

CANONICAL BASES AND THE CONJUGATING REPRESENTATION OF A SEMISIMPLE GROUP

PIERRE BAUMANN

Let G be a semisimple simply connected affine algebraic group over an algebraically closed field k of characteristic zero, let $A(G)$ be the k -algebra of regular functions of G , and let $C(G)$ be the subalgebra consisting of class functions. We explain how Lusztig's work on canonical bases affords a constructive proof of the fact, due to Richardson, that $A(G)$ is a free $C(G)$ -module.

1. Introduction

Fix an algebraically closed field k of characteristic zero. Let G be a reductive affine algebraic group over k and let V be an affine G -variety over k . We denote by $A(G)$ and $A(V)$ the k -algebras of regular functions on G and V respectively. The action of G on V gives rise to a rational representation of G on $A(V)$. A natural question is to investigate whether the algebra $A(V)$ is a free module over its subalgebra $A(V)^G$ of invariant elements. The case where V is a k -vector space on which G acts linearly has been investigated by Chevalley [Ch, Bo], Kostant [Ko], Popov [Po], Schwarz [Sc], and Littelmann [Li]. In the general case, only examples have been studied, for instance by Richardson [Ri1, Ri2] or Schwarz and Wilson [SW].

We will investigate the case where the variety V is the group G , acting on itself by inner automorphisms. Then the subalgebra of invariant elements $C(G) = A(G)^G$ is the set of regular class functions. We assume in the remainder of the paper that G is semisimple and simply connected. Richardson proved in [Ri1] that the following result holds under these assumptions.

Theorem 1. *There exists a G -stable vector subspace E of $A(G)$ such that the product map of $A(G)$ induces a vector space isomorphism from $C(G) \otimes_k E$ onto $A(G)$.*

Richardson's proof is based on a study of the geometric properties of the conjugacy classes of G and relies on heavy results of commutative algebra like the Giller-Suslin theorem. Furthermore, as Richardson himself observed, his method gives only the existence of a subspace E , and does not tell how to choose an explicit E . One can ask for instance (see Sect. 12.1 in loc. cit.) if it is possible to find a subspace E which behaves nicely in relation to the

Peter-Weyl decomposition of $A(G)$, that is, the decomposition into isotypical components for the left regular representation of G .

The aim of this paper is to provide an alternate proof of Richardson's theorem. Our method gives a more rigid choice for E , which satisfies the condition stated above. It relies on canonical bases, which are a quite recent tool in representation theory. The source of this method can be traced back to a paper of Joseph and Letzter [JL], who acknowledge an idea of Flo. Our main reference for canonical bases will be Lusztig's book [Lu2], whose notations will be recalled but not explained.

2. A graded quantized model and its canonical basis

In this section, tensor products and linear duals are taken over the field $\mathbb{Q}(v)$ of rational functions in one indeterminate.

2.1. Notations. We choose a maximal torus T in G . The weight lattice X is the character group of T . The coroot lattice Y is the dual lattice of X , the duality pairing between X and Y being denoted by $\langle \cdot, \cdot \rangle : Y \times X \rightarrow \mathbb{Z}$. The choice of a Borel subgroup B containing T affords a set $I \subseteq Y$ of simple coroots and an injection $(I \rightarrow X, i \mapsto i')$ that gives the corresponding simple roots. The dominant integral weights form a cone X^+ in the weight lattice. The set I is a basis of the lattice Y . We assume that a symmetric bilinear form $(\nu, \nu') \mapsto \nu \cdot \nu'$ is given on Y so that $i \cdot i$ is a positive even integer and $2(i \cdot j)/(i \cdot i) = \langle i, j' \rangle$ for all i, j in I .

We define on X^+ two order relations. For any ν, ν' in X^+ , we say that $\nu \leq \nu'$ whenever $\nu' - \nu \in \sum_{i \in I} \mathbb{N} i'$ and that $\nu \preceq \nu'$ whenever $\nu' - \nu \in X^+$. The poset (X^+, \preceq) is a distributive lattice.

Let v be an indeterminate. From the data above, one can define the $\mathbb{Q}(v)$ -algebra \mathfrak{f} , generated by the symbols $(\theta_i)_{i \in I}$ submitted to the quantized Serre relations ([Lu2], Chap 1 and §3.1). One then defines as in Chapter 3 of [Lu2] the quantized enveloping $\mathbb{Q}(v)$ -algebra \mathbf{U} and its involutive automorphism ω . Following §§3.4–3.5 in [Lu2], we denote the category of weight \mathbf{U} -modules by \mathcal{C} and its full subcategory of integrable \mathbf{U} -modules by \mathcal{C}' . Given a dominant integral weight λ , there is a unique simple object Λ_λ in \mathcal{C}' with highest weight λ and highest weight vector η_λ , and a unique simple object ${}^\omega \Lambda_\lambda$ in \mathcal{C}' with lowest weight $-\lambda$ and lowest weight vector $\xi_{-\lambda}$ ([Lu2], §3.5). In §14.4 of [Lu2], Lusztig defines the canonical basis \mathbf{B} of \mathfrak{f} and its family of subsets $\mathbf{B}(\lambda)$, where $\lambda \in X^+$. An immediate consequence of these definitions is the following fact.

Lemma 2. *For any $b \in \mathbf{B}$, there is a dominant integral weight $\varepsilon(b)$ such that $\{\lambda \in X^+ \mid b \in \mathbf{B}(\lambda)\} = \varepsilon(b) + X^+$.*

Proof. With the notations of loc. cit., b belongs to $\mathbf{B}(\lambda)$ if and only if the inequality $\langle i, \lambda \rangle \geq \min \{n \mid b \in {}^\sigma \mathbf{B}_{i,n}\}$ holds true for all $i \in I$. It is therefore sufficient to set $\varepsilon(b)$ so that for all $i \in I$, $\langle i, \varepsilon(b) \rangle = \min \{n \mid b \in {}^\sigma \mathbf{B}_{i,n}\}$. \square

2.2. A graded quantized model for $A(G)$. By §25.1 in [Lu2], for any dominant integral weights $\lambda, \mu \in X^+$, there are unique maps of \mathbf{U} -modules $i_{\lambda, \mu} : \Lambda_{\lambda+\mu} \rightarrow \Lambda_\lambda \otimes \Lambda_\mu$ and ${}^\omega i_{\lambda, \mu} : {}^\omega \Lambda_{\lambda+\mu} \rightarrow {}^\omega \Lambda_\mu \otimes {}^\omega \Lambda_\lambda$ such that $i_{\lambda, \mu}(\eta_{\lambda+\mu}) = \eta_\lambda \otimes \eta_\mu$ and ${}^\omega i_{\lambda, \mu}(\xi_{-\lambda-\mu}) = \xi_{-\mu} \otimes \xi_{-\lambda}$.

Using the antipode of \mathbf{U} , the dual vector space M^* of a \mathbf{U} -module M can be viewed as a \mathbf{U} -module. If M and N are \mathbf{U} -modules and if one of them is finite-dimensional, then the \mathbf{U} -modules $(M \otimes N)^*$ and $N^* \otimes M^*$ are naturally isomorphic. The dual of a finite-dimensional object of \mathcal{C}' belongs to \mathcal{C}' .

For any dominant integral weight λ , we define the \mathbf{U} -module $H^\lambda = (\omega \Lambda_\lambda \otimes \Lambda_\lambda)^*$. Also set $H = \bigoplus_{\lambda \in X^+} H^\lambda$. The family of maps

$$(1) \quad \left(\begin{array}{l} H^\lambda \otimes H^\mu \rightarrow H^{\lambda+\mu} \\ (\Lambda_\lambda)^* \otimes (\omega \Lambda_\lambda)^* \otimes (\Lambda_\mu)^* \otimes (\omega \Lambda_\mu)^* \rightarrow (\Lambda_{\lambda+\mu})^* \otimes (\omega \Lambda_{\lambda+\mu})^* \\ p \otimes q \otimes r \otimes s \mapsto (i_{\lambda, \mu})^*(r \otimes p) \otimes ({}^\omega i_{\lambda, \mu})^*(q \otimes s) \end{array} \right)$$

induces a product $m : H \otimes H \rightarrow H$ which endows H with the structure of a X^+ -graded algebra. One can easily show that this algebra is associative and has a unit.

By Proposition 25.1.4 (a) in [Lu2], for any dominant integral weight λ there is a unique \mathbf{U} -linear map $\delta_\lambda : \omega \Lambda_\lambda \otimes \Lambda_\lambda \rightarrow \mathbb{Q}(v)$ such that $\delta_\lambda(\xi_{-\lambda} \otimes \eta_\lambda) = 1$, where $\mathbb{Q}(v)$ is considered as a \mathbf{U} -module via the co-unit of \mathbf{U} . This form δ_λ is a \mathbf{U} -invariant element in H^λ .

For any two dominant integral weights λ and μ , Lusztig defines in §25.1.5 of [Lu2] the map $t_\lambda : \omega \Lambda_{\lambda+\mu} \otimes \Lambda_{\lambda+\mu} \rightarrow \omega \Lambda_\mu \otimes \Lambda_\mu$ as the composition

$$\omega \Lambda_{\lambda+\mu} \otimes \Lambda_{\lambda+\mu} \xrightarrow{{}^\omega i_{\lambda, \mu} \otimes i_{\lambda, \mu}} \omega \Lambda_\mu \otimes \omega \Lambda_\lambda \otimes \Lambda_\lambda \otimes \Lambda_\mu \xrightarrow{\text{id} \otimes \delta_\lambda \otimes \text{id}} \omega \Lambda_\mu \otimes \mathbb{Q}(v) \otimes \Lambda_\mu.$$

Lemma 3. (a) The dual map $(t_\lambda)^* : H^\mu \rightarrow H^{\lambda+\mu}$ is injective and coincides with the left multiplication by δ_λ in the algebra H .

(b) In the algebra H , one has $\delta_\lambda \delta_\mu = \delta_{\lambda+\mu}$ for any dominant integral weights λ and μ .

Proof. The injectivity of $(t_\lambda)^*$ follows from the surjectivity of t_λ , which is shown in [Lu2], Lemma 25.1.6 (c). Let us write $\delta_\lambda = \sum_i p_i \otimes q_i$ in $(\Lambda_\lambda)^* \otimes (\omega \Lambda_\lambda)^*$. Then for any elements $\sum_j r_j \otimes s_j \in (\Lambda_\mu)^* \otimes (\omega \Lambda_\mu)^*$ and $\sum_k t_k \otimes u_k \in \omega \Lambda_{\lambda+\mu} \otimes \Lambda_{\lambda+\mu}$, we have

$$\begin{aligned} & \langle \delta_\lambda \times (\sum_j r_j \otimes s_j), \sum_k t_k \otimes u_k \rangle \\ &= \sum_{i, j, k} \langle (i_{\lambda, \mu})^*(r_j \otimes p_i) \otimes ({}^\omega i_{\lambda, \mu})^*(q_i \otimes s_j), t_k \otimes u_k \rangle \\ &= \sum_{i, j, k} \langle r_j \otimes p_i \otimes q_i \otimes s_j, {}^\omega i_{\lambda, \mu}(t_k) \otimes i_{\lambda, \mu}(u_k) \rangle \end{aligned}$$

$$\begin{aligned}
&= \sum_{j,k} \langle r_j \otimes \delta_\lambda \otimes s_j, ({}^\omega i_{\lambda,\mu} \otimes i_{\lambda,\mu})(t_k \otimes u_k) \rangle \\
&= \langle \sum_j r_j \otimes s_j, t_\lambda(\sum_k t_k \otimes u_k) \rangle.
\end{aligned}$$

This calculation proves (a).

Now the linear form $\delta_\lambda \delta_\mu$ on ${}^\omega \Lambda_{\lambda+\mu} \otimes \Lambda_{\lambda+\mu}$ is \mathbf{U} -linear and takes the value 1 on the element $\xi_{-\lambda-\mu} \otimes \eta_{\lambda+\mu}$, since it can be written as $(t_\lambda)^*(\delta_\mu) = \delta_\mu \circ t_\lambda$. Therefore it coincides with $\delta_{\lambda+\mu}$, which proves (b). \square

2.3. Dual-based modules and isotypical decompositions. The simple objects of the category \mathcal{C}' are the \mathbf{U} -modules Λ_σ , where σ is a dominant integral weight; they are pairwise non-isomorphic. Given an object M in \mathcal{C}' and a dominant integral weight σ , we denote the sum of the simple subobjects of M isomorphic to Λ_σ by $M[\sigma]$. By complete reducibility we have $M = \bigoplus_{\sigma \in X^+} M[\sigma]$. Given $P \subseteq X^+$, we denote the subspace $\bigoplus_{\sigma \in P} M[\sigma]$ by $M[P]$. For short, we will write $\tau \geq \sigma$ instead of $\{\tau \in X^+ \mid \tau \geq \sigma\}$, $\not\geq \sigma$ instead of $\{\tau \in X^+ \mid \tau \not\geq \sigma\}$, and so on.

In Chapter 27 of his book [Lu2], Lusztig defines the notion of a based module. A based module is a pair (M, B) consisting of a finite-dimensional \mathbf{U} -module M which belongs to \mathcal{C}' and a $\mathbb{Q}(v)$ -basis B of M satisfying several properties stated in loc. cit. Based modules are the objects of a category: a morphism from the based module (M, B) to the based module (M', B') is a morphism $f : M \rightarrow M'$ of \mathbf{U} -modules such that $f(B) \subseteq B' \cup \{0\}$ and such that the set $B \cap \ker f$ is a basis of $\ker f$.

We define a dual-based module as a pair (M, B) consisting of a finite-dimensional \mathbf{U} -module M which belongs to \mathcal{C}' and a $\mathbb{Q}(v)$ -basis B of M such that the dual module M^* together with the basis B^* dual to B is a based module. Dual-based modules form a category, the morphisms between two dual-based modules being defined in the same way as morphisms between based modules.

For any dual-based module (M, B) and any dominant integral weight σ , we put

$$B[\sigma] = (B \cap M[\leq \sigma]) \setminus (B \cap M[< \sigma]).$$

The following properties of dual-based modules are direct consequences of similar properties of based modules.

Proposition 4. *Let (M, B) be a dual-based module and let σ be a dominant integral weight.*

(a) *The subspaces $M[\leq \sigma]$ and $M[< \sigma]$ are spanned over $\mathbb{Q}(v)$ by their intersection with B .*

(b) *The restriction of the canonical surjection $p : M[\leq \sigma] \rightarrow M[\leq \sigma]/M[< \sigma]$ to $B[\sigma]$ is injective and the pair $(M[\leq \sigma]/M[< \sigma], p(B[\sigma]))$ is a dual-based module.*

(c) *When σ runs over X^+ , the sets $B[\sigma]$ form a partition of B .*

(d) Let (M', B') be a sub-dual-based module of (M, B) and assume that M has only one non-zero isotypical component. Then the $\mathbb{Q}(v)$ -vector space M'' spanned by $B \setminus B'$ is a complementary sub- \mathbf{U} -module of M' in M and the pair $(M'', B \setminus B')$ is a dual-based module.

(e) Let (M', B') be a dual-based module and assume that \mathbf{U} acts trivially on M or on M' . Then $(M \otimes M', B \otimes B')$ is a dual-based module, where $B \otimes B'$ denotes the set $\{b \otimes b' \mid b \in B, b' \in B'\}$.

Proof. Proposition 27.1.8 in [Lu2] asserts that for any dominant integral weight τ and any based module (N, C) , the submodule $N[\geq \tau]$ is spanned over $\mathbb{Q}(v)$ by its intersection with C . One deduces from this fact that the submodule $N[P]$ is spanned over $\mathbb{Q}(v)$ by its intersection with C for any subset $P \subseteq X^+$ such that $P + (\sum_i \mathbb{N}i') \subseteq P$. In particular, this property holds for $N[\leq \sigma^*]$ and $N[\not\leq \sigma^*]$, where σ^* is the highest weight of $(\Lambda_\sigma)^*$. Applying this result to the case of the based module (M^*, B^*) and taking orthogonals, we obtain Property (a).

Property (a) proves that the restriction of p defines a bijection from $B[\sigma]$ onto a basis of the $\mathbb{Q}(v)$ -vector space $M[\leq \sigma]/M[\lt \sigma]$. To check that the pair $(M[\leq \sigma]/M[\lt \sigma], p(B[\sigma]))$ satisfies all the axioms of a dual-based module, it suffices to use duality as in the proof of Property (a) and to refer to the definition of based modules in §27.1.2 of [Lu2]. Property (b) is proved

Choose any x in B . We can find $\sigma \in X^+$ such that $x \in M[\leq \sigma]$ and such that σ is minimal for this property with respect to the order \leq . Since B is a basis of M , the element x does not belong to the span of $\bigcup_{\tau < \sigma} (B \cap M[\leq \tau])$. By Property (a), one deduces that x does not belong to $M[\lt \sigma]$ and therefore that x belongs to $B[\sigma]$. We have proved that B is the union of its subsets $B[\sigma]$, and it remains us to show that these sets $B[\sigma]$ are pairwise disjoint. Suppose that $B[\sigma]$ and $B[\tau]$ share a certain element x . Then $M[\leq \sigma]$ and $M[\leq \tau]$ intersect non-trivially. This implies that $\sigma - \tau$ belongs to the root lattice $\sum_i \mathbb{Z}i'$, and thus there exists a weight ρ less than or equal to σ and τ such that $M[\leq \sigma] \cap M[\leq \tau] = M[\leq \rho]$. Since x belongs to $M[\leq \rho]$ but not to $M[\lt \sigma]$, we cannot have $\rho < \sigma$. Therefore $\rho = \sigma$, and similarly $\rho = \tau$. Therefore $\sigma = \tau$, which completes the proof of Property (c).

Finally Property (d) is a consequence of the proof of Proposition 27.1.7 in [Lu2], and Property (e) follows by dualizing the construction given in §27.3 and Theorem 27.3.2 of [Lu2]. □

It is of course possible to extend the notion of (dual-) based module to the case of an infinite-dimensional \mathbf{U} -module which is graded with finite-dimensional graded components. In this case, the basis is required to be compatible with the decomposition of the module as the direct sum of its graded components. □

2.4. The basis of H . By §§24.3 and 27.3.4 in [Lu2], each module ${}^\omega\Lambda_\lambda \otimes \Lambda_\lambda$ has a canonical basis, with which it forms a based module. By Proposition 27.3.5 (a) in [Lu2], the map $t_\lambda : {}^\omega\Lambda_{\lambda+\mu} \otimes \Lambda_{\lambda+\mu} \rightarrow {}^\omega\Lambda_\mu \otimes \Lambda_\mu$ is a morphism of based modules.

Each module $H^\lambda = ({}^\omega\Lambda_\lambda \otimes \Lambda_\lambda)^*$ comes therefore with the dual basis B_λ , so that the pair (H^λ, B_λ) is a dual-based module. By Lemma 3 (a), the left multiplication by δ_λ defines an injective morphism of dual-based modules from (H^μ, B_μ) to $(H^{\lambda+\mu}, B_{\lambda+\mu})$.

In particular, we get an injective map from B_μ to $B_{\lambda+\mu}$. By Lemma 3 (b) these maps form a directed system of injective maps between sets B_λ , and we denote its limit¹ by $B_\infty = \varinjlim B_\lambda$. We denote the canonical injective map $B_\lambda \rightarrow B_\infty$ by ι_λ . By Proposition 27.2.2 in [Lu2], this directed system is compatible with the decompositions $B_\lambda = \bigsqcup_{\sigma \in X^+} B_\lambda[\sigma]$, which yields a similar decomposition $B_\infty = \bigsqcup_{\sigma \in X^+} B_\infty[\sigma]$.

Lemma 5. *Given $x \in B_\infty$, there is a dominant integral weight $\varepsilon(x)$ such that $\{\lambda \in X^+ \mid x \in \iota_\lambda(B_\lambda)\} = \varepsilon(x) + X^+$.*

Proof. By duality the assertion is equivalent to the following fact: for any $\lambda, \mu, \nu \in X^+$ such that $\lambda \preceq \nu$ and $\mu \preceq \nu$ and any y in the canonical basis of ${}^\omega\Lambda_\nu \otimes \Lambda_\nu$, the non-vanishing of both $t_\lambda(y)$ and $t_\mu(y)$ implies that of $t_{\sup(\lambda, \mu)}(y)$, where $\sup(\cdot, \cdot)$ is the supremum in the distributive lattice (X^+, \preceq) . In turn, this fact is a direct consequence of Proposition 25.1.10 in [Lu2] and Lemma 2. □

Lemma 6. *The set $B_\lambda[0]$ is reduced to the element δ_λ .*

Proof. The space $H^\lambda[0] = \text{Hom}_{\mathbb{U}}({}^\omega\Lambda_\lambda \otimes \Lambda_\lambda, \mathbb{Q}(v))$ has dimension at most one, since ${}^\omega\Lambda_\lambda \otimes \Lambda_\lambda$ is generated by a single element, namely $\xi_{-\lambda} \otimes \eta_\lambda$. Therefore $B_\lambda[0]$ has at most one element and it suffices to show that $\delta_\lambda \in B_\lambda$. We observe that the kernel of δ_λ is $({}^\omega\Lambda_\lambda \otimes \Lambda_\lambda)[> 0]$, which by Proposition 27.1.8 in [Lu2] is spanned over $\mathbb{Q}(v)$ by its intersection with the canonical basis of ${}^\omega\Lambda_\lambda \otimes \Lambda_\lambda$. Therefore δ_λ vanishes on all elements of this canonical basis but one. The exception is the vector $\xi_{-\lambda} \otimes \eta_\lambda$: it belongs to the canonical basis by Theorem 24.3.3 in [Lu2] and δ_λ evaluates to 1 on it. This shows that δ_λ belongs to the basis dual to the canonical basis of ${}^\omega\Lambda_\lambda \otimes \Lambda_\lambda$, that is to say δ_λ belongs to B_λ . □

The direct sum of the dual-based modules (H^λ, B_λ) will be denoted by (H, B) . Lemma 6 tells that $B[0] = \{\delta_\lambda \mid \lambda \in X^+\}$ and Proposition 4 (a) implies that the pair $(H[0], B[0])$ is a dual-based module. By Lemma 3 (a), for any $\lambda \in X^+$, the left multiplication by δ_λ is an injective morphism from the dual-based module (H, B) into itself.

¹This limit B_∞ is, in a certain sense, the basis dual to the canonical basis of the subspace $\tilde{\mathbb{U}}_{1_0}$ of Lusztig's modified quantized enveloping algebra, see Chap. 23 of [Lu2].

2.5. A filtration of H and the freeness theorem for its associated graded. The dual-based module (H, B) is filtered by the family of submodules $(H[\leq \sigma], B \cap H[\leq \sigma])$, the indexing set being the poset (X^+, \leq) . The associated graded dual-based module is $\bigoplus_{\sigma \in X^+} (\text{gr}^\sigma(H), \mathcal{B}[\sigma])$, where $\text{gr}^\sigma(H) = H[\leq \sigma]/H[\prec \sigma]$ and $\mathcal{B}[\sigma]$ is the image of $B[\sigma] = \bigsqcup_{\lambda \in X^+} B_\lambda[\sigma]$ under the canonical surjection $p: H[\leq \sigma] \rightarrow \text{gr}^\sigma(H)$.

We view H as the regular left H -module. The subspace $H[0]$ acts by morphisms of U -modules; therefore its action stabilizes each isotypical component of H and induces an action on any $\text{gr}^\sigma(H)$.

We now fix a dominant integral weight σ . We define

$$B[\sigma]^{\text{prim}} = \{\iota_{\varepsilon(x)}^{-1}(x) \mid x \in B_\infty[\sigma]\} = \bigsqcup_{\lambda \in X^+} \{x \in B_\lambda[\sigma] \mid \varepsilon(\iota_\lambda(x)) = \lambda\},$$

and we call $\mathcal{B}[\sigma]^{\text{prim}}$ its image under the canonical surjection p . We denote by K^σ the $\mathbb{Q}(v)$ -vector subspace spanned in $\text{gr}^\sigma(H)$ by $\mathcal{B}[\sigma]^{\text{prim}}$.

Proposition 7. (a) The action of δ_λ on $\text{gr}^\sigma(H)$ induces an injective morphism from the dual-based module $(\text{gr}^\sigma(H), \mathcal{B}[\sigma])$ into itself.

(b) The family of sets $(\delta_\lambda \cdot \mathcal{B}[\sigma]^{\text{prim}})_{\lambda \in X^+}$ form a partition of $\mathcal{B}[\sigma]$.

(c) The pair $(K^\sigma, \mathcal{B}[\sigma]^{\text{prim}})$ is a dual-based module.

Proof. Assertion (a) follows from the fact that the left multiplication by δ_λ is an injective morphism from the dual-based module (H, B) into itself.

As for Assertion (b), we consider an element $x \in B_\mu[\sigma]$. Let $\nu = \varepsilon(\iota_\mu(x))$. By Lemma 5, $\lambda = \mu - \nu$ belongs to X^+ and there exists $y \in B_\nu[\sigma]$ such that $\iota_\nu(y) = \iota_\mu(x)$. By construction, $y \in B[\sigma]^{\text{prim}}$ and $p(x)$ is the image of $p(y)$ under the action of δ_λ . This proves that $\mathcal{B}[\sigma] = \bigcup_{\lambda \in X^+} (\delta_\lambda \cdot \mathcal{B}[\sigma]^{\text{prim}})$. A similar reasoning based on Lemma 5 and on Assertion (a) shows that the union is disjoint.

To prove Assertion (c), it is enough to show that for all dominant integral weight λ , the pair $(K^\sigma \cap \text{gr}^\sigma(H^\lambda), \mathcal{B}[\sigma]^{\text{prim}} \cap \text{gr}^\sigma(H^\lambda))$ is a dual-based module. This is trivial for $\lambda = 0$. The case of a general λ will be proved by induction on $\sum_i \langle i, \lambda \rangle$. Assume that $\lambda \neq 0$ is given. By the induction hypothesis, we can assume that the pair $(K^\sigma \cap \text{gr}^\sigma(H^\mu), \mathcal{B}[\sigma]^{\text{prim}} \cap \text{gr}^\sigma(H^\mu))$ is a dual-based module for all $\mu \in X^+$ such that $\mu \prec \lambda$. Assertion (b) then says that the pair

$$\left(\bigoplus_{\mu \in X^+, \mu \prec \lambda} \delta_{\lambda-\mu} \cdot (K^\sigma \cap \text{gr}^\sigma(H^\mu)), \bigsqcup_{\mu \in X^+, \mu \prec \lambda} \delta_{\lambda-\mu} \cdot (\mathcal{B}[\sigma]^{\text{prim}} \cap \text{gr}^\sigma(H^\mu)) \right)$$

is a sub-dal-based mdle of $(\text{gr}^\sigma(H^\lambda), \mathcal{B}[\sigma] \cap \text{gr}^\sigma(H^\lambda))$ and that

$$\mathcal{B}[\sigma]^{\text{prim}} \cap \text{gr}^\sigma(H^\lambda) = \left(\mathcal{B}[\sigma] \cap \text{gr}^\sigma(H^\lambda) \right) \setminus \left(\bigsqcup_{\mu \in X^+, \mu \prec \lambda} \delta_{\lambda-\mu} \cdot (\mathcal{B}[\sigma]^{\text{prim}} \cap \text{gr}^\sigma(H^\mu)) \right).$$

Now Assertion (c) follows from Proposition 4 (d). \square

We now have three dal-based mdles $(\text{gr}^\sigma(H), \mathcal{B}[\sigma])$, $(H[0], B[0])$, and $(K^\sigma, \mathcal{B}[\sigma]^{\text{prim}})$. By Proposition 4 (e), the pair $(H[0] \otimes K^\sigma, B[0] \otimes \mathcal{B}[\sigma]^{\text{prim}})$ is a dal-based mdle.

Theorem 8. *The action of $H[0]$ on $\text{gr}^\sigma(H)$ gives rise to an isomorphism from $(H[0] \otimes K^\sigma, B[0] \otimes \mathcal{B}[\sigma]^{\text{prim}})$ onto $(\text{gr}^\sigma(H), \mathcal{B}[\sigma])$.*

Proof. Since \mathbf{U} acts trivially on $H[0]$, the \mathbf{U} -linear action of $H[0]$ on $\text{gr}^\sigma(H)$ induces a morphism of \mathbf{U} -mdles from $H[0] \otimes \text{gr}^\sigma(H)$ to $\text{gr}^\sigma(H)$. By Proposition 7 (a) and (b), this morphism restricts to a bijection from $B[0] \otimes \mathcal{B}[\sigma]^{\text{prim}}$ onto $\mathcal{B}[\sigma]$. The theorem follows. \square

3. Specialization to the classical case

3.1. Specialization of \mathbf{U} -modules. Let \mathcal{A} be the ring $\mathbb{Z}[v, v^{-1}]$. The field k is an \mathcal{A} -algebra on which v acts as the identity. For any \mathcal{A} -mdle ${}_{\mathcal{A}}T$, we denote by ${}_kT$ the k -mdle $k \otimes_{\mathcal{A}} {}_{\mathcal{A}}T$ obtained by base ring change.

We call \mathfrak{g} the Lie algebra of the group G and we choose Chevalley generators $E_1, \dots, E_\ell, F_1, \dots, F_\ell, H_1, \dots, H_\ell$ in it.

In §3.1.13 of [Lu2] (see also Theorem 4.5 in [Lu1]), Lusztig defines an \mathcal{A} -form ${}_{\mathcal{A}}\mathbf{U}$ of \mathbf{U} . Formulas in §§3.1.5 and 3.3.3 of [Lu2] show that ${}_{\mathcal{A}}\mathbf{U}$ inherits from \mathbf{U} the structure of a Hopf algebra over \mathcal{A} . Therefore ${}_k\mathbf{U}$ is a Hopf algebra over k . Furthermore, since the quantized Serre relations are verified by the simple root vectors in ${}_{\mathcal{A}}\mathbf{U}$, there is a natural morphism of Hopf algebras $c: U(\mathfrak{g}) \rightarrow {}_k\mathbf{U}$. Thanks to c , every ${}_k\mathbf{U}$ -mdle has a natural structure of a $U(\mathfrak{g})$ -mdle.

We use the standard strategy to specialize a finite-dimensional \mathbf{U} -mdle M : we first choose a $\mathbb{Q}(v)$ -basis B of M such that the \mathcal{A} -submdle ${}_{\mathcal{A}}M$ spanned by B in M is stable under the action of ${}_{\mathcal{A}}\mathbf{U}$, and then ${}_kM$ is a $U(\mathfrak{g})$ -mdle. So what we really specialize is the pair (M, B) . Thanks to Condition (b) in Definition 27.1.2 of [Lu2], based mdles satisfy the required condition to be specializable. One can also construct new specializable pairs by standard procedures like dualization, tensor product, or twist with ω , and then the specialization commutes with these constructions. We extend this framework to infinite-dimensional \mathbf{U} -mdles provided that they are graded with finite-dimensional graded components and that the \mathbf{U} -ir bases consist of homogeneous elements.

Let $\lambda \in X^+$. In Theorem 4.4.11 of [Lu2], Lusztig constructs a $\mathbb{Q}(v)$ -basis $\mathbf{B}(\Lambda_\lambda)$ of Λ_λ so that $(\Lambda_\lambda, \mathbf{B}(\Lambda_\lambda))$ is a based module. Lusztig shows in §3.1.2 of [Lu2] that the specialized module ${}_k(\Lambda_\lambda)$ is a simple highest weight module with highest weight λ . The basis $\mathbf{B}(\Lambda_\lambda)$ embeds ${}_k(\Lambda_\lambda)$ with a preferred highest weight vector ${}_k\eta_\lambda$. For another $\mu \in X^+$. By Proposition 25.1.2 in [Lu2], the map $i_{\lambda,\mu} : \Lambda_{\lambda+\mu} \rightarrow \Lambda_\lambda \otimes_{\mathbb{Q}(v)} \Lambda_\mu$ sends the \mathcal{A} -submodule spanned by $\mathbf{B}(\Lambda_{\lambda+\mu})$ in $\Lambda_{\lambda+\mu}$ into the \mathcal{A} -submodule spanned by $\mathbf{B}(\Lambda_\lambda) \otimes \mathbf{B}(\Lambda_\mu)$ in $\Lambda_\lambda \otimes_{\mathbb{Q}(v)} \Lambda_\mu$. It therefore specializes to the morphism of $U(\mathfrak{g})$ -modules ${}_k(i_{\lambda,\mu}) : {}_k(\Lambda_{\lambda+\mu}) \rightarrow {}_k(\Lambda_\lambda) \otimes_k {}_k(\Lambda_\mu)$ that sends ${}_k\eta_{\lambda+\mu}$ to ${}_k\eta_\lambda \otimes_k \eta_\mu$. Similarly, the U -module ${}^\omega\Lambda_\lambda$ comes with a canonical basis ${}^\omega\mathbf{B}(\Lambda_\lambda)$. Therefore it can be specialized to the $U(\mathfrak{g})$ -module ${}_k({}^\omega\Lambda_\lambda)$, which is a simple lowest weight module with lowest weight $-\lambda$ and lowest weight vector ${}_k\xi_{-\lambda}$. The specialization of ${}^\omega i_{\lambda,\mu} : {}^\omega\Lambda_{\lambda+\mu} \rightarrow {}^\omega\Lambda_\mu \otimes_{\mathbb{Q}(v)} {}^\omega\Lambda_\lambda$ is the morphism of $U(\mathfrak{g})$ -modules ${}_k({}^\omega i_{\lambda,\mu}) : {}_k({}^\omega\Lambda_{\lambda+\mu}) \rightarrow {}_k({}^\omega\Lambda_\mu) \otimes_k {}_k({}^\omega\Lambda_\lambda)$ that sends ${}_k\xi_{-\lambda-\mu}$ to ${}_k\xi_{-\mu} \otimes_k \xi_{-\lambda}$.

The family $({}_k(\Lambda_\sigma))_{\sigma \in X^+}$ affords a complete set of pairwise non-isomorphic finite-dimensional simple $U(\mathfrak{g})$ -modules. Given a finite-dimensional $U(\mathfrak{g})$ -module M and a dominant integral weight σ , we denote its isotypical component of type ${}_k(\Lambda_\sigma)$ by $M[\sigma]$. Given $P \subseteq X^+$, we denote the subspace $\bigoplus_{\sigma \in P} M[\sigma]$ by $M[P]$.

Proposition 9. *Let (M, B) be a dual-based module and ${}_kM$ its specialization. Then for any $\sigma \in X^+$, the dual-based modules $(M[\leq \sigma], B \cap M[\leq \sigma])$ and $(M[< \sigma], B \cap M[< \sigma])$ specialize to $({}_kM)[\leq \sigma]$ and $({}_kM)[< \sigma]$, respectively. In particular $(M[0], B \cap M[0])$ specializes to $({}_kM)[0]$.*

Proof. We will only prove the case of $(M[\leq \sigma], B \cap M[\leq \sigma])$. We can enumerate the weights in $\leq \sigma$ as a finite sequence τ_1, \dots, τ_n such that $\tau_i \leq \tau_j \Rightarrow i \leq j$. The dual-based module $(M[\leq \sigma], B \cap M[\leq \sigma])$ is then filtered by the composition series $(M[\{\tau_1, \dots, \tau_i\}], B \cap M[\{\tau_1, \dots, \tau_i\}])_{0 \leq i \leq n}$. As U -modules, the quotient modules are isotypical of type Λ_{τ_i} and specialize therefore to isotypical modules of type ${}_k(\Lambda_{\tau_i})$, by the dual version of Proposition 27.1.7 in [Lu2]. Thus the specialization of $(M[\leq \sigma], B \cap M[\leq \sigma])$ has a filtration with quotients isomorphic to ${}_k(\Lambda_{\tau_1}), \dots, \alpha {}_k(\Lambda_{\tau_n})$, which shows that ${}_k(M[\leq \sigma]) \subseteq ({}_kM)[\leq \sigma]$. A similar reasoning shows that the specialization of $M/M[\leq \sigma]$ has a filtration with quotients isomorphic to modules of the form ${}_k(\Lambda_\tau)$ with $\tau \not\leq \sigma$, where

$$(({}_kM)/{}_k(M[\leq \sigma]))[\leq \sigma] = ({}_k(M/M[\leq \sigma]))[\leq \sigma] = 0.$$

Therefore the equality ${}_k(M[\leq \sigma]) = ({}_kM)[\leq \sigma]$ holds. \square

3.2. Specialization of H . We are now in a position where we can specialize the U -module H , the multiplication map $m : H \otimes_{\mathbb{Q}(v)} H \rightarrow H$, and the freeness result from Theorem 8.

First observe that by Theorem 24.3.3 in [Lu2], the \mathcal{A} -lattice spanned in H^λ by the basis B_λ is the same as the \mathcal{A} -lattice spanned by the basis dual to the basis ${}^\omega \mathbf{B}(\Lambda_\lambda) \otimes \mathbf{B}(\Lambda_\lambda)$ of ${}^\omega \Lambda_\lambda \otimes_{\mathbb{Q}(v)} \Lambda_\lambda$. Therefore the multiplication map m sends the \mathcal{A} -submodule spanned in $H \otimes_{\mathbb{Q}(v)} H$ by $B \otimes B$ into the \mathcal{A} -submodule spanned in H by B . It gives rise to a multiplication map ${}_k m : {}_k H \otimes_k {}_k H \rightarrow {}_k H$.

Proposition 10. *The specialization ${}_k H$ is the $U(\mathfrak{g})$ -module*

$$\bigoplus_{\lambda \in X^+} ({}_k(\Lambda_\lambda)^* \otimes_k ({}^\omega \Lambda_\lambda)^*).$$

The multiplication map ${}_k m$ is given by Formula (1) in which the maps $(i_{\lambda,\mu})^*$ and $({}^\omega i_{\lambda,\mu})^*$ are replaced by their specializations ${}_k(i_{\lambda,\mu})^*$ and ${}_k({}^\omega i_{\lambda,\mu})^*$.

Now fix a dominant integral weight σ . By Proposition 9, the isotypical component $({}_k H)[\sigma]$ is naturally isomorphic to the specialization of the dual-based module $(\text{gr}^\sigma(H), \mathcal{B}[\sigma])$. The specialization ${}_k(K^\sigma)$ of $(K^\sigma, \mathcal{B}[\sigma]^{\text{prim}})$ is then seen as a $U(\mathfrak{g})$ -submodule of $({}_k H)[\sigma]$. By Theorem 8 and Proposition 9, we get the following result.

Theorem 11. *The map ${}_k m$ induces an isomorphism of $U(\mathfrak{g})$ -modules from $({}_k H)[0] \otimes_k {}_k(K^\sigma)$ onto $({}_k H)[\sigma]$.*

3.3. The Cartan filtration on $A(G)$. To complete the proof of Theorem 1, it only remains to relate the specialized algebra ${}_k H$ to the algebra $A(G)$. We first describe this latter.

Let M be a rational G -module. Then for any $v \in M$ and $f \in M^*$, the function on G

$$c_{f,v}^M : g \mapsto \langle f, g \cdot v \rangle$$

is regular. The map from $M^* \otimes_k M$ to $A(G)$ which sends $f \otimes v$ to $c_{f,v}^M$ is a morphism of G -modules; it is injective if M is simple. By definition, its image is the coefficient space $C(M)$ of the module M . Then the Peter-Weyl decomposition

$$A(G) = \bigoplus_{\lambda \in X^+} C({}_k(\Lambda_\lambda))$$

holds. The filtration of $A(G)$ indexed by the poset (X^+, \leq) and given by the submodules

$$A_\lambda(G) = \bigoplus_{\mu \in X^+, \mu \leq \lambda} C({}_k(\Lambda_\mu))$$

is a filtration of algebra. The associated graded is

$$\text{gr}(A(G)) = \bigoplus_{\lambda \in X^+} \text{gr}^\lambda(A(G)),$$

where

$$\mathrm{gr}^\lambda(A(G)) = A_\lambda(G) / \sum_{\mu < \lambda} A_\mu(G) \simeq C(k(\Lambda_\lambda)) \simeq k(\Lambda_\lambda)^* \otimes_k k(\Lambda_\lambda).$$

For any $\lambda, \mu \in X^+$, there is a k -isomorphism

Remark. The author does not understand the relation between the point of view presented in this paper and the extension by Donkin [Do] of Richardson's work to the case where the ground field has a positive characteristic.

Acknowledgements. The author has the pleasure to thank N. Reshetikin for a three-months long invitation at the University of California at Berkeley.

References

- [Bo] N. Bourbaki, *Groupes et algèbres de Lie, Chapitres 4, 5 et 6*, Masson, Paris, 1981, MR 39 #1590, Zbl 0186.33001 and 0483.22001.
- [Ch] C. Chevalley, *Invariants of finite groups generated by reflections*, Amer. J. Math. 77 (1955), 778–782, MR 17,345d, Zbl 0065.26103.
- [Do] S. Donkin, *On conjugating representations and adjoint representations of semisimple groups*, Invent. Math. 91 (1988), 137–145, MR 89a:20047, Zbl 0639.20021.
- [JL] A. Joseph and G. Letzter, *Separation of variables for quantized enveloping algebras*, Amer. J. Math. 116 (1994), 127–177, MR 95e:17017, Zbl 0811.17007.
- [Ko] B. Kostant, *Lie group representations on polynomial rings*, Amer. J. Math. 85 (1963), 327–404, MR 28 #1252, Zbl 0124.26802.
- [Li] P. Littelmann, *Koreguläre und äquidimensionale Darstellungen*, J. Algebra 123 (1989), 193–222, MR 90e:20039, Zbl 0688.14042.
- [Lu1] G. Lusztig, *Finite dimensional Hopf algebras arising from quantized universal enveloping algebras*, J. Amer. Math. Soc. 3 (1990), 257–296, MR 91e:17009, Zbl 0695.16006.
- [Lu2] G. Lusztig, *Introduction to quantum groups*, Progress in Mathematics, 110, Birkhäuser Boston, Boston, MA, 1993, MR 94m:17016, Zbl 0788.17010.
- [Po] V. L. Popov, *Representations with a free module of covariants*, Funktsional. Anal. i Prilozhen. 10 (1976), 91–92 (= Funct. Anal. Appl. 10 (1976), 242–243), MR 54 #5255, Zbl 0365.20053.
- [Ri1] R. W. Richardson, *The conjugating representation of a semisimple group*, Invent. Math. 54 (1979), 229–245, MR 81a:14023, Zbl 0424.20035.
- [Ri2] R. W. Richardson, *An application of the Serre conjecture to semisimple algebraic groups*, in 'Algebra, Carbondale 1980', R. K. Amayo, editor, Lecture Notes in Math., 848, Springer-Verlag, Berlin-Heidelberg-New York, 1981, pp. 141–151, MR 83j:20047, Zbl 0457.14022.
- [Sc] G. W. Schwarz, *Representations of simple Lie groups with a free module of covariants*, Invent. Math. 50 (1978), 1–12, MR 80c:14008, Zbl 0391.20033.
- [SW] G. W. Schwarz and D. L. Wehlau, *Invariants of four subspaces*, Ann. Inst. Fourier (Grenoble) 48 (1998), 667–697, MR 99i:14055, Zbl 0899.20024.

Received date / Revised version date

INSTITUT DE RECHERCHE MATHÉMATIQUE AVANCÉE
UNIVERSITÉ LOUIS PASTEUR ET CNRS
7, RUE RENÉ DESCARTES
F-67084 STRASBOURG CEDEX
FRANCE

E-mail address: baumann@math.u-strasbg.fr

The author acknowledges partial financial support from the European TMR network "Algebraic Lie representations", contract no. ERB FMRX-CT97-0100.