

Yann BUGEAUD (Strasbourg)

Abstract. *Let α in $(0, 1]$ and $\beta > 1$ be algebraic numbers. We study the asymptotic behaviour of the function that counts the number of digit changes in the β -expansion of α .*

1. Introduction

Let $\beta > 1$ be a real number. The β -transformation T_β is defined on $[0, 1]$ by $T_\beta : x \mapsto \beta x \bmod 1$. In 1957, Rényi [12] introduced the β -expansion of a real x in $[0, 1]$, denoted by $d_\beta(x)$ and defined by

$$d_\beta(x) = 0.x_1x_2\dots x_k\dots,$$

where $x_k = \lfloor \beta T_\beta^{k-1}(x) \rfloor$ for $k \geq 1$, except when β is an integer and $x = 1$, in which case $d_\beta(1) := 0.(\beta - 1)\dots(\beta - 1)\dots$. Here and throughout the present paper, $\lfloor \cdot \rfloor$ denotes the integer part function. Clearly, we have

$$x = \sum_{k \geq 1} \frac{x_k}{\beta^k}.$$

For $x < 1$, this expansion coincides with the representation of x computed by the ‘greedy algorithm’. If β is an integer b , then the digits x_i of x lie in the set $\{0, 1, \dots, b - 1\}$ and, if $x < 1$, then $d_b(x)$ corresponds to the b -ary expansion of x . If β is not an integer, then the digits x_i lie in the set $\{0, 1, \dots, \lfloor \beta \rfloor\}$. We direct the reader to [2] and to the references quoted therein for more on β -expansions. Throughout this note, we say that $d_\beta(x)$ is finite (resp. infinite) if there are only finitely many (resp. there are infinitely many) non-zero digits in the β -expansion of x .

We stress that the β -expansion of 1 has been extensively studied, for it yields a lot of information on the β -shift. In particular, Blanchard [5] proposed a classification of the β -shifts according to the properties of the (finite or infinite) word given by $d_\beta(1)$, see Section 4 of [2]. The occurrences of consecutive 0’s in $d_\beta(1)$ play a crucial role in Blanchard’s classification of the β -shifts. This motivates the following problem first investigated in [17].

Let $\beta > 1$ be a real number such that $d_\beta(1)$ is infinite and let $(a_k)_{k \geq 1}$ be the β -expansion of 1. Assume that there exist a sequence of positive integers $(r_n)_{n \geq 1}$ and an increasing sequence of positive integers $(s_n)_{n \geq 1}$ such that

$$a_{s_n+1} = a_{s_n+2} = \dots = a_{s_n+r_n} = 0, \quad a_{s_n+r_n+1} \neq 0,$$

and $s_{n+1} > s_n + r_n$ for every positive integer n . The problem is then to estimate the gaps between two consecutive non-zero digits in $d_\beta(1)$, that is, to estimate the asymptotic behaviour of the ratio r_n/s_n .

The main result of [17], quoted as Theorem VG below, mainly shows that $d_\beta(1)$ cannot be ‘too lacunary’ when β is an algebraic number. Recall that the Mahler measure of a real algebraic number θ , denoted by $M(\theta)$, is, by definition, equal to the product

$$M(\theta) := a \prod_{i=1}^d \max\{1, |\theta_i|\},$$

where $\theta = \theta_1, \theta_2, \dots, \theta_d$ are the complex conjugates of θ and a is the leading coefficient of the minimal defining polynomial of θ over the integers.

Theorem VG. *Let $\beta > 1$ be a real algebraic number. Then, with the above notation, we have*

$$\limsup_{n \rightarrow \infty} \frac{r_n}{s_n} \leq \frac{\log M(\beta)}{\log \beta} - 1.$$

Theorem VG was extended in [2], where, roughly speaking, repetitions of arbitrary (finite) blocks in the β -expansion of an algebraic number (where $\beta > 1$ is algebraic) are studied, see Theorem 2 from [2] for a precise statement.

The purpose of the present note is to study the β -expansion of an algebraic number α from another point of view, introduced in [8]. We aim at estimating the asymptotic behaviour of the number of digit changes in $d_\beta(\alpha)$. For α in $(0, 1]$, write

$$d_\beta(\alpha) = 0.a_1a_2\dots,$$

and define the function nbdc_β , ‘number of digit changes in the β -expansion’, by

$$\text{nbdc}_\beta(n, \alpha) = \text{Card}\{1 \leq k \leq n : a_k \neq a_{k+1}\},$$

for any positive integer n . This function was first studied in [8] when β is an integer, see also [9] for an improvement of the main result of [8]. The short Section 6 of [8] is devoted to the study of nbdc_β for β algebraic, but it contains some little mistakes (see below) and its main result can be strengthened (see Theorem 2 below).

The present note is organized as follows. Our results on the behaviour of the function nbdc_β when β is an algebraic number are stated in Section 2 and proved in Section 4. New results on values of lacunary series at algebraic points are discussed in Section 3.

2. Results

We begin by stating a consequence of Theorem 2 from [2], that can also be obtained with the tools used in [17].

Theorem 1. *Let $\beta > 1$ be a real algebraic number. Let α be an algebraic number in $(0, 1]$. If $d_\beta(\alpha)$ is infinite, then*

$$\liminf_{n \rightarrow +\infty} \frac{\text{nbdc}_\beta(n, \alpha)}{\log n} \geq \left(\log \left(\frac{\log M(\beta)}{\log \beta} \right) \right)^{-1}. \quad (2.1)$$

For sake of completeness, Theorem 1 is established in Section 4 along the proof of Theorem 2.

A Pisot (resp. Salem) number is an algebraic integer greater than 1 whose conjugates are of modulus less than 1 (resp. less than or equal to 1, with at least one conjugate on the unit circle). In particular, an algebraic number $\beta > 1$ is a Pisot or a Salem number if, and only if, $M(\beta) = \beta$. In that case, Theorem 1 implies that

$$\frac{\text{nbdc}_\beta(n, \alpha)}{\log n} \xrightarrow{n \rightarrow +\infty} +\infty. \quad (2.2)$$

The main purpose of the present note is to show how the use of a suitable version of the Quantitative Subspace Theorem allows us to strengthen (2.2).

Theorem 2. *Let β be a Pisot or a Salem number. Let α be an algebraic number in $(0, 1]$ such that $d_\beta(\alpha)$ is infinite and write*

$$d_\beta(\alpha) = 0.a_1a_2 \dots a_k \dots$$

Then, there exists an effectively computable constant $c(\alpha, \beta)$, depending only on α and β , such that

$$\text{nbdc}_\beta(n, \alpha) \geq c(\alpha, \beta) (\log n)^{3/2} \cdot (\log \log n)^{-1/2}, \quad (2.3)$$

for every positive integer n .

We stress that the exponent of $(\log n)$ in (2.3) is independent of β , unlike in Theorem 3 of [8]. This is a consequence of the use of the Parametric Subspace Theorem, exactly as in Theorem 3.1 of [9]. Note that Theorem 3 of [8] is not correctly stated: indeed, it claims a result valid for all expansions, whereas in the proof we are led to construct good algebraic approximations to α and to use one property of the β -expansion (see (4.14) below) to ensure that, roughly speaking, all these approximations are different.

We display two immediate corollaries of Theorem 2. A first one is concerned with the number of non-zero digits in the β -expansion of an algebraic number for β being a Pisot or a Salem number.

Corollary 1. *Let ε be a positive real number. Let β be a Pisot or a Salem number. Let α be an algebraic number in $(0, 1]$ whose β -expansion is infinite. Then, for n large enough, there are at least*

$$(\log n)^{3/2} \cdot (\log \log n)^{-1/2-\varepsilon}$$

non-zero digits among the first n digits of the β -ary expansion of α .

For $\beta = 2$, Corollary 1 gives a much weaker result than the one obtained by Bailey, Borwein, Crandall, and Pomerance [3], who proved that, among the first n digits of the binary expansion of a real irrational algebraic number ξ of degree d , there are at least $c(\xi)n^{1/d}$ occurrences of the digit 1, where $c(\xi)$ is a suitable positive constant (see also Rivoal [14]).

Recall that β is called a *Parry number* if $d_\beta(1)$ is finite or eventually periodic. Every Pisot number is a Parry number [15, 4] and K. Schmidt [15] conjectured that all Salem numbers are Parry numbers. This was proved for all Salem numbers of degree 4 by Boyd [6], who gave in [7] a heuristic suggesting the existence of Salem numbers of degree 8 that are not Parry numbers.

We highlight the special case of the β -expansion of 1 in a base β that is a Salem number.

Corollary 2. *Let ε be a positive real number. Let β be a Salem number. Assume that $d_\beta(1)$ is infinite and write*

$$d_\beta(1) = 0.a_1a_2\dots$$

For any sufficiently large integer n , we have

$$a_1 + \dots + a_n > (\log n)^{3/2} \cdot (\log \log n)^{-1/2-\varepsilon},$$

and there are at least $(\log n)^{3/2} (\log \log n)^{-1/2-\varepsilon}$ indices j with $1 \leq j \leq n$ and $a_j \neq 0$.

In view of Theorem 2, our Corollaries 1 and 2 can be (very) slightly improved.

3. On values of lacunary series at algebraic points

The following problem was posed in Section 7 of [8].

Problem 1. *Let $\mathbf{n} = (n_j)_{j \geq 1}$ be a strictly increasing sequence of positive integers and set*

$$f_{\mathbf{n}}(z) = \sum_{j \geq 1} z^{n_j}. \tag{3.1}$$

If the sequence \mathbf{n} increases sufficiently rapidly, then the function $f_{\mathbf{n}}$ takes transcendental values at every non-zero algebraic point in the open unit disc.

By a clever use of the Schmidt Subspace Theorem, Corvaja and Zannier [10] proved that the conclusion of the Problem holds for $f_{\mathbf{n}}$ given by (3.1) when the strictly increasing sequence \mathbf{n} is lacunary, that is, satisfies

$$\liminf_{j \rightarrow +\infty} \frac{n_{j+1}}{n_j} > 1.$$

Under the weaker assumption that

$$\limsup_{j \rightarrow +\infty} \frac{n_{j+1}}{n_j} > 1,$$

it follows from the Ridout Theorem that the function $f_{\mathbf{n}}$ given by (3.1) takes transcendental values at every point $1/b$, where $b \geq 2$ is an integer (see e.g. Satz 7 from Schneider's monograph [16]), and even at every point $1/\beta$, where β is a Pisot or a Salem number [1] (see also Theorem 3 of [10]).

The latter result can be improved with the methods of the present paper. Namely, we extend Corollary 4 of [8] and Corollary 3.2 of [9] as follows.

Corollary 3. *Let β be a Pisot or a Salem number. For any real number $\eta > 2/3$, the sum of the series*

$$\sum_{j \geq 1} \beta^{-n_j}, \quad \text{where } n_j = 2^{\lfloor j^\eta \rfloor} \text{ for } j \geq 1, \quad (3.2)$$

is transcendental.

The growth of the sequence $(n_j)_{j \geq 1}$ defined in (3.2) shows that our Corollary 3 is not a consequence of the results of [10].

To establish Corollary 3, it is enough to check that, for any positive integer N , the number of positive integers j such that $2^{\lfloor j^\eta \rfloor} \leq N$ is less than some absolute constant times $(\log N)^{1/\eta}$, and to apply Theorem 2 to conclude.

To be precise, to establish Corollary 3, we prove that any real number α having an expansion in base β given by (3.2) is transcendental. We do not need to assume (or to prove) that (3.2) is the β -expansion of α . Namely, this assumption is used in the proof to guarantee that the approximants α_j constructed in the proof of Theorem 2 are (essentially) all different. Under the assumption of Corollary 3, this condition is automatically satisfied.

4. Proofs

The proof of Theorem 2 follows the same lines as that of Theorem 1 of [8]. For convenience, we first explain the case where β is an integer $b \geq 2$. Then, we point out which changes have to be made to treat the case of a real algebraic number $\beta > 1$.

The key point for our argument is the following result of Ridout [13].

For a prime number ℓ and a non-zero rational number x , we set $|x|_\ell := \ell^{-u}$, where u is the exponent of ℓ in the prime decomposition of x . Furthermore, we set $|0|_\ell = 0$. With this notation, the main result of [13] reads as follows.

Theorem (Ridout, 1957). *Let S_1 and S_2 be disjoint finite sets of prime numbers. Let θ be a real algebraic number. Let ε be a positive real number. Then there are only finitely many rational numbers p/q with $q \geq 1$ such that*

$$0 < \left| \theta - \frac{p}{q} \right| \cdot \prod_{\ell \in S_1} |p|_\ell \cdot \prod_{\ell \in S_2} |q|_\ell < \frac{1}{q^{2+\varepsilon}}. \quad (4.1)$$

More precisely, we need a quantitative version of Ridout's Theorem, namely an explicit upper bound for the number of solutions to (4.1). In this direction, Locher [11] proved that, if $\varepsilon < 1/4$, the degree of θ is at most d and its Mahler measure at most H , then (4.1) has at most

$$\mathcal{N}_1(\varepsilon) := c_1(d) e^{7s} \varepsilon^{-s-4} \log(\varepsilon^{-1}) \quad (4.2)$$

solutions p/q with $q \geq \max\{H, 4^{4/\varepsilon}\}$, where s denotes the cardinality of the set $S_1 \cup S_2$, and $c_1(d)$ depends only on d .

Actually, as will be apparent below, in the present application of the quantitative Ridout's Theorem, S_1 is the empty set and we have actually to estimate the total number of solutions to the system of inequalities

$$0 < \left| \theta - \frac{p}{q} \right| < \frac{c}{q^{1+\varepsilon}}, \quad \prod_{\ell \in S_2} |q|_\ell < \frac{c}{q}, \quad (4.3)$$

where c is a positive integer. Every solution to (4.3) with q large is a solution to (4.1), with ε replaced by 2ε , but the converse does not hold. Furthermore, the best known upper bound for the total number of large solutions to (4.3) does not depend on the set S_2 . Namely, if $\varepsilon < 1/4$, then there exists an explicit number $c_2(d)$, depending only on the degree d of θ , such that (4.3) has at most

$$\mathcal{N}_2(\varepsilon) := c_2(d) \varepsilon^{-3} \log(\varepsilon^{-1}) \quad (4.4)$$

solutions p/q with $q \geq \max\{2H, 4^{4/\varepsilon}\}$; see Corollary 5.2 of [9]. Since there is no dependence on s in (4.4), unlike in (4.2), this explains the improvement obtained in [9] on the result from [8].

After these preliminary remarks, let us explain the method of the proof. Let α be an irrational (otherwise, the result is clearly true) real number in $(0, 1)$ and write

$$\alpha = \sum_{k \geq 1} \frac{a_k}{b^k} = 0.a_1 a_2 \dots$$

Define the increasing sequence of positive integers $(n_j)_{j \geq 1}$ by $a_1 = \dots = a_{n_1}$, $a_{n_1} \neq a_{n_1+1}$ and $a_{n_j+1} = \dots = a_{n_{j+1}}$, $a_{n_{j+1}} \neq a_{n_{j+1}+1}$ for every $j \geq 1$. Observe that

$$\text{nbdc}_b(n, \alpha) = \max\{j : n_j \leq n\} \quad (4.5)$$

for $n \geq n_1$, and that $n_j \geq j$ for $j \geq 1$. To construct good rational approximations to α , we simply truncate its b -ary expansion at rank a_{n_j+1} and then complete with repeating the digit a_{n_j+1} . Precisely, for $j \geq 1$, we define the rational number

$$\alpha_j = \sum_{k=1}^{n_j} \frac{a_k}{b^k} + \sum_{k=n_j+1}^{+\infty} \frac{a_{n_j+1}}{b^k} = \sum_{k=1}^{n_j} \frac{a_k}{b^k} + \frac{a_{n_j+1}}{b^{n_j}(b-1)} =: \frac{p_j}{b^{n_j}(b-1)}.$$

Set $q_j := b^{n_j}(b-1)$ and take for S_2 the set of prime divisors of b . Observe that

$$0 < |\alpha - \alpha_j| < \frac{1}{b^{n_{j+1}}}, \quad \prod_{\ell \in S_2} |q_j|_\ell = \frac{b-1}{q_j}. \quad (4.6)$$

On the other hand, the Liouville inequality as stated by Waldschmidt [18], p. 84, asserts that there exists a positive constant c , depending only on α , such that

$$\left| \alpha - \frac{p}{q} \right| \geq \frac{c}{q^d}, \quad \text{for all positive integers } p, q,$$

where d is the degree of α . Consequently, we have

$$n_{j+1} \leq 2dn_j, \quad (4.7)$$

for every sufficiently large integer j , say for $j \geq j_0$.

It then follows from (4.6) and (4.7) that

$$0 < \left| \alpha - \frac{p_j}{q_j} \right| < \frac{(b-1)^{2d}}{q_j^{n_{j+1}/n_j}}, \quad \prod_{\ell \in S_2} |q_j|_\ell = \frac{b-1}{q_j}. \quad (4.8)$$

Note that all the p_j/q_j 's are different. We are in position to apply the quantitative form of the Ridout Theorem to (4.8). Let ε be a real number with $0 < \varepsilon < 1/4$. Let $J > j_0$ be a large positive integer. It follows from Corollary 5.2 of [9] that there exist at most $\mathcal{N}_2(\varepsilon)$ positive integers $j > J$ such that $n_{j+1} \geq (1 + \varepsilon)n_j$. Consequently, we infer from (4.4) and (4.7) that

$$\frac{n_J}{n_{j_0}} = \frac{n_J}{n_{J-1}} \times \dots \times \frac{n_{j_0+1}}{n_{j_0}} \leq (1 + \varepsilon)^J (2d)^{\mathcal{N}_2(\varepsilon)},$$

and

$$\log n_J \ll J\varepsilon + \varepsilon^{-4},$$

where the numerical constant implied in \ll depends only on α . Selecting $\varepsilon = J^{-1/5}$, we get that

$$\log n_J \ll J^{4/5}. \quad (4.9)$$

By (4.5), this implies a lower bound for $\text{nbdc}_b(n, \alpha)$. Here, to get (4.9), we have used a rather crude upper bound for $\mathcal{N}_2(\varepsilon)$. A further refinement can be obtained by means of the trick that allowed us to prove Theorem 3.1 of [9], which is similar to Theorem 2 for $\beta = b$.

Replacing b by an algebraic number $\beta > 1$, everything goes along the same lines, except that we have to apply a suitable extension of Ridout's Theorem, and several technical difficulties arise.

We proceed exactly as above, keep the same notation, and set

$$\alpha_j = \sum_{k=1}^{n_j} \frac{a_k}{\beta^k} + \sum_{k=n_j+1}^{+\infty} \frac{a_{n_j+1}}{\beta^k} = \sum_{k=1}^{n_j} \frac{a_k}{\beta^k} + \frac{a_{n_j+1}}{\beta^{n_j}(\beta-1)} =: \frac{p_j}{\beta^{n_j}(\beta-1)}. \quad (4.10)$$

Here, p_j is an element of the number field generated by β . We have to prove that α_j is distinct from α : unlike when β is an integer, this is not straightforward.

Recall that $a_{n_j+1} = \dots = a_{n_{j+1}}$ and $a_{n_j+1} \neq a_{n_{j+1}+1}$. Assume first that

$$a_{n_j+1} > a_{n_{j+1}+1}. \quad (4.11)$$

Then, using (4.10), we have

$$\alpha_j - \sum_{k=1}^{n_{j+1}} \frac{a_k}{\beta^k} \geq \frac{a_{n_j+1}}{\beta^{n_{j+1}+1}} + \frac{a_{n_j+1}}{\beta^{n_{j+1}+2}}, \quad (4.12)$$

while

$$\alpha - \sum_{k=1}^{n_{j+1}} \frac{a_k}{\beta^k} \leq \frac{a_{n_{j+1}+1}}{\beta^{n_{j+1}+1}} + \frac{1}{\beta^{n_{j+1}+1}} \leq \frac{a_{n_j+1}}{\beta^{n_{j+1}+1}}, \quad (4.13)$$

since, by the property of the β -expansion,

$$\sum_{k \geq r+1} \frac{a_k}{\beta^k} \leq \frac{1}{\beta^r}, \quad \text{for every } r \geq 0. \quad (4.14)$$

Note that $a_{n_j+1} \geq 1$, by (4.11). Combining this with (4.12) and (4.13), we get that

$$\alpha_j - \alpha \geq \frac{1}{\beta^{n_{j+1}+2}}. \quad (4.15)$$

Assume now that

$$a_{n_j+1} < a_{n_{j+1}+1}. \quad (4.16)$$

Then, we have

$$\alpha_j - \sum_{k=1}^{n_{j+1}} \frac{a_k}{\beta^k} = \frac{a_{n_j+1}}{\beta^{n_{j+1}}(\beta-1)}, \quad (4.17)$$

while

$$\alpha - \sum_{k=1}^{n_{j+1}} \frac{a_k}{\beta^k} > \frac{a_{n_{j+1}+1}}{\beta^{n_{j+1}+1}} \geq \frac{a_{n_j+1} + 1}{\beta^{n_{j+1}+1}}, \quad (4.18)$$

by (4.16). Since $a_{n_j+1} < \beta - 1$, we infer from (4.17) that

$$\alpha_j - \sum_{k=1}^{n_{j+1}} \frac{a_k}{\beta^k} < \frac{1}{\beta^{n_{j+1}+1}},$$

and then from (4.18) that

$$\alpha - \alpha_j > 0.$$

Note also that, by (4.18), at least one of the following statements holds:

$$n_{j+2} = n_{j+1} + 1 \quad \text{and} \quad 0 = a_{n_{j+2}+1} < a_{n_{j+1}+1} \quad (4.19)$$

or

$$\alpha - \alpha_j > \frac{1}{\beta^{n_{j+1}+2}}. \quad (4.20)$$

On the other hand, we check that

$$|\alpha - \alpha_j| \ll \frac{1}{\beta^{n_{j+1}}}, \quad \text{for } j \geq 1. \quad (4.21)$$

Here, and throughout the end of the paper, the constants implied by \ll depend only on α and β . Disregarding the indices j for which we are in case (4.19) (and this concerns at most one index in every pair $(j, j+1)$), we infer from (4.15), (4.20) and (4.21) that the number of occurrences of a given element in the sequence $(\alpha_j)_{j \geq 1}$ is bounded by an absolute constant.

Now, we apply the extension to number fields of the aforementioned results of Ridout and Locher. We keep the notation from Section 6 of [2], noticing that the r_n (resp. s_n) in that paper corresponds to our n_j (resp. to 1). In particular, the height function H is defined as in [2].

Let \mathbf{K} be the number field generated by α and β and denote by D its degree. We consider the following linear forms, in two variables and with algebraic coefficients. For the place v_0 corresponding to the embedding of \mathbf{K} defined by $\beta \mapsto \beta$, set $L_{1,v_0}(x, y) = x$ and $L_{2,v_0}(x, y) = \alpha(\beta - 1)x + y$. It follows from (4.10) and (4.21) that

$$|L_{2,v_0}(\beta^{n_j}, -p_j)|_{v_0} \ll \frac{1}{\beta^{(n_{j+1}-n_j)/D}},$$

where we have chosen the continuation of $|\cdot|_{v_0}$ to $\overline{\mathbf{Q}}$ defined by $|x|_{v_0} = |x|^{1/D}$.

Denote by S'_∞ the set of all other infinite places on \mathbf{K} and by S_0 the set of all finite places v on \mathbf{K} for which $|\beta|_v \neq 1$. For any v in $S_0 \cup S'_\infty$, set $L_{1,v}(x, y) = x$ and $L_{2,v}(x, y) = y$. Denote by S the union of S_0 and the infinite places on \mathbf{K} . Clearly, for any v in S , the linear forms $L_{1,v}$ and $L_{2,v}$ are linearly independent.

To simplify the exposition, set

$$\mathbf{x}_j = (\beta^{n_j}, -p_j).$$

We wish to estimate the product

$$\Pi_j := \prod_{v \in S} \prod_{i=1}^2 \frac{|L_{i,v}(\mathbf{x}_j)|_v}{|\mathbf{x}_j|_v}$$

from above. Arguing exactly as in [2], we get that

$$\begin{aligned}\Pi_j &\ll n_j^D \beta^{-n_{j+1}/D} M(\beta)^{n_j/D} \prod_{v \in S} |\mathbf{x}_j|_v^{-2} \\ &\ll n_j^D \beta^{-n_{j+1}/D} M(\beta)^{n_j/D} H(\mathbf{x}_j)^{-2},\end{aligned}\tag{4.22}$$

since $|\mathbf{x}_j|_v = 1$ if v does not belong to S .

Note that

$$\beta^{n_j} \ll H(\mathbf{x}_j) \ll n_j^D M(\beta)^{n_j}.\tag{4.23}$$

Let ρ denote a positive real number that is strictly smaller than the right-hand side of (2.1). Assume that there are arbitrarily large integers n such that

$$\text{nbdc}_\beta(n, \alpha) \leq \rho \log n.$$

Consequently, there must be infinitely many indices j with

$$n_{j+1} \geq \exp\{\rho^{-1}\}n_j.$$

It then follows from (4.22) and (4.23) that there are a positive real number ε and arbitrarily large integers j such that

$$\Pi_j \ll H(\mathbf{x}_j)^{-2-\varepsilon}.$$

We then get infinitely many indices j such that p_j/β^{n_j} takes the same value. This contradicts the fact that the number of occurrences of a given element in the sequence $(\alpha_j)_{j \geq 1}$ is bounded by an absolute constant, and proves Theorem 1.

From now on, we assume that $M(\beta) = \beta$. We infer from (4.22) and (4.23) that

$$\Pi_j \ll H(\mathbf{x}_j)^{-2-(n_{j+1}/n_j-1)/(2D)}$$

as soon as j is sufficiently large.

We need a suitable extension of Corollary 5.2 from [9] to conclude (unfortunately, the notations used in [9] differ from ours). Exactly as in the case when β is an integer, we do not have to consider a product of linear forms, but rather a system

$$\begin{aligned}|L_{1,v_0}(\mathbf{x}_j)|_{v_0} &\leq \kappa H(\mathbf{x}_j), & |L_{2,v_0}(\mathbf{x}_j)|_{v_0} &\leq H(\mathbf{x}_j)^{-\delta}, \\ |L_{1,v}(\mathbf{x}_j)|_v &\leq \kappa H(\mathbf{x}_j)^{-c_v}, & |L_{2,v}(\mathbf{x}_j)|_v &\leq H(\mathbf{x}_j)^\eta, \quad (v \in S'_\infty), \\ |L_{1,v}(\mathbf{x}_j)|_v &\leq \kappa H(\mathbf{x}_j)^{-c_v}, & |L_{2,v}(\mathbf{x}_j)|_v &\leq 1, \quad (v \in S_0).\end{aligned}$$

Here, κ is a positive real number, the c_v are defined by $|\beta|_v = \beta^{-c_v/D}$ and $\delta = (s' + 1)\eta$, where s' is the cardinality of S'_∞ . Observe that $\sum_{v \in S'_\infty \cup S_0} c_v = 1$. We do not work out the technical details. Everything goes along the same lines as in [9]. It remains to note that, by the general form of the Liouville inequality (as in [18], p. 83), we get that

$$|\alpha - \alpha_j| \gg \beta^{-Dn_j}.$$

This provides us with the needed extension of (4.7) and completes the sketch of the proof of Theorem 2.

References

- [1] B. Adamczewski, *Transcendance « à la Liouville » de certains nombres réels*, C. R. Acad. Sci. Paris 338 (2004), 511–514.
- [2] B. Adamczewski and Y. Bugeaud, *Dynamics for β -shifts and Diophantine approximation*, Ergod. Th. & Dynam. Sys. 27 (2007), 1695–1711.
- [3] D. H. Bailey, J. M. Borwein, R. E. Crandall and C. Pomerance, *On the binary expansions of algebraic numbers*, J. Théor. Nombres Bordeaux 16 (2004), 487–518.
- [4] A. Bertrand-Mathis, *Développements en base θ , répartition modulo un de la suite $(x\theta^n)_{n \geq 0}$, langages codés et θ -shift*, Bull. Soc. Math. France 114 (1986), 271–323.
- [5] F. Blanchard, *β -expansions and symbolic dynamics*, Theoret. Comput. Sci. 65 (1989), 131–141.
- [6] D. Boyd, *Salem numbers of degree four have periodic expansions*. In: Théorie des nombres (Quebec, PQ, 1987), pp. 57–64, de Gruyter, 1989.
- [7] D. Boyd, *On the beta expansion for Salem numbers of degree 6*, Math. Comp. 65 (1996), 861–875.
- [8] Y. Bugeaud, *On the b -ary expansion of an algebraic number*, Rend. Sem. Mat. Univ. Padova 118 (2007), 217–233.
- [9] Y. Bugeaud and J.-H. Evertse, *On two notions of complexity of algebraic numbers*, Acta Arith. 133 (2008), 221–250.
- [10] P. Corvaja and U. Zannier, *Some new applications of the subspace theorem*, Compositio Math. 131 (2002), 319–340.
- [11] H. Locher, *On the number of good approximations of algebraic numbers by algebraic numbers of bounded degree*, Acta Arith. 89 (1999), 97–122.
- [12] A. Rényi, *Representations for real numbers and their ergodic properties*, Acta Math. Acad. Sci. Hung. 8 (1957), 477–493.
- [13] D. Ridout, *Rational approximations to algebraic numbers*, Mathematika 4 (1957), 125–131.
- [14] T. Rivoal, *On the bits counting function of real numbers*, J. Austral. Math. Soc. 85 (2008), 95–111.
- [15] K. Schmidt, *On periodic expansions of Pisot and Salem numbers*, Bull. London Math. Soc. 12 (1980), 269–278.
- [16] Th. Schneider, Einführung in die transzendenten Zahlen. Springer-Verlag, Berlin-Göttingen-Heidelberg, 1957.
- [17] J.-L. Verger-Gaugry, *On gaps in Rényi β -expansions of unity and β -integers for $\beta > 1$ an algebraic number*, Ann. Inst. Fourier (Grenoble) 56 (2006), 2565–2579.

- [18] M. Waldschmidt, *Diophantine Approximation on Linear Algebraic Groups. Transcendence properties of the exponential function in several variables*. Grundlehren der Mathematischen Wissenschaften 326. Springer–Verlag, Berlin, 2000.

Yann Bugeaud
Université Louis Pasteur
U. F. R. de mathématiques
7, rue René Descartes
67084 STRASBOURG Cedex (FRANCE)
bugeaud@math.u-strasbg.fr