

SIGNED WORDS AND PERMUTATIONS, I; A FUNDAMENTAL TRANSFORMATION

DOMINIQUE FOATA AND GUO-NIU HAN

This paper is dedicated to the memory of Percy Alexander MacMahon.

ABSTRACT. The statistics major index and inversion number, usually defined on ordinary words, have their counterparts in signed words, namely the so-called flag-major index and flag-inversion number. We give the construction of a new transformation on those signed words that maps the former statistic onto the latter one. It is proved that the transformation also preserves two other set-statistics: the inverse line of route and the lower records.

1. INTRODUCTION

The *second fundamental transformation*, as it was called later on (see [16], chap. 10 or [15], ex. 5.1.1.19), was described in these proceedings [8]. Let $w = x_1x_2 \dots x_m$ be a (finite) word, whose letters x_1, x_2, \dots, x_m are integers. The integer-valued statistics *Inversion Number* “inv” and *Major Index* “maj” attached to the word w are defined by

$$(1.1) \quad \text{inv } w := \sum_{1 \leq i \leq m-1} \sum_{i < j} \chi(x_i > x_j);$$

$$(1.2) \quad \text{maj } w := \sum_{1 \leq i \leq m-1} i \chi(x_i > x_{i+1});$$

making use of the χ -notation that maps each statement A to the value $\chi(A) = 1$ or 0 depending on whether A is true or not.

If $\mathbf{m} = (m_1, m_2, \dots, m_r)$ is a sequence of r nonnegative integers, the rearrangement class of the nondecreasing word $1^{m_1}2^{m_2} \dots r^{m_r}$, that is, the class of all the words than can be derived from $1^{m_1}2^{m_2} \dots r^{m_r}$ by permutation of the letters, is denoted by $R_{\mathbf{m}}$. The second fundamental transformation, denoted by Φ , maps each word w on another word $\Phi(w)$ and has the following properties:

- (a) $\text{maj } w = \text{inv } \Phi(w)$;
- (b) $\Phi(w)$ is a rearrangement of w and the restriction of Φ to each rearrangement class $R_{\mathbf{m}}$ is a bijection of $R_{\mathbf{m}}$ onto itself.

Further properties were proved later on by Foata, Schützenberger [10] and Björner, Wachs [5], in particular, when the transformation is restricted to act on rearrangement classes $R_{\mathbf{m}}$ such that $m_1 = \dots = m_r = 1$, that is, on symmetric groups \mathfrak{S}_r .

Received by the editors March 7, 2005.

1991 *Mathematics Subject Classification*. Primary 05A15, 05A30, 05E15.

Key words and phrases. Flag-inversion number, flag-major index, length function, signed permutations, signed words.

The purpose of this paper is to construct an analogous transformation not simply on words, but on *signed words*, so that new equidistribution properties on classical statistics, such as the (Coxeter) *length function* (see [7, p. 9], [14, p. 12]), defined on the group B_n of the *signed permutations* can be derived. By *signed word* we understand a word $w = x_1 x_2 \dots x_m$, whose letters are positive or negative integers. If $\mathbf{m} = (m_1, m_2, \dots, m_r)$ is a sequence of nonnegative integers such that $m_1 + m_2 + \dots + m_r = m$, let $B_{\mathbf{m}}$ be the set of all rearrangements $w = x_1 x_2 \dots x_m$ of the sequence $1^{m_1} 2^{m_2} \dots r^{m_r}$, with the convention that some letters i may be replaced by their opposite values $-i$. For typographical reasons we shall use the notation $\bar{i} := -i$ in the sequel. The class $B_{\mathbf{m}}$ contains $2^m \binom{m}{m_1, m_2, \dots, m_r}$ signed words. When $m_1 = m_2 = \dots = m_r = 1$, $m = r$, the class $B_{\mathbf{m}}$ is simply the group B_m of the signed permutations of order m .

Next, the statistics “inv” and “maj” must be adapted to signed words and correspond to classical statistics when applied to signed permutations. Let

$$(\omega; q)_n := \begin{cases} 1, & \text{if } n = 0; \\ (1 - \omega)(1 - \omega q) \dots (1 - \omega q^{n-1}), & \text{if } n \geq 1; \end{cases}$$

denote the usual q -ascending factorial in a ring element ω and

$$\begin{bmatrix} m_1 + \dots + m_r \\ m_1, \dots, m_r \end{bmatrix}_q := \frac{(q; q)_{m_1 + \dots + m_r}}{(q; q)_{m_1} \dots (q; q)_{m_r}}$$

be the q -multinomial coefficient. Back to MacMahon [17, 18, 19] it was known that the above q -multinomial coefficient, which is the true q -analog of the cardinality of $R_{\mathbf{m}}$, was the generating function for the class $R_{\mathbf{m}}$ by either one of the statistics “inv” or “maj.” Consequently, the generating function for $B_{\mathbf{m}}$ by the new statistics that are to be introduced on $B_{\mathbf{m}}$ must be a *plausible* q -analog of the cardinality of $B_{\mathbf{m}}$. The most natural q -analog we can think of is certainly $(-q; q)_m \begin{bmatrix} m_1 + \dots + m_r \\ m_1, \dots, m_r \end{bmatrix}_q$, that tends to $2^m \binom{m_1 + \dots + m_r}{m_1, \dots, m_r}$ when q tends to 1. As a substitute for “inv” we are led to introduce the following statistic “finv,” called the *flag-inversion number*, which will be shown to meet our expectation, that is,

$$(1.3) \quad (-q; q)_m \begin{bmatrix} m_1 + \dots + m_r \\ m_1, \dots, m_r \end{bmatrix}_q = \sum_{w \in B_{\mathbf{m}}} q^{\text{finv } w}.$$

This identity is easily proved by induction on r . Let $w = x_1 x_2 \dots x_m$ be a signed word from the class $B_{\mathbf{m}}$. To define $\text{finv } w$ we use “inv” defined in (1.1), together with

$$(1.4) \quad \bar{\text{inv}} w := \sum_{1 \leq i \leq m-1} \sum_{i < j} \chi(\bar{x}_i > x_j)$$

and define

$$(1.5) \quad \text{finv } w := \text{inv } w + \bar{\text{inv}} w + \sum_{1 \leq j \leq m} \chi(x_j < 0).$$

The salient feature of this definition of “finv” is the fact that it does *not* involve the *values* of the letters, but only the *comparisons* between letters, so that it can be applied to each *arbitrary* signed word. Moreover, the definition of “finv” is similar to that of “fmaaj” given below in (1.7). Finally, its restriction to the group B_m of the *signed permutations* is the traditional *length function*:

$$(1.6) \quad \text{finv}|_{B_m} = \ell.$$

This is easily shown, for instance, by using the formula derived by Brenti [6] for the length function ℓ over B_n , that reads.

$$\ell w = \text{inv } w + \sum_{1 \leq j \leq m} |x_j| \chi(x_j < 0).$$

Next, the statistic “maj” is to be replaced by “fmaj”, the *flag-major index*, introduced by Adin and Roichman [1] for *signed permutations*. The latter authors (see also [2]) showed that “fmaj” was equidistributed with the length function ℓ over B_n . Their definition of “fmaj” can be used *verbatim* for signed words, as well as their definition of “fdes.” For a signed word $w = x_1 x_2 \dots x_m$ those definitions read:

$$(1.7) \quad \text{fmaj } w := 2 \text{maj } w + \sum_{1 \leq j \leq m} \chi(x_j < 0);$$

$$(1.8) \quad \text{fdes } w := 2 \text{des } w + \chi(x_1 < 0);$$

where “des” is the usual *number of descents* $\text{des } w := \sum_i \chi(x_i > x_{i+1})$. We postpone the construction of our transformation Ψ on signed words to the next section. The main purpose of this paper is to prove the following theorem.

Theorem 1.1. *The transformation Ψ constructed in section 2 has the following properties:*

- (a) $\text{fmaj } w = \text{finv } \Psi(w)$ for every signed word w ;
- (b) the restriction of Ψ to each rearrangement class $B_{\mathbf{m}}$ of signed words is a bijection of $B_{\mathbf{m}}$ onto itself, so that “fmaj” and “finv” are equidistributed over each class $B_{\mathbf{m}}$.

Definition 1.1. Let $w = x_1 x_2 \dots x_m \in B_{\mathbf{m}}$ be a signed word. We say that a nonnegative integer i belongs to the *inverse ligne of route*, $\text{Iligne } w$, of w , if one of the following two conditions holds:

- (1) $i = 0$, $m_1 \geq 1$ and the rightmost letter x_k satisfying $|x_k| = 1$ is equal to $\bar{1}$;
- (2) $i \geq 1$, $m_i = m_{i+1} = 1$ and the rightmost letter that belongs to $\{i, \bar{i}, i+1, \overline{i+1}\}$ is equal to i or $\overline{i+1}$.

For example, with $w = \bar{4}4\bar{1}325\bar{5}6\bar{7}$ we have: $\text{Iligne } w = \{0, 2, 6\}$.

Remark. The expression “line of route” was used by Foulkes [11,12]. We have added the letter “g” making up “ligne of route,” thus bringing a slight touch of French. Notice that 0 may or may not belong to the inverse ligne of route. The *ligne of route* of a signed word $w = x_1 x_2 \dots x_m$ is defined to be the set, denoted by $\text{Ligne } w$, of all the i ’s such that either $1 \leq i \leq m-1$ and $x_i > x_{i+1}$, or $i = 0$ and $x_1 < 0$. In particular, $\text{maj } w = \sum_{0 \leq i \leq m-1} i \chi(i \in \text{Ligne } w)$. Finally, if w is a *signed permutation*, then $\text{Iligne } w = \text{Ligne } w^{-1}$. For *ordinary permutations*, some authors speak of *descent set* and *descent set of the inverse*, instead of ligne of route and inverse ligne of route, respectively.

Theorem 1.2. *The transformation Ψ constructed in section 2 preserves the inverse ligne of route:*

- (c) $\text{Iligne } \Psi(w) = \text{Iligne } w$ for every signed word w .

Definition 1.2. Let $w = x_1x_2 \dots x_m$ be a signed word of length m . A letter x_i is said to be a *lower record* of w , if either $i = m$, or $1 \leq i \leq m-1$ and $|x_i| < |x_j|$ for all j such that $i+1 \leq j \leq m$. When reading the lower records of w from left to right, we get a *signed subword* $x_{i_1}x_{i_2} \dots x_{i_k}$, called the *lower record subword*, denoted by $\text{Lower } w$, which has the property that: $\min_i x_i = |x_{i_1}| < |x_{i_2}| < \dots < |x_{i_k}| = |x_m|$. The notion of lower record is classical in the statistical literature. In combinatorics the expression “strict right-to-left minimum” is also used.

With our previous example $w = \bar{4}4\bar{1}325\bar{5}6\bar{7}$ we get $\text{Lower } w = \bar{1}2\bar{5}6\bar{7}$. Our third goal is to prove the following result.

Theorem 1.3. *The transformation Ψ constructed in section 2 preserves all the lower records:*

(d) $\text{Lower } \Psi(w) = \text{Lower } w$ for every signed word w .

For each signed permutation $w = x_1x_2 \dots x_m$ let

$$(1.11) \quad \text{ifmaj } w := 2 \sum_{1 \leq j \leq m} i \chi(j \in \text{Iligne } w) + \sum_{1 \leq j \leq m} \chi(x_j < 0);$$

$$(1.12) \quad \text{ifdes } w := 2 \sum_{1 \leq j \leq m} \chi(j \in \text{Iligne } w) + \chi(x_i = -1 \text{ for some } i).$$

It is immediate to verify that

$$\text{finv } w = \text{finv } w^{-1}, \quad \text{ifmaj } w = \text{fmaj } w^{-1}, \quad \text{ifdes } w = \text{fdes } w^{-1},$$

where w^{-1} denotes the inverse of the signed permutation w (written as a linear word $w^{-1} = w^{-1}(1) \dots w^{-1}(m)$).

Let $\mathbf{i}w := w^{-1}$; then the chain

$$\begin{array}{ccccccc} w & \xrightarrow{\mathbf{i}} & w_1 & \xrightarrow{\Psi} & w_2 & \xrightarrow{\mathbf{i}} & w_3 \\ \left(\begin{smallmatrix} \text{fdes} \\ \text{ifmaj} \end{smallmatrix} \right) & & \left(\begin{smallmatrix} \text{ifdes} \\ \text{fmaj} \end{smallmatrix} \right) & & \left(\begin{smallmatrix} \text{ifdes} \\ \text{finv} \end{smallmatrix} \right) & & \left(\begin{smallmatrix} \text{fdes} \\ \text{finv} \end{smallmatrix} \right) \end{array}$$

shows that the four generating polynomials $\sum t^{\text{fdes } w} q^{\text{ifmaj } w}$, $\sum t^{\text{ifdes } w} q^{\text{fmaj } w}$, $\sum t^{\text{ifdes } w} q^{\text{finv } w}$ and $\sum t^{\text{fdes } w} q^{\text{finv } w}$ ($w \in B_m$) are identical. Their analytic expression will be derived in a forthcoming paper [9].

2. THE CONSTRUCTION OF THE TRANSFORMATION

For each signed word $w = x_1x_2 \dots x_m$ the first or leftmost (resp. last or rightmost) letter x_1 (resp. x_m) is denoted by $F(w)$ (resp. $L(w)$). Next, define $\mathbf{s}_1 w := \bar{x}_1x_2 \dots x_m$. The transformation \mathbf{s}_1 changes the sign of the first letter. Together with \mathbf{s}_1 the main ingredients of our transformation are the bijections γ_x and δ_x defined for each integer x , as follows.

If $L(w) \leq x$ (resp. $L(w) > x$), then w admits the unique factorization

$$(v_1y_1, v_2y_2, \dots, v_py_p),$$

called its *x-right-to-left factorisation* having the following properties:

- (i) each y_i ($1 \leq i \leq p$) is a *letter* verifying $y_i \leq x$ (resp. $y_i > x$);
- (ii) each v_i ($1 \leq i \leq p$) is a factor which is either empty or has all its letters greater than (resp. smaller than or equal to) x .

Then, γ_x is defined to be the bijection that maps $w = v_1 y_1 v_2 y_2 \dots v_p y_p$ onto the signed word

$$(2.1) \quad \gamma_x(w) := y_1 v_1 y_2 v_2 \dots y_p v_p.$$

In a dual manner, if $F(w) \geq x$ (resp. $F(w) < x$) the signed word w admits the unique factorization

$$(z_1 w_1, z_2 w_2, \dots, z_q w_q)$$

called its *x-left-to-right factorisation* having the following properties:

- (i) each z_i ($1 \leq i \leq q$) is a *letter* verifying $z_i \geq x$ (resp. $z_i < x$);
- (ii) each w_i ($1 \leq i \leq q$) is a factor which is either empty or has all its letters less than (resp. greater than or equal to) x .

Then, δ_x is defined to be the bijection that sends $w = z_1 w_1 z_2 w_2 \dots z_q w_q$ onto the signed word

$$(2.2) \quad \delta_x(w) := w_1 z_1 w_2 z_2 \dots w_q z_q.$$

Next, if $(v_1 y_1, v_2 y_2, \dots, y_p v_p)$ is the *x-right-to-left factorization* of w , we define

$$(2.3) \quad \beta_x(w) := \begin{cases} \delta_{\bar{x}} \gamma_x(w), & \text{if either } \bar{x} \leq y_1 \leq x, \text{ or } x < y_1 < \bar{x}; \\ \delta_{\bar{x}} \mathbf{s}_1 \gamma_x(w), & \text{otherwise.} \end{cases}$$

The fundamental transformation Ψ on signed words that is the main object of this paper is defined as follows: if w is a one-letter signed word, let $\Psi(w) := w$; if it has more than one letter, write the word as wx , where x is the last letter. By induction determine $\Psi(w)$, then apply β_x to $\Psi(w)$ and define $\Psi(wx)$ to be the juxtaposition product:

$$(2.4) \quad \Psi(wx) := \beta_x(\Psi(w))x.$$

The proof of Theorem 1.1 is given in section 3. It is useful to notice the following relation

$$(2.5) \quad y_1 \leq x \Leftrightarrow L(w) \leq x$$

and the identity

$$(2.6) \quad \Psi(wx) = \Psi(w)x, \quad \text{whenever } x < -\max\{|x_i|\} \text{ or } x \geq \max\{|x_i|\}.$$

Example. Let $w = 3\bar{2}1\bar{3}\bar{4}3$. The factorizations used in the definitions of γ_x and $\delta_{\bar{x}}$ are indicated by vertical bars. First, $\Psi(3) = 3$. Then

$$\begin{aligned} |3| &\xrightarrow{\gamma_{\bar{2}}} 3 \xrightarrow{\mathbf{s}_1} |\bar{3}| \xrightarrow{\delta_2} \bar{3}, \text{ so that } \Psi(3\bar{2}) = \bar{3}\bar{2}; \\ |\bar{3}|\bar{2}| &\xrightarrow{\gamma_1} \bar{3}\bar{2} \xrightarrow{\mathbf{s}_1} |3\bar{2}| \xrightarrow{\delta_{\bar{1}}} \bar{2}3, \text{ so that } \Psi(3\bar{2}1) = \bar{2}31; \\ |\bar{2}|3|1| &\xrightarrow{\gamma_{\bar{3}}} |\bar{2}3|1| \xrightarrow{\delta_3} 3\bar{2}1, \text{ so that } \Psi(3\bar{2}1\bar{3}) = 3\bar{2}1\bar{3}; \\ &\text{and } \Psi(3\bar{2}1\bar{3}\bar{4}) = 3\bar{2}1\bar{3}\bar{4}, \text{ because of (2.6);} \\ |3|\bar{2}|1|\bar{3}|\bar{4}| &\xrightarrow{\gamma_3} |3|\bar{2}|1|\bar{3}\bar{4}| \xrightarrow{\delta_{\bar{3}}} 3\bar{2}1\bar{4}\bar{3}. \end{aligned}$$

Thus, with $w = 3\bar{2}1\bar{3}\bar{4}3$ we get $\Psi(w) = 3\bar{2}1\bar{4}\bar{3}3$. We verify that $\text{fmaj } w = \text{finv } \Psi(w) = 19$, $\text{lligne } w = \text{lligne } \Psi(w) = \{1\}$, $\text{Lower } w = \text{Lower } \Psi(w) = 13$.

3. PROOF OF THEOREM 1.1

Before proving the theorem we state a few properties involving the above statistics and transformations. Let $|w|$ be the number of letters of the signed word w and $|w|_{>x}$ be the number of its letters greater than x with analogous expressions involving the subscripts “ $\geq x$ ”, “ $< x$ ” and “ $\leq x$ ”. We have:

$$(3.1) \quad \text{fmaj } wx = \text{fmaj } w + \chi(x < 0) + 2|w| \chi(L(w) > x);$$

$$(3.2) \quad \text{finv } wx = \text{finv } w + |w|_{>x} + |w|_{<\bar{x}} + \chi(x < 0);$$

$$(3.3) \quad \text{finv } \gamma_x(w) = \text{finv } w + |w|_{\leq x} - |w| \chi(L(w) \leq x);$$

$$(3.4) \quad \text{finv } \delta_x(w) = \text{finv } w + |w|_{\geq x} - |w| \chi(F(w) \geq x).$$

Next, let y_1 denote the first letter of the signed word w'' . Then

$$(3.5) \quad \text{finv } \mathbf{s}_1 w'' = \text{finv } w'' + \chi(y_1 > 0) - \chi(y_1 < 0);$$

$$(3.6) \quad \begin{aligned} |\mathbf{s}_1 w''|_{>x} &= |w''|_{>x} + \chi(x > 0)(\chi(y_1 < \bar{x}) - \chi(x < y_1)) \\ &\quad + \chi(x < 0)(\chi(y_1 \leq x) - \chi(\bar{x} \leq y_1)). \end{aligned}$$

Theorem 1.1 is now proved by induction on the word length. Assume that $\text{fmaj } w = \text{finv } \Psi(w)$ for a given w . Our purpose is to show that

$$(3.7) \quad \text{fmaj } wx = \text{finv } \Psi(wx)$$

holds for all letters x . Let $w' = \Psi(w)$, so that by (2.4) the words w and w' have the same rightmost letter. Denote the x -right-to-left factorization of w' by $(v_1 y_1, \dots, v_p y_p)$. By (2.3) the signed word $v := \beta_x(w')$ is defined by the chain

$$(3.8) \quad \begin{aligned} w' = v_1 y_1 \dots v_p y_p &\xrightarrow{\gamma_x} w'' = y_1 v_1 \dots y_p v_p = z_1 w_1 \dots z_q w_q \\ &\xrightarrow{\delta_{\bar{x}}} v = w_1 z_1 \dots w_q z_q \end{aligned}$$

if either $\bar{x} \leq y_1 \leq x$, or $x < y_1 < \bar{x}$, and by the chain

$$(3.9) \quad \begin{aligned} w' = v_1 y_1 \dots v_p y_p &\xrightarrow{\gamma_x} w'' = y_1 v_1 \dots y_p v_p \\ &\xrightarrow{\mathbf{s}_1} w''' = \bar{y}_1 v_1 \dots y_p v_p = z_1 w_1 \dots z_q w_q \\ &\xrightarrow{\delta_{\bar{x}}} v = w_1 z_1 \dots w_q z_q, \end{aligned}$$

otherwise. Notice that $(z_1 w_1, \dots, z_q w_q)$ designates the \bar{x} -left-to-right factorization of w'' in chain (3.8) and of w''' in chain (3.9).

(i) Suppose that one of the conditions $\bar{x} \leq y_1 \leq x$, $x < y_1 < \bar{x}$ holds, so that (3.8) applies. We have

$$\begin{aligned} \text{finv } \Psi(wx) &= \text{finv } vx = \text{finv } v + |v|_{>x} + |v|_{<\bar{x}} + \chi(x < 0) && [\text{by (3.2)}] \\ \text{finv } v &= \text{finv } \delta_{\bar{x}}(w'') = \text{finv } w'' + |w''|_{\geq \bar{x}} - |w''| \chi(F(w'') \geq \bar{x}) && [\text{by (3.4)}] \\ \text{finv } w'' &= \text{finv } \gamma_x(w') = \text{finv } w' + |w'|_{\leq x} - |w'| \chi(L(w') \leq x) && [\text{by (3.3)}] \\ \text{finv } w' &= \text{fmaj } w && [\text{by induction}] \\ \text{fmaj } w &= \text{fmaj } wx - \chi(x < 0) - 2|w| \chi(L(w) > x). && [\text{by (3.1)}] \end{aligned}$$

By induction,

$$(3.10) \quad L(w) = L(w') \text{ and } \chi(L(w') > x) = 1 - \chi(L(w') \leq x).$$

Also $F(w'') = y_1$. As w' , w'' , v are true rearrangements of each other, we have $|v|_{>x} + |w'|_{\leq x} = |w|$, $|v|_{<\bar{x}} + |w''|_{\geq \bar{x}} = |w|$. Hence,

$$\text{finv } \Psi(wx) = \text{fmaj } wx + |w|[\chi(L(w') \leq x) - \chi(y_1 \geq \bar{x})].$$

By (2.5), if $\bar{x} \leq y_1 \leq x$ holds, then $L(w') \leq x$ and the expression between brackets is null. If $x < y_1 < \bar{x}$ holds, then $L(w') > x$ and the same expression is also null. Thus (3.7) holds.

(ii) Suppose that none of the conditions $\bar{x} \leq y_1 \leq x$, $x < y_1 < \bar{x}$ holds, so that (3.9) applies. We have

$$\begin{aligned} \text{finv } \Psi(wx) &= \text{finv } vx = \text{finv } v + |v|_{>x} + |v|_{<\bar{x}} + \chi(x < 0) && [\text{by (3.2)}] \\ \text{finv } v &= \text{finv } \delta_{\bar{x}}(w''') = \text{finv } w''' + |w'''|_{\geq \bar{x}} - |w'''| \chi(F(w''') \geq \bar{x}) && [\text{by (3.4)}] \\ \text{finv } w''' &= \text{finv } \mathbf{s}_1 w'' = \text{finv } w'' + \chi(y_1 > 0) - \chi(y_1 < 0) && [\text{by (3.5)}] \\ \text{finv } w'' &= \text{finv } \gamma_x(w') = \text{finv } w' + |w'|_{\leq x} - |w'| \chi(L(w') \leq x) && [\text{by (3.3)}] \\ \text{finv } w' &= \text{fmaj } w && [\text{by induction}] \\ \text{fmaj } w &= \text{fmaj } wx - \chi(x < 0) - 2|w| \chi(L(w) > x). && [\text{by (3.1)}] \end{aligned}$$

Moreover, $F(w''') = \bar{y}_1$, so that $\chi(F(w''') \geq \bar{x}) = \chi(\bar{y}_1 \geq \bar{x}) = \chi(y_1 \leq x) = \chi(L(w') \leq x) = \chi(L(w) \leq x)$. As v and w''' are rearrangements of each other, we have $|v|_{<\bar{x}} + |w'''|_{\geq \bar{x}} = |w|$. Using (3.6) since $|v|_{>x} = |w'''|_{>x} = |\mathbf{s}_1 w''|_{>x}$ we have:

$$\begin{aligned} \text{finv } \Psi(wx) &= |w''|_{>x} + \chi(x > 0)(\chi(y_1 < \bar{x}) - \chi(x < y_1)) \\ &\quad + \chi(x < 0)(\chi(y_1 \leq x) - \chi(\bar{x} \leq y_1)) + \chi(x < 0) \\ &\quad + |w| - |w| \chi(y_1 \leq x) + \chi(y_1 > 0) - \chi(y_1 < 0) \\ &\quad + |w'|_{\leq x} - |w'| \chi(y_1 \leq x) \\ &\quad + \text{fmaj } wx - \chi(x < 0) - 2|w| + 2|w| \chi(y_1 \leq x) \\ &= \text{fmaj } wx + \chi(x > 0)(\chi(y_1 < \bar{x}) - \chi(x < y_1)) \\ &\quad + \chi(x < 0)(\chi(y_1 \leq x) - \chi(\bar{x} \leq y_1)) \\ &\quad + \chi(y_1 > 0) - \chi(y_1 < 0) \\ &= \text{fmaj } wx, \end{aligned}$$

for, if none of the conditions $\bar{x} \leq y_1 \leq x$, $x < y_1 < \bar{x}$ holds, then one of the following four ones holds: (a) $y_1 > x > 0$; (b) $\bar{y}_1 > x > 0$; (c) $y_1 \leq x < 0$; (d) $\bar{y}_1 \leq x < 0$; and in each case the sum of the factors in the above sum involving χ is zero.

The construction of Ψ is perfectly reversible. First, note that \mathbf{s}_1 is an involution and the maps γ_x , $\delta_{\bar{x}}$ send each class $B_{\mathbf{m}}$ onto itself, so that their inverses are perfectly defined. They can also be described by means of left-to-right and right-to-left factorizations. Let us give the construction of the inverse Ψ^{-1} of Ψ . Of course, $\Psi^{-1}(v) := v$ if v is a one-letter word. If vx is a signed word, whose last letter is x , determine $v' := \delta_{\bar{x}}^{-1}(v)$ and let z_1 be its first letter. If one of the conditions $\bar{x} \leq z_1 \leq x$ or $x < z_1 < \bar{x}$ holds, the chain (3.8) is to be used in reverse order, so that $\Psi^{-1}(vx) := (\Psi^{-1} \gamma_x^{-1} \delta_{\bar{x}}^{-1}(v))x$. If none of those two conditions holds, then $\Psi^{-1}(vx) := (\Psi^{-1} \gamma_x^{-1} \mathbf{s}_1 \delta_{\bar{x}}^{-1}(v))x$. \square

4. PROOFS OF THEOREMS 1.2 AND 1.3

Before proving Theorem 1.2 we note the following two properties.

Property 4.1. *Let $I_x = \{i \in \mathbb{Z} : i < -|x|\}$ (resp. $J_x = \{i \in \mathbb{Z} : -|x| < i < |x|\}$, resp. $K_x = \{i \in \mathbb{Z} : |x| < i\}$) and w be a signed word. Then, the bijections γ_x and $\delta_{\bar{x}}$ do not modify the mutual order of the letters of w that belong to I_x (resp. J_x , resp. K_x).*

Proof. Let $y \in I_x$ and $z \in I_x$ (resp. $y \in J_x$ and $z \in J_x$, resp. $y \in K_x$ and $z \in K_x$) be two letters of w with y to the left of z . In the notations of (2.1) (resp. of (2.2)) both y, z are, either among the y_i 's (resp. the z_i 's), or letters of the v_i 's (resp. the w_i 's). Accordingly, y remains to the left of z when γ_x (resp. $\delta_{\bar{x}}$) is applied to w . \square

Property 4.2. *Let $w = x_1x_2 \dots x_m \in B_{\mathbf{m}}$ be a signed word and i be a positive integer such that $m_i = m_{i+1} = 1$. Furthermore, let x be an integer such that $x \notin \{i, \bar{i}, i+1, \bar{i}+1\}$. Then the following conditions are equivalent:*

- (a) $i \in \text{Iligne } w$; (b) $i \in \text{Iligne } \mathbf{s}_1 w$; (c) $i \in \text{Iligne } \gamma_x w$; (d) $i \in \text{Iligne } \delta_{\bar{x}} w$.

Proof. (a) \Leftrightarrow (b) holds by definition 1.1, because \mathbf{s}_1 has no action on the rightmost letter belonging to $\{i, \bar{i}, i+1, \bar{i}+1\}$. For the other equivalences we can say the following. If the two letters of w that belong to $\{i, \bar{i}, i+1, \bar{i}+1\}$ are in I_x (resp. J_x , resp. K_x), Property 4.1 applies. Otherwise, if $i \in \text{Iligne } w$, then w is either of the form $\dots \bar{i}+1 \dots i \dots$ or $\dots i \dots \bar{i}+1 \dots$ and the order of those two letters is immaterial. \square

Theorem 1.2 holds for each one-letter signed word. Let $w = x_1x_2 \dots x_m \in B_{\mathbf{m}}$ be a signed word, x a letter and i a positive integer. Assume that $\text{Iligne } w = \text{Iligne } \Psi(w)$.

If $x = i$, then $i \in \text{Iligne } wx$ if and only if w contains no letter equal to $\pm i$ and exactly one letter equal to $\pm(i+1)$. As $\beta_x \Psi(w)$ is a rearrangement of w with possibly sign changes for some letters, the last statement is equivalent to saying that $\beta_x \Psi(w)$ has no letter equal to $\pm i$ and exactly one letter equal to $\pm(i+1)$. This is also equivalent to saying that $i \in \text{Iligne } \beta_x \Psi(w)x = \text{Iligne } \Psi(wx)$. In the same manner, we can show that

- if $x = \bar{i}+1$, then $i \in \text{Iligne } wx$ if and only if $i \in \text{Iligne } \Psi(wx)$;
- if $x = \bar{i}$, then $i \notin \text{Iligne } wx$ and $i \notin \text{Iligne } \Psi(wx)$;
- if $x = i+1$, then $i \notin \text{Iligne } wx$ and $i \notin \text{Iligne } \Psi(wx)$.

Now, let i be such that none of the integers $i, \bar{i}+1, \bar{i}, i+1$ is equal to x . There is nothing to prove if $m_i = m_{i+1} = 1$ does not hold, as i does not belong to any of the sets $\text{Iligne } w, \text{Iligne } \beta_x \Psi(w)$. Otherwise, the result follows from Property 4.2 because Ψ is a composition product of β_x, \mathbf{s}_1 and $\gamma_{\bar{x}}$. Finally, the equivalence $[0 \in \text{Iligne } \beta_x(w)x] \Leftrightarrow [0 \in \text{Iligne } wx]$ follows from Proposition 4.1 when $|x| > 1$ and the result is evident when $|x| = 1$. \square

The proof of Theorem 1.3 also follows from Property 4.1. By definition the lower records of w , other than x , belong to J_x . As the bijections γ_x and $\delta_{\bar{x}}$ do not modify the mutual order of the letters of w that belong to J_x , we have $\text{Lower } wx = \text{Lower } \beta_x(w)x$ when the chain (3.8) is used. When (3.9) is applied, so that $y_1 \notin J_x$, we also have $z_1 = \bar{y}_1 \notin J_x$. Thus, neither y_1 , nor z_1 can be lower records for each word ending with x . Again, $\text{Lower } wx = \text{Lower } \beta_x(w)x$. \square

5. CONCLUDING REMARKS

Since the works by MacMahon, much attention has been given to the study of statistics on the symmetric group or on classes of word rearrangements, in particular by the M.I.T. school ([25, 26, 27, 13, 6]). It was then natural to extend those studies to other classical Weyl groups, as was done by Reiner [20, 21, 22, 23, 24] for the signed permutation group. Today the work has been pursued by the Israeli and Roman schools [1, 2, 3, 4]. The contribution of Adin, Roichman [1] has been essential with their definition of the *flag major index* for signed permutations. In our forthcoming paper [9] we will derive new analytical expressions, in particular for several *multivariable* statistics involving “fma_j,” “finv” and the number of lower records.

REFERENCES

1. Ron M. Adin and Yuval Roichman, *The flag major index and group actions on polynomial rings*, Europ. J. Combin. **22** (2001), 431–446.
2. Ron M. Adin, Francesco Brenti and Yuval Roichman, *Descent Numbers and Major Indices for the Hyperoctahedral Group*, Adv. in Appl. Math. **27** (2001), 210–224.
3. Riccardo Biagioli, *Major and descent statistics for the even-signed permutation group*, Adv. in Appl. Math. **31** (2003), 163–179.
4. Riccardo Biagioli and Fabrizio Caselli, *Invariant algebras and major indices for classical Weyl groups*, Proc. London Math. Soc. **88** (2004), 603–631.
5. Anders Björner and Michelle L. Wachs, *Permutation Statistics and Linear Extensions of Posets*, J. Combin. Theory, Ser. A **58** (1991), 85–114.
6. N. Bourbaki, *Groupes et algèbres de Lie, Chapitres 4, 5 et 6*, Hermann, Paris, 1968.
7. Francesco Brenti, *q-Eulerian Polynomials Arising from Coxeter Groups*, Europ. J. Combinatorics **15** (1994), 417–441.
8. Dominique Foata, *On the Netto inversion number of a sequence*, Proc. Amer. Math. Soc. **19** (1968), 236–240.
9. Dominique Foata and Guo-Niu Han, *Signed words and permutations, III; the MacMahon Verfahren, (preprint)* (2005).
10. Dominique Foata and Marcel-Paul Schützenberger, *Major Index and Inversion number of Permutations*, Math. Nachr. **83** (1978), 143–159.
11. Herbert O. Foulkes, *Tangent and secant numbers and representations of symmetric groups*, Discrete Math. **15** (1976), 311–324.
12. Herbert O. Foulkes, *Eulerian numbers, Newcomb’s problem and representations of symmetric groups*, Discrete Math. **30** (1980), 3–49.
13. Ira Gessel, *Generating functions and enumeration of sequences*, Ph. D. thesis, Dept. Math., M.I.T., Cambridge, Mass., 111 p., 1977.
14. James E. Humphreys, *Reflection Groups and Coxeter Groups*, Cambridge Univ. Press, Cambridge, (Cambridge Studies in Adv. Math., **29**), 1990.
15. Donald E. Knuth, *The Art of Computer Programming*, vol.3, *Sorting and Searching*, Addison-Wesley, Reading, 1973.
16. M. Lothaire, *Combinatorics on Words*, Addison-Wesley, London (Encyclopedia of Math. and its Appl., **17**), 1983.
17. Percy Alexander MacMahon, *The indices of permutations and the derivation therefrom of functions of a single variable associated with the permutations of any assemblage of objects*, Amer. J. Math. **35** (1913), 314–321.
18. Percy Alexander MacMahon, *Combinatory Analysis*, vol. 1 and 2, Cambridge, Cambridge Univ. Press, 1915 (Reprinted by Chelsea, New York, 1995).
19. Percy Alexander MacMahon, *“Collected Papers, vol. 1” [G.E. Andrews, ed.]*, Cambridge, Mass., The M.I.T. Press, 1978.
20. V. Reiner, *Signed permutation statistics*, Europ. J. Combinatorics **14** (1993), 553–567.
21. V. Reiner, *Signed permutation statistics and cycle type*, Europ. J. Combinatorics **14** (1993), 569–579.

- 22. V. Reiner, *Upper binomial posets and signed permutation statistics*, Europ. J. Combinatorics **14** (1993), 581–588.
- 23. V. Reiner, *Descents and one-dimensional characters for classical Weyl groups*, Discrete Math. **140** (1995), 129–140.
- 24. V. Reiner, *The distribution of descents and length in a Coxeter group*, Electronic J. Combinatorics **2**, # **R25** (1995).
- 25. Richard P. Stanley, *Ordered structures and partitions*, Mem. Amer. Math. Soc. vol. 119, Amer. Math. Soc., Providence, RI, 1972.
- 26. Richard P. Stanley, *Binomial posets, Möbius inversion, and permutation enumeration*, J. Combinatorial Theory Ser. A **20** (1976), 336–356.
- 27. John Stembridge, *Eulerian numbers, tableaux, and the Betti numbers of a toric variety*, Discrete Math. **99** (1992), 307–320.

INSTITUT LOTHAIRE, 1 RUE MURNER, F-67000 STRASBOURG, FRANCE
E-mail address: foata@math.u-strasbg.fr

I.R.M.A. UMR 7501, UNIVERSITÉ LOUIS PASTEUR ET CNRS, 7 RUE RENÉ-DESCARTES,
F-67084 STRASBOURG, FRANCE
E-mail address: guoniu@math.u-strasbg.fr