

Lorentzian Kleinian groups

Charles Frances

4 December 2003

Abstract. In this article, we introduce some basic tools for the study of Lorentzian Kleinian groups. These groups are discrete subgroups of the Lorentzian Möbius group $O(2, n)$, acting properly discontinuously on some non empty open subset of Einstein's universe, the Lorentzian analogue of the conformal sphere.

Mathematics Subject Classification: 22E40, 30F40, 53A30, 53C50.

Key words: Lorentzian conformal geometry, Einstein's universe, conformal dynamics, Kleinian groups.

1 Introduction

To understand an hyperbolic manifold \mathbf{H}^{n+1}/Γ (\mathbf{H}^{n+1} denotes here the $(n+1)$ -hyperbolic space and $\Gamma \subset O(1, n+1)$ is a discrete group of hyperbolic isometries), a nice and powerful tool is the dynamical study of the conformal action of Γ on the sphere \mathbf{S}^n . This deep relationship between hyperbolic and conformally flat geometry has a counterpart in Lorentzian geometry, often quoted by physicists as *AdS/CFT correspondence*. Let us first recall what is the Lorentzian analogue of the pair $(\mathbf{H}^{n+1}, \mathbf{S}^n)$. The $(n+1)$ -dimensional Lorentzian model space of constant curvature -1 is called *anti-de Sitter space*, denoted \mathbf{AdS}_{n+1} (precisely, we are speaking here of the quotient of the simply connected model $\widetilde{\mathbf{AdS}}_{n+1}$ by the center of its isometry group, see [O'N], [Wo]). This space, like the hyperbolic space, has a conformal boundary. It is called Einstein's universe, denoted \mathbf{Ein}_n , and it can be defined, up to a two-sheeted covering, as the product $\mathbf{S}^1 \times \mathbf{S}^{n-1}$ endowed with the conformal class of the metric $-dt^2 \times g_{\mathbf{S}^{n-1}}$. From the conformal viewpoint, Einstein's universe has a lot of properties reminiscent of those of the sphere. In particular, the group $O(2, n)$ of isometries of \mathbf{AdS}_{n+1} turns out to be also the group of conformal transformations of \mathbf{Ein}_n . The understanding of an anti-de Sitter manifold $\mathbf{AdS}_{n+1}/\Gamma$ thanks to the conformal dynamics of Γ on \mathbf{Ein}_n is one of the motivations for studying *Lorentzian Kleinian groups*, that we define by analogy with the classical theory, as discrete subgroups

of $O(2, n)$ acting freely properly discontinuously on some non empty open subset of \mathbf{Ein}_n .

Since the works of Poincaré and Klein at the end of the nineteenth century, the classical theory of Kleinian groups has generated a great amount of works, and progressed very far (we refer the reader to [A], [Ka], [Ma], [MK] for an historical account and good expositions on the subject).

Other notions of Kleinian groups also appeared in other geometric contexts, such as complex hyperbolic and projective geometry (see for example [Go], [SV]).

To our knowledge, nothing systematic has been done for studying Lorentzian Kleinian groups, so that the aim of this article is to lay some basis for the theory. In particular, our first task is to build and study nontrivial examples of such groups.

The first part of the paper (sections 3 and 4) is devoted to what could be called *Lorentzian Möbius dynamics*, namely the dynamical study of divergent sequences of $O(2, n)$, acting on \mathbf{Ein}_n . This dynamics appears richer than that of classical Möbius transformations on the sphere. This is essentially due to the fact that $O(2, n)$ has rank two, and the different ways to reach infinity in $O(2, n)$ induce different dynamical patterns for the action on \mathbf{Ein}_n . These patterns, which are essentially three, are described in section 3, propositions 3, 4 and 5. Let us mention here two new phenomena (with respect to the Riemannian context) illustrating the dynamical complications we are confronted with. Firstly, the Lorentzian Möbius group $O(2, n)$ is not a convergence group for its action on \mathbf{Ein}_n (roughly speaking, a group G acting by homeomorphisms on a manifold X is a convergence group if any sequence (g_i) of G tending to infinity admits a subsequence with a "north-south" dynamics, i.e a dynamics with an attracting pole p^+ and a repelling one p^- . See for example [A] p.40 for a precise definition). Secondly, a discrete subgroup $\Gamma \subset O(2, n)$ does not always act properly on AdS_{n+1} .

In spite of these differences with respect to the classical theory, it is still possible to define the *limit set* of a discrete subgroup $\Gamma \subset O(2, n)$ (see section 4). This is a closed Γ -invariant subset $\Lambda_\Gamma \subset \mathbf{Ein}_n$, such that the action on the complement Ω_Γ is proper. It is moreover a union of lightlike geodesics, so that it defines naturally a Γ -invariant closed subset $\hat{\Lambda}_\Gamma$ of \mathbf{L}_n , the space of lightlike geodesics of \mathbf{Ein}_n (this space is described in section 2.5). Unfortunately, the nice properties of the limit set in the classical case of groups of conformal transformations of the sphere, are generally no longer satisfied in our situation. For example, the limit set that we define is not, in general, a minimal set for the action of Γ on \mathbf{Ein}_n (although $\hat{\Lambda}_\Gamma$ is sometimes minimal for the action of Γ on \mathbf{L}_n , see theorem 1 below). Groups $\Gamma \subset O(2, n)$ acting properly on AdS_{n+1} are those whose behaviour is closest to that of classical Kleinian groups. They will be called *groups of the first type*. For them, we get nice properties for the limit set:

Theorem 1. *Let Γ be a Kleinian group of the first type and Λ_Γ its limit set.*

- (i) *The action of Γ is proper on $\Omega_\Gamma \cup \mathbf{AdS}_{n+1} \subset \mathbf{Ein}_{n+1}$.*
- (ii) *Ω_Γ is the unique maximal element among the open sets $\Omega \subset \mathbf{Ein}_n$ such that Γ acts properly on $\Omega \cup \mathbf{AdS}_{n+1}$.*
- (iii) *If moreover Γ is Zariski dense in $O(2, n)$, then Ω_Γ is the unique maximal open subset of \mathbf{Ein}_n on which Γ acts properly, and $\hat{\Lambda}_\Gamma$ is a minimal set for the action of Γ on \mathbf{L}_n .*

In section 5, we give several examples of families of Lorentzian Kleinian groups. These basic examples being constructed, it is natural to try to combine two of them to get other more complicated examples. It is the aim of section 6, where we prove the following result (an analogue of the celebrated Klein's combination theorem):

Theorem 2. *Let Γ_1 and Γ_2 be two cocompact Lorentzian Kleinian groups, with fundamental domains D_1 and D_2 . Suppose that both D_1 and D_2 contain a lightlike geodesic. Then one can construct from Γ_1 and Γ_2 another cocompact Kleinian group, isomorphic to the free product $\Gamma_1 * \Gamma_2$.*

By *cocompact Kleinian group*, we mean a group acting properly on some open subset of \mathbf{Ein}_n , with compact quotient.

We then use theorem 2 in section 7 to construct *Lorentzian Schottky groups*. The study of such groups can be carried out quite completely. The limit set Λ_Γ and the topology of the conformally flat Lorentz manifold obtained as the quotient Ω_Γ/Γ of the domain of properness, are made explicit in this case, and we get:

Theorem 3. *Let $\Gamma = \langle s_1, \dots, s_g \rangle$ ($g \geq 2$) be a Lorentzian Schottky group.*

- (i) *The group Γ is of the first type.*
- (ii) *The limit set Λ_Γ is a lamination by lightlike geodesics. Topologically, it is a product of \mathbf{RP}^1 with a Cantor set.*
- (iii) *The action of Γ is minimal on the set of lightlike geodesics of Λ_Γ .*
- (iv) *The quotient manifold Ω_Γ/Γ is diffeomorphic to the product $\mathbf{S}^1 \times (\mathbf{S}^1 \times \mathbf{S}^{n-1})^{(g-1)\sharp}$, where $(\mathbf{S}^1 \times \mathbf{S}^{n-1})^{(g-1)\sharp}$ is the connected sum of $(g-1)$ copies of $\mathbf{S}^1 \times \mathbf{S}^{n-1}$.*

2 Geometry of Einstein's universe

A detailed description of the geometry of Einstein's universe can be found in [Fr1], [Fr2] and [CK]. Also, for the reader who are not very familiar with Lorentzian space-times of constant curvatures, good expositions can be found in [Wo] chapter 11, and [O'N] chapter 8. In this section, we briefly recall (without any proof) the main properties which will be useful in this article.

2.1 Projective model for Einstein's universe

Let $\mathbf{R}^{2,n}$ be the space \mathbf{R}^{n+2} , endowed with the quadratic form $q^{2,n}(x) = -x_1^2 - x_2^2 + x_3^2 + \dots + x_{n+2}^2$. The isotropic cone of $q^{2,n}$ is the subset of $\mathbf{R}^{2,n}$ on which $q^{2,n}$ vanishes. We call $C^{2,n}$ this isotropic cone, with the origin removed. Throughout this article, we will denote by π the projection from $\mathbf{R}^{2,n}$ minus the origin, on $\mathbf{R}P^{n+1}$. The set $\pi(C^{2,n})$ is a smooth hypersurface Σ of $\mathbf{R}P^{n+1}$. This hypersurface turns out to be endowed with a natural Lorentzian conformal structure. Indeed, for any $x \in C^{2,n}$, the restriction of $q^{2,n}$ to the tangent space $T_x C^{2,n}$, that we call $\hat{q}_x^{2,n}$, is degenerate. Its kernel is just the kernel of the tangent map $d_x \pi$. Thus, pushing $\hat{q}_x^{2,n}$ by $d_x \pi$, we get a well defined Lorentzian metric on $T_{\pi(x)} \Sigma$. If $\pi(x) = \pi(y)$ the two Lorentzian metrics on $T_{\pi(x)} \Sigma$ obtained by pushing $\hat{q}_x^{2,n}$ and $\hat{q}_y^{2,n}$ are in the same conformal class. Thus, the form $q^{2,n}$ determines a well defined conformal class of Lorentzian metrics on Σ . One calls *Einstein's universe* the hypersurface Σ , together with this canonical conformal structure.

The intersection of $C^{2,n}$ with the euclidean sphere defined by $x_1^2 + x_2^2 + \dots + x_{n+2}^2 = 1$ is a smooth hypersurface $\hat{\Sigma} \subset \mathbf{R}^{2,n}$. One can check that $q^{2,n}$ has lorentzian signature when restricted to $\hat{\Sigma}$, and in fact, $(\hat{\Sigma}, q_{|\hat{\Sigma}}^{2,n})$ is isometric to the product $(\mathbf{S}^1 \times \mathbf{S}^{n-1}, -dt^2 + g_{\mathbf{S}^{n-1}})$. Now Einstein's universe is conformally equivalent to the quotient of $(\mathbf{S}^1 \times \mathbf{S}^{n-1}, -dt^2 + g_{\mathbf{S}^{n-1}})$ by an involution (induced by the map $x \mapsto -x$ of $\mathbf{R}^{2,n}$).

2.2 Conformal group

In the previous projective model for Einstein's universe, the subgroup $O(2, n) \subset GL_{n+2}(\mathbf{R})$ preserving $q^{2,n}$, acts conformally on \mathbf{Ein}_n . In fact, the conformal group $\text{Conf}(\mathbf{Ein}_n)$ of \mathbf{Ein}_n is exactly $PO(2, n)$. Let us now recall the following result, which is an extension to Einstein's universe of a classical theorem of Liouville in Euclidean conformal geometry (see for example [CK] [Fr3]):

Theorem 4. *Any conformal transformation between two open sets of \mathbf{Ein}_n is the restriction of a unique element of $PO(2, n)$.*

2.3 Lightlike geodesics and lightcones

It is a remarkable fact of pseudo-Riemannian geometry that all the metrics of a given conformal class have the same lightlike geodesics (as sets but not as parametrized curves). In the case of Einstein's universe, the lightlike geodesics are the projections on \mathbf{Ein}_n of 2-planes $P \subset \mathbf{R}^{2,n}$ such that $q_P^{2,n} = 0$. So, lightlike geodesics of \mathbf{Ein}_n are copies of $\mathbf{R}P^1$.

Given a point p in \mathbf{Ein}_n , the *lightcone with vertex p* , denoted by $C(p)$, is the set of lightlike geodesics containing p . In the projective model, if

$p = \pi(u)$, with u some isotropic vector of $\mathbf{R}^{2,n}$, then $C(p)$ is just $\pi(P \cap C^{2,n})$, where P is the degenerate hyperplane $P = u^\perp$ (the orthogonal is taken for the form $q^{2,n}$). The lightcones are not smooth submanifolds of \mathbf{Ein}_n . The only singular point of $C(p)$ is p , and $C(p) \setminus \{p\}$ is topologically $\mathbf{R} \times \mathbf{S}^{n-2}$.

2.4 Homogeneous open subsets

We will deal in this paper with several interesting open subsets of \mathbf{Ein}_n , all obtained by removing to \mathbf{Ein}_n the projectivization of peculiar linear subspaces of $\mathbf{R}^{2,n}$. We will be very brief here, and we refer to [Wo] for a more detailed study (especially concerning de Sitter and anti- de Sitter spaces).

- *Minkowski components.*

Given a point $p \in \mathbf{Ein}_n$, the complement of $C(p)$ in \mathbf{Ein}_n is an homogeneous open subset of \mathbf{Ein}_n , which is conformally equivalent to Minkowski space $\mathbf{R}^{1,n-1}$. We say that this is the Minkowski component associated to p . In fact, we have an explicit formula for the stereographic projection identifying $\mathbf{Ein}_n \setminus C(p)$ and $\mathbf{R}^{1,n-1}$ (see [CK] [Fr1]).

- *De Sitter and anti-de Sitter components.*

Just as Minkowski space arises by removing to \mathbf{Ein}_n the projectivization of a lightlike hyperplane, one also gets interesting open subsets by removing the projectivization of other (i.e nondegenerate) hyperplanes.

If P is some hyperplane of $\mathbf{R}^{2,n}$, with Lorentzian signature, then $\pi(P \cap C^{2,n})$ is a Riemannian sphere S of codimension one. The canonical conformal structure of \mathbf{Ein}_n induces on this sphere the canonical Riemannian conformal structure. The stabilizer of S in $O(2, n)$ is a group G isomorphic to $O(1, n)$. The complement of S in \mathbf{Ein}_n is an homogeneous open subset of \mathbf{Ein}_n , conformally equivalent to the de Sitter space \mathbf{dS}_n . So, \mathbf{S}^{n-1} , with its canonical conformal structure, appears as the conformal boundary of \mathbf{dS}_n .

If P is some hyperplane of $\mathbf{R}^{2,n}$, with signature $(2, n - 1)$, then the projection $\pi(P \cap C^{2,n})$ is a codimension one Einstein's universe E . The stabilizer of E in $O(2, n)$ is a subgroup isomorphic to $O(2, n - 1)$. The complement of E is an homogeneous open subset of \mathbf{Ein}_n which is conformally equivalent to the anti-de Sitter space \mathbf{AdS}_n . In this way, we see \mathbf{Ein}_{n-1} as the conformal boundary of \mathbf{AdS}_n .

- *Complement of a lightlike geodesic.*

What do we get if we remove from \mathbf{Ein}_n the projectivization of a maximal isotropic subspace of $\mathbf{R}^{2,n}$? Such subspaces are 2-planes, so that the resulting open set is the complement Ω_Δ of a lightlike geodesic $\Delta \subset \mathbf{Ein}_n$. Open sets like Ω_Δ admit a natural foliation by degenerate hypersurfaces, and this foliation \mathcal{H}_Δ is preserved by the whole conformal group of Ω_Δ . This foliation can be described as follows: given a point $p \in \Delta$, we consider the lightcone

The interest of this definition for the study of actions of discrete groups can be illustrated by the following: let Γ be a discrete group of $\text{Homeo}(X)$ acting on some open subset $\Omega \subset X$. Then, one proves easily:

Proposition 1. *The group Γ acts properly on Ω iff no two points of Ω are dynamically related.*

Assuming that the action of Γ on Ω is proper, we also have:

Proposition 2. *If the action of Γ on Ω has compact quotient, then every $x \in \partial\Omega$ must be dynamically related to some point y of Ω (depending on x).*

Now, let (g_k) be a divergent sequence of $O(2, n)$. We define $\lambda_k = \lambda(g_k)$, $\mu_k = \mu(g_k)$ and $\delta_k = \lambda_k - \mu_k$. We say that the sequence (g_k) **tends simply to infinity** when:

- a) the three sequences (λ_k) , (μ_k) and (δ_k) converge respectively to some λ_∞ , μ_∞ and δ_∞ in $\overline{\mathbf{R}}$.
- b) compact factors in the Cartan decomposition of (g_k) both admit a limit in K .

Of course, every sequence tending to infinity admits some subsequence tending simply to infinity, so that we will restrict our study to these last ones. The sequences tending simply to infinity split into three categories:

(i) Sequences with balanced distortions

This name denotes the sequences (g_k) for which $\lambda_\infty = \mu_\infty = +\infty$ and δ_∞ is finite.

(ii) Sequences with bounded distortion

This denotes the sequences (g_k) for which $\mu_\infty \neq +\infty$.

(iii) Sequences with mixed distortions

This denotes the sequences (g_k) for which $\lambda_\infty = \mu_\infty = \delta_\infty = +\infty$.

To each type corresponds, as we will see soon, distinct dynamical behaviours.

Notations 1. *In the following, we will use notations such as $C(p)$, \mathcal{H}_Δ ... We invite the reader to look at section 2, where these notation were introduced.*

For any set E in $\mathbf{R}^{2,n}$, we use the notation $\tilde{\pi}(E)$ for $\pi(E \cap C^{2,n})$. If y and ϵ are two real numbers, we write $I_\epsilon(y)$ for the closed interval $[y - \epsilon, y + \epsilon]$. For every $x = (x_1, x_2, \dots, x_{n+2})$ in $\mathbf{R}^{2,n}$, we define the ϵ -box centered at x as:

$$B_\epsilon(x) = I_\epsilon(x_1) \times I_\epsilon(x_2) \times \dots \times I_\epsilon(x_{n+2})$$

For a sequence (g_k) of $O(2, n)$ tending simply to infinity, we call $B_\epsilon^\infty(x)$ the compact set obtained as the limit (for the Hausdorff topology) of the sequence of compact sets $g_k \circ \tilde{\pi}(B_\epsilon(x))$ (this limit will always exist in the examples we will deal with).

Lastly, we will often denote in the same way an element of $O(2, n)$ and the conformal transformation of \mathbf{Ein}_n that it induces.

3.2.1 Dynamics with balanced distortions

Proposition 3. *Let (g_k) be a sequence of $O(2, n)$ with balanced distortions. Then we can associate to (g_k) two lightlike geodesics Δ^+ and Δ^- called **attracting and repelling circles** of (g_k) , and two submersions $\pi_+ : \mathbf{Ein}_n \setminus \Delta^- \rightarrow \Delta^+$ (resp. $\pi_- : \mathbf{Ein}_n \setminus \Delta^+ \rightarrow \Delta^-$), whose fibers are the leaves of \mathcal{H}_{Δ^-} (resp. \mathcal{H}_{Δ^+}), such that :*

For every compact subset K of $\mathbf{Ein}_n \setminus \Delta^-$ (resp. $\mathbf{Ein}_n \setminus \Delta^+$), $D_{(g_k)}(K) = \pi_+(K)$ (resp. $D_{(g_k^{-1})}(K) = \pi_-(K)$).

Remark 1. *Before beginning the proof, let us remark that if (g_k) has balanced distortions (resp. bounded distortion, resp. mixed distortions), it will be so for any compact perturbation of (g_k) , i.e any sequence $(l_k^{(1)} g_k l_k^{(2)})$ for $(l_k^{(1)})$ and $(l_k^{(2)})$ two converging sequences of $O(2, n)$. In the same way, the conclusions of the above proposition are not modified by a compact perturbation, even if of course, π_{\pm} and Δ^{\pm} are. So in the following (and also for sections 3.2.2 3.2.3), we will restrict the proofs to the case where (g_k) is a sequence of A^+ .*

Proof : We restrict the proof to the case $\lambda_k = \mu_k$, so that $\delta_{\infty} = 0$.

We begin by defining Δ^{\pm} and π^{\pm} . Let us call P^+ (resp. P^-) the 2- plane spanned by e_1 and e_2 (resp. e_{n+1} and e_{n+2}), and Δ^+ (resp. Δ^-) the projection on \mathbf{Ein}_n of these 2- planes. The space $\mathbf{R}^{2, n}$ splits as a direct sum $P_+ \oplus P_0 \oplus P_-$, where P_0 is the span of e_3, \dots, e_n . This splitting defines a projection $\tilde{\pi}_+$ (resp. $\tilde{\pi}_-$) from $\mathbf{R}^{2, n}$ to the plane P^+ (resp. P^-). The image $\tilde{\pi}_+(x)$ is nonzero as soon as x is an isotropic vector of $q^{2, n}$ which is not in P^- . Thus $\tilde{\pi}_+$ induces a projection π_+ of $\mathbf{Ein}_n \setminus \Delta^-$ on Δ^+ whose fibers are the projections on \mathbf{Ein}_n of the fibers of $\tilde{\pi}_+$. These are degenerate hyperplanes of $\mathbf{R}^{2, n}$, defined as $q^{2, n}$ -orthogonals of vectors of P^- . So, the fibers of π_+ are the intersections of $\mathbf{Ein}_n \setminus \Delta^-$ with the lightcones with vertex on Δ^- , i.e the leaves of \mathcal{H}_{Δ^-} .

Now, let us choose x such that $\pi(x) \notin \Delta^-$. Since $g_k \circ \tilde{\pi}(B_{\epsilon}(x)) = \tilde{\pi}(I_{e^{\lambda_k \epsilon}}(e^{\lambda_k x_1}) \times I_{e^{\mu_k \epsilon}}(e^{\mu_k x_2}) \times I_{\epsilon}(x_3) \times \dots \times I_{\epsilon}(x_n) \times I_{e^{-\mu_k \epsilon}}(e^{-\mu_k x_{n+1}}) \times I_{e^{-\lambda_k \epsilon}}(e^{-\lambda_k x_{n+2}}))$, we get that for ϵ sufficiently small, we have $B_{\epsilon}^{\infty}(x) = \tilde{\pi}(I_{\epsilon}(x_1) \times I_{\epsilon}(x_2) \times \{0\} \times \dots \times \{0\})$

We thus have $B_{\epsilon}^{\infty}(x) \subset \Delta^+$. Since ϵ is arbitrarily close to 0, for any sequence (x_k) such that $\pi(x_k)$ tends to $\pi(x)$, we have $\lim_{k \rightarrow \infty} g_k \circ \pi(x_k) = \pi(x_1, x_2, 0, \dots, 0)$. This concludes the proof. \square

3.2.2 Dynamics with bounded distortion

Proposition 4. *Let (g_k) be a sequence of $O(2, n)$ with bounded distortions. Then we can associate to (g_k) two points p^+ and p^- of \mathbf{Ein}_n called **attracting and repelling poles** of (g_k) , and a diffeomorphism \hat{g}_{∞} from the space*

3.2.3 Mixed dynamics

Proposition 5. *Let (g_k) be a sequence of $O(2, n)$ with mixed distortions. Then we can associate to (g_k) two points p^+ and p^- called **attracting** and **repelling poles** of the sequence, as well as two lightlike geodesics Δ^+ et Δ^- (called **attracting** and **repelling circles**), with the inclusions $p^+ \in \Delta^+ \subset C^+ = C(p^+)$ and $p^- \in \Delta^- \subset C^- = C(p^-)$, such that the following properties hold:*

- (i) *For every compact subset K inside $\mathbf{Ein}_n \setminus C^-$, the set $D_{(g_k)}(K)$ is $\{p^+\}$.*
- (ii) *If x is a point of C^- , not on Δ^- , then $D_{(g_k)}(x)$ is the lightlike geodesic Δ^+ .*
- (iii) *If x is a point of Δ^- , distinct from p^- , then $D_{(g_k)}(x)$ is the attracting cone C^+ .*
- (iv) *The set $D_{(g_k)}p^-$ is the whole \mathbf{Ein}_n .*

The cones C^+ and C^- are called **attracting** and **repelling cones** of the sequence (g_k) .

Proof: Once again, we suppose that (g_k) is in A^+ .

We define $p^+ = \pi(e_1)$, $p^- = \pi(e_{n+2})$, $C^+ = \tilde{\pi}((e_1)^\perp)$, $C^- = \tilde{\pi}((e_{n+2})^\perp)$. The circle Δ^+ (resp. Δ^-) is the projection of the 2-plane spanned by e_1 and e_2 (resp. e_{n+1} and e_{n+2}). We don't prove points (i) and (iv), the proof being exactly the same as for proposition 4.

If $\pi(x) \in C^-$ but $\pi(x) \notin \Delta^-$, it means that $x_1 = 0$ but $x_2 \neq 0$. In this case, we get for $B_\epsilon^\infty(x) = \tilde{\pi}(\mathbf{R} \times I_\epsilon(x_2) \times \{0\} \times \dots \times \{0\})$, that is to say Δ^+ .

The intersection of all the $B_\epsilon^\infty(x)$ is $\tilde{\pi}(\mathbf{R} \times \{x_2\} \times \{0\} \times \dots \times \{0\})$, i.e the lightlike geodesic Δ^+ . The fact that $D_{(g_k)}(\pi(x)) = \Delta^+$ is proved exactly as in proposition 4.

When $\pi(x) \in \Delta^-$, only x_{n+1} and x_{n+2} don't vanish and by the assumption $\pi(x) \neq p^-$, we get $x_{n+1} \neq 0$. Hence, we have that $B_\epsilon^\infty(x)$ is $\tilde{\pi}(\mathbf{R} \times \dots \times \mathbf{R} \times I_\epsilon(x_{n+1}) \times \{0\})$, that is to say C^+ .

As previously, we get $D_{(g_k)}(\pi(x)) = C^+$.

□

Remark 3. *Notice that different configurations for the dynamical elements described above can occur. For example, attracting and repelling circles of a dynamics with balanced or mixed distortions can intersect, or even be the same. In fact, all the possible configurations can occur.*

4 About the limit set of a Lorentzian Kleinian group

4.1 Definition of the limit set

Given a Kleinian group Γ on a manifold X , it is quite natural to ask if there is in some sense a “canonical” open set $\Omega \subset X$ on which Γ acts properly. For example, any Kleinian group Γ on the sphere \mathbf{S}^n admits a limit set Λ_Γ and the open set $\Omega_\Gamma = \mathbf{S}^n \setminus \Lambda_\Gamma$ is distinguished, since it is the only maximal open subset on which Γ acts properly. The nice properties of the limit set of a Kleinian group on \mathbf{S}^n rest essentially on the fact that the Möbius group $O(1, n+1)$ is a convergence group on \mathbf{S}^n . We just saw in the previous section that $O(2, n)$ is quite far from being a convergence group on \mathbf{Ein}_n but nevertheless, we would like to define a limit set Λ_Γ associated to a given discrete group $\Gamma \subset O(2, n)$. We require that such a limit set have at least the two following properties:

- (i) Λ_Γ is a Γ -invariant closed subset of \mathbf{Ein}_n .
- (ii) The action of Γ on $\Omega_\Gamma = \mathbf{Ein}_n \setminus \Lambda_\Gamma$ is properly discontinuous.

Definition 2. *Given Γ discrete in $O(2, n)$, we define \mathcal{S}_Γ (resp. \mathcal{T}_Γ) the set of sequences (γ_k) of Γ , tending simply to infinity, with mixed or balanced distortions (resp. with bounded distortion). If (γ_k) is a sequence of \mathcal{S}_Γ (resp. \mathcal{T}_Γ), we call $\Delta^+(\gamma_k)$ and $\Delta^-(\gamma_k)$ (resp. $C^+(\gamma_k)$ and $C^-(\gamma_k)$) its attracting and repelling circles (resp. attracting and repelling cones).*

Definition 3. *We define the limit set of a discrete $\Gamma \subset O(2, n)$ as:*

$$\Lambda_\Gamma = \Lambda_\Gamma^{(1)} \cup \Lambda_\Gamma^{(2)} \text{ where}$$

$$\Lambda_\Gamma^{(1)} = \overline{\bigcup_{(\gamma_k) \in \mathcal{S}_\Gamma} \Delta^+(\gamma_k) \cup \Delta^-(\gamma_k)}$$

and

$$\Lambda_\Gamma^{(2)} = \overline{\bigcup_{(\gamma_k) \in \mathcal{T}_\Gamma} C^+(\gamma_k) \cup C^-(\gamma_k)}$$

Notation 1. *The complement of Λ_Γ in \mathbf{Ein}_n is denoted by Ω_Γ .*

It is clear that Λ_Γ is closed and Γ -invariant. Let us remark that Λ_Γ is a union of lightlike geodesics, so that it also defines a closed Γ -invariant subset $\hat{\Lambda}_\Gamma \subset \mathbf{L}_n$.

From the dynamical properties stated in the previous section, one checks easily that no pair of points in Ω_Γ can be dynamically related, so that the action of Γ on Ω_Γ is proper.

4.2 Lorentzian Kleinian groups of the first and the second type

Until now, we didn't focus on a fundamental difference between the action of $O(1, n+1)$ on \mathbf{S}^n and that of $O(2, n)$ on \mathbf{Ein}_n . Although any discrete group $\Gamma \subset O(1, n+1)$ automatically acts properly on \mathbf{H}^{n+1} , it is not true in general that a discrete $\Gamma \subset O(2, n)$ does so on \mathbf{AdS}_{n+1} . This motivates the following distinction between subgroups of $O(2, n)$:

Definition 4. A discrete group Γ of $O(2, n)$ is of the **first type** if it acts properly on \mathbf{AdS}_{n+1} . If not, it is said to be of the **second type**.

Notice that this terminology has no connection with the denomination *first kind* and *second kind* for the standard Kleinian groups on the sphere.

The previous dichotomy has a nice translation into dynamical terms thanks to the:

Proposition 6. A Kleinian group Γ of $O(2, n)$ is of the first type if and only if it doesn't admit any sequence (γ_k) with bounded distortion.

Proof: We endow $\mathbf{R}^{2, n+1}$ with the quadratic form $q^{2, n+1}(x) = -2x_1x_{n+2} + 2x_2x_{n+1} + x_3^2 + \dots + x_n^2 + x_{n+3}^2$ and call e_1, \dots, e_{n+3} the canonical basis. The subgroup of $O(2, n+1)$ leaving invariant the subspace spanned by the first $n+2$ basis vectors can be canonically identified with $O(2, n)$. This identification defines an embedding j from $O(2, n)$ into $O(2, n+1)$. The action of $j(O(2, n))$ on \mathbf{Ein}_{n+1} leaves invariant a codimension 1 Einstein universe that we call \mathbf{Ein}_n . As we saw in the introduction, the complement of \mathbf{Ein}_n in \mathbf{Ein}_{n+1} is conformally equivalent to the anti-de Sitter space \mathbf{AdS}_{n+1} .

Let us consider some g in $O(2, n)$. In the basis e_1, \dots, e_{n+3} , $j(g) = \begin{pmatrix} g & \\ & 1 \end{pmatrix}$, so that when we perform the Cartan decomposition of $j(g)$, we find the same distortions as for g .

Suppose now that Γ admits some sequence (γ_k) with bounded distortion. By the remark above, $j(\gamma_k)$ has also bounded distortion as a sequence of $O(2, n+1)$. We call C^+ and C^- its attracting and repelling cones in \mathbf{Ein}_{n+1} . By proposition 4, $D_{(g_k)}(C^- \cap \mathbf{AdS}_{n+1}) = C^+ \cap \mathbf{AdS}_{n+1}$. Therefore, we can find two points of \mathbf{AdS}_{n+1} which are dynamically related, so that the action of (γ_k) on \mathbf{AdS}_{n+1} can not be proper (proposition 1).

Conversely, let us consider some sequence (γ_k) tending simply to infinity and with balanced or mixed distortions. Then the sequence $j(\gamma_k)$ has the same properties. Let us call Δ^+ and Δ^- the attracting and repelling circles of this latter sequence. Looking at the matrix expressions, it is clear that $\Delta^+ \subset \mathbf{Ein}_n$ and $\Delta^- \subset \mathbf{Ein}_n$. By propositions 3 and 5, $D_{(g_k)}(x) \subset \mathbf{Ein}_n$ for any point $x \in \mathbf{AdS}_{n+1}$. So, if we assume that Γ has no sequence with bounded distortion, we get $D_\Gamma(x) \subset \mathbf{Ein}_n$ for any point $x \in \mathbf{AdS}_{n+1}$. Using proposition 1, we get that Γ acts properly on \mathbf{AdS}_{n+1} . \square

4.3 Limit set of a group of the first type: proof of Theorem 1

Since Γ is of the first type, Λ_Γ is also the limit set of Γ , seen as a subgroup of $O(2, n+1)$ acting on \mathbf{Ein}_{n+1} . The complement of this limit set in \mathbf{Ein}_{n+1} is precisely $\Omega_\Gamma \cup \mathbf{AdS}_{n+1}$, so that point (i) of the theorem is clear.

To prove point (ii), let us suppose that Γ acts properly on some $\Omega \cup \mathbf{AdS}_{n+1}$ with Ω not included in Ω_Γ . Then there is a sequence (γ_k) of Γ (with balanced or mixed distortions) such that $\Delta^-(\gamma_k)$ meets Ω .

Lemma 1. *Let Γ be a discrete group of $O(2, n)$ acting properly on some open set Ω . Then for any sequence (γ_k) of Γ with balanced distortions, neither $\Delta^+(\gamma_k)$ nor $\Delta^-(\gamma_k)$ meets Ω .*

Proof. Suppose on the contrary that for some (γ_k) with balanced distortions, we have $\Delta^+(\gamma_k) \cap \Omega \neq \emptyset$. From proposition 3, we infer that the set $D_{(\gamma_k)}(\Delta^+(\gamma_k) \cap \Omega)$ contains a lightlike geodesic Δ in its interior. So, there is a tubular neighbourhood U of Δ contained in $Ext(\Omega)$ ($Ext(\Omega)$ denotes the complement of Ω in \mathbf{Ein}_n). But we also infer from proposition 3 that for any Δ not meeting $\Delta^-(\gamma_k)$, we have $\lim_{k \rightarrow +\infty} \gamma_k(\Delta) = \Delta^+(\gamma_k)$. As a consequence, any lightlike geodesic of $Ext(\Omega)$ has to cut $\Delta^-(\gamma_k)$. Since all the lightlike geodesics included in U can't all meet $\Delta^-(\gamma_k)$, we get a contradiction. \square

The lemma above tells us that the sequence (γ_k) has mixed distortions. For any point $x \in \Delta^-(\gamma_k) \cap \Omega$, we have $D_{(\gamma_k)}(x) = C^+(\gamma_k)$. Since $C^+(\gamma_k)$ meets \mathbf{AdS}_{n+1} , we get pairs of points in $\Omega \cup \mathbf{AdS}_{n+1}$ which are dynamically related, and the action can't be proper, by proposition 1.

Remark 4. *For Γ Kleinian of the first type, the manifold Ω_Γ/Γ appears as the conformal boundary of the complete anti-de Sitter manifold $\mathbf{AdS}_{n+1}/\Gamma$ (see [Fr4] for more details on this point).*

To prove point (iii), we begin by showing that $\hat{\Lambda}_\Gamma \subset \mathbf{L}_n$ is a minimal set. This is in fact a particular case of a general result of Benoist ([B]), but we give a simple proof.

Let $\hat{\Lambda}$ be a closed Γ -invariant subset of \mathbf{L}_n . Any sequence (γ_k) tending simply to infinity in Γ has either mixed or balanced distortions. As a simple consequence of propositions 3 and 5, we get that if Δ is a lightlike geodesic of \mathbf{Ein}_n which does not meet $\Delta^-(\gamma_k)$, then $\lim_{k \rightarrow +\infty} \gamma_k(\Delta) = \Delta^+(\gamma_k)$. So, if for any sequence (γ_k) as above, no geodesic of $\hat{\Lambda}$ meets $\Delta^-(\gamma_k)$, we have $\Lambda_\Gamma \subset \Lambda$, and we are done.

On the contrary, if for some (γ_k) , all the geodesics of $\hat{\Lambda}$ meet $\Delta^-(\gamma_k)$, we claim that Γ can't be Zariski dense. Indeed, by Zariski density, Γ can't leave $\Delta^-(\gamma_k)$ invariant. So, let us choose $\gamma \in \Gamma$ such that $\gamma(\Delta^-(\gamma_k)) \neq \Delta^-(\gamma_k)$. If $\gamma(\Delta^-(\gamma_k))$ and $\Delta^-(\gamma_k)$ are disjoint, the set of lightlike geodesics meeting

both $g(\Delta^-(\gamma_k))$ and $\Delta^-(\gamma_k)$ is contained in a two dimensional Einstein's universe, which have to be fixed by Γ : a contradiction with the Zariski density of Γ .

If $g(\Delta^-(\gamma_k))$ and $\Delta^-(\gamma_k)$ meet in one point p , then any lightlike geodesic meeting both $g(\Delta^-(\gamma_k))$ and $\Delta^-(\gamma_k)$ has to contain p . Indeed, due to the fact that the quadratic form $q^{2,n}$ can't have some 3 dimensional isotropic subspace, there is no nontrivial triangle of \mathbf{Ein}_n , whose edges are pieces of lightlike geodesics. We infer that Γ has to fix the lightcone $C(p)$ and we get once again a contradiction.

We can now prove that Ω_Γ is the maximal open set on which the action of Γ is proper. Suppose that Γ acts properly on Ω which is not included in Ω_Γ . We call Λ the complement of Ω in \mathbf{Ein}_n . Since $\Lambda_\Gamma \not\subset \Lambda$, there is a sequence (γ_k) tending simply to infinity in Γ with $\Delta^+(\gamma_k) \cap \Omega \neq \emptyset$.

Lemma 2. *If an infinite Kleinian group $\Gamma \subset O(2, n)$ acts properly on some open subset Ω , then the complement Λ of Ω in \mathbf{Ein}_n contains a lightlike geodesic.*

Proof. Let us pick a sequence (γ_k) tending simply to infinity in Γ . Suppose first that (γ_k) has mixed dynamics. Suppose that $\Delta^-(\gamma_k)$ meets Ω at a point x (if $\Delta^-(\gamma_k) \cap \Omega = \emptyset$, we are done). By properness, $D_{(g_k)}(x) \cap \Omega = \emptyset$. But $D_{(g_k)}(x) = C^+(g_k)$, which contains infinitely many lightlike geodesics, and the conclusion holds.

Also, if (g_k) has balanced (resp. bounded) distortions, the dynamic set $D_{(g_k)}x$ of $x \in \Delta^-(\gamma_k)$ (resp. $x \in C^-(\gamma_k)$) contains infinitely many lightlike geodesics. The proof works thus in the same way. □

Now, let us look at the lightlike geodesics of Λ . Since by Zariski density, Γ can't fix a finite family of lightlike geodesics, there are infinitely many lightlike geodesics in Λ . But all these geodesics have to meet $\Delta^-(\gamma_k)$, because if some Δ does not, $\lim_{k \rightarrow +\infty} \gamma_k(\Delta) = \Delta^+(\gamma_k)$. A contradiction with $\Delta^+(\gamma_k) \cap \Omega \neq \emptyset$. Now, we conclude as for proving the minimality property of $\hat{\Lambda}_\Gamma$: all the lightlike geodesics of Λ are in the same Γ -invariant Einstein torus, or the same Γ -invariant lightcone and we get a contradiction with the Zariski density of Γ .

5 Some examples of Lorentzian Kleinian groups

5.1 Examples arising from structures with constant curvature

In Lorentzian geometry, a completeness result ensures that any compact Lorentzian manifold with constant sectional curvature is obtained as a quo-

tient $\mathbf{R}^{1,n-1}/\Gamma$ or $\widetilde{\mathbf{AdS}}_n/\Gamma$, where Γ is a discrete group of Lorentzian isometries. This deep theorem was first proved for the case of curvature zero by Carrière in [Ca], and generalized by Klingler in [Kl] (note that compact Lorentzian manifolds can not have curvature $+1$). Another result, known as *finiteness of level* (see [KR], [Ze]), ensures that any compact quotient $\widetilde{\mathbf{AdS}}_n/\tilde{\Gamma}$ (where $\tilde{\Gamma}$ is a discrete group of isometries) is in fact, up to finite cover, a quotient \mathbf{AdS}_n/Γ . Since we saw in section 2 that $\mathbf{R}^{1,n-1}$ and \mathbf{AdS}_n both embed conformally into \mathbf{Ein}_n , and thanks to theorem 4, we get that any compact Lorentzian structure with constant curvature is (up to finite cover) uniformized by a Lorentzian Kleinian groups. Moreover, in this case, the structure of the groups involved is fairly well understood, thanks to [CaD], [Sa] and [Ze].

5.2 Examples arising from flat CR -geometry

Let us consider the complex vector space \mathbf{C}^{n+1} , endowed with the hermitian form $h^{1,n-1}(z) = -|z_1|^2 + |z_2|^2 + |z_3|^2 + \dots + |z_{n+1}|^2$. We consider $C_{\mathbf{C}}^{1,n}$, the lightcone defined as $\{z \in \mathbf{C}^{n+1} \mid h^{1,n}(z) = 0\}$, and call Ω^- the open set $\{z \in \mathbf{C}^{n+1} \mid h^{1,n}(z) < 0\}$. If we project Ω^- on the complex projective space \mathbf{CP}^n , we get the complex hyperbolic space $\mathbf{H}_{\mathbf{C}}^n$. If we project $C_{\mathbf{C}}^{1,n}$ minus the origin on \mathbf{CP}^n , we get a sphere \mathbf{S}^{2n-1} , naturally endowed with a CR -structure. This CR -sphere can be seen at the infinity of $\mathbf{H}_{\mathbf{C}}^n$. If, instead of looking at the complex directions of $C_{\mathbf{C}}^{1,n}$, we consider the quotient $C_{\mathbf{C}}^{1,n}/\mathbf{R}^*$ of $C_{\mathbf{C}}^{1,n}$ by the real homotheties, then the space that we get is Einstein's universe of dimension $2n$. In other words, there is a fibration $f : \mathbf{Ein}_{2n} \rightarrow \mathbf{S}^{2n-1}$ whose fibers are circles. The fibration is preserved by the group $U(1, n)$, which acts on \mathbf{Ein}_{2n} as a subgroup of $O(2, 2n)$. If Z denotes the center of $U(1, n)$ (homotheties by complex numbers of modulus 1), then the fibers of f are exactly the orbits of Z on \mathbf{Ein}_{2n} . These orbits are lightlike geodesics.

Proposition 7. *If $\Gamma \in U(1, n)$ is a discrete group, whose projection $\hat{\Gamma}$ on $PU(1, n)$ acts properly discontinuously on $\hat{\Omega} \subset \mathbf{S}^{2n-1}$, then Γ is a Kleinian group of \mathbf{Ein}_{2n} and acts properly discontinuously on $\Omega = f^{-1}(\hat{\Omega})$. If \hat{G} acts with compact quotient on $\hat{\Omega}$, so does Γ on Ω .*

Remark 5. *The group $PU(1, n)$ acting on \mathbf{S}^{2n-1} is a convergence group, and there is a good notion of limit set for a discrete group \hat{G} as above (see for example [A]). In fact, it is not difficult to check that the Lorentzian Kleinian groups Γ built as in proposition 7 are of the first type. Their limit set is just the preimage by f of the limit set $\hat{\Lambda}_{\hat{\Gamma}}$ of $\hat{\Gamma}$ on \mathbf{S}^{2n-1} .*

To illustrate this case, let us mention the two following examples:

Example 1.

We write each $z \in \mathbf{C}^{n+1}$ as $z = (x, y)$ with x and y in \mathbf{R}^n . We identify the real hyperbolic space $\mathbf{H}_{\mathbb{R}}^n$ with the set of points $(x, 0)$ with $-x_1^2 + x_2^2 + \dots + x_{n+1}^2 = -1$ and $x_1 > 0$. If (x, y) is moreover in the unit tangent bundle of $\mathbf{H}_{\mathbb{R}}^n$, it satisfies the two extra equations:

$$\begin{aligned} -x_1y_1 + x_2y_2 + \dots + x_ny_n &= 0 \\ -y_1^2 + y_2^2 + \dots + y_{n+1}^2 &= 1 \end{aligned}$$

Projectivising, we get an open subset $\hat{\Omega} \subset \mathbf{S}^{2n-1}$. In fact $\hat{\Omega}$ is precisely \mathbf{S}^{2n-1} minus a $(n-1)$ -dimensional sphere Σ (the projection on \mathbf{S}^{2n-1} of the set $\{z = (x, 0) \mid -x_1^2 + x_2^2 + \dots + x_{n+1}^2 = 0\}$).

Now, the subgroup $G = O(1, n)$ of real matrices in $U(1, n)$ acts on \mathbf{S}^{2n-1} , and preserves $\hat{\Omega}$. Identifying $\hat{\Omega}$ with $T^1\mathbf{H}_{\mathbb{R}}^n$, we get that G acts properly and transitively on $\hat{\Omega}$. As a consequence, we have the following:

Fact 1. *Any discrete group Γ in $O(1, n)$ acts properly discontinuously on $\hat{\Omega}$. Seen as a subgroup of $O(2, 2n)$ it yields a Kleinian group acting on \mathbf{Ein}_{2n} .*

The Kleinian manifold Ω/Γ obtained in this way are circle bundles over $T^1(N)$, where N is the hyperbolic manifold $\mathbf{H}_{\mathbb{R}}^n/\Gamma$.

Example 2.

Inside $U(1, n)$, there is a group G isomorphic to the Heisenberg group of dimension $2n-1$. The group G fixes a point p_{∞} on \mathbf{S}^{2n-1} and acts simply transitively on the complement of this point. By proposition 7, any discrete group in G will yield a Lorentzian Kleinian group, acting properly on the complement of a lightlike geodesic. The Kleinian manifolds obtained in this way will be circle bundles over nilmanifolds.

5.3 Subgroups of $O(1, r) \times O(1, s)$

We still endow $\mathbf{R}^{2,n}$ with the quadratic form $q^{2,n}(x) = -2x_1x_{n+2} + 2x_2x_{n+1} + x_3^2 + \dots + x_n^2$, and we consider an orthogonal splitting $\mathbf{R}^{2,n} = E_1 + E_2$ with E_1 and E_2 two spaces of signature $(1, r)$ and $(1, s)$ respectively ($r \neq 0, s \neq 0$ and $r + s = n$). We suppose also $r \leq s$. For example, we take $E_1 = (e_1, e_3, \dots, e_s, e_{n+2})$ and $E_2 = (e_2, e_{r+1}, \dots, e_{n+1})$. The subgroup G of $O(2, n)$ preserving this splitting is isomorphic to the product $O(1, r) \times O(1, s)$. Before describing some examples of Kleinian groups in G , let us say a few words about the geometric meaning of this splitting on \mathbf{Ein}_n .

Lemma 3. *We can write \mathbf{Ein}_n as a union $\Omega_1 \cup \Omega_2 \cup \Sigma$. The set Ω_1 (resp. Ω_2) is open, G -invariant, homogeneous under the action of G , and conformally equivalent to the product $\mathbf{dS}_r \times \mathbf{H}^s$ (resp. $\mathbf{H}^r \times \mathbf{dS}_s$). Σ is a singular, degenerate G -invariant hypersurface.*

Proof: We call π_1 and π_2 the projections of $\mathbf{R}^{2,n}$ on E_1 and E_2 respectively. The projection of vectors $u = (v, w)$ of $\mathbf{R}^{2,n}$ for which both $v = \pi_1(u)$ and

$w = \pi_2(u)$ are isotropic gives the hypersurface Σ . We will say more about it later.

The vectors $u = (v, w)$ for which neither v nor w is isotropic are of two kinds.

- *Those for which $q^{2,n}(v) > 0$.*

Since we work projectively, we can suppose that $q^{2,n}(v) = 1$ and $q^{2,n}(w) = -1$. In a further quotient by $-Id$, these vectors project on the product $\mathbf{dS}_r \times \mathbf{H}^s$. They constitute the open set Ω_1 .

- *Those for which $q^{2,n}(v) < 0$.*

These vectors project on a product $\mathbf{H}^r \times \mathbf{dS}_s$, and constitute the open set Ω_2 .

□

The hypersurface Σ can be seen as the conformal boundary of the spaces $\mathbf{dS}_r \times \mathbf{H}^s$ and $\mathbf{H}^r \times \mathbf{dS}_s$. Let us describe it more precisely. The isotropic vectors (v, w) of $\mathbf{R}^{2,n}$ for which v and w are isotropic split themselves into two sets. Those for which either v or w is zero. Their projectivisation gives two Riemannian spheres Σ_1 and Σ_2 of dimension $(r - 1)$ and $(s - 1)$ respectively.

Those for which v and w are non zero project on the product of the projectivisation of the lightcone of E_1 by the lightcone of E_2 , namely $\mathbf{S}^{r-1} \times \mathbf{C}^{1,s}$. So Σ minus $\Sigma_1 \cup \Sigma_2$ has two connected components, each of which is diffeomorphic to $\mathbf{S}^{r-1} \times \mathbf{S}^{s-1} \times \mathbf{R}$. One can check that Σ is obtained as the union of the lightlike geodesics intersecting both Σ_1 and Σ_2 .

We now give some examples of Kleinian groups in G .

Example 3.

Let us take a discrete group $\hat{\Gamma}$ inside $O(1, r)$ and any representation ρ of $\hat{\Gamma}$ inside $O(1, s)$. We call $\Gamma_\rho = \text{Graph}(\hat{\Gamma}, \rho) = \{(\hat{\gamma}, \rho(\hat{\gamma})) | \hat{\gamma} \in \hat{\Gamma}\}$. Then Γ_ρ is a Lorentzian Kleinian group of $O(2, n)$. Indeed, its action on $\Omega_2 = \mathbf{H}^r \times \mathbf{dS}_s$ is clearly proper. Let us say a little bit more about the limit set of these groups. We call $\Lambda_{\hat{\Gamma}}$ the limit set of the group $\hat{\Gamma}$ on the sphere Σ_1 .

Case a): ρ is injective with discrete image.

A sequence (γ_k) of Γ_ρ can be written as a matrix $\begin{pmatrix} \hat{\gamma}_k & \\ & \rho(\hat{\gamma}_k) \end{pmatrix}$. If (γ_k) tends simply to infinity, so does the sequence $(\hat{\gamma}_k)$ (resp. $\rho(\hat{\gamma}_k)$) in $O(1, r)$ (resp. in $O(1, s)$). We thus see that (γ_k) has either mixed or balanced distortions. In particular, the group Γ_ρ is always of the first type in this case. The attracting and repelling circles of (γ_k) can be describe as follows. Since the sequence $(\hat{\gamma}_k)$ (resp. $\rho(\hat{\gamma}_k)$) tends simply to infinity in $O(1, r)$ (resp. $O(1, s)$), it has two attracting and repelling poles $p^+(\hat{\gamma}_k)$ and $p^-(\hat{\gamma}_k)$ (resp. $p^+(\rho(\hat{\gamma}_k))$ and $p^-(\rho(\hat{\gamma}_k))$) on Σ_1 (resp. Σ_2). Then $\Delta^+(\gamma_k)$ (resp. $\Delta^-(\gamma_k)$) is simply the lightlike geodesic of \mathbf{Ein}_n joining $p^+(\hat{\gamma}_k)$ and $p^+(\rho(\hat{\gamma}_k))$ (resp.

$p^-(\hat{\gamma}_k)$ and $p^-(\rho(\hat{\gamma}_k))$). In particular, the limit set Λ_{Γ_ρ} is a closed subset of Σ (strictly included in Σ if $\Lambda_{\hat{\Gamma}} \neq \Sigma_1$).

An interesting subcase arises when we take for $\hat{\Gamma}$ a cocompact lattice in $O(1, 2)$, and a quasi-fuchsian representation $\rho : \hat{\Gamma} \rightarrow O(1, s)$ ($s \geq 2$). The limit set of $\rho(\hat{\Gamma})$ on Σ_2 is a topological circle, and we get for the limit set Λ_{Γ_ρ} a topological torus. One can prove moreover (but we don't do it here) that the action of Γ_ρ is cocompact on the complement of its limit set.

Cas b): ρ is not injective with discrete image.

In this case, there is a sequence (γ_k) tending simply to infinity in Γ_ρ such that $\rho(\hat{\gamma}_k)$ is bounded. Such a sequence (γ_k) has bounded distortion and the group Γ_ρ is of the second type. The attracting and repelling poles $p^+(\gamma_k)$ and $p^-(\gamma_k)$ are both on Σ_1 . In fact they are the attracting and repelling poles of $(\hat{\gamma}_k)$ (acting as a sequence of $O(1, r)$ on Σ_1). In this case, the limit set Λ_{Γ_ρ} is just the union of lightcones with vertex on $\Lambda_{\hat{\Gamma}}$.

6 About Klein's combination theorem

Until now, the examples of Kleinian groups we gave could not appear completely satisfactory, since they arise from geometrical contexts such as Lorentzian spaces with constant curvature or flat CR -geometry, and in some way are not "typical" of conformally flat Lorentzian geometry. For example, we still don't have examples of Zariski dense Kleinian groups on \mathbf{Ein}_n . One way to construct other classes of examples, is to combine two existing Lorentzian Kleinian groups to get a third one. In the theory of Kleinian groups on the sphere, this kind of construction is achieved thanks to the celebrated Klein's combination theorem ([A] [Ma]). We state now a generalized version of this theorem. Before this, we need the:

Definition 5. *Let X be a manifold. A Kleinian group on X is a discrete subgroup of diffeomorphisms Γ acting properly discontinuously on some nonempty open set $\Omega \subset X$. We say that an open set $D \subset \Omega$ is a fundamental domain for the action of Γ on Ω if D does not contain two points of the same Γ -orbit and if moreover $\bigcup_{\gamma \in \Gamma} \gamma(\overline{D}) = \Omega$.*

Notation 2. *For any subset D of the manifold X , we call $Ext(D)$ the complement of D in X .*

Theorem 5 (Klein). *Let Γ_i ($i = 1, \dots, m$) a finite family of Kleinian groups on a compact connected manifold X . We suppose that each Γ_i acts cocompactly on some open subset Ω_i of X , with fundamental domain D_i . We assume moreover that for each $i \neq j$, $Ext(D_i) \subset D_j$, and that $D = \bigcap_{i=1}^m D_i \neq \emptyset$. Then:*

(i) *The group Γ generated by the Γ_i 's is isomorphic to the free product $\Gamma_1 * \dots * \Gamma_m$.*

(ii) The group Γ is Kleinian. More precisely, $\Omega = \bigcup_{\gamma \in \Gamma} \gamma(\overline{D})$ is an open subset of X , and Γ acts properly discontinuously and cocompactly on Ω , with fundamental domain D .

Proof: We do the proof for two groups Γ_1 and Γ_2 , the final result being then obtained by induction. Let $\gamma = \gamma_s \gamma_{s-1} \dots \gamma_2 \gamma_1$ be a word of Γ such that $\gamma_i \in G_{j_i}$ ($j_i \in \{1, 2\}$) and $j_i \neq j_{i+1}$. Then, the first condition on the fundamental domains yields the inclusions $\gamma_s \gamma_{s-1} \dots \gamma_2 \gamma_1(D) \subset \gamma_s \gamma_{s-1} \dots \gamma_2(\text{Ext}(\overline{D}_{j_1})) \subset \dots \subset \gamma_s(\text{Ext}(\overline{D}_{j_{s-1}})) \subset \text{Ext}(\overline{D}_{j_s})$. So, for any non trivial reduced g , $\gamma(D) \cap D = \emptyset$. This proves that γ can't be the identity and the point (i) follows. In the same way, we prove that $\gamma(\overline{D}) \cap \overline{D} = \emptyset$ as soon as $s > 1$. Since \overline{D} is compact in Ω_1 and Ω_2 and the action of Γ_1 and Γ_2 is proper, we get the:

Lemma 4. *The intersection $\gamma(\overline{D}) \cap \overline{D}$ is empty for all but a finite number of γ 's.*

Lemma 5. *There is a finite family $\gamma_1, \dots, \gamma_s$ of elements of Γ such that $\overline{D} \cup \gamma_1(\overline{D}) \cup \dots \cup \gamma_m(\overline{D})$ contains \overline{D} in its interior.*

Proof: We choose some open neighbourhood U_1 of ∂D_1 such that $U_1 \subset \Omega_1$ and \overline{U}_1 is a compact subset of Ω_1 . Since D_1 is a fundamental domain of Γ_1 , for each $x \in U_1$, there exists a $\gamma_x \in \Gamma_1$ such that $x \in \gamma_x(\overline{D}_1)$. But since the action of Γ_1 is proper $\gamma(\overline{D}_1) \cap U_1$ is nonempty only for a finite number of elements $\gamma_1^{(1)}, \dots, \gamma_s^{(1)}$ of Γ_1 . Thus $\overline{D}_1 \cup U_1$ is included in $\overline{D}_1 \cup \gamma_1^{(1)}(\overline{D}_1) \cup \dots \cup \gamma_s^{(1)}(\overline{D}_1)$ and \overline{D}_1 is contained in the interior of $\overline{D}_1 \cup \gamma_1^{(1)}(\overline{D}_1) \cup \dots \cup \gamma_s^{(1)}(\overline{D}_1)$. But if $D'_1 = D_1 \setminus K$, where K is a compact subset of D_1 , then we also have $\overline{D}'_1 \cup U_1 \subset \overline{D}'_1 \cup \gamma_1^{(1)}(\overline{D}'_1) \cup \dots \cup \gamma_s^{(1)}(\overline{D}'_1)$. In particular, when K is the exterior of D_2 , we get that $\overline{D} \cup U_1 \subset \overline{D} \cup \gamma_1^{(1)}(\overline{D}) \cup \dots \cup \gamma_s^{(1)}(\overline{D})$. Now, we can apply the same argument for a neighbourhood U_2 of ∂D_2 in Ω_2 . We get a finite family $\gamma_1^{(2)}, \dots, \gamma_t^{(2)}$ of Γ_2 such that $\overline{D} \cup U_2 \subset \overline{D} \cup \gamma_1^{(2)}(\overline{D}) \cup \dots \cup \gamma_t^{(2)}(\overline{D})$. Setting $m = s + t$, $\gamma_i = \gamma_i^{(1)}$ for $i = 1, \dots, s$ and $\gamma_{s+i} = \gamma_i^{(2)}$ for $i = 1, \dots, t$, we get the lemma. \square

As a consequence of this lemma, we get that the set $\Omega = \bigcup_{\gamma \in \Gamma} \gamma(\overline{D})$ is an open set.

It remains to prove that the action of Γ on Ω is proper. Indeed, since Γ is not *a priori* a convergence group, the fact that Γ acts discontinuously on Ω no longer ensures that the action is proper. That is why our assumptions (in particular the assumption of cocompactness) are stronger as for the classical Klein's theorem on the sphere.

Suppose, on the contrary, that there is a sequence (x_i) of Ω converging to $x_\infty \in \Omega$, and a sequence (γ_i) tending to infinity in Γ , such that $y_i = \gamma_i(x_i)$ converges to $y_\infty \in \Omega$. We can assume that $x_\infty \in \overline{D}$. On the other hand, by definition of Ω , there is a γ_0 such that $y_\infty \in \gamma_0(\overline{D})$. The lemma 5 ensures

We now take some $g \in \text{Conf}(\mathbf{Ein}_n)$ as in the lemma above. Let us choose V_1 (resp. V_2) an open tubular neighbourhood of Δ_1 (resp. Δ_2) such that $\overline{V_1} \subset D_1$ (resp. $\overline{V_2} \subset D_2$). The complement of V_i ($i = 1, 2$) in \mathbf{Ein}_n is denoted by $\text{Ext}(V_i)$. It follows from proposition 5 (points (i) and (ii)) that the set dynamically associated to $\text{Ext}(V_2)$ with respect to (g^k) is included in Δ^+ . Since $\text{Ext}(V_2)$ contains a lightlike geodesic, it is exactly Δ^+ . Hence, for k_0 sufficiently large, $g^{k_0}(\text{Ext}(V_2)) \subset V_1$. We call $\Gamma'_2 = g^{k_0}\Gamma_2g^{-k_0}$. The group Γ'_2 is a cocompact Lorentzian Kleinian group, with fundamental domain $D'_2 = g^{k_0}(D_2)$. But $g^{k_0}(D_2)$ contains $g^{k_0}(\text{Int}(V_2))$, and as we just saw, $\text{Ext}(V_1) \subset g^{k_0}(\text{Int}(V_2))$. So $\text{Ext}(D'_2) \subset D_1$. We can then apply theorem 5, and we get that the group generated by Γ'_2 and Γ_1 is still Kleinian, cocompact, and isomorphic to $\Gamma_1 * \Gamma'_2$, i.e $\Gamma_1 * \Gamma_2$.

Example 4

All the cocompact Lorentzian Kleinian groups of the examples 1 and 2 of section 5 satisfy the hypothesis of theorem 2. This is also the case of most instances of example 3, when ρ is injective with discrete image. Thus, such groups can be combined and give new examples. Notice that in the proof of theorem 2, the gluing element g can be chosen in many ways. In particular, starting from two groups of the examples 1, 2 or 3, suitable choices of g will give combined groups which are Zariski dense in $O(2, n)$.

6.2 Lorentzian surgery

Theorem 2 is in fact the group theoretical aspect of a slightly more general process of conformal Lorentzian surgery.

Let M_1 and M_2 be two conformally flat Lorentzian manifolds (we don't make any compactness assumption). Suppose that M_1 contains a closed lightlike geodesic Δ_1 , admitting some open neighbourhood U_1 which embeds conformally, via a certain embedding ϕ_1 , into \mathbf{Ein}_n . Suppose moreover that the same property is satisfied by M_2 , for a closed lightlike geodesic Δ_2 , an open neighbourhood U_2 , and a conformal embedding ϕ_2 . We can suppose that $\phi_1(\Delta_1)$ and $\phi_2(\Delta_2)$ are disjoint in \mathbf{Ein}_n . By lemma 6, $\phi_1(\Delta_1)$ and $\phi_2(\Delta_2)$ are the attracting and repelling circles of some element $g \in \text{Conf}(\mathbf{Ein}_n)$. As in the proof of theorem 2, there exist two open neighbourhoods V_1 and V_2 of Δ_1 and Δ_2 respectively, such that $V_1 \subset U_1$, $V_2 \subset U_2$, and $g(\text{Ext}(\phi_2(V_2))) = \phi_1(V_1)$. In particular $g(\partial(\phi_2(V_2))) = \partial(\phi_1(V_1))$ (recall that ∂ denotes the boundary). so, the element g provides a gluing map f between ∂V_1 and ∂V_2 . We denote by \dot{M}_1 (resp. \dot{M}_2) the manifold M_1 (resp. M_2) with V_1 (resp. V_2) removed. We call $M = \dot{M}_1 \#_f \dot{M}_2$ the manifold obtained from $\dot{M}_1 \cup \dot{M}_2$, after identification of ∂V_1 and ∂V_2 by means of the map f . Since $g \in \text{Conf}(\mathbf{Ein}_n)$, the "surgered manifold" M is still endowed with a conformally flat Lorentzian structure. Theorem 2 ensures that if one starts with two compact Kleinian structures M_1 and M_2 , the conformally flat structure on $\dot{M}_1 \#_f \dot{M}_2$ is still Kleinian.

Remark 6. *This surgery process is reminiscent of Kulkarni's construction of a conformally flat Riemannian structure on the connected sum of two conformally flat Riemannian manifolds ([K1]). We don't know if the connected sum of two conformally flat Lorentzian manifolds can still be endowed with a conformally flat Lorentzian structure.*

7 Lorentzian Schottky groups

As an application of the former sections, we study here the Lorentzian Schottky groups. These groups are interesting since we can completely determine their limit set and the Kleinian manifolds they uniformize. Moreover, they can be used to construct examples of conformally flat manifolds with some peculiar properties (see [Fr2]).

Let us consider a family $\{(\Delta_1^-, \Delta_1^+), \dots, (\Delta_g^-, \Delta_g^+)\}$ of pairs of lightlike geodesics in \mathbf{Ein}_n . We suppose moreover that the Δ_i^\pm are all disjoint. By lemma 6, there exists a family s_1, \dots, s_g of elements of $\text{Conf}(\mathbf{Ein}_n)$ with mixed dynamics such that the attracting and repelling circles of s_i are precisely Δ_i^+ and Δ_i^- . Looking if necessary at suitable powers $s_i^{k_i}$ of s_i , we can find open tubular neighbourhoods U_i^\pm of the Δ_i^\pm such that:

- (i) The $\overline{U_i^\pm}$ are all disjoint.
- (ii) $s_i(\text{Ext}(U_i^-)) = \overline{U_i^+}$ for all $i = 1, \dots, g$.

Such a group $\Gamma = \langle s_1, \dots, s_g \rangle$ is called a *Lorentzian Schottky group*. Properties (i) and (ii) are classically known as *ping-pong dynamics* (see for example [dlH]). For each i , the group $\langle s_i \rangle$ acts properly cocompactly on the open set $\mathbf{Ein}_n \setminus \{\Delta_i^- \cup \Delta_i^+\}$, and a fundamental domain is just given by $D_i = \mathbf{Ein}_n \setminus \{\overline{U_i^+} \cup \overline{U_i^-}\}$. Now, since the $\overline{U_i^\pm}$ are disjoint, we get that $\text{Ext}(D_i) \subset D_j$ for all $i \neq j$. If we call $D = \bigcap_{i=1}^g D_i$, it is clear that $D \neq \emptyset$. We then apply theorem 5 to obtain:

Proposition 8. *A Lorentzian Schottky group $\Gamma = \langle s_1, \dots, s_g \rangle$ is a free group of $\text{Conf}(\mathbf{Ein}_n)$. Moreover, Γ is Kleinian: it acts properly and cocompactly on $\Omega = \bigcup_{\gamma \in \Gamma} \gamma(\overline{D})$. A fundamental domain for this action is given by $D = \bigcap_{i=1}^g D_i$.*

We are now going to describe Ω , and its complement $\Lambda \subset \mathbf{Ein}_n$ more precisely.

Let us recall that in a finitely generated free group, each element γ can be written in a unique way as a reduced word in the generators. We denote by $|\gamma|$ the length of this word. Let us also recall that we can define the boundary $\partial\Gamma$ of Γ as the set of totally reduced words of infinite length. So, the elements of the boundary can be written $s_{i_1}^{\epsilon_1} \dots s_{i_k}^{\epsilon_k} \dots$ with $\epsilon_j \in \{\pm 1\}$ and $i_j \epsilon_j \neq -i_{j+1} \epsilon_{j+1}$ for all $j \geq 1$. Since we supposed $g \geq 2$, the boundary $\partial\Gamma$ is a compact metrizable space, homeomorphic to a Cantor set (see [GdlH]).

For each $k \in \mathbb{N}$, we call $F_k = \bigcup_{|\gamma| \leq k} \gamma(\overline{D})$, with the convention $F_0 = \overline{D}$. It's not difficult to check that $F_{k-1} \subset F_k$, and $\Omega = \bigcup_{k \in \mathbb{N}} F_k$. So, $\Lambda = \bigcap_{k \in \mathbb{N}} \text{Ext}(F_k)$. For each k , we set $\Lambda_k = \text{Ext}(F_k)$, and thus, we also have $\Lambda = \bigcap_{k \in \mathbb{N}} \overline{\Lambda}_k$. The set $\overline{\Lambda}_k$ is a disjoint union of exactly $2g \cdot (2g-1)^k$ connected components, in one to one correspondence with the words of length $k+1$ in Γ . For example, to the word $s_{i_1}^{\epsilon_1} \dots s_{i_{k+1}}^{\epsilon_{k+1}}$ corresponds the component $s_{i_1}^{\epsilon_1} \dots s_{i_k}^{\epsilon_k} (\overline{U}_{i_{k+1}}^{\epsilon_{k+1}})$ of $\overline{\Lambda}_k$. We can now state:

Lemma 7. *There is an homeomorphism K between the boundary $\partial\Gamma$ and the space of connected components of Λ (endowed with the Hausdorff topology for the compact subsets of \mathbf{Ein}_n).*

Proof: Let $\gamma_\infty = s_{i_1}^{\epsilon_1} \dots s_{i_k}^{\epsilon_k} \dots$ be an element of $\partial\Gamma$. We call $\gamma_k = s_{i_1}^{\epsilon_1} \dots s_{i_k}^{\epsilon_k}$ and we look at the decreasing sequence of compact subsets $K(\gamma_k) = s_{i_1}^{\epsilon_1} \dots s_{i_{k-1}}^{\epsilon_{k-1}} (\overline{U}_{i_k}^{\epsilon_k})$. This decreasing sequence of compact sets tends to a limit compact set $K(\gamma_\infty)$ for the Hausdorff topology. Since the U_i^\pm are connected, so are the $K(\gamma_k)$, and $K(\gamma_\infty)$ is itself connected. Let us remark that if γ_∞ and γ'_∞ are distinct in $\partial\Gamma$, then $K(\gamma_k)$ and $K(\gamma'_k)$ are disjoint for k large (they represent two distinct components of $\overline{\Lambda}_k$), so that $K(\gamma_\infty)$ and $K(\gamma'_\infty)$ are disjoint.

Reciprocally, choose $x_\infty \in \Lambda$. Since $\Lambda = \bigcap_{k \in \mathbb{N}} \overline{\Lambda}_k$ with $\overline{\Lambda}_{k+1} \subset \overline{\Lambda}_k$, x_∞ must be an element of some connected component $C_k \subset \overline{\Lambda}_k$ for each k . Moreover $C_{k+1} \subset C_k$. But C_k is then a decreasing sequence of compact subsets of the form $s_{i_1}^{\epsilon_1} \dots s_{i_{k-1}}^{\epsilon_{k-1}} (\overline{U}_{i_k}^{\epsilon_k})$, and thus converges to a limit compact set $K(\gamma_\infty)$ for $\gamma_\infty = s_{i_1}^{\epsilon_1} \dots s_{i_k}^{\epsilon_k} \dots$.

We have proved that the mapping K between $\partial\Gamma$ and the set of connected components of Λ is a bijection. It remains to prove that it is an homeomorphism, and for this, it is sufficient to show that K is continuous. Let us consider a sequence $\gamma_\infty^{(n)}$ of elements of Γ , converging to some γ_∞ . It means that there is a sequence (r_n) of integers which tends to infinity, such that $\gamma_\infty^{(n)}$ and γ_∞ have the same r_n first letters. For each $n \in \mathbb{N}$, $K(\gamma_\infty^{(n)})$ is a decreasing sequence of compact sets $C_k^{(n)}$, where each $C_k^{(n)}$ is a connected component of $\overline{\Lambda}_k$. On the other hand, $K(\gamma_\infty)$ is the limit of a decreasing sequence of C_k , where each C_k is a connected component of $\overline{\Lambda}_k$. Since $\gamma_\infty^{(n)}$ and γ_∞ have the same r_n first letters, we have $C_{r_n-1}^{(n)} = C_{r_n-1}$ for all n . Thus, the limit, as n tends to infinity, of $C_{r_n-1}^{(n)}$ is $K(\gamma_\infty)$. But since $K(\gamma_\infty^{(n)}) \subset C_{r_n-1}^{(n)}$, we get that $\lim_{n \rightarrow \infty} K(\gamma_\infty^{(n)}) = K(\gamma_\infty)$ and we are done. \square

The next step is the:

Lemma 8. *The connected components of Λ are lightlike geodesics.*

Proof: Let us consider $\gamma_\infty = s_{i_1}^{\epsilon_1} \dots s_{i_k}^{\epsilon_k} \dots$ in the boundary of Γ . We know that $K(\gamma_\infty)$ is the limit of the sequence $s_{i_1}^{\epsilon_1} \dots s_{i_{k-1}}^{\epsilon_{k-1}} (\overline{U}_{i_k}^{\epsilon_k})$. Since the sequence is

decreasing, the limit remains the same if we consider a subsequence. Thus, we can make the extra assumption that $K(\gamma_\infty)$ is the limit of a sequence $\gamma_k(\overline{U}_{j_0}^{\epsilon_{j_0}})$, such that (γ_k) tends simply to infinity and the first and last letters of γ_k are always the same, namely $s_{i_1}^{\epsilon_1}$ and $s_{j_1}^{\epsilon_{j_1}}$. Let us precise that $j_1 \epsilon_{j_1} \neq -j_0 \epsilon_{j_0}$. We are going to discuss the different possible dynamics for (γ_k) , and we first prove that (γ_k) can't have bounded distortion.

Suppose that it is the case. We call p^+ (resp. p^-) and C^+ (resp. C^-) the attracting (resp. repelling) pole and cone of (γ_k) . If x is a point of D , then for all $k \in \mathbb{N}$, $\gamma_k(x) \in U_{i_1}^{\epsilon_1}$ and $\gamma_k^{-1}(x) \in U_{j_1}^{-\epsilon_{j_1}}$. So, we must have $p^+ \in U_{i_1}^{\epsilon_1}$ and $p^- \in U_{j_1}^{-\epsilon_{j_1}}$. In particular, p^- is not in $U_{j_0}^{\epsilon_{j_0}}$. On the other hand, this is a general fact that in \mathbf{Ein}_n , any lightlike cone meets any lightlike geodesic (just because degenerate hyperplanes always meet null 2-planes in $\mathbf{R}^{2,n}$). In particular, the cone C^- meets $\Delta_{j_0}^{\epsilon_{j_0}}$, and thus $U_{j_0}^{\epsilon_{j_0}}$. We call $V_{j_0}^{\epsilon_{j_0}} = C^- \cap U_{j_0}^{\epsilon_{j_0}}$. Since $U_{j_0}^{\epsilon_{j_0}}$ does not contain p^- , we infer from proposition 4 (points (i) and (ii)) that $K(\gamma_\infty) = D_{(\gamma_k)}(\overline{V}_{j_0}^{\epsilon_{j_0}})$. More precisely, if $\hat{V}_{j_0}^{\epsilon_{j_0}}$ is the set of lightlike geodesics of C^- meeting $V_{j_0}^{\epsilon_{j_0}}$, then $K(\gamma_\infty)$ is the closure of the union the lightlike geodesics of $\hat{\gamma}_\infty(\hat{V}_{j_0}^{\epsilon_{j_0}})$ (see proposition 4 for the notation $\hat{\gamma}_\infty$). In particular, $K(\gamma_\infty)$ contains a lightlike geodesic. Now, some lightlike geodesic of C^- does not meet $V_{j_0}^{\epsilon_{j_0}}$. Indeed, if it is not the case, the point (ii) of proposition 4 ensures that $K(\gamma_\infty) = C^-$. But if we take $\gamma'_\infty \neq \gamma_\infty$, $K(\gamma'_\infty)$ contains some lightlike geodesic by the remark above, and since any lightlike geodesic meets C^- , we get a contradiction with the fact that $K(\gamma_\infty)$ and $K(\gamma'_\infty)$ have to be disjoint.

Now, let us perturb slightly the sets $U_{j_0}^{\epsilon_{j_0}}$ and $U_{j_0}^{-\epsilon_{j_0}}$ into some sets $U'_{j_0}{}^{\epsilon_{j_0}}$ and $U'_{j_0}{}^{-\epsilon_{j_0}}$, in order to get another fundamental domain D' , very close to D . Since it is very near D , \overline{D}' is included in some F_k for k sufficiently large, and so $\bigcup_{\gamma \in \Gamma} \gamma(\overline{D}') = \bigcup_{\gamma \in \Gamma} \gamma(D)$. We prove as above that the limit of the compact sets $\gamma_k(U'_{j_0}{}^{\epsilon_{j_0}})$ is still a connected component of Λ , and consequently of the form $K(\gamma'_\infty)$. We just saw that some lightlike geodesics of C^- don't meet $\overline{V}_{j_0}^{\epsilon_{j_0}}$, so that $\hat{V}_{j_0}^{\epsilon_{j_0}}$ is not the whole \mathbf{S}^{n-2} . It is thus possible to choose $U'_{j_0}{}^{\epsilon_{j_0}}$ in such a way that some points of $\hat{V}'_{j_0}{}^{\epsilon_{j_0}}$ are not in $\hat{V}_{j_0}^{\epsilon_{j_0}}$. But then, $K(\gamma'_\infty)$ and $K(\gamma_\infty)$ will be two different components, hence disjoint. On the other hand, since the intersection of $U_{j_0}^{\epsilon_{j_0}}$ and $U'_{j_0}{}^{\epsilon_{j_0}}$ is not empty ($\Delta_{j_0}^{\epsilon_{j_0}}$ is inside), $K(\gamma_\infty)$ and $K(\gamma'_\infty)$ must have some common points. We thus get a contradiction.

It remains to deal with the case where (γ_k) has mixed or balanced distortions. Once again, if x is a point of D then for all $k \in \mathbb{N}$, $\gamma_k(x) \in U_{i_1}^{\epsilon_1}$ and $\gamma_k^{-1}(x) \in U_{j_1}^{-\epsilon_{j_1}}$. Hence, the attracting circle Δ^+ is in $U_{i_1}^{\epsilon_1}$ and the repelling one Δ^- is in $U_{j_1}^{-\epsilon_{j_1}}$. In particular $\overline{U}_{j_0}^{\epsilon_{j_0}}$ does not meet Δ^- . We infer from proposition 5 and proposition 3 that $\lim_{k \rightarrow \infty} \gamma_k(\overline{U}_{j_0}^{\epsilon_{j_0}}) \subset \Delta^+$, but since $\overline{U}_{j_0}^{\epsilon_{j_0}}$

contains a lightlike geodesic, we have the equality $\lim_{k \rightarrow \infty} \gamma_k(\overline{U}_{j_0}^{\epsilon_{j_0}}) = \Delta^+$. We finally obtain that $K(\gamma_\infty) = \Delta^+$. □

7.1 Proof of Theorem 3

We begin by proving that the group Γ is of the first type. Suppose on the contrary that there is some sequence (γ_k) in Γ with bounded distortion. Then D meets the repelling cone C^- . Indeed, if not, C^- would be included in some U_i^\pm , for example U_1^+ . But since Δ_1^- meets C^- , the intersection between Δ_1^- and U_1^+ would be non empty, a contradiction. By the point (ii) of proposition 4, $\lim_{k \rightarrow \infty} \gamma_k(\overline{D})$ is a compact subset containing infinitely many lightlike geodesics. But $\lim_{k \rightarrow \infty} \gamma_k(\overline{D})$ is also a connected subset of Λ . This contradicts the fact that the connected components of Λ are lightlike geodesics.

Now, we claim that the equality $\Lambda_\Gamma = \Lambda$ holds. Indeed, for any sequence (γ_k) of Γ tending simply to infinity, (γ_k) tends to $\Delta^+(\gamma_k)$. We thus see that $\Lambda_\Gamma \subset \Lambda$. Now, it is a general fact that if a group Γ acts properly co-compactly on some open set Ω , it can't act properly on some open set Ω' strictly containing Ω . So, Ω can't be strictly contained in $\mathbf{Ein}_n \setminus \Lambda_\Gamma$ and we get $\Lambda_\Gamma = \Lambda$.

Thanks to the homeomorphism K , we get that since the action of Γ on its boundary is minimal (see for instance [GdlH]), the action of Γ on the space of lightlike geodesics of Λ_Γ is also minimal.

We now prove that Λ_Γ is the product of \mathbf{RP}^1 with a Cantor set. The space \mathbf{Ein}_n is the quotient of $\mathbf{S}^1 \times \mathbf{S}^{n-1}$ by the product of antipodal maps, so that there is a fibration $f : \mathbf{Ein}_n \rightarrow \mathbf{RP}^1$. The fibers of f are conformal Riemannian spheres of codimension one. In the projective model, they are obtained as the projection of the intersection between $C^{2,n}$ and some hyperplanes $P \subset \mathbf{R}^{2,n}$ of Lorentzian signature. As a consequence, any lightlike geodesic is transverse to any fiber of f . Let us choose a fiber \mathcal{F}_0 above a point t_0 of \mathbf{RP}^1 . From lemmas 7 and 8, Λ (and thus Λ_Γ) is transverse to \mathcal{F}_0 and intersects it along a Cantor set \mathcal{C} . For each $x \in \mathcal{C}$, we call $x(t)$ the unique element of $f^{-1}(t) \cap \Lambda_\Gamma$ such that x and $x(t)$ are on the same lightlike geodesic of Λ . Then, the lemma 7 ensures that the following mapping is an homeomorphism.

$$\begin{aligned} \mathbf{RP}^1 \times \mathcal{C} &\rightarrow \Lambda \\ (t, x) &\mapsto x(t) \end{aligned}$$

The point (ii) is then proved.

Thanks to the homeomorphism K , we get that since the action of Γ on its boundary is minimal (see for instance [GdlH]), the action of Γ on the space of lightlike geodesics of Λ_Γ is also minimal, what proves (iii).

For the proof of (iv), we refer to the theorem 5 of [Fr2] (in fact, in [Fr2], we considered only particular cases of Schottky groups, but the proof of theorem 5 includes the general case).

Acknowledgments: I would like to thank Thierry Barbot for his useful remarks on this text, the ACI "Structures géométriques et trous noirs", as well as the referee, who made a lot of valuable suggestions to improve the original version. Thank you also to the *Institut für Mathematik* of Humboldt University in Berlin, where part of this work was done in autumn 2003.

References

- [A] B.N.APANASOV - *Conformal geometry of discrete groups and manifolds*. de Gruyter Expositions in Mathematics, 32. Walter de Gruyter and Co., Berlin, 2000.
- [B] Y.BENOIST - *Propriétés asymptotiques des groupes linéaires*. *Geom. Funct. Anal.* 7 (1997), no. 1, 1–47.
- [BTZ1] M.BAÑADOS; C.TEITELBOIM; J.ZANELLI - *Black hole in three-dimensional spacetime*. *Phys. Rev. Lett.* 69 (1992), no. 13, 1849–1851
- [BTZ2] M.BAÑADOS; M.HENNEAUX; C.TEITELBOIM; J.ZANELLI - *Geometry of the 2 + 1 black hole*. *Phys. Rev. D* (3) 48 (1993), no. 4, 1506–1525.
- [CK] M.CAHEN; Y.KERBRAT - *Domaines symétriques des quadriques projectives*. *J. Math. Pures Appl.* (9) 62 (1983), no. 3, 327–348.
- [Ca] Y.CARRIÈRE - *Autour de la conjecture de Markus sur les variétés affines*. *Invent. Math.*, vol 95, n°3, (1989), pp. 615-628.
- [CaD] Y.CARRIÈRE; F.DAL'BO - *Generalisations du premier théorème de Bieberbach sur les groupes cristallographiques*. *Enseign. Math.* (2) 35 (1989), no. 3-4, 245–262.
- [dlH] P.DE LA HARPE - *Free groups in linear groups*. *Enseign. Math.* (2) 29 (1983), no. 1-2, 129–144.
- [DGo] T.A.DRUMM; W.M.GOLDMAN - *The geometry of crooked planes*. *Topology* 38 (1999), no. 2, 323–351.
- [Fr1] C.FRANCES - *Géométrie et dynamique lorentziennes conformes*. Thèse. ENS-Lyon (2002). Available at <http://www.math.u-psud.fr/~frances/>.

- [Fr2] C.FRANCES - *Sur les variétés lorentziennes dont le groupe conforme est essentiel.* (2003) . To appear in *Mathematische Annalen*.
- [Fr3] C.FRANCES - *Une preuve du théorème de Liouville en géométrie conforme dans le cas analytique.* *Enseign. Math.* (2) 49 (2003), no. 1-2, 95–100.
- [Fr4] C.FRANCES - *The conformal boundary of anti-de Sitter spacetimes.* To appear in “AdS-CFT Correspondance: Einstein Metrics and their Conformal Boundaries” (Olivier Biquard, ed.), IRMA Lectures in Mathematics and Theoretical Physics, Volume 8.
- [GdlH] E.GHYS; P.DE LA HARPE - *Sur les groupes hyperboliques d’après Mikhael Gromov.* *Progress in Mathematics*, 83. Birkhuser Boston, Inc., Boston, MA, 1990.
- [Go] W.GOLDMAN - *Complex hyperbolic kleinian groups.* in: *Complex geometry, Proceedings of the Osaka International Conference*, Marcel Dekker, Inc., New York, G. Komatsu and Y. Sakane (Eds.) (1992), 31–52.
- [HE] S.HAWKING; G.ELLIS - *The large scale structure of universe*-Cambridge University Press, 1973.
- [IW] A.IOZZI; D.WITTE - *Cartan-decomposition subgroups of $SU(2, n)$.* *J. Lie Theory* 11 (2001), no. 2, 505–543.
- [Kl] B.KLINGLER - *Complétude des variétés lorentziennes à courbure constante.*- *Math. Ann.*, vol 306, n°2, (1996), pp.353-370.
- [Ka] M.KAPOVICH - *Hyperbolic manifolds and discrete groups.* *Progress in Mathematics*, 183. Birkhuser Boston, Inc., Boston, MA, 2001.
- [K1] R.S.KULKARNI - *The principle of uniformization.* *J. Diff. Geo.* no. 13 (1978), 109-138.
- [K2] R.S.KULKARNI - *Groups with domains of discontinuity.* *Math. Ann.* 237 (1978), no. 3, 253–272.
- [KP] R.S.KULKARNI; U.PINKALL - *Uniformization of geometric structures with applications to conformal geometry.* *Differential geometry*, Pescola 1985, 190–209, *Lecture Notes in Math.*, 1209, Springer, Berlin, 1986.
- [KR] R.S.KULKARNI; F.RAYMOND - *Three-dimensional Lorentz space forms and Seifert fiber spaces-* *J.Diff.Geo* 21, 231-268 (1985)

- [Ma] B.MASKIT - *Kleinian groups*. Grundlehren der Mathematischen Wissenschaften , 287. Springer-Verlag, Berlin, 1988.
- [MK] K.MATSUZAKI; M.TANIGUCHI - *Hyperbolic manifolds and Kleinian groups*. Oxford Mathematical Monographs. Oxford Science Publications. The Clarendon Press, Oxford University Press, New York, 1998.
- [O’N] B.O’NEILL - *Semi-riemannian geometry with applications to relativity*- New York, Academic Press Inc, 1983.
- [P] H.POINCARÉ - *Mémoire sur les groupes Kleinéens*. Acta Mathematica. t3 (1883), 49–92.
- [PR] R.PENROSE ; W.RINDLER - *Vol.1. Spinors and space-time. Vol. 2. Spinor and twistor methods in space-time geometry*. Second edition. Cambridge Monographs on Mathematical Physics. Cambridge University Press, Cambridge, 1988.
- [Sa] F.SALEIN - *Variétés anti- de Sitter de dimension 3*- Thèse de Doctorat de l’école normale sup. de Lyon, 1999.
- [SV] J.SEADE; A.VERJOVSKY - *Higher dimensional complex Kleinian groups*. Math. Ann. 322 (2002), no. 2, 279–300.
- [Th] W.THURSTON - *Three dimensional geometry and topology. Vol 1.-* Princeton University Press, 1997. Edited by Silvio Levy.
- [Wo] J.WOLF - *Spaces of constant curvature* - Houston, Tex., Publish or Perish Inc., 1984, fifth edition.
- [Ze] A.ZEGHIB - *On closed anti de Sitter spacetimes*- Math. Ann. 310 695-716 (1998)

Charles FRANCES
 Laboratoire de Topologie et Dynamique
 Université de Paris-Sud, Bât. 430
 91405 ORSAY
 FRANCE
 email: Charles.Frances@math.u-psud.fr