

## A combinatorial interpretation of the Seidel generation of $q$ -derangement numbers

Robert J. Clarke

Pure Mathematics Department, University of Adelaide, Adelaide, South Australia 5005  
rclarke@maths.adelaide.edu.au

Guo-Niu Han

I.R.M.A., Université Louis-Pasteur et C.N.R.S., 67084 Strasbourg Cedex, France  
guoniu@math.u-strasbg.fr

Jiang Zeng

Institut Girard Desargues, Université Claude Bernard (Lyon I), F-69622 Villeurbanne Cedex  
zeng@desargues.univ-lyon1.fr

Received November 5, 1997

AMS Subject Classification: 05A15, 05A30

**Abstract.** In [8] Dumont and Randrianarivony have given several combinatorial interpretations for the coefficients of the Euler-Seidel matrix associated to  $n!$ . In this paper we consider a  $q$ -analogue of their results, which leads to the discovery of a new mahonian statistic “maf” on the symmetric group. We then give new proofs and generalizations of some results of Gessel and Reutenauer [12] and Wachs [17].

*Keywords:* mahonian statistics, permutations,  $q$ -derangement numbers, Seidel matrices

### 1. Introduction

Euler (see [8]) considered the *difference table*  $(d_n^k)_{0 \leq k \leq n}$ , where the generic coefficients  $d_n^k$  are defined by

$$d_n^n = n! \quad \text{and} \quad d_n^k = d_n^{k+1} - d_{n-1}^k \quad (1 \leq k \leq n-1). \quad (1.1)$$

Let  $a_n^k = d_{n+k}^k$  ( $n, k \geq 0$ ). Then the above relations can be written as

$$a_0^k = k! \quad \text{and} \quad a_n^k = a_n^{k-1} + a_{n+1}^{k-1} \quad (n, k \geq 0).$$

The matrix  $(a_n^k)_{n,k \geq 0}$  is also called the *Seidel matrix* associated to the sequence  $a_n^0$  in the literature (see [7, 9]). The first terms of these matrices are as follows:

$$\begin{array}{c|cccccc}
 n \setminus k & 0 & 1 & 2 & 3 & 4 & 5 \\
 \hline
 0 & 1 & & & & & \\
 1 & 0 & 1 & & & & \\
 2 & 1 & 1 & 2 & & & \\
 3 & 2 & 3 & 4 & 6 & & \\
 4 & 9 & 11 & 14 & 18 & 24 & \\
 5 & 44 & 53 & 64 & 78 & 96 & 120 \\
 \hline
 & & & & & & (d_n^k)
 \end{array}
 \quad \longrightarrow \quad
 \begin{array}{c|cccccc}
 k \setminus n & 0 & 1 & 2 & 3 & 4 & 5 \\
 \hline
 0 & 1 & 0 & 1 & 2 & 9 & 44 \\
 1 & 1 & 1 & 3 & 11 & 53 & \\
 2 & 2 & 4 & 14 & 64 & & \\
 3 & 6 & 18 & 78 & & & \\
 4 & 24 & 96 & & & & \\
 5 & 120 & & & & & \\
 \hline
 & & & & & & (a_n^k)
 \end{array}$$

Iterating the difference equation (1.1) we derive

$$a_n^0 = d_n^0 = n! \left( 1 - \frac{1}{1!} + \frac{1}{2!} - \cdots + (-1)^n \frac{1}{n!} \right), \quad (1.2)$$

which is the *classical derangement number*  $d_n$ , that is, the number of derangements on  $\{1, 2, \dots, n\}$  (cf. [16, p. 67]).

In several recent papers [4, 6, 12, 17], the  $q$ -maj counting of the derangements on  $\{1, 2, \dots, n\}$  has been studied. Consider the  $q$ -derangement numbers  $d_n(q)$  defined by

$$d_n(q) = \sum_{\sigma \in \mathcal{D}_n} q^{\text{maj}\sigma}, \quad (1.3)$$

where  $\mathcal{D}_n$  is the set of all derangements on  $\{1, 2, \dots, n\}$ . Then the following  $q$ -analogue of equation (1.2) has been obtained:

$$d_n(q) = [n]_q! \sum_{i=0}^n (-1)^i \frac{q^{\binom{i}{2}}}{[i]_q!} \quad (n \geq 1). \quad (1.4)$$

Here,  $[n]_q = 1 + q + \cdots + q^{n-1}$  is the  $q$ -analogue of the nonnegative integer  $n$  and  $[n]_q! = [1]_q [2]_q \cdots [n]_q$  is the  $q$ -analogue of  $n!$ .

In this paper, we shall put the  $q$ -derangement numbers in the context of a Seidel matrix as Dumont and Randrianarivony [8] did for the ordinary derangement numbers. To this end, in section 2 we introduce the notion of  $q$ -Seidel matrix. In section 3 we define a new statistic “maf” on permutations and then prove bijectively that this is a mahonian statistic. In section 4 we consider the  $q$ -Seidel matrix associated to the  $q$ -derangement numbers and give combinatorial interpretations for all of the coefficients in this matrix in terms of the new statistic “maf”. As a consequence we get a new proof of a formula of Gessel and Reutenauer [12] and of Wachs [17]. Finally we close this paper with some remarks and open questions.

We will need the following notations and results of  $q$ -calculus (see [11]). The  $q$ -binomial coefficients are defined by

$$\binom{n}{k}_q = \frac{[n]_q!}{[k]_q! [n-k]_q!} \quad (n \geq k \geq 0).$$

Define also  $(t; q)_n = (1-t)(1-qt) \cdots (1-q^{n-1}t)$  and  $(t; q)_\infty = \lim_{n \rightarrow \infty} (t; q)_n$ . Then the two  $q$ -analogues of the exponential series  $e^t = \sum_{n \geq 0} t^n/n!$  are defined by

$$e_q(t) = \sum_{n \geq 0} \frac{t^n}{[n]_q!} = \frac{1}{((1-q)t; q)_\infty}, \quad (1.5)$$

$$E_q(t) = \sum_{n \geq 0} \frac{q^{\binom{n}{2}} t^n}{[n]_q!} = (- (1-q)t; q)_\infty. \quad (1.6)$$

Notice that  $e_q(t) \cdot E_q(-t) = 1$ .

## 2. $q$ -Seidel matrices

Let us introduce the following generalization of Seidel matrix.

**Definition 1.** Given a sequence  $(a_n(x, q))$  ( $n \geq 0$ ) of elements in a commutative ring, we call the  $q$ -Seidel matrix associated to  $(a_n(x, q))$  the double sequence  $(a_n^k(x, q))$  ( $n \geq 0, k \geq 0$ ) given by the recurrence

$$\begin{cases} a_n^0(x, q) = a_n(x, q), & (n \geq 0) \\ a_n^k(x, q) = xq^n a_n^{k-1}(x, q) + a_{n+1}^{k-1}(x, q). & (k \geq 1, n \geq 0) \end{cases} \quad (2.7)$$

Moreover  $(a_n^0(x, q))$  is called the initial sequence and  $(a_0^n(x, q))$  the final sequence of the  $q$ -Seidel matrix.

**Lemma 1.** We have

$$a_n^k(x, q) = \sum_{i=0}^k (xq^n)^{k-i} \binom{k}{i}_q a_{n+i}^0(x, q). \quad (2.8)$$

**Proof:** Recall that

$$\binom{n}{k}_q = q^{n-1} \binom{n-1}{k-1}_q + \binom{n-1}{k}_q.$$

We proceed by recurrence on  $k$ . Clearly (2.8) is valid for  $k = 1$ . Suppose (2.8) is true for  $k - 1$ . We then have

$$\begin{aligned} a_n^k(x, q) &= \sum_{i=0}^{k-1} \binom{k-1}{i}_q \left( (xq^n)^{k-i} a_{n+i}^0(x, q) + (xq^{n+1})^{k-1-i} a_{n+1+i}^0(x, q) \right) \\ &= (xq^n)^k a_n^0(x, q) + \sum_{i=1}^{k-1} (xq^n)^{k-i} \binom{k-1}{i}_q a_{n+i}^0(x, q) \\ &\quad + \sum_{i=0}^{k-2} (xq^{n+1})^{k-1-i} \binom{k-1}{i}_q a_{n+1+i}^0(x, q) + a_{n+k}^0(x, q) \\ &= (xq^n)^k a_n^0(x, q) + \sum_{i=1}^{k-1} (xq^n)^{k-i} \binom{k}{i}_q a_{n+i}^0(x, q) + a_{n+k}^0(x, q). \end{aligned}$$

Thus completes the proof.  $\square$

In particular we pass from the initial sequence to the final sequence and conversely by the *Gauss inversion formula* [2, p. 96]:

$$a_0^n(x, q) = \sum_{i=0}^n x^{n-i} \binom{n}{i}_q a_i^0(x, q), \quad (2.9)$$

$$a_n^0(x, q) = \sum_{i=0}^n (-x)^{n-i} q^{\binom{n-i}{2}} \binom{n}{i}_q a_i^0(x, q). \quad (2.10)$$

Define the generating functions as follows:

$$a(t) = \sum_{n \geq 0} a_n^0(x, q) t^n, \quad \bar{a}(t) = \sum_{n \geq 0} a_0^n(x, q) t^n,$$

and

$$A(t) = \sum_{n \geq 0} a_n^0(x, q) \frac{t^n}{[n]_q!}, \quad \bar{A}(t) = \sum_{n \geq 0} a_0^n(x, q) \frac{t^n}{[n]_q!}.$$

**Proposition 2.** *The generating functions of the initial and final sequences are related by the following equations:*

$$\bar{a}(t) = \sum_{n \geq 0} a_n^0(x, q) \frac{t^n}{(xt; q)_{n+1}}; \quad (2.11)$$

$$\bar{A}(t) = e_q(xt)A(t). \quad (2.12)$$

**Proof:** Note that

$$\frac{1}{(t; q)_{n+1}} = \sum_{k=0}^{\infty} \binom{n+k}{k}_q t^k.$$

Hence

$$\begin{aligned} \sum_{n \geq 0} a_n^0(x, q) \frac{t^n}{(xt; q)_{n+1}} &= \sum_{n, k \geq 0} \binom{n+k}{k}_q a_n^0(x, q) x^k t^{n+k} \\ &= \sum_{m \geq 0} t^m \sum_{n=0}^m \binom{m}{n}_q x^{m-n} a_n^0(x, q) \\ &= \sum_{m \geq 0} a_0^m(x, q) t^m. \end{aligned}$$

By (1.5) we have

$$\begin{aligned} e_q(xt)A(t) &= \sum_{i, j \geq 0} \frac{a_i^0(x, q) t^i}{[i]_q!} \cdot \frac{x^j t^j}{[j]_q!} \\ &= \sum_{i, j \geq 0} \binom{i+j}{i}_q a_i^0(x, q) x^j \frac{t^{i+j}}{[i+j]_q!} \\ &= \sum_{n \geq 0} \left( \sum_{i=0}^n x^{n-i} \binom{n}{i}_q a_i^0(x, q) \right) \frac{t^n}{[n]_q!}, \end{aligned}$$

which completes the proof of (2.12) in view of (2.9).  $\square$

**Remark:** If  $x = q = 1$  we get the classical formulas [7, 9]:

$$\bar{a}(t) = \frac{1}{1-t} a\left(\frac{t}{1-t}\right) \quad \text{and} \quad \bar{A}(t) = e^t A(t).$$

If  $x = 0$  we have  $\bar{A}(t) = A(t)$ .

### 3. A new mahonian statistic “maf”

Let  $\mathcal{S}_n$  be the set of permutations on  $[n] = \{1, 2, \dots, n\}$ . Recall that  $i \in [n]$  is a *fixed point* of  $\sigma \in \mathcal{S}_n$  if  $\sigma(i) = i$ . Let  $\text{fix } \sigma$  denote the number of fixed points of  $\sigma$ . The permutation  $\sigma$  has a *descent* at  $i \in \{1, 2, \dots, n-1\}$  if  $\sigma(i) > \sigma(i+1)$  and we call  $i$  the *descent place* of  $\sigma$ . The *major index* of  $\sigma$ , denoted  $\text{maj } \sigma$ , is the sum of all the descent places of  $\sigma$ . Let  $\text{FIX}(\sigma) = \{i \mid \sigma(i) = i\}$  be the set of all fixed points of  $\sigma$  and  $\tilde{\sigma}$  the *restriction* of  $\sigma$  to  $\{1, 2, \dots, n\} \setminus \text{FIX}(\sigma)$ .

**Definition 2.** If  $\sigma \in \mathcal{S}_n$  with  $\text{FIX}(\sigma) = \{i_1, i_2, \dots, i_l\}$ , then the statistic “maf” is defined by

$$\text{maf } \sigma = \sum_{j=1}^l (i_j - j) + \text{maj } \tilde{\sigma}.$$

**Example 1.** Let  $\sigma = 321659487$ . Then  $\text{FIX}(\sigma) = \{2, 5, 8\}$  and  $\tilde{\sigma} = 316947$ . Hence  $\text{fix } \sigma = 3$ ,  $\text{maj } \sigma = 1 + 2 + 4 + 6 + 8 = 21$  and  $\text{maf } \sigma = (2 - 1) + (5 - 2) + (8 - 3) + (1 + 4) = 14$ .

We now show that the bivariate statistics  $(\text{fix}, \text{maf})$  and  $(\text{fix}, \text{maj})$  are equidistributed on the symmetric group  $\mathcal{S}_n$  (Corollary 7). In particular, this shows that maf is a Mahonian statistic.

Let  $\sigma = x_1 x_2 \dots x_n \in \mathcal{S}_n$ . For convenience we put  $x_0 = -\infty$  and  $x_{n+1} = +\infty$ . For  $0 \leq i \leq n$ , a pair  $(i, i+1)$  of positions is the  $j$ -th slot of  $\sigma$  provided that  $x_i \neq i$ , i.e.,  $i$  is not a fixed point of  $\sigma$  and that  $\sigma$  has  $i - j$  fixed points  $f$  such that  $f < i$ . Clearly we can insert a fixed point into the  $j$ -th slot to obtain the permutation

$$(\sigma, j) = x'_1 x'_2 \dots x'_i (i+1) x'_{i+1} \dots x'_n, \quad (3.13)$$

where  $x' = x$  if  $x \leq i$  and  $x' = x + 1$  if  $x > i$ .

More generally, if  $\sigma$  is a derangement in  $\mathcal{S}_n$  and  $(i_1, i_2, \dots, i_m)$  a sequence of integers such that  $0 \leq i_1 \leq i_2 \leq \dots \leq i_m \leq n$ , we can insert  $m$  fixed points into the derangement  $\sigma$  successively, finally obtaining

$$(\sigma, i_1, \dots, i_m) = ((\sigma, i_1, \dots, i_{m-1}), i_m).$$

Note that the fixed points of this last permutation are  $i_1 + 1, i_2 + 2, \dots, i_m + m$ .

**Example 2.** Let  $\sigma = 2143$  and  $(i_1, i_2, \dots, i_m) = (0, 1, 1, 4)$ . Then we have  $(\sigma, 0) = 13254$ ,  $(\sigma, 0, 1) = 143265$ ,  $(\sigma, 0, 1, 1) = 1534276$  and  $(\sigma, 0, 1, 1, 4) = 15342768$ .

We can of course undertake the reverse operation. That is, if a permutation  $\sigma$  in  $\mathcal{S}_{m+n}$  has  $m$  fixed points we can find a unique derangement  $\text{dp}(\sigma) \in \mathcal{S}_n$ , called (following Wachs [17]) the *derangement part* of  $\sigma$ , and a unique sequence of integers  $i_1 \leq \dots \leq i_m$ , which we call the *fixed point sequence* of  $\sigma$ , such that

$$\sigma = (\text{dp}(\sigma), i_1, \dots, i_m).$$

It is easy to see that

$$\text{maj}\sigma = \text{maj}\text{dp}(\sigma) + i_1 + \dots + i_m. \quad (3.14)$$

Consider a permutation  $\sigma$  with  $n$  slots. The  $j$ -th slot  $(i, i+1)$  of  $\sigma$  is said to be *green* if  $\text{des}(\sigma, j) = \text{des}\sigma$ , *red* if  $\text{des}(\sigma, j) = \text{des}\sigma + 1$ . We assign *values* to the green slots of  $\sigma$  from right to left, from 0 to  $g$ , and to the red slots from left to right, from  $g+1$  to  $n$ . Denote the value of the  $j$ -th slot by  $g_j$ . (When we refer to the "largest" slot, we will mean largest in terms of  $j$ .)

**Example 3.** Let  $\sigma = 2143$ . Then  $(\sigma, 0) = 13254$ ,  $(\sigma, 1) = 32154$ ,  $(\sigma, 2) = 21354$ ,  $(\sigma, 3) = 21543$ ,  $(\sigma, 4) = 21435$ . Hence slots 0, 2 and 4 are green, while 1 and 3 are red. Therefore

$$(g_0, \dots, g_n) = (2, 3, 1, 4, 0). \quad (3.15)$$

It is easy to see that every slot is either green or red. In fact, one can see that  $(i, i+1)$  is green if either  $x_{i+1} < x_i \leq i$ , or  $i < x_{i+1} < x_i$ , or  $x_i \leq i < x_{i+1}$ . So  $(i, i+1)$  is red if either  $x_{i+1} \leq i < x_i$ , or  $i < x_i < x_{i+1}$  or  $x_i < x_{i+1} \leq i$ . (Expressed in terms of cyclic intervals (cf. [13]), slot  $(i, i+1)$  is green if  $i+1 \in \llbracket x_i, x_{i+1} \rrbracket$ .)

Denote by  $d_j$  the number of descents of  $(\sigma, j)$  that lie to the right of  $x'_i$  in (3.13).

**Lemma 3.** Let  $\sigma$  be a permutation in  $\mathcal{S}_n$ . If the  $j$ -th slot  $(i, i+1)$  is green then  $\text{maj}(\sigma, j) - \text{maj}\sigma = d_j$ , if  $(i, i+1)$  is red then  $\text{maj}(\sigma, j) - \text{maj}\sigma = d_j + i$ .

**Proof:** Let  $(i, i+1)$  be a green slot. Since no new descents are formed by inserting a fixed point into the  $j$ -th slot of  $\sigma$ ,  $\text{maj}(\sigma, j) - \text{maj}\sigma$  equals the number of descents of  $\sigma$  that are displaced to the right when this fixed point is inserted. This number equals  $d_j$ . The case in which  $(i, i+1)$  is red is dealt with similarly.  $\square$

**Remark:** If  $\sigma$  is a derangement in  $\mathcal{S}_n$ , the  $j$ -th slot of  $\sigma$  is just  $(j, j+1)$  for  $0 \leq j \leq n$ .

**Lemma 4.** If  $\sigma$  is a derangement in  $\mathcal{S}_n$ , then

$$\text{maj}(\sigma, j) = \text{maj}\sigma + g_j \quad \text{for } 0 \leq j \leq n.$$

**Proof:** Let  $i$  and  $j$  be slots. It follows from Lemma 3 that if  $i$  and  $j$  are both green and  $i < j$  then  $\text{maj}(\sigma, i) \geq \text{maj}(\sigma, j)$ , while if  $i$  and  $j$  are both red and  $i < j$  then  $\text{maj}(\sigma, i) \leq \text{maj}(\sigma, j)$ . Therefore 0 is the green slot of  $\sigma$  of highest value, and if  $i$  is red and  $j$  is green we have  $\text{maj}(\sigma, i) \geq \text{maj}(\sigma, j)$ . This is because for any red slot  $i$  we have  $\text{maj}(\sigma, i) \geq \text{maj}(\sigma, 0)$  by Lemma 3. Hence, if  $m$  is the largest red slot of  $\sigma$ , i.e.,  $g_m = n$ , for any two slots  $i$  and  $j$  with  $g_i < g_j$  we have

$$\text{maj}\sigma \leq \text{maj}(\sigma, i) \leq \text{maj}(\sigma, j) \leq \text{maj}(\sigma, m).$$

It therefore suffices to show that

$$\text{maj}(\sigma, m) = \text{maj} \sigma + n.$$

Now, consider a green slot  $(i, i + 1)$ . If  $i + 1$  is a non-excedance place, i.e.,  $x_{i+1} \leq i + 1$ , then, as  $\sigma$  is a derangement,  $x_{i+1} \leq i$ . Hence  $x_{i+1} < x_i \leq i$ . Thus  $i$  is a non-excedance place. Since  $n$  is a non-excedance place and  $m + 1, m + 2, \dots, n$  are green slots, we have

$$m + 1 > x_{m+1} > \dots > x_n.$$

As the slot  $m$  is red, either  $m$  is a non-excedance place and  $m$  is a non-descent or  $m$  is an excedance place and  $m$  is a descent. In each case, inserting a fixed point into the  $m$ -th slot introduces a new descent for  $i = m + 1$  and moves  $n - (m + 1)$  descents one place further to the right. Hence

$$\text{maj}(\sigma, m) = \text{maj} \sigma + (m + 1) + (n - m - 1) = \text{maj} \sigma + n,$$

as required.  $\square$

**Remark:** Suppose that  $\sigma$  is a derangement in  $\mathcal{S}_n$  and  $0 \leq i \leq n$ . It follows from Lemmas 3 and 4 that  $d_i = g_i$  if  $i$  is green and  $d_i = g_i - i$  if  $i$  is red. If  $i$  is green then

$$\text{maj}(\sigma, i, i) = \text{maj}(\sigma, i) + g_i.$$

Hence, if  $j \leq i$ , it follows from Lemma 4 that

$$\text{maj}(\sigma, j, i) = \text{maj}(\sigma, i) + g_j.$$

On the other hand, if  $i$  is red, then

$$\text{maj}(\sigma, i, i) = \text{maj}(\sigma, i) + g_i - i.$$

Now one can easily see that, if  $k$  is the largest green slot to the left of slot  $i$ ,  $g_k = g_i - i$ . Hence, if  $j < i$ , it follows again from Lemma 4 that

$$\text{maj}(\sigma, j, i) = \text{maj}(\sigma, i) + g_j + 1.$$

We are now ready to state the key result of this section. Let  $S(\sigma, m)$  denote the set of permutations in  $\mathcal{S}_{n+m}$  with derangement part  $\sigma \in \mathcal{D}_n$ . Note that

$$S(\sigma, m) = \{(\sigma, \mathbf{i}) \mid \mathbf{i} = (i_1, \dots, i_m) \text{ and } 0 \leq i_1 \leq i_2 \leq \dots \leq i_m \leq n\}.$$

**Theorem 5.** *There is a bijection  $\Psi$  on  $S(\sigma, m)$  such that if  $\Psi(\sigma, \mathbf{i}) = (\sigma, \mathbf{j})$  then*

$$\text{maj}(\sigma, \mathbf{i}) = \text{maj}(\sigma, \mathbf{j}). \tag{3.16}$$

**Proof:** We divide the proof into two parts.

**The definition of  $\Psi$ .** We will define such a bijection  $\Psi$  by induction on  $m \geq 0$ .

First,  $\Psi$  is the identity mapping on  $S(\sigma, 0)$ . Next, we define  $\Psi$  on  $S(\sigma, 1)$  by

$$\Psi(\sigma, i) = (\sigma, g_i).$$

Then using equation (3.14) and Lemma 4 we see that  $\Psi$  satisfies equation (3.16).

Let  $m > 1$  and suppose that  $\Psi$  has been defined on  $S(\sigma, k)$  for  $0 \leq k \leq m-1$ . Consider  $\tau = (\sigma, i_1, \dots, i_m)$ . Suppose that the  $i_m$ -th slot of  $(\sigma, i_1, \dots, i_{m-1})$  is green. Then, if

$$\Psi(\sigma, i_1, \dots, i_{m-1}) = (\sigma, j_2, \dots, j_m),$$

we define

$$\Psi(\tau) = (\sigma, g_{i_m}, j_2, \dots, j_m).$$

Suppose that the  $i_m$ -th slot of  $(\sigma, i_1, \dots, i_{m-1})$  is red. Then the slots  $i_1, \dots, i_m$  cannot be all the same. Let  $k$  be the smallest positive integer such that  $i_{m-k} < i_m$ . Thus  $i_{m-k} < i_{m-k+1} = \dots = i_m$ . Then, if

$$\Psi(\sigma, i_1, \dots, i_{m-k}) = (\sigma, j_1, \dots, j_{m-k}),$$

we define

$$\Psi(\tau) = (\sigma, \underbrace{g_{i_m - i_m}, \dots, g_{i_m - i_m}}_{k-1 \text{ terms}}, j_1 + 1, \dots, j_{m-k} + 1, g_{i_m}).$$

The following lemma is easily proved by induction.

**Lemma 6.** *Let  $\tau = (\sigma, i_1, \dots, i_m)$  and  $\Psi(\tau) = (\sigma, j_1, \dots, j_m)$ .*

*Suppose that at least one of the slots  $i_1, \dots, i_m$  is either green or is repeated. Let  $i_l$  be the largest such slot. If  $i_l$  is green then  $j_1 = g_{i_l}$ . If  $i_l$  is red and is repeated then  $j_1 = g_{i_l} - i_l$ .*

*If on the other hand all of the slots  $i_1, \dots, i_m$  are red and are distinct, then  $j_1 = g_{i_1}$ .*

*Suppose that at least one of the slots  $i_1, \dots, i_m$  is red. If  $i_l$  is the largest red slot then  $j_m = g_{i_l}$ .*

*If on the other hand all of these slots are green then  $j_m = g_{i_1}$ .*

It follows from this lemma that  $j_1, \dots, j_m$  as defined above are in ascending order.

We now show by induction on  $m$  that  $\Psi$  satisfies equation (3.16).

If  $i_m$  is green, then using Lemma 4 we have

$$\begin{aligned} \text{maj}(\sigma, i_1, \dots, i_m) &= \text{maj}(\sigma, i_1, \dots, i_{m-1}) + g_{i_m} \\ &= \text{maf}(\sigma, j_2, \dots, j_m) + g_{i_m} \\ &= \text{maf}(\sigma, g_{i_m}, j_2, \dots, j_m). \end{aligned}$$

If  $i_m$  is red, let  $k$  be the smallest positive integer such that  $i_{m-k} < i_m$ , then

$$\begin{aligned} \text{maj}(\sigma, i_1, \dots, i_m) &= \text{maj}(\sigma, i_1, \dots, i_{m-k}) + (m-k) + (k-1)(g_{i_m} - i_m) + g_{i_m} \\ &= \text{maf}(\sigma, j_1, \dots, j_{m-k}) + (m-k) + (k-1)(g_{i_m} - i_m) + g_{i_m} \\ &= \text{maf}(\sigma, \underbrace{g_{i_m - i_m}, \dots, g_{i_m - i_m}}_{k-1 \text{ terms}}, j_1 + 1, \dots, j_{m-k} + 1, g_{i_m}). \end{aligned}$$



This is because inserting the first fixed point  $i_m$  into  $(\sigma, i_1, \dots, i_{m-k})$  adds a descent and increases  $\text{maj}$  by  $g_{i_m} + (m - k)$ . Inserting each of the remaining fixed points  $i_m$  has the same affect as inserting a fixed point into a green slot of value  $g_{i_m} - i_m$ .

**$\Psi$  is a bijection.** It remains to show that  $\Psi$  is a bijection on  $S(\sigma, m)$ . It suffices to show that  $\Psi$  is an injection.

We use induction on  $m$ . The result is clearly true for  $m = 0$  and  $m = 1$ .

Let  $\tau = (\sigma, i_1, \dots, i_m)$  and  $\Psi(\tau) = (\sigma, j_1, \dots, j_m)$ . Suppose that  $\Psi(\tau) = \Psi(\tau')$ , where  $\tau' = (\sigma, i'_1, \dots, i'_m)$ .

If both  $i_m$  and  $i'_m$  are green or red then it is easy to show using the induction hypothesis that  $\tau = \tau'$ . So suppose that  $i_m$  is green and  $i'_m$  is red. Thus  $j_1 = g_{i_m}$ ,  $j_m = g_{i'_m}$ .

Suppose that  $i_1, \dots, i_m$  are all green. Then  $j_m = g_{i_1}$ . Hence  $i_1 = i'_m$ , contradiction.

Let  $i_u$  be the largest red slot amongst  $i_1, \dots, i_m$ . Then  $j_m = g_{i_u}$ . Hence  $i'_m = i_u < i_m$ .

*Case 1:* Suppose that one of the slots  $i'_1, \dots, i'_m$  is either green or is repeated. Let  $i'_v$  be the largest such slot. If  $i'_v$  is green, then

$$\Psi(\tau') = (\sigma, g_{i'_v} + (m - v), \dots, g_{i'_m}).$$

Hence  $j_1 = g_{i_m} = g_{i'_v} + (m - r)$ . Since  $i_m$  and  $i'_v$  are both green, this means that  $i_m \leq i'_v < i'_m$ , contradiction.

If  $i'_v$  is red, then

$$\Psi(\tau') = (\sigma, g_{i'_v} - i'_v + (m - v), \dots, g_{i'_m}).$$

Hence  $j_1 = g_{i_m} = g_{i'_v} - i'_v + (m - r)$ . But  $g_{i'_v} - i'_v$  is the value of the largest green slot  $i_w$  less than  $i'_v$ . As  $i_m$  is green this means that  $i_m \leq i_w < i'_v \leq i'_m$ , contradiction.

*Case 2:* Suppose that all of the slots  $i'_1, \dots, i'_m$  are red and distinct. Then

$$\Psi(\tau') = (\sigma, g_{i'_1} + (m - 1), \dots, i'_m).$$

Hence  $j_1 = g_{i_m} = g_{i'_1} + (m - 1) > g_{i'_{-1}}$ . This is a contradiction, since  $i_m$  is green and  $i'_1$  is red.  $\square$

**Example 4.** Let  $\sigma = 2143$  and consider  $(\sigma, 0, 1, 1, 4) \in S(\sigma, 4)$ . Then the values of the slots of  $\sigma$  have been calculated in (3.15). The bijection  $\Psi$  goes as follows: since slot 0 is green in  $\sigma$  we have

$$\Psi(\sigma, 0) = (\sigma, g_0) = (\sigma, 2);$$

since slot 1 is red we have

$$\Psi(\sigma, 0, 1) = (\sigma, 2 + 1, g_1) = (\sigma, 3, 3);$$

again, since slot 1 is red we have

$$\Psi(\sigma, 0, 1, 1) = (\sigma, g_1 - 1, 2 + 1, g_1) = (\sigma, 2, 3, 3);$$

Finally, since slot 4 is green we obtain

$$\Psi(\sigma, 0, 1, 1, 4) = (\sigma, g_4, 2, 3, 3) = (\sigma, 0, 2, 3, 3) \in S(\sigma, 4).$$

Let  $\tau = (\sigma, 0, 1, 1, 4)$  and  $\tau' = (\sigma, 0, 2, 3, 3)$ . Then  $\tau = 15342768$  and  $\tau' = 13248675$ . It is easy to see that  $\text{maj } \tau = 12$  and  $\text{maf } \tau' = 12$ . Hence we have checked equation (3.16).

Using theorem 5, we obtain the following result.

**Corollary 7.** (a) *There is a bijection  $\phi: S_n \rightarrow S_n$  such that for any  $\sigma \in S_n$  we have*

$$(\text{fix}, \text{maf})\sigma = (\text{fix}, \text{maj})\phi(\sigma).$$

(b) *The bi-statistic  $(\text{fix}, \text{maf})$  is equidistributed with the bi-statistic  $(\text{fix}, \text{maj})$  on the symmetric group  $S_n$ .*

The following result was first proved by Wachs [17, corollary 3].

**Corollary 8.** *Let  $\sigma$  be a derangement in  $S_n$  and  $m \geq 0$ . We have*

$$\sum_{\pi \in S(\sigma, m)} q^{\text{maj}\pi} = q^{\text{maj}\sigma} \binom{m+n}{n}_q.$$

**Proof:** By theorem 5 we have

$$\begin{aligned} \sum_{\pi \in S(\sigma, m)} q^{\text{maj}\pi} &= \sum_{\pi \in S(\sigma, m)} q^{\text{maf}\pi} \\ &= q^{\text{maj}\sigma} \sum_{0 \leq i_1 \leq \dots \leq i_m \leq n} q^{i_1 + i_2 + \dots + i_m} \\ &= q^{\text{maj}\sigma} \binom{m+n}{n}_q. \end{aligned}$$

The last line follows from a well-known result [1, p. 33].  $\square$

#### 4. $q$ -derangement matrices

We first prove the following result.

**Proposition 9.** *Let  $(a_n^k(x, q))$  be a  $q$ -Seidel matrix. Then the following three conditions are equivalent:*

$$a_n^0(x, q) = [n]_q! \sum_{i=0}^n (-1)^i \frac{q^{\binom{i}{2}}}{[i]_q!}, \quad (4.17)$$

$$a_0^n(x, q) = [n]_q! \left( 1 + \sum_{i=1}^n \frac{(x-1)(x-q)\cdots(x-q^{i-1})}{[i]_q!} \right), \quad (4.18)$$

$$a_0^n(1, q) = [n]_q! \quad \text{and} \quad a_n^0(x, q) \text{ is independent of } x. \quad (4.19)$$

**Proof:** By the  $q$ -binomial formula [11, p.7]

$$\sum_{n=0}^{\infty} \frac{(a; q)_n}{(q; q)_n} t^n = \frac{(at; q)_{\infty}}{(t; q)_{\infty}},$$

we have in view of (1.5) and (1.6),

$$1 + \sum_{n=1}^{\infty} \frac{(x-1)(x-q)\cdots(x-q^{n-1})}{[n]_q!} t^n = e_q(xt) E_q(-t).$$

Therefore the generating functions of (4.17), (4.18) and (4.19) are respectively the following:

$$A(t) = \sum_{n \geq 0} a_n^0(x, q) \frac{t^n}{[n]_q!} = \frac{E_q(-t)}{1-t}, \quad (4.20)$$

$$\bar{A}(t) = \sum_{n \geq 0} a_0^n(x, q) \frac{t^n}{[n]_q!} = \frac{e_q(xt)E_q(-t)}{1-t}, \quad (4.21)$$

$$\bar{A}(t)|_{x=1} = \sum_{n \geq 0} a_0^n(1, q) \frac{t^n}{[n]_q!} = \frac{1}{1-t}. \quad (4.22)$$

So, it suffices to prove that the equivalence of (4.20), (4.21) and (4.22). Indeed,

(4.20)  $\iff$  (4.21): this follows from proposition 2;

(4.21)  $\implies$  (4.22): this is obvious;

(4.22)  $\implies$  (4.20): since  $A(t)$  is independent of  $x$ , equation (4.20) follows then from (2.12) by setting  $x = 1$ .  $\square$

**Definition 3.** A  $q$ -derangement matrix is the  $q$ -Seidel matrix satisfying any of the three conditions of proposition 8.

If  $x = 1$ , then  $a_0^n(x, q) = [n]_q!$  and the  $q$ -derangement matrix is as follows :

$k \setminus n$	0	1	2	3	4
0	1	0	$q$	$q + q^2$	$\binom{q+2q^2+2q^3}{+2q^4+q^5+q^6}$
1	1	$q$	$q + q^2 + q^3$	$\binom{q+2q^2+2q^3}{+3q^4+2q^5+q^6}$	
2	$1 + q$	$q + 2q^2 + q^3$	$\binom{q+2q^2+3q^3}{+4q^4+3q^5+q^6}$		
3	$[3]_q!$	$\binom{q+3q^2+5q^3}{+5q^4+3q^5+q^6}$			
4	$[4]_q!$				

$$(a_n^k(1, q))$$

Denote by  $\mathcal{S}_n^k$  the set of permutations on  $[n + k]$  of which all the fixed points are included in  $\{n + 1, n + 2, \dots, n + k\}$ . In particular  $\mathcal{S}_n^0$  is the set of permutations without fixed points on  $[n]$  and  $\mathcal{S}_0^k$  the set of all permutations on  $[k]$ . The following result generalizes a result of Dumont and Randrianarivony [8].

**Theorem 10.** The coefficients  $a_n^k(x, q)$  ( $n, k \geq 0$ ) in a  $q$ -derangement matrix have the following combinatorial interpretation:

$$a_n^k(x, q) = \sum_{\sigma \in \mathcal{S}_n^k} x^{\text{fix} \sigma} q^{\text{maf} \sigma}. \quad (4.23)$$

**Proof:** Notice that  $\mathcal{S}_{n+1}^{k-1} \subset \mathcal{S}_n^k$ . Set

$$\Delta_n^k = \mathcal{S}_n^k \setminus \mathcal{S}_{n+1}^{k-1} = \{\sigma \in \mathcal{S}_n^k \mid \sigma(n+1) = n+1\}.$$

We construct a bijection  $\varphi : \Delta_n^k \rightarrow \mathcal{S}_n^{k-1}$  such that for all  $\sigma \in \Delta_n^k$ ,

$$\begin{aligned} \text{maf } \sigma &= n + \text{maf}(\varphi(\sigma)), \\ \text{fix } \sigma &= 1 + \text{fix}(\varphi(\sigma)). \end{aligned}$$

Indeed, if  $\sigma \in \Delta_n^k$  we define  $\varphi(\sigma)$  as the word obtained from  $\sigma$  by deleting  $n+1$  and reduce all the values strictly bigger than  $n+1$ . It is readily verified that  $\varphi$  is the desired bijection. Therefore

$$\sum_{\sigma \in \mathcal{S}_n^k} x^{\text{fix } \sigma} q^{\text{maf } \sigma} = xq^n \sum_{\sigma \in \mathcal{S}_n^{k-1}} x^{\text{fix } \sigma} q^{\text{maf } \sigma} + \sum_{\sigma \in \mathcal{S}_{n+1}^{k-1}} x^{\text{fix } \sigma} q^{\text{maf } \sigma}, \quad (4.24)$$

which is the recurrence (2.7). So it remains to check the initial condition. Now  $\mathcal{S}_0^n = \mathcal{S}_n$  and it is well-known [14] that  $\sum_{\sigma \in \mathcal{S}_n} q^{\text{maj } \sigma} = [n]_q!$ , so it follows from corollary 7 that

$$a_0^n(1, q) = \sum_{\sigma \in \mathcal{S}_n} q^{\text{maf } \sigma} = \sum_{\sigma \in \mathcal{S}_n} q^{\text{maj } \sigma} = [n]_q!.$$

The theorem follows then from proposition 9, since  $a_n^0(x, q)$  is clearly independent of  $x$ .  $\square$

**Remark:** Since  $(\text{fix}, \text{maf})$  and  $(\text{fix}, \text{maj})$  are not equidistributed on  $\mathcal{S}_1^2$  we cannot replace  $\text{maf}$  by  $\text{maj}$  in the above theorem.

From Corollary 7, proposition 9 and theorem 10 we derive the following result.

**Corollary 11.** *The final sequence of the  $q$ -derangement matrix has the following interpretation:*

$$a_0^n(x, q) = \sum_{\sigma \in \mathcal{S}_n} x^{\text{fix } \sigma} q^{\text{maf } \sigma} \quad (4.25)$$

$$= \sum_{\sigma \in \mathcal{S}_n} x^{\text{fix } \sigma} q^{\text{maj } \sigma} \quad (4.26)$$

$$= [n]_q! \left( 1 + \sum_{i=1}^n \frac{(x-1)(x-q) \cdots (x-q^{i-1})}{[i]_q!} \right). \quad (4.27)$$

Note that the last equation has been obtained by Gessel and Reutenauer [12] and by Wachs [17] in the special  $x=0$  case using different methods.

## 5. An open problem about $q$ -succession numbers

Let  $\sigma$  be a permutation in  $\mathcal{S}_n$ . For convenience put  $\sigma(0) = 0$ . We say that an element  $p$  (with  $1 \leq p \leq n$ ) is a *succession* of  $\sigma$  if  $\sigma(p) = \sigma(p-1) + 1$ . The  $p$  is called the *succession position*, while  $\sigma(p)$  is called the *succession value*. Let  $\text{SUC}(\sigma)$  be the set of succession values of  $\sigma$  and let  $\text{suc } \sigma$  be the number of successions of  $\sigma$ . For example, if

$$\sigma = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 \\ 1 & 4 & 3 & 8 & 9 & 5 & 6 & 7 & 2 \end{pmatrix},$$

then  $\text{SUC}(\sigma) = \{1, 9, 6, 7\}$  and  $\text{suc } \sigma = 4$ .

We use a variant of Foata's first fundamental transformation [10] to show that the statistics  $\text{fix}$  and  $\text{suc}$  are equidistributed on  $\mathcal{S}_n$ .

Given a permutation  $\sigma = \sigma(1)\sigma(2)\cdots\sigma(n) \in \mathcal{S}_n$  we set  $\sigma^d = \sigma(2)\cdots\sigma(n)\sigma(1)$ . We call the *standard form* of the factorization into cycles of  $\sigma$  the unique writing  $\bar{\sigma}$  such that in each cycle  $(a, \sigma(a), \dots, \sigma^l(a))$  the maximum  $\sigma^l(a)$  is in the last position and the cycles of  $\sigma$  are decreasingly ordered according to their maxima. (Note that this is *not* the usual definition of standard form.) We define  $\varphi(\sigma)$  as the permutation obtained by erasing the parentheses in the standard form of  $\bar{\sigma}^d$ .

The following lemma is easy to verify.

**Lemma 12.** *The mapping  $\varphi$  is a bijection on  $\mathcal{S}_n$  such that for all  $\sigma \in \mathcal{S}_n$ ,  $\text{FIX}(\sigma) = \text{SUC}(\varphi(\sigma))$  and  $\text{fix } \sigma = \text{suc } \varphi(\sigma)$ . Hence the statistics  $\text{fix}$  and  $\text{suc}$  are equidistributed on  $\mathcal{S}_n$ .*

For example, if  $\sigma = 142836759 \in \mathcal{S}_9$ , then

$$\sigma^d = 428367591 \quad \text{and} \quad \bar{\sigma}^d = (14389)(567)(2).$$

Erasing the parentheses we obtain the permutation  $\varphi(\sigma) = 143895672$ . We have

$$\text{FIX}(\sigma) = \text{SUC}(\varphi(\sigma)) = \{1, 6, 7, 9\}.$$

Define the statistic

$$\text{suc}' \sigma = \begin{cases} \text{suc } \sigma, & \text{if } \sigma(1) \neq 1, \\ \text{suc } \sigma - 1, & \text{if } \sigma(1) = 1; \end{cases}$$

and let

$$F_n(x) = \sum_{\sigma \in \mathcal{S}_n} x^{\text{fix } \sigma}, \quad S_n(x) = \sum_{\sigma \in \mathcal{S}_n} x^{\text{suc}' \sigma}.$$

Then, using lemma 12, we obtain a bijective proof of the following known results (See [3, 15]).

**Proposition 13.** *We have*

$$S_{n+1}(x) = F_{n+1}(x) + (1-x)F_n(x), \tag{5.28}$$

and in particular

$$S_{n+1}(0) = d_{n+1} + d_n. \tag{5.29}$$

Setting  $q = 1$  in (4.20) we see that

$$\sum_{n \geq 0} F_n(x) \frac{t^n}{n!} = \frac{e^{(x-1)t}}{1-t}. \tag{5.30}$$

Hence, from equation (5.28), we have

$$\sum_{n \geq 0} S_n(x) \frac{t^n}{n!} = \frac{e^{(x-1)t}}{1-t} + (1-x) \sum_{n \geq 1} F_{n-1}(x) \frac{t^n}{n!}, \tag{5.31}$$

in which by convention  $S_0(x) = F_0(x) = 1$ . Thus

$$\sum_{n \geq 0} S_n(x) \frac{t^n}{n!} = \frac{e^{(x-1)t}}{1-t} + (1-x) \int_0^t \frac{e^{(x-1)z}}{1-z} dz. \quad (5.32)$$

Let  $\mathcal{L}$  be the formal Laplace transformation on the ring of formal power series, that is,  $\mathcal{L}(\sum a_n x^n / n!) = \sum a_n x^n$ . Then

$$\sum_{n \geq 0} F_n(x) t^n = \mathcal{L} \left( \frac{e^{(x-1)t}}{1-t} \right) = \sum_{n \geq 0} \frac{n! t^n}{[1 - (x-1)t]^{n+1}}. \quad (5.33)$$

Therefore

$$\begin{aligned} \sum_{n \geq 0} S_n(x) t^n &= \sum_{n \geq 0} F_n(x) t^n + (1-x) \sum_{n \geq 0} F_n(x) t^{n+1} \\ &= [1 - (x-1)t] \sum_{n \geq 0} F_n(x) t^n \\ &= \sum_{n \geq 0} \frac{n! t^n}{[1 - (x-1)t]^n}. \end{aligned} \quad (5.34)$$

In the case of  $q = 1$ , using lemma 11, we can restate theorem 9 in terms of successions. Unfortunately, since the mapping  $\phi$  does not keep track of the maj statistic, we do not have a full interpretation in the last model.

The distribution of our statistics on  $\mathcal{S}_3$  is as follows:

$\sigma \backslash \text{stat}$	maf	maj	suc	fix
1 2 3	0	0	3	3
1 3 2	1	2	1	1
2 1 3	3	1	0	1
2 3 1	2	2	1	0
3 1 2	1	1	1	0
3 2 1	2	3	0	1

**Statistic distributions on  $\mathcal{S}_3$**

Finally we record two open problems related to our work.

- 1) Find a mahonian statistic “mag” such that  $(\text{suc}, \text{mag})$  is equidistributed with  $(\text{fix}, \text{maj})$  on the symmetric group  $\mathcal{S}_n$ .
- 2) Generalize the statistic “maf” on permutations to *words* as in [5, 13] for other mahonian statistics.

## References

1. ANDREWS (G.). *The Theory of Partitions*, Addison-Wesley Publishing Company, 1976.
2. AIGNER (M.). *Combinatorial Theory*, Springer-Verlag, Berlin, 1979.

3. CHEN (W.). *The skew, relative, and classical derangements*, Disc. Math., **160** (1996), 235-239.
4. CHEN (W. Y. C.) and ROTA (G.-C.).  *$q$ -Analogues of the inclusion-exclusion principle and permutations with restricted position*, Disc. Math., **104** (1992), 7-22
5. CLARKE (R.) and FOATA (D.). *Eulerian Calculus, II: An Extension of Han's Fundamental Transformation*, Europ. J. Combinatorics, **16** (1995), 221-252.
6. DÉARMÉNIEN (J.) and WACHS (M. L.). *Descent classes of permutations with a Given Number of Fixed Points*, J. Combin. Theory, Ser. A. **64** (1993), 311-328.
7. DUMONT (D.). *Matrices d'Euler-Seidel*, Séminaire Lotharingien de Combinatoire, **B05c** (1981), <http://cartan.u-strasbg.fr/~slc>.
8. DUMONT (D.) et RANDRIANARIVONY (A.). *Dérangements et nombres de Genocchi*, Disc. Math., **132** (1994), 37-49.
9. DUMONT (D.) and VIENNOT (G.). *A combinatorial interpretation of the Seidel generation of Genocchi numbers*, Ann. Disc. Math., **6** (1980), 77-87.
10. FOATA (D.). *Rearrangements of words*, in: Combinatorics on words, Encyclopedia of Mathematics, Vol. 17, ed., M. Lothaire, Addison-Wesley, Reading, MA, 1981, pp. 184-212.
11. GASPER (G.) and RAHMAN (M.). *Basic Hypergeometric Series*, Cambridge University Press, 1990.
12. GESSEL (I.) and REUTENAUER (C.). *Counting Permutations with Given Cycle Structure and Descent Set*, J. Combin. Theory Ser. A, **64** (1993), 189-215.
13. HAN (G-N.). *Une transformation fondamentale sur les réarrangements de mots*, Advances in Math., **105** (1994), 26-41.
14. MACMAHON (P. A.). *The indices of permutations and the derivation therefrom of functions of a single variable associated with the permutations of any assemblage of objects*, Amer. J. Math., **35** (1913), 314-321.
15. ROSELLE (D. P.). *Permutations by number of rises and successions*, Proc. Amer. Math. Soc., **19** (1968), 8-16.
16. STANLEY (R.). *Enumerative Combinatorics I*, Cambridge University Press, 1997.
17. WACHS (M.). *On  $q$ -derangement numbers*, Proc. Amer. Math. Soc., **106** (1989), 273-278.