

DIFFERENCE OPERATORS FOR PARTITIONS UNDER THE LITTLEWOOD DECOMPOSITION

PAUL-OLIVIER DEHAYE, GUO-NIU HAN, AND HUAN XIONG*

ABSTRACT. The concept of t -difference operator for functions of partitions is introduced to prove a generalization of Stanley's theorem on polynomiality of Plancherel averages of symmetric functions related to contents and hook lengths. Our extension uses a generalization of the notion of Plancherel measure, based on walks in the Young lattice with steps given by the addition of t -hooks. It is well-known that the hook lengths of multiples of t can be characterized by the Littlewood decomposition. Our study gives some further information on the contents and hook lengths of other congruence classes modulo t .

1. INTRODUCTION

1.1. Goal. The purpose of this paper is to compute explicit averages over partitions of some statistics of their invariants, such as contents and hook lengths, and establish polynomiality of these averages in the size of the partitions considered. In contrast with previous results, we restrict our attention to subsets of partitions classified by the divisibility properties of those invariants, and we modify the averaging measures accordingly. Our main result is stated in Theorem 1.3.

1.2. Basic definitions. A *partition* is a finite weakly decreasing sequence of positive integers $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_r)$. The *size* of the partition λ is defined by the integer $|\lambda| = \sum_{1 \leq i \leq r} \lambda_i$. A partition λ could be represented by its *Young diagram*, which is a collection of boxes arranged in left-justified rows with λ_i boxes in the i -th row. The *content* of a box $\square = (i, j)$ in a partition λ is defined by $c_{\square} = j - i$. With each (i, j) -box is associated its *hook length*, denoted by $h_{(i,j)}$, which is the number of boxes exactly to the right, or exactly below, including the box itself. The hook length multi-set of λ is denoted by $\mathcal{H}(\lambda)$. In Figure 1, the Young diagram and hook lengths of the partition $(6, 3, 2, 2)$ are represented. For example, $h_{(1,2)} = 8$ and $h_{(3,1)} = 3$.

Let t be a positive integer. A partition is called a *t -core partition* if none of its hook lengths is divisible by t . For example, from Figure 1 we can see that $\lambda = (6, 3, 2, 2)$ is a 7-core partition. Let

$$\mathcal{H}_t(\lambda) = \{h \in \mathcal{H}(\lambda) : h \equiv 0 \pmod{t}\}$$

be the multi-set of hook lengths of multiples of t .

2010 *Mathematics Subject Classification.* 05A15, 05A17, 05A19, 05E05, 05E10, 11P81.

Key words and phrases. Littlewood decomposition, partition, Plancherel measure, Difference operator.

* Huan Xiong is the corresponding author.

9	8	5	3	2	1
5	4	1			
3	2				
2	1				

FIGURE 1. The Young diagram of the partition $(6, 3, 2, 2)$ and the hook lengths of corresponding boxes.

1.3. 01-sequences, Littlewood decomposition and Plancherel averages.

Each partition λ can be associated with its *01-sequence* $z(\lambda) = (z_{\lambda,i})_{i \in \mathbb{Z}} = (z_i)_{i \in \mathbb{Z}}$ as follows. First, consider the edges on the boundary of λ , starting at the bottom and ending to the right. Label the vertical (resp. horizontal) edges 0 (resp. 1). We obtain a bi-infinite 01-sequence $(z_i)_{i \in \mathbb{Z}}$ beginning with infinitely many 0's and ending with infinitely many 1's. The index 0 of the bi-infinite 01-sequence is defined by the condition $\#\{z_i = 0 : i \geq 0\} = \#\{z_i = 1 : i < 0\}$. Every partition and its 01-sequence are uniquely determined by each other (see Figure 2 for a running example of the constructions presented in this section).




FIGURE 2. This is our running example for the 01-sequence and the Littlewood decomposition of a partition. The Young diagram of the partition $\lambda = (18, 7, 6)$ is drawn, together with its 01-sequence $\cdots 0111|111010111111111101 \cdots$ (the $|$ indicates the unique location where the number of 1's to the left equals the number of 0's to the right, which is between index -1 and 0 in the 01-sequence; visually this corresponds to the main diagonal in the Young diagram). Notice the mnemonic “RGB” for the colourings, starting after the $|$ mark. The grey shapes indicate a sequence of 3-hooks that can be removed from λ , to finally obtain the 3-core $(3, 1)$ of λ . One could start with the yellow shape for instance, and then make various choices for the order of the removals. The process always ends with the same 3-core $(3, 1)$, irrespective of the order of removals. Quotients will be $\cdots 01|101111 \cdots$ (partition (2)), $\cdots 0|111111 \cdots$ (empty partition \emptyset) and $\cdots 011|011101 \cdots$ (partition $(5, 2)$).

The 01-sequences contain a lot of information about partitions (see [2, 14]). By the construction of 01-sequences the following lemma is obvious.

Lemma 1.1. *For a partition λ and its 01-sequence $z(\lambda) = (z_i)_{i \in \mathbb{Z}}$, we have*

$$|\lambda| = \#\{(i, j) : i < j, z_i = 1, z_j = 0\}.$$

Actually, each box in λ is uniquely determined by such a pair (i, j) .¹ The hook length of such box is $j - i$. Also, for an edge labeled by $z_i = 0$, the box to the left has content i ; for an edge labeled by $z_i = 1$, the box above has content $i + 1$.

By Lemma 1.1 a partition λ is a t -core partition if and only if there are no $i \in \mathbb{Z}$ satisfying $z_i = 1$, $z_{i+t} = 0$ in the 01-sequence $z(\lambda)$. If $z_i = 1$, $z_{i+t} = 0$ (resp. $z_i = 0$, $z_{i+t} = 1$) in a 01-sequence $z(\lambda) = (z_j)_{j \in \mathbb{Z}}$, by exchanging the values of z_i and z_{i+t} we obtain a new 01-sequence and thus a new partition. We say that this new partition is obtained by removing (resp. adding) a t -hook from (resp. to) λ .

Starting with a partition λ , we can always remove several t -hooks until no more t -hooks could be removed. The final partition we get by this operation is called the t -core of λ and denoted by $\lambda_{t\text{-core}}$, which is actually independent of the ordering of removing t -hooks from λ (see [2, 14]). For $0 \leq i \leq t - 1$, the i -th t -quotient of the partition λ is defined to be the partition associated with the 01-sequence obtained by taking the subsequence $(z_{tj+i})_{j \in \mathbb{Z}}$ of $z(\lambda)$, which is denoted by λ^i . For each positive integer t , we actually build a map

$$\{\text{partitions}\} \rightarrow \{t\text{-core partitions}\} \times \{\text{partitions}\}^t$$

which sends a partition λ to $(\lambda_{t\text{-core}}; \lambda^0, \lambda^1, \dots, \lambda^{t-1})$. This map is a bijection, called the *Littlewood decomposition of partitions at t* [16]. In this paper, we always set that a partition λ is identical with its image under the Littlewood decomposition, i.e., we always write $\lambda = (\lambda_{t\text{-core}}; \lambda^0, \lambda^1, \dots, \lambda^{t-1})$. The Littlewood decomposition has the following two fundamental properties (see [16]):

$$|\lambda| = |\lambda_{t\text{-core}}| + t(|\lambda^0| + |\lambda^1| + \dots + |\lambda^{t-1}|)$$

and

$$\{h/t \mid h \in \mathcal{H}_t(\lambda)\} = \mathcal{H}(\lambda^0) \cup \mathcal{H}(\lambda^1) \cup \dots \cup \mathcal{H}(\lambda^{t-1}).$$

Let f_λ (resp. $f_{\lambda/\mu}$) be the number of standard Young tableaux of shape λ (resp. λ/μ) and $H_\lambda = \prod_{\square \in \lambda} h_\square$ be the product of all hook lengths of boxes in λ . For convenience we set $f_\emptyset = 1$ and $H_\emptyset = 1$ for the empty partition \emptyset . It is well known that (see [5, 8, 15, 24])

$$(1.1) \quad f_\lambda = \frac{|\lambda|!}{H_\lambda} \quad \text{and} \quad \frac{1}{n!} \sum_{|\lambda|=n} f_\lambda^2 = 1.$$

The latter identity defines a measure on partitions of size n , called the Plancherel measure. Alternatively, we can see f_λ as counting “walks” among Young diagrams up to the shape λ (so-called *hook walks*, see [6, 7]).

1.4. Polynomiality of Plancherel averages and modified hook walks. Expanding on a formula of Nekrasov and Okounkov on hook lengths [17], the second author conjectured [9] that

$$P(n) = \frac{1}{n!} \sum_{|\lambda|=n} f_\lambda^2 \sum_{\square \in \lambda} h_\square^{2k}$$

is always a polynomial of n for any $k \in \mathbb{N}$. This conjecture was generalized and proved by Stanley [22] (see also [1, 4, 10, 20]), and later generalized in [11, 12, 13]. Let Q be a symmetric function in infinitely many variables and E be a finite set

¹ Notice that here (i, j) is not the coordinate of the box in λ .

with n elements. The symbol $Q(x : x \in E)$ means that n of the variables are substituted by x for $x \in E$, and all other variables by 0.

Theorem 1.2 (Stanley [22]). *Let Q be a symmetric function in infinitely many variables. Then*

$$(1.2) \quad P(n) = \frac{1}{n!} \sum_{|\lambda|=n} f_\lambda^2 Q(c_\square : \square \in \lambda)$$

and

$$(1.3) \quad P^*(n) = \frac{1}{n!} \sum_{|\lambda|=n} f_\lambda^2 Q(h_\square^2 : \square \in \lambda)$$

are polynomials in n .

Olshanski [19] also proved the content case of Theorem 1.2.

Let λ and μ be two partitions. We say that $\lambda \geq_t \mu$ if λ could be obtained by adding some t -hooks to μ . Then, the set of all partitions becomes a partially ordered set under this relation. If $\lambda \geq_t \mu$, define $F_{\lambda/\mu}$ to be the number of ways of removing t -hooks:

$$(1.4) \quad F_{\mu/\mu} := 1 \quad \text{and} \quad F_{\lambda/\mu} := \sum_{\substack{\lambda \geq_t \lambda^- \geq_t \mu \\ |\lambda/\lambda^-|=t}} F_{\lambda^-/\mu} \quad (\text{for } \lambda \neq \mu).$$

It is easy to see that $F_{\lambda/\mu} = \#\{(P_0, P_1, \dots, P_{t-1}) : P_i \text{ is a Young tableau of shape } \lambda^i/\mu^i, \text{ the union of entries in } P_0, P_1, \dots, P_{t-1} \text{ are } 1, 2, \dots, \sum_{i=0}^{t-1} |\lambda^i/\mu^i|\}$. Hence,

$$F_{\lambda/\mu} = \left(|\lambda^0/\mu^0|, |\lambda^1/\mu^1|, \dots, |\lambda^{t-1}/\mu^{t-1}| \right) \prod_{i=0}^{t-1} f_{\lambda^i/\mu^i}.$$

These $F_{\lambda/\mu}$ can be seen as counting hook walks distributed among the quotient partitions of λ . Let

$$(1.5) \quad F_\lambda := F_{\lambda/\lambda_{t\text{-core}}} = \left(|\lambda^0|, |\lambda^1|, \dots, |\lambda^{t-1}| \right) \prod_{i=0}^{t-1} f_{\lambda^i} = n! t^n G_\lambda,$$

where $n = |\lambda^0| + |\lambda^1| + \dots + |\lambda^{t-1}|$ and

$$G_\lambda := \frac{1}{\prod_{h \in \mathcal{H}_t(\lambda)} h}.$$

When $t = 1$, we have $F_\lambda = f_\lambda$ and $G_\lambda = 1/H_\lambda$. Also, when λ is a t -core partition, we have $F_\lambda = G_\lambda = 1$.

1.5. Difference operators and main result. Let g be a function of partitions and λ a partition. The difference operator D for partitions was defined in [11] as

$$Dg(\lambda) = \sum_{|\lambda^+/\lambda|=1} g(\lambda^+) - g(\lambda).$$

In this paper, we introduce a generalization of D . For every integer t , let

$$(1.6) \quad D_t g(\lambda) = \sum_{\substack{\lambda^+ \geq_t \lambda \\ |\lambda^+/\lambda|=t}} g(\lambda^+) - g(\lambda).$$

Accordingly, $D_1 = D$.

The higher-order difference operators for D_t are defined by induction:

$$D_t^0 g := g \quad \text{and} \quad D_t^k g := D_t(D_t^{k-1} g) \quad (k \geq 1).$$

These operators also fit in Stanley's theory of differential posets [23], although this is a language we will not use here. The following theorem is our main result, which will be proved in Section 5.

Theorem 1.3. *Suppose that t is a positive integer, $u', v', j_u, j'_v, k_u, k'_v$ are non-negative integers and μ is a given t -core partition. Then, there exists some fixed $r \in \mathbb{N}$ such that*

$$D_t^r \left(G_\lambda \left(\prod_{u=1}^{u'} \sum_{\substack{\square \in \lambda \\ h_\square \equiv \pm j_u \pmod{t}}} h_\square^{2k_u} \right) \left(\prod_{v=1}^{v'} \sum_{\substack{\square \in \lambda \\ c_\square \equiv j'_v \pmod{t}}} c_\square^{k'_v} \right) \right) = 0$$

for every partition λ with $\lambda_{t\text{-core}} = \mu$. Moreover,

$$P(n) = \sum_{\substack{\lambda_{t\text{-core}} = \mu \\ |\lambda/\mu| = nt}} F_{\lambda/\mu} G_\lambda \left(\prod_{u=1}^{u'} \sum_{\substack{\square \in \lambda \\ h_\square \equiv \pm j_u \pmod{t}}} h_\square^{2k_u} \right) \left(\prod_{v=1}^{v'} \sum_{\substack{\square \in \lambda \\ c_\square \equiv j'_v \pmod{t}}} c_\square^{k'_v} \right)$$

is a polynomial of n (n is a nonnegative integer).

Let Q_1 and Q_2 be two symmetric functions. By Theorem 1.3, there exists some fixed $r \in \mathbb{N}$ such that

$$D_t^r (G_\lambda Q_1(h_\square^2 : \square \in \lambda) Q_2(c_\square : \square \in \lambda)) = 0$$

for every partition λ with $\lambda_{t\text{-core}} = \mu$. Moreover,

$$(1.7) \quad P(n) = \sum_{\substack{\lambda_{t\text{-core}} = \mu \\ |\lambda/\mu| = nt}} F_{\lambda/\mu} G_\lambda Q_1(h_\square^2 : \square \in \lambda) Q_2(c_\square : \square \in \lambda)$$

is a polynomial of n . When $t = 1, \mu = \emptyset$, we have (1.7) implies Theorem 1.2.

1.6. Specialization for the square case. We now focus on expressions of square sum of contents or hook lengths, and thereby obtain explicit results. Details of the proofs are given in Section 6. We start by considering individual partitions.

Theorem 1.4. *Let $0 \leq k \leq t-1$. For every partition λ with $\lambda_{t\text{-core}} = \emptyset$ we have*

$$(1.8) \quad \sum_{\substack{\square \in \lambda \\ h_\square \equiv k \pmod{t}}} h_\square^2 + \sum_{\substack{\square \in \lambda \\ h_\square \equiv t-k \pmod{t}}} h_\square^2 - \left(\sum_{\substack{\square \in \lambda \\ c_\square \equiv k \pmod{t}}} c_\square^2 + \sum_{\substack{\square \in \lambda \\ c_\square \equiv t-k \pmod{t}}} c_\square^2 \right) \\ = 2t^2 \left(\sum_{i=0}^{t-1-k} n_i n_{i+k} + \sum_{i=0}^{k-1} n_i n_{i+t-k} \right),$$

where $n_i = |\lambda^i|$.

By letting $t = 1$ we obtain the following well-known result (see [16]).

Corollary 1.5. *For every partition λ we have*

$$\sum_{\square \in \lambda} h_\square^2 - \sum_{\square \in \lambda} c_\square^2 = |\lambda|^2.$$

We now turn to t -Plancherel averages, introducing $F_{\lambda/\mu}$ and G_λ as weights in the previous results and summing over λ . The results are stated in two cases according to the divisibility of the hook lengths by t .

Theorem 1.6. *Suppose that μ is a given t -core partition and $1 \leq k \leq t-1$. Then we have*

$$\begin{aligned} & \sum_{\substack{\lambda_{t\text{-core}}=\mu \\ |\lambda/\mu|=nt}} F_{\lambda/\mu} G_\lambda \left(\sum_{\substack{\square \in \lambda \\ h_\square \equiv k \pmod{t}}} h_\square^2 + \sum_{\substack{\square \in \lambda \\ h_\square \equiv t-k \pmod{t}}} h_\square^2 \right) \\ &= 6t \binom{n}{2} + (2k(t-k) + 4t|\mu(k)| + 4t|\mu(t-k)|)n \\ & \quad + \left(\sum_{\substack{\square \in \mu \\ h_\square \equiv k \pmod{t}}} h_\square^2 + \sum_{\substack{\square \in \mu \\ h_\square \equiv t-k \pmod{t}}} h_\square^2 \right), \end{aligned}$$

where $\mu(k) = \{h \in \mathcal{H}(\mu) \mid h \equiv k \pmod{t}\}$, viewed as a multiset. In particular, let $\mu = \emptyset$. Then we have

$$\begin{aligned} & \sum_{\substack{\lambda_{t\text{-core}}=\emptyset \\ |\lambda|=nt}} F_\lambda G_\lambda \left(\sum_{\substack{\square \in \lambda \\ h_\square \equiv k \pmod{t}}} h_\square^2 + \sum_{\substack{\square \in \lambda \\ h_\square \equiv t-k \pmod{t}}} h_\square^2 \right) \\ &= 6t \binom{n}{2} + 2k(t-k)n. \end{aligned}$$

Theorem 1.7. *Suppose that μ is a given t -core partition. Then we have*

$$\sum_{\substack{\lambda_{t\text{-core}}=\mu \\ |\lambda/\mu|=nt}} F_{\lambda/\mu} G_\lambda \sum_{\substack{\square \in \lambda \\ h_\square \equiv 0 \pmod{t}}} h_\square^2 = nt^2 + 3t \binom{n}{2}.$$

By Theorems 1.6 and 1.7 we obtain the following result.

Corollary 1.8. *Suppose that μ is a given t -core partition. Then*

$$\sum_{\substack{\lambda_{t\text{-core}}=\mu \\ |\lambda/\mu|=nt}} F_{\lambda/\mu} G_\lambda \sum_{\square \in \lambda} h_\square^2 = \frac{3t^2 n^2}{2} + \frac{nt(t^2 - 3t - 1 + 24|\mu|)}{6} + \sum_{\square \in \mu} h_\square^2.$$

In particular, let $\mu = \emptyset$. Then we have

$$\sum_{\substack{\lambda_{t\text{-core}}=\emptyset \\ |\lambda|=nt}} F_\lambda G_\lambda \sum_{\square \in \lambda} h_\square^2 = \frac{3t^2 n^2}{2} + \frac{nt(t^2 - 3t - 1)}{6}.$$

Similar results for contents can also be obtained.

Theorem 1.9. *Let $0 \leq k \leq t-1$. We have*

$$\sum_{\substack{\lambda_{t\text{-core}}=\emptyset \\ |\lambda|=nt}} F_\lambda G_\lambda \sum_{\substack{\square \in \lambda \\ c_\square \equiv k \pmod{t}}} c_\square^2 = t \binom{n}{2} + k(t-k)n.$$

Theorem 1.9 implies the following identity.

$$(1.9) \quad \sum_{\substack{\lambda_{t\text{-core}}=\emptyset \\ |\lambda|=nt}} F_\lambda G_\lambda \sum_{\square \in \lambda} c_\square^2 = t^2 \binom{n}{2} + \frac{(t^3 - t)n}{6}.$$

In fact we will derive a more general result in Theorem 6.3 by replacing \emptyset by a general t -core partition.

2. PARTITIONS AND t -DIFFERENCE OPERATOR

We now turn our attention to preliminary lemmas on the t -difference operators D_t^k , and in particular a theorem outlining how polynomiality of t -Plancherel averages can be deduced from the vanishing of D_t^k for large enough k . It is known [11] that

$$(2.1) \quad D\left(\frac{1}{H_\lambda}\right) = 0.$$

We establish a similar result for the t -difference operator.

Lemma 2.1. *Suppose that λ is a partition. Then*

$$D_t(G_\lambda) = 0.$$

In other words,

$$G_\lambda = \sum_{\substack{\lambda^+ \geq_t \lambda \\ |\lambda^+/\lambda|=t}} G_{\lambda^+}.$$

Proof. Suppose that

$$\lambda = (\lambda_{t\text{-core}}; \lambda^0, \lambda^1, \dots, \lambda^{i-1}, \lambda^i, \lambda^{i+1}, \dots, \lambda^{t-1})$$

and

$$\lambda^+ = (\lambda_{t\text{-core}}; \lambda^0, \lambda^1, \dots, \lambda^{i-1}, (\lambda^i)^+, \lambda^{i+1}, \dots, \lambda^{t-1})$$

where $|(\lambda^i)^+/\lambda^i| = 1$. Then, by the definition of G_λ we have

$$\frac{G_{\lambda^+}}{G_\lambda} = \frac{H_{\lambda^i}}{tH_{(\lambda^i)^+}}.$$

By (2.1) we have

$$\sum_{|(\lambda^i)^+/\lambda^i|=1} \frac{G_{(\lambda_{t\text{-core}}; \lambda^0, \lambda^1, \dots, \lambda^{i-1}, (\lambda^i)^+, \lambda^{i+1}, \dots, \lambda^{t-1})}}{G_\lambda} = \frac{1}{t}.$$

Summing the above equality over $0 \leq i \leq t-1$, we prove our claim. \square

We now prove a commutation relation between the averaging over partitions of a given weight and the D_t operator.

Lemma 2.2. *Suppose that μ is a given partition and g is a function of partitions. For every $n \in \mathbb{N}$, let*

$$P_g(n) := \sum_{\substack{\lambda \geq_t \mu \\ |\lambda/\mu|=nt}} F_{\lambda/\mu} g(\lambda).$$

Then

$$P_g(n+1) - P_g(n) = P_{D_t g}(n).$$

Proof. The proof is straightforward and follows from expanding definitions:

$$\begin{aligned}
P_g(n+1) - P_g(n) &= \sum_{\substack{\nu \geq_t \mu \\ |\nu/\mu|=(n+1)t}} F_{\nu/\mu} g(\nu) - \sum_{\substack{\lambda \geq_t \mu \\ |\lambda/\mu|=nt}} F_{\lambda/\mu} g(\lambda) \\
&= \sum_{\substack{\nu \geq_t \mu \\ |\nu/\mu|=(n+1)t}} \sum_{\substack{\nu \geq_t \nu^- \geq_t \mu \\ |\nu/\nu^-|=t}} F_{\nu^-/\mu} g(\nu) - \sum_{\substack{\lambda \geq_t \mu \\ |\lambda/\mu|=nt}} F_{\lambda/\mu} g(\lambda) \\
&= \sum_{\substack{\lambda \geq_t \mu \\ |\lambda/\mu|=nt}} F_{\lambda/\mu} \left(\sum_{\substack{\lambda^+ \geq_t \lambda \\ |\lambda^+/\lambda|=t}} g(\lambda^+) - g(\lambda) \right) \\
&= P_{D_t g}(n). \quad \square
\end{aligned}$$

Example. Let $g(\lambda) = G_\lambda$. Then $D_t g(\lambda) = 0$ by Lemma 2.1, which means that $P_{D_t g}(n) = 0$. Consequently, $P_g(n+1) = P_g(n) = \dots = P_g(0) = G_\mu$, or

$$(2.2) \quad \sum_{\substack{\lambda \geq_t \mu \\ |\lambda/\mu|=nt}} F_{\lambda/\mu} G_\lambda = G_\mu.$$

Recall that when μ is a t -core partition, we have $G_\mu = 1$. This implies

$$(2.3) \quad \sum_{\substack{\lambda_{t\text{-core}}=\mu \\ |\lambda/\mu|=nt}} F_{\lambda/\mu} G_\lambda = 1.$$

In particular, let $\mu = \emptyset$, we obtain a generalization of the second identity of (1.1):

$$(2.4) \quad \sum_{\substack{\lambda_{t\text{-core}}=\emptyset \\ |\lambda|=nt}} F_\lambda G_\lambda = 1,$$

or equivalently,

$$(2.5) \quad \sum_{|\lambda^0|+|\lambda^1|+\dots+|\lambda^{t-1}|=n} \binom{n}{|\lambda^0|, |\lambda^1|, \dots, |\lambda^{t-1}|} \prod_{i=0}^{t-1} \frac{f_{\lambda^i}^2}{|\lambda^i|!} = t^n.$$

Formulas such as this one can be used to create new measures on partitions, called *t-Plancherel*.

Recall that the higher-order difference operators for D_t are defined by induction $D_t^0 g := g$ and $D_t^k g := D_t(D_t^{k-1} g)$ ($k \geq 1$). We write $D_t g(\nu) := D_t g(\lambda)|_{\lambda=\nu}$ for a fixed partition ν .

Theorem 2.3. *Let g be a function of partitions and μ be a given partition. Then,*

$$(2.6) \quad P_g(n) = \sum_{\substack{\lambda \geq_t \mu \\ |\lambda/\mu|=nt}} F_{\lambda/\mu} g(\lambda) = \sum_{k=0}^n \binom{n}{k} D_t^k g(\mu)$$

and

$$(2.7) \quad D_t^n g(\mu) = \sum_{k=0}^n (-1)^{n+k} \binom{n}{k} P_g(k).$$

In particular, if there exists some positive integer r such that $D_t^r g(\lambda) = 0$ for every partition $\lambda \geq_t \mu$, then $P_g(n)$ is a polynomial in n .

Proof. Identity (2.6) will be proved by induction. The case $n = 0$ is obvious. Assume that (2.6) is true for some nonnegative integer n . By Lemma 2.2 we obtain

$$\begin{aligned} P_g(n+1) &= P_g(n) + P_{D_t g}(n) \\ &= \sum_{k=0}^n \binom{n}{k} D_t^k g(\mu) + \sum_{k=0}^n \binom{n}{k} D_t^{k+1} g(\mu) \\ &= \sum_{k=0}^{n+1} \binom{n+1}{k} D_t^k g(\mu). \end{aligned}$$

Identity (2.7) follows from the famous Möbius inversion formula [21]. \square

3. CORNERS OF PARTITIONS

Now we introduce the concept of corners in a partition. For a partition λ , the *outer corners* (see [3, 11]) are the boxes which can be removed in such a way that after removal the resulting diagram is still a Young diagram. The coordinates of outer corners are denoted by $(\alpha_1, \beta_1), \dots, (\alpha_m, \beta_m)$ such that $\alpha_1 > \alpha_2 > \dots > \alpha_m$. Let $y_j := \beta_j - \alpha_j$ ($1 \leq j \leq m$) be the contents of outer corners. Set $\alpha_{m+1} = \beta_0 = 0$. The *inner corners* of λ are defined to be the boxes whose coordinates are $(\alpha_1, \beta_0), (\alpha_2, \beta_1), \dots, (\alpha_{m+1}, \beta_m)$. Let $x_i = \beta_i - \alpha_{i+1}$ be the contents of the inner corners for $0 \leq i \leq m$.

Example. For $\lambda = (6, 3, 2, 2)$ (see Figure 1) we have $m = 3$, $(\alpha_1, \beta_1) = (4, 2)$, $(\alpha_2, \beta_2) = (2, 3)$ and $(\alpha_3, \beta_3) = (1, 6)$. Therefore $(x_0, x_1, x_2, x_3) = (-4, 0, 2, 6)$ and $(y_1, y_2, y_3) = (-2, 1, 5)$. Also note that $\sum_{0 \leq i \leq 3} x_i - \sum_{1 \leq j \leq 3} y_j = 0$ and $\sum_{0 \leq i \leq 3} x_i^2 - \sum_{1 \leq j \leq 3} y_j^2 = 26 = 2|\lambda|$.

Let λ be a partition and x_i, y_j be the quantities associated to λ defined above. Define (see [11])

$$(3.1) \quad q_k(\lambda) := \sum_{0 \leq i \leq m} x_i^k - \sum_{1 \leq j \leq m} y_j^k$$

for each integer k , and

$$(3.2) \quad q_\nu(\lambda) := \prod_{k=1}^{\ell} q_{\nu_k}(\lambda)$$

for each partition $\nu = (\nu_1, \nu_2, \dots, \nu_\ell)$.²

We need the following results developed in [11] (see (6.3), (6.6), Theorem 6.1 and Lemma 6.4 in [11] respectively).

Theorem 3.1. *Let $\lambda^{i+} = \lambda \cup \{\square_i\}$ with $c_{\square_i} = x_i$ for $0 \leq i \leq m$.*

(1) *For $k = 0, 1, 2$, we have*

$$q_0(\lambda) = 1, \quad q_1(\lambda) = 0 \quad \text{and} \quad q_2(\lambda) = 2|\lambda|.$$

(2) *Suppose that k is a nonnegative integer. Then we have*

$$q_k(\lambda^{i+}) - q_k(\lambda) = \sum_{1 \leq j \leq k/2} 2 \binom{k}{2j} x_i^{k-2j}.$$

²We need to be careful here: in general, $q_\nu(\lambda) = \prod_i q_{\nu_i}(\lambda) \neq \prod_i q_{\nu^i}(\lambda)$, which was used in Theorem 3.4 and commented on in footnote 3. Writing q_ν allows for a convenient access to $|\nu|$.

(3) Let ν be a partition. Then, there exists some $b_\delta \in \mathbb{Q}$ such that

$$D\left(\frac{q_\nu(\lambda)}{H_\lambda}\right) = \sum_{|\delta| \leq |\nu|-2} b_\delta \frac{q_\delta(\lambda)}{H_\lambda}$$

for every partition λ .

(4) Let k be a nonnegative integer. Then there exists some $b_\delta \in \mathbb{Q}$ such that

$$\sum_{0 \leq i \leq m} \frac{H_\lambda}{H_{\lambda^{i+}}} x_i^k = \sum_{|\delta| \leq k} b_\delta q_\delta(\lambda)$$

for every partition λ .

The next result is easy to prove by Theorem 3.1(4).

Theorem 3.2. Let $\lambda^{i+} = \lambda \cup \{\square_i\}$ with $c_{\square_i} = x_i$ for $0 \leq i \leq m$. Then,

$$\sum_{0 \leq i \leq m} \frac{H_\lambda}{H_{\lambda^{i+}}} x_i = 0$$

and

$$\sum_{0 \leq i \leq m} \frac{H_\lambda}{H_{\lambda^{i+}}} x_i^2 = |\lambda|$$

for every partition λ .

We obtain the following result, which is a generalization of Theorem 3.1(3).

Theorem 3.3. Let ν be a given partition. Then, there exists some $b_{\delta^0, \delta^1, \dots, \delta^{t-1}} \in \mathbb{Q}$ such that

$$(3.3) \quad D_t \left(G_\lambda \prod_{i=0}^{t-1} q_{\nu^i}(\lambda^i) \right) = \sum_{(*)} b_{\delta^0, \delta^1, \dots, \delta^{t-1}} G_\lambda \prod_{i=0}^{t-1} q_{\delta^i}(\lambda^i)$$

for every partition λ , where $(*)$ ranges over all partitions $\delta^0, \delta^1, \dots, \delta^{t-1}$ such that $\sum_{i=0}^{t-1} |\delta^i| \leq \sum_{i=0}^{t-1} |\nu^i| - 2$. Notice that $b_{\delta^0, \delta^1, \dots, \delta^{t-1}}$ is independent of λ .

Furthermore, if $r \geq \frac{1}{2} \sum_{i=0}^{t-1} |\nu^i| + 1$, then

$$D_t^r \left(G_\lambda \prod_{i=0}^{t-1} q_{\nu^i}(\lambda^i) \right) = 0$$

for every partition λ .

Proof. We have

$$\begin{aligned} & D_t \left(G_\lambda \prod_{i=0}^{t-1} q_{\nu^i}(\lambda^i) \right) \\ &= \sum_{\substack{\lambda^+ \geq_t \lambda \\ |\lambda^+/\lambda|=t}} G_{\lambda^+} \prod_{i=0}^{t-1} q_{\nu^i}((\lambda^+)^i) - G_\lambda \prod_{i=0}^{t-1} q_{\nu^i}(\lambda^i) \\ &= G_\lambda \sum_{\substack{\lambda^+ \geq_t \lambda \\ |\lambda^+/\lambda|=t}} \frac{G_{\lambda^+}}{G_\lambda} \left(\prod_{i=0}^{t-1} q_{\nu^i}((\lambda^+)^i) - \prod_{i=0}^{t-1} q_{\nu^i}(\lambda^i) \right) \end{aligned}$$

$$\begin{aligned}
&= \frac{1}{t} G_\lambda \sum_{j=0}^{t-1} \prod_{\substack{0 \leq i \leq t-1 \\ i \neq j}} q_{\nu^i}(\lambda^i) \sum_{|(\lambda^j)^+ / \lambda^j|=1} \frac{H_{\lambda^j}}{H_{(\lambda^j)^+}} (q_{\nu^j}((\lambda^j)^+) - q_{\nu^j}(\lambda^j)) \\
&= \frac{1}{t} G_\lambda \sum_{j=0}^{t-1} \prod_{\substack{0 \leq i \leq t-1 \\ i \neq j}} q_{\nu^i}(\lambda^i) H_{\lambda^j} D \left(\frac{q_{\nu^j}(\lambda^j)}{H_{\lambda^j}} \right) \\
&= \sum_{(*)} b_{\delta^0, \delta^1, \dots, \delta^{t-1}} G_\lambda \prod_{i=0}^{t-1} q_{\delta^i}(\lambda^i)
\end{aligned}$$

for some $b_{\delta^0, \delta^1, \dots, \delta^{t-1}} \in \mathbb{Q}$, where $(*)$ ranges over all partitions $\delta^0, \delta^1, \dots, \delta^{t-1}$ such that $\sum_{i=0}^{t-1} |\delta^i| \leq \sum_{i=0}^{t-1} |\nu^i| - 2$. The last equality is due to Theorem 3.1(3). \square

Definition 3.1. Let μ be a t -core partition. A function g of partitions is called μ -admissible, if there exist some functions of partitions η_s (here η_s depends on t, g, i, μ) such that $g(\lambda^+) - g(\lambda) = \sum_{s=0}^{s'} \eta_s(\lambda) c_{\square_i}^s$ is a polynomial of c_{\square_i} for every pair of partitions

$$\lambda = (\mu; \lambda^0, \lambda^1, \dots, \lambda^{i-1}, \lambda^i, \lambda^{i+1}, \dots, \lambda^{t-1})$$

and

$$\lambda^+ = (\mu; \lambda^0, \lambda^1, \dots, \lambda^{i-1}, \lambda^i \cup \{\square_i\}, \lambda^{i+1}, \dots, \lambda^{t-1}),$$

where each coefficient $\eta_s(\lambda)$ is a linear combination of some $\prod_{j=0}^{t-1} q_{\tau^j}(\lambda^j)$ for some partition τ ,³ and for every partition λ with $\lambda_{t\text{-core}} = \mu$.

By Theorem 3.1(2), $\lambda \mapsto \prod_{i=0}^{t-1} q_{\nu^i}(\lambda^i)$ is μ -admissible for any partition ν and any t -core partition μ . As can be seen in Section 5, the following two functions of partitions

$$g(\lambda) = \sum_{\substack{\square \in \lambda \\ h_{\square} \equiv \pm j \pmod{t}}} h_{\square}^{2k}$$

and

$$g(\lambda) = \sum_{\substack{\square \in \lambda \\ c_{\square} \equiv j \pmod{t}}} c_{\square}^k$$

are μ -admissible for any t -core partition μ , any nonnegative integer k , and $0 \leq j \leq t-1$. This is a consequence of Lemmas 5.2 and 5.4.

Theorem 3.4. Let t be a given integer, ν be a partition, and μ be a t -core partition. Suppose that g_1, g_2, \dots, g_v are μ -admissible functions of partitions. Then, there exists some $r \in \mathbb{N}$ such that

$$D_t^r \left(G_\lambda \prod_{u=1}^v g_u(\lambda) \prod_{i=0}^{t-1} q_{\nu^i}(\lambda^i) \right) = 0$$

³We really need here a t -tuple of partitions. We decide to reuse the quotient notation for this purpose. Different τ with the same t -core would give the same tuple, but the surjectivity is inconsequential here.

for every partition λ with $\lambda_{t\text{-core}} = \mu$. Furthermore, by Theorem 2.3,

$$(3.4) \quad P(n) = \sum_{\substack{\lambda_{t\text{-core}} = \mu \\ |\lambda/\mu| = nt}} F_{\lambda/\mu} G_{\lambda} \prod_{u=1}^v g_u(\lambda)$$

is a polynomial in n .

Proof of Theorem 3.4. Notice that this theorem is true for $v = 0$ by Theorem 3.3. We will prove it by induction. First, we have

$$\begin{aligned} & D_t \left(G_{\lambda} \prod_{u=1}^v g_u(\lambda) \prod_{i=0}^{t-1} q_{\nu^i}(\lambda^i) \right) \\ &= G_{\lambda} \sum_{\substack{\lambda^+ \geq_t \lambda \\ |\lambda^+/\lambda| = t}} \frac{G_{\lambda^+}}{G_{\lambda}} \left(\prod_{u=1}^v g_u(\lambda^+) \prod_{i=0}^{t-1} q_{\nu^i}((\lambda^+)^i) - \prod_{u=1}^v g_u(\lambda) \prod_{i=0}^{t-1} q_{\nu^i}(\lambda^i) \right) \\ &= G_{\lambda} \sum_{\substack{\lambda^+ \geq_t \lambda \\ |\lambda^+/\lambda| = t}} \frac{G_{\lambda^+}}{G_{\lambda}} (A \cdot \Delta B + B \cdot \Delta A + \Delta A \cdot \Delta B), \end{aligned}$$

where

$$\begin{aligned} A &= \prod_{u=1}^v g_u(\lambda), \\ \Delta A &= \prod_{u=1}^v g_u(\lambda^+) - \prod_{u=1}^v g_u(\lambda), \\ B &= \prod_{i=0}^{t-1} q_{\nu^i}(\lambda^i), \\ \Delta B &= \prod_{i=0}^{t-1} q_{\nu^i}((\lambda^+)^i) - \prod_{i=0}^{t-1} q_{\nu^i}(\lambda^i). \end{aligned}$$

Next,

$$\begin{aligned} & G_{\lambda} \sum_{\substack{\lambda^+ \geq_t \lambda \\ |\lambda^+/\lambda| = t}} \frac{G_{\lambda^+}}{G_{\lambda}} (A \cdot \Delta B) \\ &= \frac{1}{t} G_{\lambda} \prod_{u=1}^v g_u(\lambda) \sum_{j=0}^{t-1} \left(\prod_{\substack{0 \leq i \leq t-1 \\ i \neq j}} q_{\nu^i}(\lambda^i) \right) H_{\lambda^j} D \left(\frac{q_{\nu^j}(\lambda^j)}{H_{\lambda^j}} \right), \\ & G_{\lambda} \sum_{\substack{\lambda^+ \geq_t \lambda \\ |\lambda^+/\lambda| = t}} \frac{G_{\lambda^+}}{G_{\lambda}} (B \cdot \Delta A) \\ &= \frac{1}{t} G_{\lambda} \prod_{i=0}^{t-1} q_{\nu^i}(\lambda^i) \sum_{j=0}^{t-1} \sum_{|(\lambda^j)^+/\lambda^j| = 1} \frac{H_{\lambda^j}}{H_{(\lambda^j)^+}} \end{aligned}$$

$$\begin{aligned}
& \left(\prod_{u=1}^v g_u((\mu; \lambda^0, \dots, (\lambda^j)^+, \dots, \lambda^{t-1})) - \prod_{u=1}^v g_u(\lambda) \right), \\
G_\lambda & \sum_{\substack{\lambda^+ \geq t\lambda \\ |\lambda^+/\lambda|=t}} \frac{G_{\lambda^+}}{G_\lambda} (\Delta A \cdot \Delta B) \\
&= \frac{1}{t} G_\lambda \sum_{j=0}^{t-1} \sum_{|(\lambda^j)^+/\lambda^j|=1} \frac{H_{\lambda^j}}{H_{(\lambda^j)^+}} \left(\prod_{\substack{0 \leq i \leq t-1 \\ i \neq j}} q_{\nu^i}(\lambda^i) \right) \left(q_{\nu^j}((\lambda^j)^+) - q_{\nu^j}(\lambda^j) \right) \\
& \left(\prod_{u=1}^v g_u((\mu; \lambda^0, \dots, (\lambda^j)^+, \dots, \lambda^{t-1})) - \prod_{u=1}^v g_u(\lambda) \right).
\end{aligned}$$

By Theorems 3.1 and 3.3 each of the last three terms could be written as a linear combination of some $G_\lambda \prod_{u'=1}^{v'} g'_{u'}(\lambda) \prod_{i=0}^{t-1} q_{(\nu')^i}(\lambda^i)$ for some μ -admissible functions $g'_{u'}$, where either $v' = v$ and simultaneously $\sum_{i=0}^{t-1} |(\nu')^i| \leq \sum_{i=0}^{t-1} |\nu^i| - 2$, or $v' < v$. Then, by Theorem 3.3 and induction we prove the claim. \square

4. FURTHER PROPERTIES OF LITTLEWOOD DECOMPOSITION

In this section, we always assume that μ is a given t -core partition with 01-sequence $z(\mu) = (z_{\mu,j})_{j \in \mathbb{Z}}$. For $0 \leq i \leq t-1$, we have $\mu^i = \emptyset$. Thus, μ^i has a 01-sequence equal to $\cdots 0000|11111\cdots$, which means that $z_{\mu^i,j} = 0$ for $j < 0$ and $z_{\mu^i,j} = 1$ for $j \geq 0$. Let

$$b_i := b_i(\mu) = \min\{j \in \mathbb{Z} : j \equiv i \pmod{t}, z_{\mu,j} = 1\}.$$

Then, there exist some $d_i \in \mathbb{Z}$ such that $b_i = td_i + i$ for $0 \leq i \leq t-1$. Therefore, $z_{\mu,tj+i} = 0$ for $j < d_i$ and $z_{\mu,tj+i} = 1$ for $j \geq d_i$.

Example. The 01-sequence of the 3-core partition $(5, 3, 1, 1)$ is

$$\cdots 001001|1011011\cdots.$$

Therefore, $b_0 = 0$, $b_1 = 7$, $b_2 = -4$, $d_0 = 0$, $d_1 = 2$, and $d_2 = -2$. Notice that $d_0 + d_1 + d_2 = 0$.

We define

$$B_k = \{(i, i+k) : 0 \leq i \leq t-1-k\} \cup \{(i, i+t-k) : 0 \leq i \leq k-1\},$$

viewed as multisets for $1 \leq k \leq t-1$ (i.e., if k equals $t-k$, we keep two copies). Then we have

$$(4.1) \quad \sum_{(i,j) \in B_k} (j-i)^2 = tk(t-k).$$

Lemma 4.1. *Suppose that μ is a given t -core partition. Let b_i and d_i be defined as above. Then*

$$\sum_{i=0}^{t-1} d_i = 0.$$

Proof. Let $0 \leq i \leq t-1$. First, the fact that μ is a t -core partition implies that $z_{\mu,tj+i} = 0$ for $j < d_i$ and $z_{\mu,tj+i} = 1$ for $j \geq d_i$. By the definition of 01-sequence we have $\#\{z_{\mu,j} = 0 : j \geq 0\} = \#\{z_{\mu,j} = 1 : j < 0\}$. Thus,

$$\sum_{i=0}^{t-1} d_i = \#\{z_{\mu,j} = 0 : j \geq 0\} - \#\{z_{\mu,j} = 1 : j < 0\} = 0.$$

□

For every partition λ and $0 \leq k \leq t-1$, recall that $\lambda(k) = \{h_{\square} \equiv k(\text{mod } t) : \square \in \lambda\}$, viewed as a multiset.

Lemma 4.2. *Suppose that μ is a given t -core partition and $1 \leq k \leq t-1$. Then, $|\mu(0)| = 0$ and*

$$|\mu(k)| + |\mu(t-k)| = \sum_{(i,j) \in B_k} \binom{d_i - d_j}{2}.$$

Proof. First, $|\mu(0)| = 0$ since μ is a t -core partition. By Lemma 1.1 we have

$$|\mu(k)| = \#\{(i', j') : i' < j', j' - i' \equiv k(\text{mod } t), z_{\mu,i'} = 1, z_{\mu,j'} = 0\}.$$

Now we just need to consider the right hand side of the above identity. Notice that for $0 \leq i \leq t-1$ we have $z_{\mu,tj+i} = 0$ for $j < d_i$ and $z_{\mu,tj+i} = 1$ for $j \geq d_i$. Then, for each $(i, j) \in B_k$, we have

$$\begin{aligned} & \#\{(i', j') : i' \equiv i(\text{mod } t), j' \equiv j(\text{mod } t), i' < j', z_{\mu,i'} = 1, z_{\mu,j'} = 0\} \\ & + \#\{(i', j') : i' \equiv j(\text{mod } t), j' \equiv i(\text{mod } t), i' < j', z_{\mu,i'} = 1, z_{\mu,j'} = 0\} \\ & = 1 + 2 + \cdots + (d_i - d_j - 1) = \binom{d_i - d_j}{2} = \binom{d_j - d_i + 1}{2} \end{aligned}$$

whenever $d_i > d_j$ or $d_i \leq d_j$. This means that

$$|\mu(k)| + |\mu(t-k)| = \sum_{(i,j) \in B_k} \binom{d_i - d_j}{2}.$$

□

By Lemmas 4.1 and 4.2 we obtain

$$(4.2) \quad \sum_{(i,j) \in B_k} (2j - 2i)(d_i - d_j) = t \sum_{(i,j) \in B_k} (d_i - d_j),$$

$$(4.3) \quad t \sum_{0 \leq i \leq t-1} d_i^2 = \sum_{0 \leq i < j \leq t-1} (d_i - d_j)^2,$$

$$(4.4) \quad -2 \sum_{0 \leq i \leq t-1} i d_i = \sum_{0 \leq i < j \leq t-1} (d_i - d_j),$$

and

$$\begin{aligned} (4.5) \quad |\mu| &= \sum_{0 \leq i < j \leq t-1} \binom{d_i - d_j}{2} \\ &= \frac{t}{2} \sum_{0 \leq i \leq t-1} d_i^2 + \sum_{0 \leq i \leq t-1} i d_i \end{aligned}$$

$$= \frac{1}{2t^2} \sum_{0 \leq i < j \leq t-1} ((b_i - b_j)^2 - (i - j)^2).$$

Lemma 4.3. *Suppose that μ is a given t -core partition and $0 \leq k \leq t-1$. Let λ be a partition with $\lambda_{t\text{-core}} = \mu$ and $|\lambda| = |\mu| + nt$. Then,*

$$|\lambda(k)| + |\lambda(t-k)| - |\mu(k)| - |\mu(t-k)| = 2n.$$

Proof. Adding a t -hook to a partition λ means that we change some $(z_i = 0, z_{i+t} = 1)$ to $(z_i = 1, z_{i+t} = 0)$ in the 01-sequence of λ . Then, the lemma follows directly from Lemma 1.1 and induction on the size of λ . \square

5. CONTENTS, HOOK LENGTHS AND t -DIFFERENCE OPERATORS

By the construction of 01-sequences and Lemma 1.1, the following lemma is obvious.

Lemma 5.1. *Let $z(\lambda) = (z_{\lambda,i})_{i \in \mathbb{Z}}$ be the 01-sequence of a partition λ . Then, λ has an inner (resp. outer) corner with content k if and only if $z_{\lambda,k-1} = 0$, $z_{\lambda,k} = 1$ (resp. $z_{\lambda,k-1} = 1$, $z_{\lambda,k} = 0$).*

For each partition λ let $\mathcal{C}(\lambda)$ be the multiset of its contents. We explicitly compute how contents change when adding a box to one of the quotients in the Littlewood decomposition.

Lemma 5.2. *Suppose that μ is a given t -core partition. Let*

$$\lambda = (\mu; \lambda^0, \lambda^1, \dots, \lambda^{i-1}, \lambda^i, \lambda^{i+1}, \dots, \lambda^{t-1})$$

and

$$\lambda^+ = (\mu; \lambda^0, \lambda^1, \dots, \lambda^{i-1}, \lambda^i \cup \{\square_i\}, \lambda^{i+1}, \dots, \lambda^{t-1}).$$

Then,

$$\mathcal{C}(\lambda^+) \setminus \mathcal{C}(\lambda) = \{c_{\square_i}t + b_i, c_{\square_i}t + b_i - 1, \dots, c_{\square_i}t + b_i - (t-1)\}.$$

Proof. Suppose that the 01-sequence of λ is $(z_{\lambda,j})_{j \in \mathbb{Z}}$. For each $0 \leq i \leq t-1$ let $(z_{\lambda^i,j})_{j \in \mathbb{Z}}$ be the 01-sequence of λ^i . Then, $z_{\lambda^i,j} = z_{\lambda,jt+b_i}$ since $\lambda_{t\text{-core}} = \mu$. Notice that b_i are determined by μ . Let $c = c_{\square_i}$ be the content of \square_i . Then, by Lemma 5.1 adding the box \square_i to λ^i means that $(z_{\lambda^i,c-1}, z_{\lambda^i,c})$, which is equal to $(0, 1)$ in the 01-sequence of λ^i , is changed to $(1, 0)$; or equivalently, $(z_{\lambda,(c-1)t+b_i}, z_{\lambda,ct+b_i}) = (0, 1)$ in the 01-sequence of λ is changed to $(1, 0)$. Then by Lemma 1.1 this means that we add boxes with contents

$$ct + b_i, ct + b_i - 1, \dots, ct + b_i - (t-1)$$

to λ . \square

By induction and Lemma 5.2 we obtain the following lemma.

Lemma 5.3. *Suppose that μ is a given t -core partition. Then*

$$\mathcal{C}(\lambda) \setminus \mathcal{C}(\mu) = \bigcup_{i=0}^{t-1} \{tc_{\square_i} + b_i - j : 0 \leq j \leq t-1, \square_i \in \lambda^i\}$$

for every partition λ with $\lambda_{t\text{-core}} = \mu$.

For the partition λ and $0 \leq i \leq t-1$, let $x_{i,l}$ ($0 \leq l \leq m_i$) be the contents of inner corners of λ^i and $y_{i,l}$ ($1 \leq l \leq m_i$) be the contents of outer corners of λ^i .

Lemma 5.4. Suppose that μ is a given t -core partition, r is a given integer, $1 \leq k \leq t-1$, and $0 \leq i \leq t-1$. Let

$$\lambda = (\mu; \lambda^0, \lambda^1, \dots, \lambda^{i-1}, \lambda^i, \lambda^{i+1}, \dots, \lambda^{t-1})$$

and

$$\lambda^+ = (\mu; \lambda^0, \lambda^1, \dots, \lambda^{i-1}, \lambda^i \cup \{\square_i\}, \lambda^{i+1}, \dots, \lambda^{t-1}).$$

Then

$$\sum_{\substack{\square \in \lambda^+ \\ h_{\square} \equiv 0 \pmod{t}}} h_{\square}^{2r} - \sum_{\substack{\square \in \lambda \\ h_{\square} \equiv 0 \pmod{t}}} h_{\square}^{2r} = t^{2r} + \sum_{0 \leq l \leq m_i} (t(c_{\square_i} - x_{i,l}))^{2r} - \sum_{1 \leq l \leq m_i} (t(c_{\square_i} - y_{i,l}))^{2r}$$

and

$$\begin{aligned} & \sum_{\substack{\square \in \lambda^+ \\ h_{\square} \equiv k \pmod{t}}} h_{\square}^{2r} + \sum_{\substack{\square \in \lambda^+ \\ h_{\square} \equiv t-k \pmod{t}}} h_{\square}^{2r} - \sum_{\substack{\square \in \lambda \\ h_{\square} \equiv k \pmod{t}}} h_{\square}^{2r} - \sum_{\substack{\square \in \lambda \\ h_{\square} \equiv t-k \pmod{t}}} h_{\square}^{2r} \\ &= \sum_{0 \leq l \leq m_{i'}} (tc_{\square_i} + b_i - tx_{i',l} - b_{i'})^{2r} - \sum_{1 \leq l \leq m_{i'}} (tc_{\square_i} + b_i - ty_{i',l} - b_{i'})^{2r} \\ &+ \sum_{0 \leq l \leq m_{i''}} (tc_{\square_i} + b_i - tx_{i'',l} - b_{i''})^{2r} - \sum_{1 \leq l \leq m_{i''}} (tc_{\square_i} + b_i - ty_{i'',l} - b_{i''})^{2r} \end{aligned}$$

where $0 \leq i', i'' \leq t-1$ satisfy $i' \equiv i+k \pmod{t}$ and $i'' \equiv i-k \pmod{t}$. Furthermore,

$$\begin{aligned} \sum_{\square \in \lambda^+} h_{\square}^{2r} - \sum_{\square \in \lambda} h_{\square}^{2r} &= t^{2r} + \sum_{j=0}^{t-1} \left(\sum_{0 \leq l \leq m_j} (tc_{\square_i} + b_i - tx_{j,l} - b_j)^{2r} \right. \\ &\quad \left. - \sum_{1 \leq l \leq m_j} (tc_{\square_i} + b_i - ty_{j,l} - b_j)^{2r} \right). \end{aligned}$$

Proof. Suppose that the 01-sequence of λ is $(z_{\lambda,j})_{j \in \mathbb{Z}}$. For each $0 \leq i \leq t-1$, let $(z_{\lambda^i,j})_{j \in \mathbb{Z}}$ be the 01-sequence of λ^i . Then, by the Littlewood decomposition we have $z_{\lambda^i,j} = z_{\lambda,jt+b_i}$. Let $c = c_{\square_i} = x_{i,w_i}$. By Lemma 5.1, adding the box \square_i to λ^i means that $(z_{\lambda^i,c-1}, z_{\lambda^i,c}) = (0, 1)$ in the 01-sequence of λ^i is changed to $(1, 0)$, or equivalently, $(z_{\lambda,(c-1)t+b_i}, z_{\lambda,ct+b_i}) = (0, 1)$ in the 01-sequence of λ is changed to $(z_{\lambda^+, (c-1)t+b_i}, z_{\lambda^+, ct+b_i}) = (1, 0)$. By Lemma 1.1, to see the difference between hook length sets of λ^+ and λ , we just need to consider the different pairs (j, j') where $j < j'$, $(1, 0) = (z_{\lambda,j}, z_{\lambda,j'})$ or $(z_{\lambda^+,j}, z_{\lambda^+,j'})$, and $\{j, j'\} \cap \{(c-1)t+b_i, ct+b_i\} \neq \emptyset$ in the 01-sequences of λ and λ^+ .

Case (i). First consider such pairs (j, j') satisfying $j' - j \equiv 0 \pmod{t}$. Since $(z_{\lambda,(c-1)t+b_i}, z_{\lambda,ct+b_i}) = (0, 1)$ and $(z_{\lambda^+, (c-1)t+b_i}, z_{\lambda^+, ct+b_i}) = (1, 0)$, it follows from Lemmas 1.1 and 5.1 that the hook lengths of λ^+ for such pairs (j, j') are

$$\begin{aligned} & t, t(c - x_{i,l}), t(c - x_{i,l} - 1), \dots, t(c - y_{i,l+1} + 1) \quad (0 \leq l \leq w_i - 1), \\ & t(x_{i,l} - c), t(x_{i,l} - c - 1), \dots, t(y_{i,l} - c + 1) \quad (w_i + 1 \leq l \leq m_i), \end{aligned}$$

and the hook lengths of λ for such pairs are

$$\begin{aligned} & t(c - x_{i,l} - 1), t(c - x_{i,l} - 2), \dots, t(c - y_{i,l+1}) \quad (0 \leq l \leq w_i - 1), \\ & t(x_{i,l} - c - 1), t(x_{i,l} - c - 2), \dots, t(y_{i,l} - c) \quad (w_i + 1 \leq l \leq m_i). \end{aligned}$$

Let $|a|$ be the absolute value for every real number a . Thus

$$\begin{aligned} & \{h_{\square} \in \mathcal{H}(\lambda^+) : h_{\square} \equiv 0(\text{mod } t)\} \setminus \{h_{\square} \in \mathcal{H}(\lambda) : h_{\square} \equiv 0(\text{mod } t)\} \\ &= \left(\{t\} \bigcup \{t(c - x_{i,l}) : 0 \leq l \leq w_i - 1\} \bigcup \{t(x_{i,l} - c) : w_i + 1 \leq l \leq m_i\} \right) \\ & \setminus \left(\{t(c - y_{i,l+1}) : 0 \leq l \leq w_i - 1\} \bigcup \{t(y_{i,l} - c) : w_i + 1 \leq l \leq m_i\} \right) \\ &= \left(\{t\} \bigcup \{|t(c - x_{i,l})| : 0 \leq l \leq m_i, l \neq w_i\} \right) \setminus \{|t(c - y_{i,l+1})| : 1 \leq l \leq m_i\}, \end{aligned}$$

which means that

$$\begin{aligned} \sum_{\substack{\square \in \lambda^+ \\ h_{\square} \equiv 0(\text{mod } t)}} h_{\square}^{2r} - \sum_{\substack{\square \in \lambda \\ h_{\square} \equiv 0(\text{mod } t)}} h_{\square}^{2r} = \\ t^{2r} + \sum_{0 \leq l \leq m_i} (t(c - x_{i,l}))^{2r} - \sum_{1 \leq l \leq m_i} (t(c - y_{i,l}))^{2r} \end{aligned}$$

since $c = x_{i,w_i}$.

Case (ii). For $1 \leq k \leq t-1$ consider such pairs (j, j') satisfying $j' - j \equiv k$ or $t-k(\text{mod } t)$. Let $0 \leq i', i'' \leq t-1$ satisfy $i' \equiv i+k(\text{mod } t)$ and $i'' \equiv i-k(\text{mod } t)$. Then, by the similar argument as in Case (i) we obtain

$$\begin{aligned} & (\{h_{\square} \in \mathcal{H}(\lambda^+) : h_{\square} \equiv k(\text{mod } t)\} \bigcup \{h_{\square} \in \mathcal{H}(\lambda^+) : h_{\square} \equiv t-k(\text{mod } t)\}) \\ & \setminus (\{h_{\square} \in \mathcal{H}(\lambda) : h_{\square} \equiv k(\text{mod } t)\} \bigcup \{h_{\square} \in \mathcal{H}(\lambda) : h_{\square} \equiv t-k(\text{mod } t)\}) \\ &= (\{|tc - tx_{i',l} + b_i - b_{i'}| : 0 \leq l \leq m_{i'}\} \bigcup \{|tc - tx_{i'',l} + b_i - b_{i''}| : 0 \leq l \leq m_{i''}\}) \\ & \setminus (\{|tc - ty_{i',l} + b_i - b_{i'}| : 1 \leq l \leq m_{i'}\} \bigcup \{|tc - ty_{i'',l} + b_i - b_{i''}| : 1 \leq l \leq m_{i''}\}), \end{aligned}$$

which means that

$$\begin{aligned} & \sum_{\substack{\square \in \lambda^+ \\ h_{\square} \equiv k(\text{mod } t)}} h_{\square}^{2r} + \sum_{\substack{\square \in \lambda^+ \\ h_{\square} \equiv t-k(\text{mod } t)}} h_{\square}^{2r} - \sum_{\substack{\square \in \lambda \\ h_{\square} \equiv k(\text{mod } t)}} h_{\square}^{2r} - \sum_{\substack{\square \in \lambda \\ h_{\square} \equiv t-k(\text{mod } t)}} h_{\square}^{2r} \\ &= \sum_{0 \leq l \leq m_{i'}} (tc + b_i - tx_{i',l} - b_{i'})^{2r} - \sum_{1 \leq l \leq m_{i'}} (tc + b_i - ty_{i',l} - b_{i'})^{2r} \\ &+ \sum_{0 \leq l \leq m_{i''}} (tc + b_i - tx_{i'',l} - b_{i''})^{2r} - \sum_{1 \leq l \leq m_{i''}} (tc + b_i - ty_{i'',l} - b_{i''})^{2r}. \end{aligned}$$

Finally, by Cases (i) and (ii) the proof is completed. \square

Now we are ready to prove Theorem 1.3.

Proof of Theorem 1.3. By Lemmas 5.2 and 5.4, the following two functions of partitions

$$g_j(\lambda) = \sum_{\substack{\square \in \lambda \\ h_{\square} \equiv \pm j(\text{mod } t)}} h_{\square}^{2k}$$

and

$$g'_j(\lambda) = \sum_{\substack{\square \in \lambda \\ c_{\square} \equiv j(\text{mod } t)}} c_{\square}^k$$

are μ -admissible for any t -core partition μ , nonnegative integer k , and $0 \leq j \leq t-1$. Then, the proof is achieved by Theorem 3.4. \square

6. EXPLICIT FORMULAS FOR THE SQUARE CASE

Now we are able to deduce more explicit results.

Theorem 6.1. *Suppose that μ is a given t -core partition. Recall that b_i and d_i are defined as in Section 4. Let $1 \leq k \leq t-1$. For every partition λ with $\lambda_{t\text{-core}} = \mu$ we have*

$$\begin{aligned} & \sum_{\substack{\square \in \lambda \\ h_{\square} \equiv k \pmod{t}}} h_{\square}^2 + \sum_{\substack{\square \in \lambda \\ h_{\square} \equiv t-k \pmod{t}}} h_{\square}^2 - \left(\sum_{\substack{\square \in \lambda \\ c_{\square} \equiv k \pmod{t}}} c_{\square}^2 + \sum_{\substack{\square \in \lambda \\ c_{\square} \equiv t-k \pmod{t}}} c_{\square}^2 \right) \\ = & \sum_{\substack{\square \in \mu \\ h_{\square} \equiv k \pmod{t}}} h_{\square}^2 + \sum_{\substack{\square \in \mu \\ h_{\square} \equiv t-k \pmod{t}}} h_{\square}^2 - \left(\sum_{\substack{\square \in \mu \\ c_{\square} \equiv k \pmod{t}}} c_{\square}^2 + \sum_{\substack{\square \in \mu \\ c_{\square} \equiv t-k \pmod{t}}} c_{\square}^2 \right) \\ & + \sum_{(i,j) \in B_k} \left(2t^2 n_i n_j + t(b_j + j - 2b_i) d_j n_i + t(b_i + i - 2b_j) d_i n_j \right. \\ & \left. - \frac{1}{3} t^2 (d_j q_3(\lambda^i) + d_i q_3(\lambda^j)) \right), \end{aligned}$$

where $n_i = |\lambda^i|$ for $0 \leq i \leq t-1$.

Proof. We will prove this result by induction on the size of λ . When $\lambda = \mu$, we have $\lambda^i = \emptyset$, $n_i = 0$ and thus the identity holds. Now suppose that the identity holds for

$$\lambda = (\mu; \lambda^0, \lambda^1, \dots, \lambda^{i-1}, \lambda^i, \lambda^{i+1}, \dots, \lambda^{t-1}).$$

We just need to prove it for

$$\lambda^+ = (\mu; \lambda^0, \lambda^1, \dots, \lambda^{i-1}, \lambda^i \cup \{\square_i\}, \lambda^{i+1}, \dots, \lambda^{t-1}).$$

Let $0 \leq i', i'' \leq t-1$ satisfy $i' \equiv i+k \pmod{t}$ and $i'' \equiv i-k \pmod{t}$. Set $x_{i,w_i} = c_{\square_i}$. By Lemma 5.4, we have

$$\begin{aligned} & \sum_{\substack{\square \in \lambda^+ \\ h_{\square} \equiv k \pmod{t}}} h_{\square}^2 + \sum_{\substack{\square \in \lambda^+ \\ h_{\square} \equiv t-k \pmod{t}}} h_{\square}^2 - \sum_{\substack{\square \in \lambda \\ h_{\square} \equiv k \pmod{t}}} h_{\square}^2 - \sum_{\substack{\square \in \lambda \\ h_{\square} \equiv t-k \pmod{t}}} h_{\square}^2 \\ = & \sum_{0 \leq l \leq m_{i'}} (tx_{i,w_i} + b_i - tx_{i',l} - b_{i'})^2 - \sum_{1 \leq l \leq m_{i'}} (tx_{i,w_i} + b_i - ty_{i',l} - b_{i'})^2 \\ & + \sum_{0 \leq l \leq m_{i''}} (tx_{i,w_i} + b_i - tx_{i'',l} - b_{i''})^2 - \sum_{1 \leq l \leq m_{i''}} (tx_{i,w_i} + b_i - ty_{i'',l} - b_{i''})^2 \\ = & (tx_{i,w_i} + b_i - b_{i'})^2 - 2t(tx_{i,w_i} + b_i - b_{i'}) \left(\sum_{0 \leq l \leq m_{i'}} x_{i',l} - \sum_{1 \leq l \leq m_{i'}} y_{i',l} \right) \\ & + t^2 \left(\sum_{0 \leq l \leq m_{i'}} x_{i',l}^2 - \sum_{1 \leq l \leq m_{i'}} y_{i',l}^2 \right) \\ & + (tx_{i,w_i} + b_i - b_{i''})^2 - 2t(tx_{i,w_i} + b_i - b_{i''}) \left(\sum_{0 \leq l \leq m_{i''}} x_{i'',l} - \sum_{1 \leq l \leq m_{i''}} y_{i'',l} \right) \\ & + t^2 \left(\sum_{0 \leq l \leq m_{i''}} x_{i'',l}^2 - \sum_{1 \leq l \leq m_{i''}} y_{i'',l}^2 \right) \end{aligned}$$

$$= (tx_{i,w_i} + b_i - b_{i'})^2 + (tx_{i,w_i} + b_i - b_{i''})^2 + 2t^2 n_{i'} + 2t^2 n_{i''}.$$

The last equality is due to Theorem 3.1(1). On the other hand, by Lemma 5.2 we have

$$\begin{aligned} & \sum_{\substack{\square \in \lambda^+ \\ c_{\square} \equiv k \pmod{t}}} c_{\square}^2 + \sum_{\substack{\square \in \lambda^+ \\ c_{\square} \equiv t-k \pmod{t}}} c_{\square}^2 - \left(\sum_{\substack{\square \in \lambda \\ c_{\square} \equiv k \pmod{t}}} c_{\square}^2 + \sum_{\substack{\square \in \lambda \\ c_{\square} \equiv t-k \pmod{t}}} c_{\square}^2 \right) \\ &= (tx_{i,w_i} + b_i - i')^2 + (tx_{i,w_i} + b_i - i'')^2. \end{aligned}$$

By Theorem 3.1(3) we have

$$q_3(\lambda^i \cup \{\square_i\}) - q_3(\lambda^i) = 6x_{i,w_i}.$$

Finally, after putting those identities together we know the theorem is also true for λ^+ and thus prove the claim. \square

We now turn to a theorem still focused on one partition at a time, but this time summing over contents or hook lengths divisible by t .

Theorem 6.2. *Suppose that μ is a given t -core partition. For every partition λ with $\lambda_{t\text{-core}} = \mu$ we have*

$$\begin{aligned} & \sum_{\substack{\square \in \lambda \\ h_{\square} \equiv 0 \pmod{t}}} h_{\square}^2 - \sum_{\substack{\square \in \lambda \\ c_{\square} \equiv 0 \pmod{t}}} c_{\square}^2 \\ &= t^2 \sum_{i=0}^{t-1} \left(n_i^2 - d_i^2 n_i - \frac{1}{3} d_i q_3(\lambda^i) \right) - \sum_{\substack{\square \in \mu \\ c_{\square} \equiv 0 \pmod{t}}} c_{\square}^2, \end{aligned}$$

where $n_i = |\lambda^i|$ for $0 \leq i \leq t-1$.

Proof. We will prove this result by induction on the size of λ . Notice that $\mu(0) = \{h \in \mathcal{H}(\mu) \mid h \equiv 0 \pmod{t}\} = \emptyset$ since μ is a t -core partition. When $\lambda = \mu$, we have $\lambda^i = \emptyset$, $n_i = 0$ and thus the identity holds. Now suppose that the identity holds for

$$\lambda = (\mu; \lambda^0, \lambda^1, \dots, \lambda^{i-1}, \lambda^i, \lambda^{i+1}, \dots, \lambda^{t-1}).$$

We just need to prove it for

$$\lambda^+ = (\mu; \lambda^0, \lambda^1, \dots, \lambda^{i-1}, \lambda^i \cup \{\square_i\}, \lambda^{i+1}, \dots, \lambda^{t-1}).$$

Let $x_{i,w_i} = c_{\square_i}$. By Lemma 5.4 we have

$$\begin{aligned} & \sum_{\substack{\square \in \lambda^+ \\ h_{\square} \equiv 0 \pmod{t}}} h_{\square}^2 - \sum_{\substack{\square \in \lambda \\ h_{\square} \equiv 0 \pmod{t}}} h_{\square}^2 \\ &= t^2 + \sum_{0 \leq l \leq m_i} (tx_{i,w_i} - tx_{i,l})^2 - \sum_{1 \leq l \leq m_i} (tx_{i,w_i} - ty_{i,l})^2 \\ &= t^2 + (tx_{i,w_i})^2 - 2t^2 x_{i,w_i} \left(\sum_{0 \leq l \leq m_i} x_{i,l} - \sum_{1 \leq l \leq m_i} y_{i,l} \right) \\ &\quad + t^2 \left(\sum_{0 \leq l \leq m_i} x_{i,l}^2 - \sum_{1 \leq l \leq m_i} y_{i,l}^2 \right) \\ &= t^2 + t^2 x_{i,w_i}^2 + 2t^2 n_i. \end{aligned}$$

The last equality is due to Theorem 3.1(1). Also, by Lemma 5.2 we have

$$\sum_{\substack{\square \in \lambda^+ \\ c_\square \equiv 0 \pmod{t}}} c_\square^2 - \sum_{\substack{\square \in \lambda \\ c_\square \equiv 0 \pmod{t}}} c_\square^2 = t^2(x_{i,w_i} + d_i)^2.$$

By Theorem 3.1(3) we have

$$q_3(\lambda^i \cup \{\square_i\}) - q_3(\lambda^i) = 6x_{i,w_i}.$$

Finally, putting these identities together, we prove the claim. \square

It is clear that Theorem 1.4 is a consequence of Theorems 6.1 and 6.2 by letting $\mu = \emptyset$.

Theorem 1.6 is on t -Plancherel averages of square sums of congruent hook lengths.

Proof of Theorem 1.6. For two partitions

$$\lambda = (\mu; \lambda^0, \lambda^1, \dots, \lambda^{i-1}, \lambda^i, \lambda^{i+1}, \dots, \lambda^{t-1})$$

and

$$\lambda^+ = (\mu; \lambda^0, \lambda^1, \dots, \lambda^{i-1}, \lambda^i \cup \{\square_i\}, \lambda^{i+1}, \dots, \lambda^{t-1}),$$

by Case (ii) of Lemma 5.4 we have

$$\begin{aligned} & \sum_{\substack{\square \in \lambda^+ \\ h_\square \equiv k \pmod{t}}} h_\square^2 + \sum_{\substack{\square \in \lambda^+ \\ h_\square \equiv t-k \pmod{t}}} h_\square^2 - \sum_{\substack{\square \in \lambda \\ h_\square \equiv k \pmod{t}}} h_\square^2 - \sum_{\substack{\square \in \lambda \\ h_\square \equiv t-k \pmod{t}}} h_\square^2 \\ &= \sum_{0 \leq l \leq m_{i'}} (tc_{\square_i} + b_i - tx_{i',l} - b_{i'})^2 - \sum_{1 \leq l \leq m_{i'}} (tc_{\square_i} + b_i - ty_{i',l} - b_{i'})^2 \\ & \quad + \sum_{0 \leq l \leq m_{i''}} (tc_{\square_i} + b_i - tx_{i'',l} - b_{i''})^2 - \sum_{1 \leq l \leq m_{i''}} (tc_{\square_i} + b_i - ty_{i'',l} - b_{i''})^2 \end{aligned}$$

where $i' \equiv i + k \pmod{t}$ and $i'' \equiv i - k \pmod{t}$.

Let

$$P(n) := \sum_{\substack{\lambda_{t\text{-core}} = \mu \\ |\lambda/\mu| = nt}} F_{\lambda/\mu} G_\lambda \left(\sum_{\substack{\square \in \lambda \\ h_\square \equiv k \pmod{t}}} h_\square^2 + \sum_{\substack{\square \in \lambda \\ h_\square \equiv t-k \pmod{t}}} h_\square^2 \right).$$

Then, by Lemma 2.2 and Theorems 3.1, 3.2 we have

$$\begin{aligned} & P(n+1) - P(n) \\ &= \frac{1}{t} \sum_{\substack{\lambda_{t\text{-core}} = \mu \\ |\lambda/\mu| = nt}} F_{\lambda/\mu} G_\lambda \sum_{i=0}^{t-1} \sum_{\lambda^{i+} = \lambda^i \cup \{\square_i\}} \frac{H_{\lambda^i}}{H_{\lambda^{i+}}} \\ & \quad \left(\sum_{0 \leq l \leq m_{i'}} (tc_{\square_i} + b_i - tx_{i',l} - b_{i'})^2 - \sum_{1 \leq l \leq m_{i'}} (tc_{\square_i} + b_i - ty_{i',l} - b_{i'})^2 \right. \\ & \quad \left. + \sum_{0 \leq l \leq m_{i''}} (tc_{\square_i} + b_i - tx_{i'',l} - b_{i''})^2 - \sum_{1 \leq l \leq m_{i''}} (tc_{\square_i} + b_i - ty_{i'',l} - b_{i''})^2 \right) \\ &= \frac{1}{t} \sum_{\substack{\lambda_{t\text{-core}} = \mu \\ |\lambda/\mu| = nt}} F_{\lambda/\mu} G_\lambda \sum_{i=0}^{t-1} \sum_{\lambda^{i+} = \lambda^i \cup \{\square_i\}} \frac{H_{\lambda^i}}{H_{\lambda^{i+}}} \end{aligned}$$

$$\begin{aligned}
& \left(2t^2 c_{\square_i}^2 + 2t^2 |\lambda^{i'}| + 2t^2 |\lambda^{i''}| + (b_i - b_{i'})^2 + (b_i - b_{i''})^2 \right) \\
&= 2tn + 2tn + 2tn + \frac{1}{t} \sum_{i=0}^{t-1} \left((b_i - b_{i'})^2 + (b_i - b_{i''})^2 \right).
\end{aligned}$$

On the other hand,

$$\begin{aligned}
& \frac{1}{t} \sum_{i=0}^{t-1} \left((b_i - b_{i'})^2 + (b_i - b_{i''})^2 \right) \\
&= \frac{2}{t} \sum_{(i,j) \in B_k} (b_i - b_j)^2 \\
&= \frac{2}{t} \sum_{(i,j) \in B_k} \left(t^2 (d_i - d_j)^2 + (i - j)^2 + 2t(i - j)(d_i - d_j) \right) \\
&= 2k(t - k) + 4t(|\mu(k)| + |\mu(t - k)|).
\end{aligned}$$

The last equality is due to (4.2) and Lemma 4.2. Finally,

$$P(n + 1) - P(n) = 6nt + 2k(t - k) + 4t|\mu(k)| + 4t|\mu(t - k)|.$$

Thus,

$$\begin{aligned}
P(n) &= 6t \binom{n}{2} + (2k(t - k) + 4t|\mu(k)| + 4t|\mu(t - k)|)n + P(0) \\
&= 6t \binom{n}{2} + (2k(t - k) + 4t|\mu(k)| + 4t|\mu(t - k)|)n \\
&\quad + \left(\sum_{\substack{\square \in \mu \\ h_{\square} \equiv k \pmod{t}}} h_{\square}^2 + \sum_{\substack{\square \in \mu \\ h_{\square} \equiv t - k \pmod{t}}} h_{\square}^2 \right). \quad \square
\end{aligned}$$

Theorem 1.7 is analogous to Theorem 1.6 when looking at t -Plancherel averages of square sums of hook lengths, but only the lengths divisible by t are taken into account.

Proof of Theorem 1.7. For two partitions

$$\lambda = (\mu; \lambda^0, \lambda^1, \dots, \lambda^{i-1}, \lambda^i, \lambda^{i+1}, \dots, \lambda^{t-1})$$

and

$$\lambda^+ = (\mu; \lambda^0, \lambda^1, \dots, \lambda^{i-1}, \lambda^i \cup \{\square_i\}, \lambda^{i+1}, \dots, \lambda^{t-1}),$$

by Case (i) of Lemma 5.4 we have

$$\sum_{\substack{\square \in \lambda^+ \\ h_{\square} \equiv 0 \pmod{t}}} h_{\square}^2 - \sum_{\substack{\square \in \lambda \\ h_{\square} \equiv 0 \pmod{t}}} h_{\square}^2 = t^2 + t^2 c_{\square_i}^2 + 2t^2 n_i.$$

Let

$$P(n) := \sum_{\substack{\lambda_{t\text{-core}} = \mu \\ |\lambda/\mu| = nt}} F_{\lambda/\mu} G_{\lambda} \sum_{\substack{\square \in \lambda \\ h_{\square} \equiv 0 \pmod{t}}} h_{\square}^2.$$

Then,

$$\begin{aligned}
P(n+1) - P(n) &= \frac{1}{t} \sum_{\substack{\lambda_{t\text{-core}}=\mu \\ |\lambda/\mu|=nt}} F_{\lambda/\mu} G_{\lambda} \sum_{i=0}^{t-1} \sum_{\lambda^{i+}=\lambda^i \cup \{\square_i\}} \frac{H_{\lambda}}{H_{\lambda^{i+}}} (t^2 + t^2 c_{\square_i}^2 + 2t^2 n_i) \\
&= t^2 + 3t \sum_{i=0}^{t-1} n_i \\
&= t^2 + 3nt.
\end{aligned}$$

Thus, we have

$$P(n) = nt^2 + 3t \binom{n}{2} + P(0) = nt^2 + 3t \binom{n}{2}. \quad \square$$

Finally, the next two theorems have similar concerns, looking at t -Plancherel averages, but this time for squares of contents. They are differentiated by the restriction in Theorem 6.3 or looking only at contents in one congruence class.

Theorem 6.3. *Suppose that μ is a given t -core partition and $0 \leq k \leq t-1$. For each $0 \leq i \leq t-1$ let i' satisfy $0 \leq i' \leq t-1$ and $i - i' \equiv k \pmod{t}$. Then*

$$\sum_{\substack{\lambda_{t\text{-core}}=\mu \\ |\lambda/\mu|=nt}} F_{\lambda/\mu} G_{\lambda} \sum_{\substack{\square \in \lambda \\ c_{\square} \equiv k \pmod{t}}} c_{\square}^2 = t \binom{n}{2} + \frac{1}{t} \sum_{i=0}^{t-1} (b_i - i')^2 n + \sum_{\substack{\square \in \mu \\ c_{\square} \equiv k \pmod{t}}} c_{\square}^2.$$

Proof. For two partitions

$$\lambda = (\mu; \lambda^0, \lambda^1, \dots, \lambda^{i-1}, \lambda^i, \lambda^{i+1}, \dots, \lambda^{t-1})$$

and

$$\lambda^+ = (\mu; \lambda^0, \lambda^1, \dots, \lambda^{i-1}, \lambda^i \cup \{\square_i\}, \lambda^{i+1}, \dots, \lambda^{t-1}),$$

by Lemma 5.2 we have

$$\sum_{\substack{\square \in \lambda^+ \\ c_{\square} \equiv k \pmod{t}}} c_{\square}^2 - \sum_{\substack{\square \in \lambda \\ c_{\square} \equiv k \pmod{t}}} c_{\square}^2 = (tc_{\square_i} + b_i - i')^2.$$

Let

$$P(n) := \sum_{\substack{\lambda_{t\text{-core}}=\mu \\ |\lambda/\mu|=nt}} F_{\lambda/\mu} G_{\lambda} \sum_{\substack{\square \in \lambda \\ c_{\square} \equiv k \pmod{t}}} c_{\square}^2.$$

Then, we have

$$\begin{aligned}
P(n+1) - P(n) &= \frac{1}{t} \sum_{\substack{\lambda_{t\text{-core}}=\mu \\ |\lambda/\mu|=nt}} F_{\lambda/\mu} G_{\lambda} \sum_{i=0}^{t-1} \sum_{\lambda^{i+}=\lambda^i \cup \{\square_i\}} \frac{H_{\lambda}}{H_{\lambda^{i+}}} (tc_{\square_i} + b_i - i')^2 \\
&= tn + \frac{1}{t} \sum_{i=0}^{t-1} (b_i - i')^2.
\end{aligned}$$

Finally, we obtain

$$P(n) = t \binom{n}{2} + \frac{1}{t} \sum_{i=0}^{t-1} (b_i - i')^2 n + P(0)$$

$$= t \binom{n}{2} + \frac{1}{t} \sum_{i=0}^{t-1} (b_i - i')^2 n + \sum_{\substack{\square \in \mu \\ c_{\square} \equiv k \pmod{t}}} c_{\square}^2. \quad \square$$

Theorem 6.4. *Suppose that μ is a given t -core partition. Then we have*

$$\sum_{\substack{\lambda_{t\text{-core}} = \mu \\ |\lambda/\mu| = nt}} F_{\lambda/\mu} G_{\lambda} \sum_{\square \in \lambda} c_{\square}^2 = t^2 \binom{n}{2} + \frac{(t^3 - t)n}{6} + 2tn|\mu| + \sum_{\square \in \mu} c_{\square}^2.$$

Proof. By Theorem 6.3 we have

$$\begin{aligned} & \sum_{\substack{\lambda_{t\text{-core}} = \mu \\ |\lambda/\mu| = nt}} F_{\lambda/\mu} G_{\lambda} \sum_{\square \in \lambda} c_{\square}^2 \\ &= t^2 \binom{n}{2} + \frac{1}{t} \sum_{i=0}^{t-1} \sum_{j=0}^{t-1} (b_i - j)^2 n + \sum_{\square \in \mu} c_{\square}^2 \\ &= t^2 \binom{n}{2} + \frac{(t^3 - t)n}{6} + 2tn|\mu| + \sum_{\square \in \mu} c_{\square}^2. \end{aligned}$$

The last equality is due to (4.3), (4.4) and (4.5). \square

7. ACKNOWLEDGMENTS

The authors thank the referees for carefully reading our paper and for giving helpful comments. P.-O.D. and H.X. acknowledge support from grant PP00P2_138906 of the Swiss National Science Foundation. H.X. also acknowledges support of a Forschungskredit FK-14-093 from the University of Zürich.

REFERENCES

- [1] T. Amdeberhan, Differential operators, shifted parts, and hook lengths, *Ramanujan J.* **24**(3) (2011), 259–271.
- [2] D. Armstrong, C. Hanusa and B. Jones, Results and conjectures on simultaneous core partitions, *European J. Combin.* **41** (2014), 205–220.
- [3] J. Bandlow, An elementary proof of the hook formula, *Electron. J. Combin.* **15** (2008), research paper 45.
- [4] K. Carde, J. Loubert, A. Potechin, and A. Sanborn, Proof of Han’s Hook Expansion Conjecture, preprint; [arXiv:0808.0928](#).
- [5] J. S. Frame, G. de B. Robinson, and R. M. Thrall, The hook graphs of S_n , *Canad. J. Math.* **6** (1954), 316–324.
- [6] C. Greene, A. Nijenhuis and H. S. Wilf, A probabilistic proof of a formula for the number of Young tableaux of a given shape, *Adv. in Math.* **31** (1979), no. 1, 104–109.
- [7] C. Greene, A. Nijenhuis and H. S. Wilf, Another probabilistic method in the theory of Young tableaux, *J. Combin. Theory Ser. A* **37** (1984), no. 2, 127–135.
- [8] G.-N. Han, The Nekrasov-Okounkov hook length formula: refinement, elementary proof, extension, and applications, *Ann. Inst. Fourier* **60** (2010), no. 1, 1–29.
- [9] G.-N. Han, Some conjectures and open problems on partition hook lengths, *Experimental Mathematics* **18** (2009) 97–106.
- [10] G.-N. Han, Hook lengths and shifted parts of partitions, *Ramanujan J.* **23**(1-3) (2010), 127–135.
- [11] G.-N. Han and H. Xiong, Difference operators for partitions and some applications, preprint; [arXiv:1508.00772](#).

- [12] G.-N. Han and H. Xiong, New hook-content formulas for strict partitions (extended abstract), *Proceedings of the 28th International Conference on Formal Power Series and Algebraic Combinatorics, Disc. Math. and Theoretical Computer Science, 2016*, to appear.
- [13] G.-N. Han and H. Xiong, Polynomiality of some hook-content summations for doubled distinct and self-conjugate partitions, preprint; [arXiv:1601.04369](https://arxiv.org/abs/1601.04369).
- [14] G. James and A. Kerber, *The Representation Theory of the Symmetric Group*, Encyclopedia of Mathematics and its Applications, 16. Addison-Wesley Publishing Co., Reading, Mass., 1981.
- [15] D. Knuth, *The Art of Computer Programming, Vol. 3: Sorting and Searching*, Addison-Wesley, London, 1973, pp. 54–58.
- [16] I. G. Macdonald, *Symmetric functions and Hall polynomials*, Oxford Mathematical Monographs, The Clarendon Press, Oxford University Press, New York, second edition, 1995.
- [17] N. A. Nekrasov and A. Okounkov, Seiberg-Witten theory and random partitions, in The unity of mathematics, *Progress in Mathematics* **244**, Birkhäuser Boston, 2006, pp. 525–596.
- [18] G. Olshanski, Anisotropic Young diagrams and infinite-dimensional diffusion processes with the Jack parameter, *Int. Math. Res. Not. IMRN* **6** (2010), 1102–1166.
- [19] G. Olshanski, Plancherel averages: Remarks on a paper by Stanley, *Electron. J. Combin.* **17** (2010), research paper 43.
- [20] G. Panova, Polynomiality of some hook-length statistics, *Ramanujan J.* **27** (2012), no. 3, 349–356.
- [21] G.-C. Rota, On the foundations of combinatorial theory: I. Theory of Möbius functions, *Z. Wahrscheinlichkeitstheorie und Verw. Gebiete* **2** (1964), 349–356.
- [22] R. P. Stanley, Some combinatorial properties of hook lengths, contents, and parts of partitions, *Ramanujan J.* **23(1-3)** (2010), 91–105.
- [23] R. P. Stanley, Differential posets, *J. Amer. Math. Soc.* **1** (1988), no. 4, 919–961.
- [24] R. P. Stanley, *Enumerative Combinatorics*, vol. 2, Cambridge University Press, New York/Cambridge, 1999.

(Paul-Olivier Dehaye) I-MATH, UNIVERSITÄT ZÜRICH, WINTERTHURERSTRASSE 190, ZÜRICH 8057, SWITZERLAND

E-mail address: paul-olivier.dehaye@math.uzh.ch

(Guo-Niu Han) I.R.M.A., UMR 7501, UNIVERSITÉ DE STRASBOURG ET CNRS, 7 RUE RENÉ DESCARTES, F-67084 STRASBOURG, FRANCE

E-mail address: guoniu.han@unistra.fr

(Huan Xiong) I-MATH, UNIVERSITÄT ZÜRICH, WINTERTHURERSTRASSE 190, ZÜRICH 8057, SWITZERLAND

E-mail address, Corresponding author: huan.xiong@math.uzh.ch