

WEIGHTED DERANGEMENTS AND LAGUERRE POLYNOMIALS

BY

DOMINIQUE FOATA AND DORON ZEILBERGER

1. Introduction. — One of the harbingers of the current combinatorization of the theory of special functions was a remarkable result of GILLIS and EVEN [7] that gave a certain combinatorial interpretation to the linearization coefficients of the *simple* Laguerre polynomials $L_n(x)$. Let $(L_n^{(\alpha)}(x))$ be the sequence of the (general) Laguerre polynomials that may be defined by their generating function

$$(1.1) \quad \sum_{n=0}^{\infty} L_n^{(\alpha)}(x) u^n = (1-u)^{-\alpha-1} \exp \frac{-xu}{1-u},$$

the simple Laguerre polynomials being defined by $L_n(x) = L_n^0(x)$ ($\alpha = 0$). Furthermore, for each positive integer m and each sequence (n_1, \dots, n_m) of nonnegative integers let

$$(1.2) \quad A(n_1, \dots, n_m; \alpha) = (-1)^{n_1 + \dots + n_m} \int_0^{\infty} \left(\prod_{i=1}^m L_{n_i}^{(\alpha)}(x) \right) x^{\alpha} e^{-x} dx$$

and

$$(1.3) \quad I(n_1, \dots, n_m; \alpha) = \frac{1}{\Gamma(\alpha+1)} n_1! \dots n_m! A(n_1, \dots, n_m; \alpha).$$

Then GILLIS and EVEN [7] found a combinatorial interpretation for $I(n_1, \dots, n_m; 0)$ ($\alpha = 0$) and deduced from that interpretation the fact that $I(n_1, \dots, n_m; 0)$ was positive. The positivity property was immediately reproved by ASKEY and his followers [1, 2, 3, 8] by means of analytical methods and reincluded in a more general special function set-up. As they noticed, the generating function (1.1) yields the identity :

$$(1.4) \quad \sum A(n_1, \dots, n_m; \alpha) x_1^{n_1} \dots x_m^{n_m} = \frac{\Gamma(\alpha+1)}{(1 - \sigma_2 - 2\sigma_3 - \dots - (m-1)\sigma_m)^{\alpha+1}},$$

where n_1, \dots, n_m runs over all nonnegative integers and σ_j denotes the j -th elementary symmetric function in x_1, \dots, x_m . As it was shown by ASKEY and his coauthors, the positivity of $I(n_1, \dots, n_m; \alpha)$ is an immediate consequence of (1.4) and it holds for $\alpha > -1$. In [1, p. 857–858] the authors were very close to finding a combinatorial interpretation of $I(n_1, \dots, n_m; \alpha)$ for an arbitrary α . It is the purpose of this paper to provide one by taking up again the combinatorial model introduced by GILLIS and EVEN [7] and “ α -extending” it. Let $\mathcal{P}(n_1, \dots, n_m)$ be the set of permutations on the $n_1 + \dots + n_m$ elements

$$a_{1,1}, \dots, a_{1,n_1}; \dots; a_{m,1}, \dots, a_{m,n_m}$$

and denote by $\mathcal{D}(n_1, \dots, n_m)$ the subset of $\mathcal{P}(n_1, \dots, n_m)$ consisting of what we will call (a_1, \dots, a_m) -derangements. These are permutations where no element is allowed to go to one of its kind; in other words, columns of the form

$$\begin{array}{ccc} a_{1,i} & \dots & a_{m,i} \\ a_{1,j} & \dots & a_{m,j} \end{array}$$

are forbidden in the two-line representation of the permutation. The following identity is due to GILLIS and EVENS [7] :

$$(1.5) \quad I(n_1, \dots, n_m; 0) = \text{card } \mathcal{D}(n_1, \dots, n_m).$$

In order to have an extension for *any* α introduce for each permutation π its number of cycles $\text{cyc } \pi$ and define its *weight* by

$$(1.6) \quad w(\pi) = (\alpha + 1)^{\text{cyc } \pi}.$$

The polynomial

$$D(n_1, \dots, n_m; \alpha) = \sum_{\pi} w(\pi) \quad (\pi \in \mathcal{D}(n_1, \dots, n_m))$$

reduces to $\text{card } \mathcal{D}(n_1, \dots, n_m)$ when $\alpha = 0$. Our α -analog of the Gillis-Even result now reads :

THEOREM 1. — *For each indeterminate α one has :*

$$(1.7) \quad I(n_1, \dots, n_m; \alpha) = D(n_1, \dots, n_m; \alpha).$$

As all the weights in $D(n_1, \dots, n_m; \alpha)$ are nonnegative whenever $\alpha > -1$, then the forementioned theorem implies that $I(n_1, \dots, n_m; \alpha)$ is also nonnegative, as was proved by ASKEY et al. [1, 2].

We give two proofs of THEOREM 1. The first one is based on the combinatorial interpretation of the Laguerre polynomials. The product (1.3) is expanded and the various terms interpreted in terms of generating polynomials for permutations and injections (see sections 2 and 3). The second one relies on an β -analog of the celebrated MACMAHON's Master Theorem [9, p. 97]. This β -analog has an interest of its own and will be discussed in section 4. The second proof is then completed in section 5.

2. Cycles. — We will need two results that are fundamental in the current combinatorial interpretation of special functions. First, the generating function for the set $\mathcal{P}(n)$ of all the permutations on n elements by number of cycles is given by (see, e.g., [10, p. 78])

$$(2.1) \quad w(\mathcal{P}(n)) = \sum_{\pi} w(\pi) = (\alpha+1)_n = (\alpha+1)(\alpha+2) \dots (\alpha+n) \quad (\pi \in \mathcal{P}(n)).$$

Let $1 \leq k \leq n$ and S be a $(n-k)$ -element subset of the n -element $[n]$. The set of *injections* from S into $[n]$ will be denoted by $\text{Inj}(S, n)$. An injection from S to $[n]$ consists of a (possibly empty) collection of cycles within S and some paths that wander in S , but terminate at a point outside S . Similarly, denote by $\text{cyc } \pi$ the number of cycles of π and define its *weight* by $w(\pi) = (\alpha+1)^{\text{cyc } \pi}$.

For example, if $S = \{1, 2, 3, 4, 5, 6\}$ and $n = 9$, then $(1, 3)$, (2) , $4 \rightarrow 5 \rightarrow 7$, $6 \rightarrow 8$ is an injection with weight $(\alpha+1)^2$.

The result analogous to (1.1) reads (see [6, lemma 2.1]) : if $\text{card } S = n-k$, then

$$(2.2) \quad w(\text{Inj}(S, n)) = \sum_{\pi} w(\pi) = (\alpha+1+k)_{n-k} \quad (\pi \in \text{Inj}(S, n)).$$

3. Proof of theorem 1. — By the definition of the Laguerre polynomials

$$n_i! L_{n_i}^{(\alpha)}(x) = \sum_{k_i=0}^{n_i} (-1)^{k_i} \binom{n_i}{k_i} (\alpha+1+k_i)_{n_i-k_i} x^{k_i} \quad (i = 1, \dots, m).$$

In (1.2) and (1.3) expand each Laguerre polynomial and integrate term by term using the fact that $(1/\Gamma(\alpha+1)) \int_0^\infty e^{-x} x^{n+\alpha} dx = (\alpha+1)_n$. This leads to :

$$(3.1) \quad I(n_1, \dots, n_m; \alpha) = \sum_{k_1=0}^{n_1} \dots \sum_{k_m=0}^{n_m} (\alpha+1)_{k_1+\dots+k_m} \prod_{i=1}^m (-1)^{n_i-k_i} \binom{n_i}{k_i} (\alpha+1+k_i)_{n_i-k_i}.$$

Consider now *any* permutation in $\mathcal{P}(n_1, \dots, n_m)$ and write it in *cycle form*. An element will be called *incestuous*, if it is sent to one of its own kind. Denote by $\text{Inc } \pi$ the set of all incestuous elements of π .

For example, in the permutation belonging to $\mathcal{P}(4, 5, 5)$

$$(a_1 b_1 b_2 c_3 a_2)(c_2 c_1 b_3 a_3)(c_4 b_4 a_4 c_5)$$

the elements b_1, a_2, c_2, c_5 are the incestuous elements.

Now call *marked permutation* an ordered pair (π, S) with π a permutation and S a subset of $\text{Inc } \pi$ and denote by $\mathcal{M}(n_1, \dots, n_m)$ the set of *marked permutations*.

For example,

$$(a_1 \overline{b_1} b_2 c_3 a_2)(\overline{c_2} c_1 b_3 a_3)(c_4 b_4 a_4 \overline{c_5})$$

is the marked permutation where the incestuous elements b_1, c_2, c_5 are marked ($S = \{b_1, c_2, c_5\}$) while the incestuous element a_2 is not marked.

Define the *weight* of each marked permutation (π, S) by

$$w'(\pi, S) = (-1)^{\text{card } S} (\alpha + 1)^{\text{cyc } \pi}$$

and consider the generating polynomial for marked permutations :

$$M(n_1, \dots, n_m; \alpha) = \sum_{\pi} w'(\pi) \quad (\pi \in \mathcal{M}(n_1, \dots, n_m)).$$

This generating polynomial will be computed in two different ways. One of these ways yields

$$(3.2) \quad M(n_1, \dots, n_m; \alpha) = D(n_1, \dots, n_m; \alpha).$$

The other way will give :

$$(3.3) \quad M(n_1, \dots, n_m; \alpha) = I(n_1, \dots, n_m; \alpha).$$

Consider the following weight preserving sign changing involution on marked permutations. Look at the first incestuous element. If it is marked, unmark it ; if it is unmarked, mark it. Of course, the number of cycles of the permutation does not change (since there is no change in the underlying permutation!). Only the parity of the number of marks changes, reversing the sign of the weight. Accordingly, all the terms in $M(n_1, \dots, n_m; \alpha)$ corresponding to permutations with incestuous elements can be arranged in mutually cancelling pairs and their sum is therefore zero. All that remains in $M(n_1, \dots, n_m; \alpha)$ are the terms corresponding to those marked permutations (π, S) containing no incestuous elements and so $S = \emptyset$.

But those marked permutations are simply the pairs (π, \emptyset) with $\pi \in \mathcal{D}(n_1, \dots, n_m)$ and they satisfy : $w'(\pi, \emptyset) = w(\pi)$. Therefore (3.2) holds.

Now compute $M(n_1, \dots, n_m; \alpha)$ in a different way. For each $i = 1, \dots, m$ let S_i be a certain subset of $\{a_{i,1}, \dots, a_{i,n_i}\}$ of cardinality $n_i - k_i$ and denote by $\mathcal{M}(\mathbf{S})$ the subset of $\mathcal{M}(n_1, \dots, n_m)$ consisting of marked permutations (π, S) with $S = S_1 \cup \dots \cup S_m$. Also let $\mathcal{P}(\mathbf{S})$ the set of all π in $\mathcal{P}(n_1, \dots, n_m)$ such that $(\pi, S) \in \mathcal{M}(\mathbf{S})$. Clearly,

$$(3.4) \quad w'(\mathcal{M}(\mathbf{S})) = (-1)^{n_1 - k_1 + \dots + n_m - k_m} w(\mathcal{P}(\mathbf{S})).$$

LEMMA. — One has

$$(3.5) \quad w(\mathcal{P}(\mathbf{S})) = (\alpha + 1)_{k_1 + \dots + k_m} \prod_{i=1}^m (\alpha + 1 + k_i)_{n_i - k_i}.$$

Proof. — From (2.1) and (2.2) it follows that the right-hand side of (3.5) is the generating function for the product

$$\mathcal{P}(k_1, \dots, k_m) \times \prod_{i=1}^m \text{Inj}(S_i, n_i)$$

by w . To prove the lemma it then suffices to construct a w -weight preserving bijection $\pi \mapsto (\pi_1, \dots, \pi_m, \sigma)$ of $\mathcal{P}(\mathbf{S})$ onto that product. Write π in cycle form. Then in each cycle of π delete all the elements of $S = S_1 \cup \dots \cup S_m$. What remains is a permutation written in cycle form. Call it σ .

To get π_i take all the cycles of π consisting *only* of elements of S_i . Take also the connected portions of S_i lying in other cycles. Doing this will result in a certain number of *paths* that wander through S_i but terminate in an element not in S_i .

Clearly, σ belongs to $\mathcal{P}(k_1, \dots, k_m)$ and each π_i is an injection of S_i into $\{a_{i,1}, \dots, a_{i,n_i}\}$. Moreover, the total number of cycles of $\sigma, \pi_1, \dots, \pi_m$ is equal to $\text{cyc } \pi$. Thus, the mapping is w -preserving. The reverse construction is immediate. \square

Example. — Take

$$\begin{aligned} n_1 &= 6 & n_2 &= 6 & n_3 &= 6 \\ k_1 &= 3 & k_2 &= 3 & k_3 &= 3 \\ S_1 &= \{a_1, a_2, a_3\} & S_2 &= \{b_1, b_2, b_3\} & S_3 &= \{c_1, c_2, c_3\} \\ \pi &= (a_1 a_2)(a_4 b_1 b_5 a_5 a_3)(b_2 b_4 c_1 c_4 c_3 c_2 c_6)(c_5 a_6)(b_3)(b_6) \end{aligned}$$

Then

$$\begin{aligned}\sigma &= (a_4 b_5 a_5)(b_4 c_4 c_6)(c_5 c_6)(b_6) \\ \pi_1 &= (a_1 a_2), \quad a_3 \rightarrow a_4 \\ \pi_2 &= (b_3), \quad b_1 \rightarrow b_5, \quad b_2 \rightarrow b_4 \\ \pi_3 &= c_1 \rightarrow c_4, \quad c_3 \rightarrow c_2 \rightarrow c_6\end{aligned}$$

It follows from (3.4) and (3.5) that

$$w'(\mathcal{M}(\mathbf{S})) = (-1)^{n_1 - k_1 + \dots + n_m - k_m} (\alpha + 1)_{k_1 + \dots + k_m} \prod_{i=1}^m (\alpha + 1 + k_i)_{n_i - k_i}.$$

Hence

$$\begin{aligned}M(n_1, \dots, n_m; \alpha) &= \sum w'(\mathcal{M}(\mathbf{S})) \\ &= \sum_{k_1=0}^{n_1} \dots \sum_{k_m=0}^{n_m} \prod_{i=1}^m \binom{n_i}{k_i} \sum w'(\mathcal{M}(\mathbf{S})) \\ &= \sum_{k_1=0}^{n_1} \dots \sum_{k_m=0}^{n_m} (\alpha + 1)_{k_1 + \dots + k_m} \\ &\quad \times \prod_{i=1}^m (-1)^{n_i - k_i} \binom{n_i}{k_i} (\alpha + 1 + k_i)_{n_i - k_i},\end{aligned}$$

which is the expression found for $I(n_1, \dots, n_m; \alpha)$ in (3.1). Therefore, (3.3) is proved and also THEOREM 1.

4. The β -analog of the MacMahon Master Theorem. — Let D be the determinant $\det(\delta_{ij} - b(i, j)x_j)$ ($1 \leq i, j \leq m$). The MacMahon Master Theorem asserts that the coefficient of $x_1^{n_1} \dots x_m^{n_m}$ in the expansion of D^{-1} is equal to the coefficient of $x_1^{n_1} \dots x_m^{n_m}$ in the product

$$(4.1) \quad (b(1, 1)x_1 + \dots + b(1, m)x_m)^{n_1} \dots (b(m, 1)x_1 + \dots + b(m, m)x_m)^{n_m}.$$

It will be convenient to restate this statement in a slightly different form. Let $\mathcal{R}(n_1, \dots, n_m)$ denote the set of all the rearrangements

$$r = r(1, 1) \dots r(1, n_1) \dots r(m, 1) \dots r(m, n_m)$$

of the word $1^{n_1} \dots m^{n_m}$ and let

$$v(r) = \prod_{i,j} b(i, r(i, j)) \quad (1 \leq i \leq m; 1 \leq j \leq n_i).$$

Clearly, the coefficient of $x_1^{n_1} \dots x_m^{n_m}$ in (4.1) is equal to the sum of all the $v(r)$ with r running over all the rearrangements of $1^{n_1} \dots m^{n_m}$.

Next, consider a permutation π belonging to $\mathcal{P}(n_1, \dots, n_m)$ (defined in section 1). If π sends (i, j) over (i', j') , write $i' = c\pi(i, j)$. Furthermore, define

$$v(\pi) = \prod_{i,j} b(i, c\pi(i, j)) \quad (1 \leq i \leq m; 1 \leq j \leq n_i).$$

To each rearrangement r in $\mathcal{R}(n_1, \dots, n_m)$ there correspond exactly $n_1! \dots n_m!$ permutations π in $\mathcal{P}(n_1, \dots, n_m)$ with the property that $v(\pi) = v(r)$. Therefore, the coefficient of $x_1^{n_1} \dots x_m^{n_m}$ in (4.1) is also equal to

$$(4.2) \quad n_1! \dots n_m! \sum_{\pi} v(\pi) \quad (\pi \in \mathcal{P}(n_1, \dots, n_m)).$$

The MacMahon Master identity can then be restated as

$$(4.3) \quad \sum \frac{x_1^{n_1}}{n_1!} \dots \frac{x_m^{n_m}}{n_m!} v(\mathcal{P}(n_1, \dots, n_m)) = D^{-1}.$$

Next define the β -weight $v(\beta; \pi)$ of each permutation π in $\mathcal{P}(n_1, \dots, n_m)$ by

$$v(\beta; \pi) = \beta^{\text{cyc } \pi} v(\pi).$$

THEOREM (β -analog of the MacMahon Master Theorem). — *The following identity holds :*

$$(4.4) \quad \sum \frac{x_1^{n_1}}{n_1!} \dots \frac{x_m^{n_m}}{n_m!} v(\beta; \mathcal{P}(n_1, \dots, n_m)) = D^{-\beta}.$$

Proof. — Consider the partitional complex (see, e.g., [4, 5]) of the permutations and denote by $\mathcal{CP}(n_1, \dots, n_m)$ the subset of the *connected* permutations in $\mathcal{P}(n_1, \dots, n_m)$. As the weight $v(\beta; \cdot)$ is *multiplicative* [4, 5], the following identity holds :

$$(4.5) \quad \begin{aligned} \sum \frac{x_1^{n_1}}{n_1!} \dots \frac{x_m^{n_m}}{n_m!} v(\beta; \mathcal{P}(n_1, \dots, n_m)) \\ = \exp \sum \frac{x_1^{n_1}}{n_1!} \dots \frac{x_m^{n_m}}{n_m!} v(\beta; \mathcal{CP}(n_1, \dots, n_m)). \end{aligned}$$

From (4.3) and (4.5) with $\beta = 1$

$$\sum \frac{x_1^{n_1}}{n_1!} \dots \frac{x_m^{n_m}}{n_m!} v(\mathcal{CP}(n_1, \dots, n_m)) = -\text{Log } D.$$

But $v(\beta; \pi) = \beta v(\pi)$ for every connected permutation, so that

$$\sum \frac{x_1^{n_1}}{n_1!} \cdots \frac{x_m^{n_m}}{n_m!} v(\beta; \mathcal{CP}(n_1, \dots, n_m)) = -\beta \text{Log } D,$$

and finally (4.4) holds because of (4.5). \square

5. Second proof of theorem 1. — As was noted by ASKEY [1, 2], (1.4) is an immediate consequence of (1.1). Rewriting (1.4) for $\alpha = 0$ using GILLIS-EVEN's result (1.5) yields :

$$(5.1) \quad \sum \text{card } \mathcal{D}(n_1, \dots, n_m) \frac{x_1^{n_1}}{n_1!} \cdots \frac{x_m^{n_m}}{n_m!} = (1 - \sigma_2 - 2\sigma_3 - \cdots - (m-1)\sigma_m)^{-1}.$$

Take again the v -weight of section 4 with $b(i, j) = \delta_{ij}$. Then,

$$(5.2) \quad \text{card } \mathcal{D}(n_1, \dots, n_m) = v(\mathcal{P}(n_1, \dots, n_m)).$$

Thus for this particular v identity (4.4) holds with $\beta = \alpha + 1$ and

$$D = (1 - \sigma_2 - 2\sigma_3 - \cdots - (m-1)\sigma_m).$$

On the account of (1.2), (1.3), (1.4) and (4.4) with this particular weight v we conclude that (1.7) also holds.

REFERENCES

- [1] ASKEY (Richard) and ISMAIL (Mourad E.H.). — Permutation problems and special functions, *Canad. J. Math.*, t. **28**, 1976, p. 853–874.
- [2] ASKEY (Richard), ISMAIL (Mourad E.H.) and KOORNWINDER (T.). — Weighted permutation problems and Laguerre polynomials, *J. Combinatorial Theory, Ser. A*, t. **25**, 1978, p. 277–287.
- [3] ASKEY (Richard) and GASPER (G.). — Convolution structures for Laguerre polynomials, *J. d'Anal. Math.*, t. **31**, 1977, p. 46–48.
- [4] FOATA (Dominique). — *La série génératrice exponentielle dans les problèmes d'énumération*. — Montréal, Presses de l'Université de Montréal, 1974.
- [5] FOATA (Dominique) et SCHÜTZENBERGER (Marcel-Paul). — *Théorie géométrique des polynômes eulériens*. — Berlin, Springer-Verlag, 1970 (*Lecture Notes in Math.*, **138**).
- [6] FOATA (Dominique) and STREHL (Volker). — Combinatorics of Laguerre polynomials, *Enumeration and Design* [Waterloo. June–July 1982 : D.M. Jackson and S.A. Vanstone, eds.], p. 123–140. — Toronto, Academic Press, 1984.
- [7] GILLIS (J.) and EVEN (S.). — Derangements and Laguerre polynomials, *Proc. Cambridge Phil. Soc.*, t. **79**, 1976, p. 135–143.
- [8] ISMAIL (Mourad E.H.) and TAMHANKAR (M.V.). — A combinatorial approach to some positivity problems, *S.I.A.M. J. Math. Anal.*, t. **10**, 1979, p. 478–485.

WEIGHTED DERANGEMENTS

- [9] MACMAHON (Percy Alexander). — *Combinatory Analysis*, vol. 1. — Cambridge, Univ. Press, 1915. (Reprinted by Chelsea, New York, 1955).
- [10] RIORDAN (John). — *An Introduction to Combinatorial Analysis*. — New York, John Wiley, 1959.

The present paper has been completed and updated to become :

Dominique Foata and Doron Zeilberger, Laguerre polynomials, weighted derangements and positivity, *SIAM J. Discrete Math.*, vol. **1**, 1988, p. 425–433.

Dominique FOATA,
Département de mathématique,
Université Louis-Pasteur,
7, rue René-Descartes,
F-67084 Strasbourg, France.

and

Doron ZEILBERGER,
Department of Mathematics,
Drexel University,
Philadelphia, Pennsylvania 19104, U.S.A.
now at the address :
Temple University,
Philadelphia, Pennsylvania 19122, U.S.A.