

**q-SERIES IN COMBINATORICS;
PERMUTATION STATISTICS**

(Preliminary version)

initiated in 2003 ; revised version : March 2021

Dominique Foata and Guo-Niu Han

Dominique Foata
Institut Lothaire
1, Rue Murner
F-67000 Strasbourg, France
foata@unistra.fr

or

Guo-Niu Han
I.R.M.A. UMR 7501
Université Louis Pasteur et CNRS
7, rue René-Descartes
F-67084 Strasbourg, France
guoniu@unistra.fr

© These Lecture Notes cannot be reproduced without permission of the authors. Please, send remarks and corrections to: foata@unistra.fr or guoniu@unistra.fr.

Table of Contents

- Introduction
- 1. The q -binomial theorem
- 2. Mahonian Statistics
 - 2.1. The inv-coding
 - 2.2. The maj-coding
 - 2.3. The den-coding
- 3. The algebra of q -binomial coefficients
- 4. The q -binomial structures
 - 4.1. Partitions of integers
 - 4.2. Nondecreasing sequences of integers
 - 4.3. Binary words
 - 4.4. Ordered Partitions into two blocks
- 5. The q -multinomial coefficients
- 6. The MacMahon Verfahren
- 7. A refinement of the MacMahon Verfahren
- 8. The Euler-Mahonian polynomials
 - 8.1. A finite difference q -calculus
 - 8.2. A q -iteration method
- 9. The Insertion technique
- 10. The two classes of q -Eulerian polynomials
- 11. Major Index and Inversion Number
 - 11.1. How to construct a bijection
 - 11.2. The binary case
 - 11.3. From the binary to the general case
 - 11.4. Further properties of the transformation
 - 11.5. Applications to permutations
- 12. Major and Inverse Major Indices
 - 12.1. The biword expansion
 - 12.2. Another application of the MacMahon Verfahren
- 13. A four-variable distribution
- 14. Symmetric Functions
 - 14.1. Partitions of integers
 - 14.2. The algebra of symmetric functions
 - 14.3. The classical bases
- 15. The Schur Functions

16. The Cauchy Identity
17. The Combinatorial definition of the Schur Functions
18. The inverse ligne route of a standard tableau
19. The Robinson-Schensted correspondence
 - 19.1. The Schensted-Knuth algorithm
 - 19.2. A combinatorial proof of the Cauchy identity
 - 19.3. Geometric properties of the correspondence
 - 19.4. A permutation statistic distribution
20. Eulerian Calculus; the first extensions
 - 20.1. The signed permutations
 - 20.2. Pairs of permutations
 - 20.3. The q -extension
 - 20.4. The t, q -maj extension for signed permutations
 - 20.5. A first inversion number for signed permutations
21. Eulerian Calculus; the analytic choice
 - 21.1. Inversions for signed permutations
 - 21.2. Basic Bessel Functions
 - 21.3. The iterative method
22. Eulerian Calculus; finite analogs of Bessel functions
 - 22.1. Signed biwords
 - 22.2. Signed bipermutations
 - 22.3. Signed biwords and compatible bipermutations
 - 22.4. The last specializations
23. Eulerian Calculus; multi-indexed polynomials
 - 23.1. The bi-indexed Eulerian polynomials
 - 23.2. The Désarménien Verfahren
 - 23.3. Congruences for bi-indexed polynomials
 - 23.4. The signed Eulerian Numbers
24. The basic and bibasic trigonometric Functions
 - 24.1. The basic and bibasic tangent and secant functions
 - 24.2. Alternating permutations
 - 24.3. Combinatorics of the bibasic secant and tangent
25. MacMahon's Master Theorem revisited
 - 25.1. The "Master Theorem" identity
 - 25.2. Circuit Calculus
 - 25.3. "Master Theorem" within the algebra \mathcal{A}^{MM}
 - 25.4. Further matricial properties of the algebras \mathcal{A}^{CF} and \mathcal{A}^{GLZ}
26. The decrease value theorem
 - 26.1. The first fundamental transformation for arbitrary words revisited
 - 26.2. Horizontal derangements
 - 26.3. Vertical derangements
 - 26.4. End of the proof of Theorem 26.1

27. The Decrease Value Theorem; from words to permutations

27.2. Lyndon words

27.3. The Gessel-Reutenauer standardization

27.4. A further property of the bijection $w \mapsto (c, \sigma)$

27.5. Generating functions for permutations

27.6. A quadruple q -series for permutations

Examples and Exercises

Answers to the Exercises

Notes

References

Introduction

The inverse Laplace transform maps each formal power series $\sum_{n \geq 0} a(n)u^n$ in one variable u into another series $\sum_{n \geq 0} (a(n)/n!)u^n$, whose coefficient of order n is *normalized* by the factor $n!$ We then obtain the so-called *exponential* generating function for the sequence $(a(n))$ ($n \geq 0$). The normalization has numerous advantages: the derivative is obtained by a simple shift of the coefficients; the exponential of a series can be explicitly calculated; there are closed formulas for the *exponential* generating functions for several classical orthogonal polynomials, . . . However the algebra of exponential generating functions cannot be regarded as the universal panacea. Further kinds of series are needed, for instance to express some generating series for the symmetric groups by certain statistics.

In the middle of the eighteenth century Heine introduced a new class of series in which the normalized factor $n!$ was replaced by a polynomial of degree n in another variable, more precisely, the series where the coefficient of order n is normalized by the polynomial denoted by $(q; q)_n$, in another variable q , defined by

$$(0.1) \quad (q; q)_n := \begin{cases} 1, & \text{if } n = 0; \\ (1 - q)(1 - q^2) \cdots (1 - q^n), & \text{if } n \geq 1. \end{cases}$$

The algebra of those series has been largely developed by Jackson in the beginning of the twentieth century. It has then fallen into disuse, except perhaps in the field of Partition Theory, but has vigorously come back in several mathematical domains, in particular in the theory of Quantum Groups and naturally in Combinatorics.

Those series have been named *q-series*. They are simply formal power series in two variables, say, u and q , where the latter variable q , used for the normalization, plays a privileged role. Let $\Omega[[u, q]]$ denote the algebra of formal series in two variables u and q , with coefficients in a ring Ω . Each element of that algebra can be expressed as a series

$$(0.2) \quad a = \sum_{n \geq 0, m \geq 0} a(n, m) u^n q^m,$$

INTRODUCTION

where, for each ordered pair (n, m) , the symbol $a(n, m)$ belongs to Ω . Such a series can be seen as a series *in one variable* u , with coefficients in the ring $\Omega[[q]]$ of series in one variable q , *i. e.*,

$$(0.3) \quad a = \sum_{n \geq 0} u^n \left(\sum_{m \geq 0} a(n, m) q^m \right).$$

For each integer $n \geq 0$ the expression $(q; q)_n$ is a polynomial in q , which is invertible in $\Omega[[q]]$, since its constant coefficient is 1. Thus the series a can be rewritten as

$$(0.4) \quad a = \sum_{n \geq 0} \frac{u^n}{(q; q)_n} b(n; q),$$

where $b(n; q)$ is the formal series in the variable q

$$(0.5) \quad b(n; q) := (q; q)_n \cdot \left(\sum_{m \geq 0} a(n, m) q^m \right).$$

Each formal series a written in the form (0.4) is called a *q-series*. The coefficient $u^n / (q; q)_n$ is then a formal series in the unique variable q .

The purpose of this memoir is to give a basic description of the algebra of those series and describe the use that has been made of them in Combinatorics, in particular for expressing the generating functions for certain statistics defined on permutations, words, multipermutations, signed permutations and other finite structures. It has been customary to regroup all the techniques that have been developed under the name of *Permutation Statistic Study*, even though the group of permutations is not the only group structure involved. The statistics themselves can be uni- or several-variable, or even set-valued. As will be seen, the *q-series* enter into the picture in a very natural way.

The *q-binomial theorem*, which is stated and proved in the first section, is the basic tool in *q-Calculus*. It opens the door to all the *q-series* identities and also gives rise to two expansions of the *q-exponential*, as a *q-series* itself, and also as an infinite product.

The polynomial $(q; q)_n$, defined above, is next studied in a combinatorial context. This leads to a discussion of the so-called *Mahonian* statistics, especially the *Major Index* and the *Inversion Number* that will play an essential role in this memoir. One of our goals, indeed, is to try to understand why the so-called *natural q-analogs* of various numbers or polynomials can be derived by means of either one of those statistics.

The Major Index is strongly related to the combinatorial theory of the representation of finite groups, particularly when dealing with various tableaux (standard, semi-standard, ...). The inversion number requires

other techniques, in particular the so-called q -iteration method, that will be used on several occasions.

The q -binomial coefficients or Gaussian polynomials that appear in many identities on q -series are studied in several combinatorial environments, as is done in section 4. With the study of the q -multinomial coefficient we are led to introduce the statistic “number of inversions” for classes of permutations with repetitions. We prefer to use the term “rearrangement” (of a given finite sequence) or “word.” This is the content of section 5.

The *MacMahon Verfahren*, introduced in section 6, is the first tool that makes possible the transcription of properties of certain statistics defined on the symmetric group or on some classes of rearrangements to the algebra of q -series. As a first application, it is shown that the Major Index has the same distribution as the number of inversions on each class of rearrangements.

A careful study of the *MacMahon Verfahren* serves to find a q -extension of the traditional Eulerian polynomials, namely the *Euler-Mahonian polynomials* $A_{\mathbf{m}}(t, q)$, that appear to be generating polynomials for classes of rearrangements by the bivariable statistic (des, maj) . The statistic “des” is the *number of descents* that has been studied in several combinatorial contexts and “maj” is the Major Index.

As shown in section 8, there are four equivalent definitions of the Euler-Mahonian polynomials. The proofs of those equivalences are based on fundamental techniques in q -Calculus, finite difference and iterative methods. The insertion technique that looks so natural when dealing with univariable statistics on the symmetric group becomes intricate for several-variable statistics. A *marked word* technique is presented in section 9 and appears to be successful for deriving a recurrence relation for the Euler-Mahonian polynomials $A_{\mathbf{m}}(t, q)$.

When the class of rearrangements is reduced to the symmetric group, the Euler-Mahonian polynomials become the so-called *q -maj-Eulerian polynomials* ${}^{\text{maj}}A_n(t, q)$, as they form a first q -analog of the traditional Eulerian polynomials $A_n(t)$ in one variable t . However, when the exponential generating function for the latter polynomials is q -analogized in a proper way, another q -analog of those polynomials, namely the q -inv Eulerian polynomial ${}^{\text{inv}}A_n(t, q)$ appears. As shown in the notations, “inv” plays for ${}^{\text{inv}}A_n(t, q)$ the role that “maj” does for ${}^{\text{maj}}A_n(t, q)$.

The *Major Index* and the *Inversion Number*, that can be defined for each rearrangement of a given finite sequence of integers, are equidistributed over each rearrangement class. Section 11 contains the construction of a bijection Φ of the class onto itself such that $\text{inv } \Phi(w) = \text{maj } w$.

INTRODUCTION

The bijection has several other properties, in particular when the underlying class is the symmetric group.

With section 12 we start the study of permutation statistics that involve both the permutation and its inverse. Besides “maj” we are led to introduce “imaj” that is nothing but the Major Index of the *inverse* permutation. We then see the first occurrence of the classical infinite product $\prod_{i \geq 1, j \geq 1} \frac{1}{1 - ux_i y_j}$, that is to be expanded, once the substitutions $x_i \leftarrow q_1^{i-1}$, $y_j \leftarrow q_2^{j-1}$ are made. The resulting series are series in two bases, normalized by denominators of the form $(q_1; q_1)_n (q_2; q_2)_n$.

In section 13 a further extension of the MacMahon *Verfahren* leads to the derivation of the distribution of a four-variable statistic on the symmetric group.

The theory of symmetric functions hides too many useful identities and too many combinatorial algorithms not to appear in this memoir. In particular, the infinite product mentioned above has a celebrated expansion in terms of products of *Schur functions*. As those functions have a handy combinatorial interpretation—as shown in section 17—it was essential to give the main properties of those functions. This is the content of sections 15, 16, and 17.

In the expansion of a Schur function we find monomials that are coded by the so-called *semi-standard tableaux*. In their turn, those tableaux can be further coded by *standard tableaux* and sequences of numbers. This coding has several applications. In particular, it serves to express a Schur function, in which variables are replaced by powers of a variable q , as a generating function for standard tableaux by a certain statistic. This is the content of section 18.

In section 19 we find an overview of the *Robinson-Schensted correspondence* that enables the transfer of geometric properties on tableaux to analogous properties on permutations. As an application, a bibasic generating function for polynomials in several variables defined on symmetric groups is derived.

The next four sections 20, 21, 22 and 23 deal with *Eulerian Calculus*, that is, the study of geometric properties of the Eulerian polynomials and its various extensions. By “extensions” we mean three aspects: (i) extension to the group of the *signed permutations*, (ii) q -extension, that is, the introduction of a suitable Mahonian statistic “inv” or “maj,” (iii) the study of generating polynomials for *pairs* or finite sequences of permutations or signed permutations. The combination of those three extensions leads to the combinatorial study of some Bessel functions, q -Bessel functions and finite analogs of Bessel functions.

The crucial step in Eulerian Calculus is to find the appropriate q -analog for the generating polynomial for the signed permutations by their number of descents. Our *analytic choice* (see section 21) forces us to find a suitable definition of inversions for signed permutations. This leads to a coherent study of all the extensions described above. Notice that the *length*, as defined in the theory of Coxeter groups, does not conduct to an elegant derivation in the algebra of q -series.

In our last section on Eulerian Calculus (section 23) we introduce the bi-indexed Eulerian polynomials and explain how the Désarménien *Verfahren* makes it possible the study of congruences of those polynomials with respect to the cyclotomic polynomials. The section ends with a short study on signed Eulerian Numbers.

Next, a combinatorial study of the basic trigonometric functions, especially the tangent and secant functions is developed. The coefficients in their q -expansions are the generating polynomials for the so-called alternating permutations by number of inversions. The same combinatorial set-up is used to interpret the coefficients in the p, q -expansions of the bibasic tangent and secant.

We end these Lecture Notes with an exposé on the celebrated “Master Theorem” in its main three versions: the commutative one à la MacMahon, the partially commutative one à la Cartier-Foata and the quantum one à la Garoufalidis-Lê-Zeilberger. Finally, the Decrease Value Theorem, that comes next, may be regarded as a true extension of the previous theorem, as it makes it possible to calculate the distributions of several multivariable permutations statistics.

1. The q -Binomial Theorem

Take up again the notations (0.3)–(0.5). When, for each $n \geq 0$, the ratio $b(n+1; q)/b(n; q)$ is a *rational fraction* in q^n , equal to 1 for $q = 0$ and such that $b(0; q) = 1$, we get what is called a *basic hypergeometric series*. In the analytic expression of such a series the following notation is used that extends the notation (0.1): for each element ω in the ring define the *q -ascending factorial in ω* by

$$(1.1) \quad (\omega; q)_n := \begin{cases} 1, & \text{if } n = 0; \\ (1 - \omega)(1 - \omega q) \dots (1 - \omega q^{n-1}), & \text{if } n \geq 1; \end{cases}$$

in its *finite* version and

$$(1.2) \quad (\omega; q)_\infty := \lim_n (\omega; q)_n = \prod_{n \geq 0} (1 - \omega q^n);$$

in its *infinite* version.

1. THE q -BINOMIAL THEOREM

When the underlying ring Ω is the complex field, the rational fraction $b(n+1; q)/b(n; q)$ can be written as

$$\frac{b(n+1; q)}{b(n; q)} = \frac{(1 - \alpha_1 q^n) \dots (1 - \alpha_r q^n)}{(1 - \beta_1 q^n) \dots (1 - \beta_s q^n)},$$

where $\alpha_1, \dots, \alpha_r, \beta_1, \dots, \beta_s$ are complex numbers. By iteration,

$$\begin{aligned} b(n; q) &= \frac{b(n; q)}{b(n-1; q)} \dots \frac{b(2; q)}{b(1; q)} \frac{b(1; q)}{b(0; q)} \\ &= \frac{(1 - \alpha_1 q^{n-1}) \dots (1 - \alpha_r q^{n-1})}{(1 - \beta_1 q^{n-1}) \dots (1 - \beta_s q^{n-1})} \\ &\quad \times \dots \times \frac{(1 - \alpha_1 q) \dots (1 - \alpha_r q)}{(1 - \beta_1 q) \dots (1 - \beta_s q)} \frac{(1 - \alpha_1) \dots (1 - \alpha_r)}{(1 - \beta_1) \dots (1 - \beta_s)}, \end{aligned}$$

that is

$$(1.3) \quad b(n; q) = \frac{(\alpha_1; q)_n \dots (\alpha_r; q)_n}{(\beta_1; q)_n \dots (\beta_s; q)_n}.$$

For an arbitrary ring Ω the expression of $b(n; q)$ given in (1.3) is taken as a definition. As each q -ascending factorial $(\beta_i; q)_n$ has a constant coefficient equal to 1, it is invertible in the ring $\Omega[[q]]$, so that $b(n; q)$ as shown in (1.3) is well-defined. Then call *basic hypergeometric series* each q -series of the form

$$(1.4) \quad {}_r\varphi_s \left(\begin{matrix} \alpha_1, \dots, \alpha_r \\ \beta_1, \dots, \beta_s \end{matrix}; q, u \right) := \sum_{n \geq 0} \frac{(\alpha_1; q)_n \dots (\alpha_r; q)_n}{(\beta_1; q)_n \dots (\beta_s; q)_n} \frac{u^n}{(q; q)_n}.$$

Such a series can be defined in each algebra $\Omega[[u, q]]$ of formal series in two variables u and q , whatever the underlying ring Ω is. When $r = 0$ (resp. $s = 0$, resp. $r = 0$ and $s = 0$), the following notations are used: ${}_0\varphi_s \left(\begin{matrix} \text{---} \\ \beta_1, \dots, \beta_s \end{matrix}; q, u \right)$ (resp. ${}_r\varphi_0 \left(\begin{matrix} \alpha_1, \dots, \alpha_r \\ \text{---} \end{matrix}; q, u \right)$, resp. ${}_0\varphi_0 \left(\begin{matrix} \text{---} \\ \text{---} \end{matrix}; q, u \right)$).

In the q -binomial theorem which is stated next, the series ${}_1\varphi_0 \left(\begin{matrix} \alpha \\ \text{---} \end{matrix}; q, u \right)$ has a closed expression in terms of an infinite product.

Theorem 1.1 (q -Binomial Theorem). *The following identity holds:*

$$(1.5) \quad \sum_{n \geq 0} (\alpha; q)_n \frac{u^n}{(q; q)_n} = \frac{(\alpha u; q)_\infty}{(u; q)_\infty} = \prod_{n \geq 0} \frac{1 - \alpha u q^n}{1 - u q^n};$$

or, equivalently:

$$(1.6) \quad {}_1\varphi_0 \left(\begin{matrix} \alpha \\ \text{---} \end{matrix}; q, u \right) = \frac{(\alpha u; q)_\infty}{(u; q)_\infty}.$$

Before giving the proof of the theorem it matters to make several remarks.

(a) The *order* $o(a)$ of a formal series $a = \sum_{n \geq 0, m \geq 0} a(n, m) u^n q^m$ is defined to be the smallest integer $k \geq 0$ such that the polynomial

$$\sum_{n+m=k} a(n, m) u^n q^m,$$

called the *homogeneous polynomial of degree k of a* (in u and q), is not zero. Consider a countable family (a_s) ($s \geq 0$) of formal series in two variables. As for the series in one variable, it is readily seen that if the order of a_s tends to infinity with s , then the infinite product $\prod_{s \geq 0} (1 - a_s)$ is a well-defined series. In the infinite product $(\alpha u; q)_\infty$, with $\alpha \neq 0$, the term $\alpha u q^n$ is a series (reduced to a monomial) of order $(n + 1)$. As $o(\alpha u q^n)$ tends to infinity with n , the infinite product $(\alpha u; q)_\infty$ is well-defined. The same property holds for the product $(u; q)_\infty$ occurring in the denominator.

(b) Within the coefficient $(\alpha; q)_n / (q; q)_n$ of u^n in formula (1.5), make the substitution $\alpha \leftarrow q^\alpha$ to obtain $(q^\alpha; q)_n / (q; q)_n$; then let q tend to 1. We get $(\alpha)_n / n!$. The ratio $(q^\alpha; q)_n / (q; q)_n$ is said to be the *q -analog* of the ascending factorial $(\alpha)_n / n!$, where

$$(\alpha)_n = \begin{cases} 1, & \text{if } n = 0; \\ \alpha(\alpha + 1) \cdots (\alpha + n - 1), & \text{if } n \geq 1. \end{cases}$$

But the series $\sum_{n \geq 0} u^n (\alpha)_n / n!$ is the *hypergeometric series* ${}_1F_0(\alpha; -; u)$, that satisfies the identity

$$(1.7) \quad {}_1F_0(\alpha; -; u) = (1 - u)^{-\alpha},$$

which is known to be the *binomial identity*. Notice that (1.7) is used to extend the definition of $(1 - u)^{-\alpha}$ when α is not an integer (positive or negative). Identity (1.6) is said to be the *q -analog* of (1.7).

The main difference between (1.6) and (1.7) is the following: when dealing with the algebra of formal series, formula (1.7) is a *definition* of $(1 - u)^{-\alpha}$ whenever α is not an integer, while (1.6) is an *identity*.

However, when considering the series ${}_1F_0(\alpha; -; u)$ and ${}_1\varphi_0(\alpha; q; u)$ as power series of the *complex* variables u and q , the two formulas (1.6) and (1.7) are *identities*, if the moduli of u and q are less than 1. The first proof of (1.6) given below is directly inspired from the classical proof of (1.7) in the analytic case. However, the end of the proof uses a topological argument on formal series and not on analytic series.

1. THE q -BINOMIAL THEOREM

(c) In view of the proof of Theorem 1.1 let us mention the following identity, easy to derive:

$$(1.8) \quad (\alpha; q)_n - (\alpha q; q)_n = (\alpha q; q)_{n-1} \alpha (q^n - 1) \quad (n \geq 1).$$

There exist numerous relations on the q -ascending factorials. Among the most frequent let us quote the *associativity* property

$$(1.9) \quad (\alpha; q)_{n+k} = (\alpha; q)_n (\alpha q^n; q)_k$$

and the *reverse formula*

$$(1.10) \quad (\alpha^{-1} q^{1-n}; q)_n = (\alpha; q)_n (-\alpha^{-1})^n q^{-n(n-1)/2}.$$

No comment for (1.9). The latter can be proved as follows. For $n \geq 1$ we have

$$\begin{aligned} (\alpha; q)_n &= (1 - \alpha)(1 - \alpha q) \cdots (1 - \alpha q^{n-1}) \\ &= (-\alpha)(1 - \alpha^{-1})(-\alpha q)(1 - \alpha^{-1} q^{-1}) \cdots (-\alpha q^{n-1})(1 - \alpha^{-1} q^{-(n-1)}) \\ &= (-\alpha)^n q^{n(n-1)/2} (1 - \alpha^{-1})(1 - \alpha^{-1} q^{-1}) \cdots (1 - \alpha^{-1} q^{1-n}) \\ &= (-\alpha)^n q^{n(n-1)/2} (\alpha^{-1} q^{1-n}; q)_n, \end{aligned}$$

a formula that still holds for $n = 0$.

Finally, notice that the relation

$$(1.11) \quad (\alpha; q)_n = \frac{(\alpha; q)_\infty}{(\alpha q^n; q)_\infty}$$

can be used to define the q -ascending factorial $(\alpha; q)_n$ for *every real number* n .

(d) Let $\alpha = q^N$ in (1.5). We get

$$(1.12) \quad \frac{(q^N u; q)_\infty}{(u; q)_\infty} = (u; q)_N^{-1} = \sum_{n \geq 0} (q^N; q)_n \frac{u^n}{(q; q)_n}.$$

But, if u and q are regarded as complex variables with modulus less than 1 and if we let q tend to 1, we obtain the identity

$$(1 - u)^{-N} = \sum_{n \geq 0} (N)_n \frac{u^n}{n!},$$

that is, the usual binomial identity.

First proof of Theorem 1.1. Start with the series ${}_1\varphi_0\left(\begin{smallmatrix} \alpha \\ - \end{smallmatrix}; q, u\right) = \sum_{n \geq 0} u^n (\alpha; q)_n / (q; q)_n$ and evaluate the q -difference

$$\begin{aligned} {}_1\varphi_0\left(\begin{smallmatrix} \alpha \\ - \end{smallmatrix}; q, u\right) - {}_1\varphi_0\left(\begin{smallmatrix} \alpha \\ - \end{smallmatrix}; q, qu\right) &= \sum_{n \geq 1} \frac{(\alpha; q)_n}{(q; q)_n} u^n (1 - q^n) = \sum_{n \geq 1} \frac{(\alpha; q)_n}{(q; q)_{n-1}} u^n \\ &= (1 - \alpha)u \left(1 + \sum_{n \geq 2} \frac{(\alpha q; q)_{n-1}}{(q; q)_{n-1}} u^{n-1}\right) \\ (1.13) \qquad \qquad \qquad &= (1 - \alpha)u {}_1\varphi_0\left(\begin{smallmatrix} \alpha q \\ - \end{smallmatrix}; q, u\right). \end{aligned}$$

By using (1.8) we get:

$$\begin{aligned} {}_1\varphi_0\left(\begin{smallmatrix} \alpha \\ - \end{smallmatrix}; q, u\right) - {}_1\varphi_0\left(\begin{smallmatrix} \alpha q \\ - \end{smallmatrix}; q, u\right) &= \sum_{n \geq 1} \frac{(\alpha; q)_n - (\alpha q; q)_n}{(q; q)_n} u^n \\ &= -\alpha \sum_{n \geq 1} \frac{(\alpha q; q)_{n-1}}{(q; q)_{n-1}} u^n \\ (1.14) \qquad \qquad \qquad &= -\alpha u {}_1\varphi_0\left(\begin{smallmatrix} \alpha q \\ - \end{smallmatrix}; q, u\right). \end{aligned}$$

From (1.13) and (1.14) it follows that

$${}_1\varphi_0\left(\begin{smallmatrix} \alpha \\ - \end{smallmatrix}; q, u\right) = \frac{1 - \alpha u}{1 - u} {}_1\varphi_0\left(\begin{smallmatrix} \alpha \\ - \end{smallmatrix}; q, qu\right),$$

and by iteration

$$(1.15) \quad {}_1\varphi_0\left(\begin{smallmatrix} \alpha \\ - \end{smallmatrix}; q, u\right) = \frac{(\alpha u; q)_m}{(u; q)_m} {}_1\varphi_0\left(\begin{smallmatrix} \alpha \\ - \end{smallmatrix}; q, q^m u\right) \quad (m \geq 0).$$

If ${}_1\varphi_0\left(\begin{smallmatrix} \alpha \\ - \end{smallmatrix}; q, u\right)$ is considered as an analytic series of the complex variable u , it suffices to say that ${}_1\varphi_0\left(\begin{smallmatrix} \alpha \\ - \end{smallmatrix}; q, u\right)$ is continuous inside the unit disk. As the series is equal to 1 for $u = 0$, identity (1.6) follows from (1.15) by letting m tend to infinity.

With the topology of formal series we may use the following argument: consider a pair (i, j) of nonnegative integers such that $i + j \geq 1$. As soon as $m \geq j + 1$ the coefficients of $u^i q^j$ in

$$\frac{(\alpha u; q)_\infty}{(u; q)_\infty} \quad \text{and in} \quad \frac{(\alpha u; q)_m}{(u; q)_m}$$

are equal. But ${}_1\varphi_0\left(\begin{smallmatrix} \alpha \\ - \end{smallmatrix}; q, q^m u\right)$ is of the form $1 + q^m u a$, where a is a formal series in the two variables u and q . It follows that for every i the coefficients of $u^i q^j$ in

$$\frac{(\alpha u; q)_m}{(u; q)_m} \quad \text{and if} \quad \frac{(\alpha u; q)_m}{(u; q)_m} (1 + q^m u a) = {}_1\varphi_0\left(\begin{smallmatrix} \alpha \\ - \end{smallmatrix}; q, u\right)$$

are the same. Hence, identity (1.6) is proved. \square

1. THE q -BINOMIAL THEOREM

Second proof of Theorem 1.1. The right-hand side of identity (1.6) is a formal series that can be written as

$$b(u, q) := \sum_{n \geq 0} c_n(q) u^n,$$

where, for every $n \geq 0$, the coefficient $c_n(q)$ is a formal series in the variable q . But

$$b(u, q) = \frac{1 - \alpha u}{1 - u} \prod_{n \geq 0} \frac{(1 - \alpha u q q^n)}{(1 - u q q^n)} = \frac{1 - \alpha u}{1 - u} b(uq, q);$$

therefore

$$(1 - \alpha u)b(uq, q) = (1 - u)b(u, q),$$

and then

$$(1 - \alpha u) \sum_{n \geq 0} c_n(q) q^n u^n = (1 - u) \sum_{n \geq 0} c_n(q) u^n.$$

Looking for the coefficient of u^{n+1} on each side provides:

$$c_{n+1}(q)q^{n+1} - \alpha q^n c_n(q) = c_{n+1}(q) - c_n(q);$$

so that

$$c_{n+1}(q) = c_n(q) \frac{1 - \alpha q^n}{1 - q^{n+1}}.$$

As $c_0(q) = 1$, the right expression

$$c_n(q) = \frac{(\alpha; q)_n}{(q; q)_n}$$

is found by induction on n . \square

Let $\alpha = 0$ in (1.6). We get

$$(1.16) \quad \sum_{n \geq 0} \frac{u^n}{(q; q)_n} = \frac{1}{(u; q)_\infty}.$$

Now consider the infinite product $(-u; q)_\infty$ and again take the argument developed in the second proof. If we let $(-u; q)_\infty = \sum_{n \geq 0} c_n(q) u^n$, we find $c_{n+1}(q)q^{n+1} + c_n(q)q^n = c_{n+1}(q)$. Hence,

$$c_{n+1}(q) = \frac{q^n c_n(q)}{(1 - q^{n+1})}.$$

As $c_0(q) = 1$, we get

$$c_{n+1}(q) = \frac{q^{(n+1)n/2}}{(q; q)_{n+1}};$$

and then the identity

$$(1.17) \quad \sum_{n \geq 0} q^{n(n-1)/2} \frac{u^n}{(q; q)_n} = (-u; q)_\infty.$$

The two series appearing in (1.16) and (1.17) are respectively denoted by $e_q(u)$ and $E_q(u)$ and are referred to as the *first* and the *second q -exponential*.

2. Mahonian Statistics

For each integer $n \geq 0$ let S_n be a set of cardinality $n!$ (for example, the permutation group \mathfrak{S}_n). By *statistic* on S_n we simply mean a mapping $f : S_n \rightarrow \mathbb{N}$ with nonnegative integral values. The polynomial

$$a(n) := \sum_{s \in S_n} q^{f(s)}$$

is called the *generating polynomial for S_n by the statistic f* ; or, sometimes, the generating polynomial for f . The series

$$(2.1) \quad a := \sum_{n \geq 0} u^n \frac{a(n)}{(q; q)_n}$$

is called the *q -generating function* for the polynomials $(a(n))$ ($n \geq 0$). If, for each $n \geq 0$, the polynomial $a(n)$ has the form

$$(2.2) \quad a(n) = \frac{(q; q)_n}{(1 - q)^n} = (1 + q + \cdots + q^{n-1}) \cdots (1 + q + q^2)(1 + q),$$

we say that f is a *Mahonian* statistic on the family (S_n) ($n \geq 0$).

With each positive integer n is associated its *q -analog* defined by

$$(2.3) \quad [n]_q := \frac{1 - q^n}{1 - q} = 1 + q + q^2 + \cdots + q^{n-1}$$

and its *q -factorial*

$$(2.4) \quad \begin{aligned} [n]_q! &:= [n]_q [n-1]_q \cdots [2]_q [1]_q \\ &= \frac{(1 - q^n)}{(1 - q)} \frac{(1 - q^{n-1})}{(1 - q)} \cdots \frac{(1 - q^2)}{(1 - q)} \frac{(1 - q)}{(1 - q)} \\ &= \frac{(q; q)_n}{(1 - q)^n} \\ &= (1 + q + \cdots + q^{n-1})(1 + q + \cdots + q^{n-2}) \cdots (1 + q). \end{aligned}$$

Consequently, the generating polynomial defined in (2.2) is equal to the q -factorial of n and the q -generating function has the simple form

$$(2.5) \quad a = \sum_{n \geq 0} u^n \frac{1}{(q; q)_n} \frac{(q; q)_n}{(1 - q)^n} = \left(1 - \frac{u}{1 - q}\right)^{-1}.$$

2. MAHONIAN STATISTICS

In this section our purpose is to introduce several Mahonian statistics that are of constant use in the study of the q -series. Some of their properties are being derived.

The first one of these statistics is denoted by “tot” (“tot” for “total”). Although its definition is straightforward, it is very useful in many calculations. For each $n \geq 0$ let SE_n denote the set of the *subexcedent* sequences $x = (x_1, x_2, \dots, x_n)$. By *subexcedent* we mean a sequence of integers x_i , of length n , that satisfy the inequalities $0 \leq x_i \leq i - 1$ for all $i = 1, 2, \dots, n$. The cardinality of SE_n is of course $n!$ For each sequence $x = (x_1, x_2, \dots, x_n) \in SE_n$, define

$$(2.6) \quad \text{tot } x := x_1 + x_2 + \cdots + x_n.$$

Proposition 2.1. *The statistic “tot” on SE_n is Mahonian, that is, for each $n \geq 1$ we have:*

$$(2.7) \quad \sum_{x \in SE_n} q^{\text{tot } x} = \frac{(q; q)_n}{(1 - q)^n}.$$

Proof. The result is banal for $n = 1$. By induction on n :

$$\begin{aligned} \sum_{x \in SE_n} q^{\text{tot } x} &= \sum_{x' \in SE_{n-1}} q^{\text{tot } x'} \sum_{0 \leq x_n \leq n-1} q^{x_n} \\ &= \frac{(q; q)_{n-1}}{(1 - q)^{n-1}} (1 + q + \cdots + q^{n-1}) \\ &= \frac{(q; q)_n}{(1 - q)^n}. \quad \square \end{aligned}$$

The next three Mahonian statistics that are being introduced are defined on the permutation group \mathfrak{S}_n ; they are called the *Inversion Number* “inv”, the *Major Index* “maj” and the *Denert statistic* “den”.

Let $\sigma = \sigma(1) \dots \sigma(n)$ be a permutation, written as linear word. It is traditional to define the *Inversion Number*, $\text{inv } \sigma$, of the permutation σ as the number of ordered pairs of integers (i, j) such that $1 \leq i < j \leq n$ and $\sigma(i) > \sigma(j)$.

The *Major Index* $\text{maj } \sigma$ of σ is defined to be the *sum of the positions i where a descent $\sigma(i) > \sigma(i + 1)$ occurs*. We can also write

$$(2.8) \quad \text{maj } \sigma := \sum_{1 \leq i \leq n-1} i \chi\{\sigma(i) > \sigma(i + 1)\},$$

by making use of the χ -notation. Remember that for each statement A we write $\chi(A) = 1$ or 0 depending on whether A is true or not.

The definition of “den” is based on the notion of *cyclic interval*. Let i, j be two positive integers; the *cyclic interval* $\llbracket i, j \rrbracket$ is then

$$\llbracket i, j \rrbracket := \begin{cases}]i, j], & \text{if } i \leq j; \\ [1, j] +]i, +\infty[, & \text{if } i > j. \end{cases}$$

The *Denert statistic*, $\text{den } \sigma$, of the permutation σ is defined as the number of ordered pairs (i, j) such that $1 \leq i < j \leq n$ and $\sigma(i) \in \llbracket \sigma(j), j \rrbracket$.

To show that those three statistics are Mahonian on \mathfrak{S}_n , we construct three *bijections* $\sigma \mapsto x$ of \mathfrak{S}_n onto SE_n having the properties

$$\text{inv } \sigma = \text{tot } x, \quad \text{maj } \sigma = \text{tot } x, \quad \text{den } \sigma = \text{tot } x,$$

respectively. The construction of those three bijections makes use of three different *codings* of the permutations. The image x of σ under each of those bijections is called the *inv-coding*, the *maj-coding* and the *den-coding* of σ , respectively.

2.1. *The inv-coding (also called Lehmer coding).* Let $\sigma = \sigma(1) \dots \sigma(n)$ be a permutation. For each $i = 1, \dots, n$ define x_i as being the number of terms $\sigma(j)$ to the left of $\sigma(i)$ which are greater than $\sigma(i)$, that is,

$$x_i := \sum_{j:1 \leq j \leq i-1} \chi(\sigma(j) > \sigma(i)).$$

The sequence $x := (x_1, \dots, x_n)$ just defined is obviously subexcedent. Furthermore, the correspondence $\sigma \mapsto x$ is bijective. Moreover, the sum $\text{tot } x$ of the x_i 's is precisely equal to the *Inversion Number* $\text{inv } \sigma$ of the permutation σ .

In the following example, under every element $\sigma(i)$ is written the corresponding x_i of the inv-coding.

$$\begin{array}{cccccccccc} \sigma = & \left(\begin{array}{cccccccccc} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 \\ 7 & 1 & 5 & 4 & 9 & 2 & 6 & 3 & 8 \end{array} \right) \\ x = & \quad 0 & 1 & 1 & 2 & 0 & 4 & 2 & 5 & 1 \end{array}$$

In particular, $\text{inv } \sigma = \text{tot } x = 16$.

To reconstruct σ from its inv-coding x , proceed as follows: first let $\sigma(n) := n - x_n$. Once the elements $\sigma(k+1), \dots, \sigma(n)$ have been obtained, eliminate all the terms of the sequence $(n, n-1, \dots, 2, 1)$ equal to one of the $\sigma(l)$'s for a certain $l \geq k+1$. Then, $\sigma(k)$ is equal to be the $(x_k + 1)$ -st term of the sequence $(n, n-1, \dots, 2, 1)$, when reading that sequence from left to right.

2. MAHONIAN STATISTICS

For example, start from the subexcedent sequence

$$x = (0, 1, 1, 2, 0, 4, 2, 5, 1)$$

of length $n = 9$; first, get $\sigma(9) := n - x_n = 9 - 1 = 8$. Then, form the sequence $(9, 7, 6, 5, 4, 3, 2, 1)$, where 8 has been deleted. Then, $\sigma(8)$ is equal to the $(x_8 + 1) = (5 + 1) = 6$ -th term of the sequence, that is, $\sigma(8) := 3$. The running sequence becomes $(9, 7, 6, 5, 4, 2, 1)$, whose $(x_7 + 1) = (2 + 1) = 3$ -rd term is 6; hence, $\sigma(7) := 6$. Next, consider $(9, 7, 5, 4, 2, 1)$ whose $(x_6 + 1) = (4 + 1) = 5$ -th term is 2; then $\sigma(6) := 2$, and so on.

2.2. The maj-coding. Starting with a permutation $\sigma' \in \mathfrak{S}_{n-1}$, written as a word $\sigma'(1) \dots \sigma'(n-1)$, we can generate n permutations $\sigma \in \mathfrak{S}_n$ by inserting the letter n to the left of the word, or between two letters $\sigma'(i)$ and $\sigma'(i+1)$ for $1 \leq i \leq n-2$, or to the right of the word, say, *in position* $i = 0, 1, 2, \dots, (n-1)$, respectively. Thus, every permutation $\sigma \in \mathfrak{S}_n$ is characterized by an ordered pair (σ', i) , where $\sigma' \in \mathfrak{S}_{n-1}$ and $0 \leq i \leq n-1$. The surjection $\psi : \sigma \mapsto \sigma'$ of \mathfrak{S}_n onto \mathfrak{S}_{n-1} consists of removing the letter n from the word $\sigma = \sigma(1)\sigma(2) \dots \sigma(n)$.

For describing the maj-coding we *relabel* the n possible positions where n can be inserted into $\sigma' = \sigma'(1) \dots \sigma'(n-1)$ in the following manner: label $j = 0$ is given to the insertion of n to the right of the word σ' ; suppose that σ' has d descents, that is to say, d positions $\sigma'(i)\sigma'(i+1)$ such that $\sigma'(i) > \sigma'(i+1)$. We label those descents $j = 1, j = 2, \dots, j = d$, *from right to left*; the insertion of n to the left of σ' is given label $j = d+1$ and the labels $j = d+2, d+3, \dots, n-1$ are given to the $(n-2-d)$ insertions into the other positions when reading the word σ' *from left to right*.

If the letter n in the original permutation σ is in position j for the *relabelling* just described, we adopt the notations:

$$(2.9) \quad \begin{aligned} \sigma_{n-1} &:= \sigma' = \psi(\sigma); & x_n &:= j; \\ \sigma_n &:= \sigma := [\sigma_{n-1}, x_n]. \end{aligned}$$

In the same manner, to σ_{n-1} there corresponds a pair $[\sigma_{n-2}, x_{n-1}]$ and by iteration we obtain a sequence of pairs $[\sigma_{n-3}, x_{n-2}], \dots, [\sigma_0, x_1]$, where σ_0 is the void permutation and $x_1 = 0$. This yields a sequence, necessarily subexcedent $x = (x_1, x_2, \dots, x_n)$, that is called the *maj-coding* of σ .

Example. Consider the permutation $\sigma = 715492638$. The permutations $\sigma_1, \sigma_2, \dots, \sigma_8, \sigma_9$ are simply the *subwords* reduced to the letter 1, to the letters 1, 2, \dots , to the letters 1, 2, $\dots, 8$, finally to the letters 1, 2, $\dots, 9$.

At each step of the construction we have to *maj-label* the inserting positions as was indicated before. In Table 2.1 the maj-labellings appear as subscripts of the permutations written in the third column. The maj-coding of σ is then $x = (0, 0, 0, 2, 3, 5, 4, 0, 2)$. Notice that $\text{maj } \sigma = \text{tot } x = 16$.

Permutation	maj-coding	maj-labelling
$\sigma_1 = 1$	$x_1 = 0$	$1_1 0$
$\sigma_2 = 12$	$x_2 = 0$	$1_1 2_2 0$
$\sigma_3 = 123$	$x_3 = 0$	$1_1 2_2 3_3 0$
$\sigma_4 = 1423$	$x_4 = 2$	$2_2 1_3 4_1 2_4 3_0$
$\sigma_5 = 15423$	$x_5 = 3$	$3_3 1_4 5_2 4_1 2_5 3_0$
$\sigma_6 = 154263$	$x_6 = 5$	$4_4 1_5 5_3 4_2 2_6 6_1 3_0$
$\sigma_7 = 7154263$	$x_7 = 4$	$5_5 7_4 1_6 5_3 4_2 2_7 6_1 3_0$
$\sigma_8 = 71542638$	$x_8 = 0$	$5_5 7_4 1_6 5_3 4_2 2_7 6_1 3_8 8_0$
$\sigma_9 = 715492638$	$x_9 = 2$	

Table 2.1

To reconstruct the permutation σ from a subexcedent sequence $x = (x_1, x_2, \dots, x_n)$, we put $\sigma_1 := 1$, then obtain $\sigma_2 := [\sigma_1, x_2]$ (with the notations (2.9)), \dots , until we reach $\sigma := \sigma_n = [\sigma_{n-1}, x_n]$.

Proposition 2.2. *Let $\psi : \sigma \mapsto \sigma'$ be the surjection of \mathfrak{S}_n onto \mathfrak{S}_{n-1} that consists of removing the letter n from the word $\sigma(1) \dots \sigma(n)$. For $n \geq 2$ and for each permutation $\sigma' \in \mathfrak{S}_{n-1}$ the generating polynomial for the class $\psi^{-1}(\sigma')$ by the Major Index “maj” is given by*

$$(2.10) \quad \sum_{\sigma \in \psi^{-1}(\sigma')} q^{\text{maj } \sigma} = q^{\text{maj } \sigma'} (1 + q + q^2 + \dots + q^{n-1}).$$

Moreover, with the notation (2.9)

$$(2.11) \quad \text{maj}[\sigma_{n-1}, x_n] = \text{maj } \sigma_{n-1} + x_n.$$

Finally, if $x = (x_1, x_2, \dots, x_n)$ is the maj-coding of σ , then

$$(2.12) \quad \text{maj } \sigma = x_1 + x_2 + \dots + x_n = \text{tot } x.$$

Proof. Identities (2.10) and (2.12) follow from (2.11) that is now being proved. When n is inserted to the right of $\sigma' = \sigma_{n-1}$, into the position maj-labelled 0, we have $\text{maj}[\sigma', 0] = \text{maj } \sigma'$. If n is inserted into the x_n -th descent ($1 \leq x_n \leq d = \text{des } \sigma'$) (labelled from right to left), the x_n descents occurring on the right are shifted one position to the right; the other descents remain alike. Hence, (2.11) holds. In the same manner, (2.11) holds for $x_n = d + 1$, since the maj-labelling corresponds to an

3. THE ALGEBRA OF THE q -BIMOMIAL COEFFICIENTS

insertion of n at the beginning of the word. Now if $\sigma'(i) < \sigma'(i+1)$ is the k -th non-descent when σ' is read from left to right ($1 \leq k \leq n-d-2$), the left factor $\sigma'(1)\sigma'(2)\dots\sigma'(i)$ contains $i-k$ descents and the right factor $\sigma'(i+1)\sigma'(i+2)\dots\sigma'(n-1)$ exactly $d-i+k$ descents. The insertion of n between $\sigma'(i)$ and $\sigma'(i+1)$, into a position maj-labelled $d+k+1$, increases the Major Index by $(i+1) + (d-i+k) = d+k+1$, since a new descent is created between $\sigma'(i)$ and $\sigma'(i+1)$ and the $(d-i+k)$ descents of the right factor $\sigma'(i+1)\sigma'(i+2)\dots\sigma'(n-1)$ are shifted one position to the right. \square

2.3. The den-coding. The Denert statistic, $\text{den } \sigma$, of a permutation $\sigma \in \mathfrak{S}_n$ can be calculated by means of its *den-coding*, defined as follows. For each integer j ($1 \leq j \leq n$) define x_j as the number of integers i such that

$$(2.13) \quad 1 \leq i \leq j-1 \quad \text{and} \quad \sigma(i) \in \llbracket \sigma(j), j \rrbracket.$$

The *den-coding* of σ is defined to be the sequence $x := (x_1, x_2, \dots, x_n)$, which is obviously subexcedent. Clearly, $\text{den } \sigma = x_1 + x_2 + \dots + x_n = \text{tot } x$ and the mapping $\sigma \mapsto x$ is injective and then bijective. Let us illustrate the calculation of the den-coding with an example.

In the example below the first row shows the integers j from 1 to 9, the second row the value of $\sigma(j)$, the third row the value of the cyclic interval $\llbracket \sigma(j), j \rrbracket$, the fourth row the value of x_j (which is the number of integers i such that $1 \leq i \leq j-1$ and $\sigma(i) \in \llbracket \sigma(j), j \rrbracket$).

j	1	2	3	4	5	6	7	8	9
$\sigma(j)$	7	1	5	4	9	2	6	3	8
$\llbracket \sigma(j), j \rrbracket$	$\{1, 8, 9\}$	$\{2\}$	$\{6, 7, 8, 9, 1, 2, 3\}$	\emptyset	$\{1, 2, 3, 4, 5\}$	$\{3, 4, 5, 6\}$	$\{7\}$	$\{4, 5, 6, 7, 8\}$	$\{9\}$
x_j	0	0	2	0	3	2	1	4	1

The den-coding of σ is then $x = (0, 0, 2, 0, 3, 2, 1, 4, 1)$. In particular, $\text{den } \sigma = 13$.

To recover σ from x , first define $\sigma(n) := n - x_n$. Suppose that $\sigma(j+1), \dots, \sigma(n)$ have been determined from x_{j+1}, \dots, x_n . Write the sequence

$$j, (j-1), \dots, 2, 1, n, (n-1), \dots, (j+1).$$

From that list remove all the elements equal to $\sigma(l)$ for a given $l \geq j+1$. Then $\sigma(j)$ is the $(x_j + 1)$ -st letter in the sequence when reading it from left to right.

Three bijections ϕ^{inv} , ϕ^{maj} , ϕ^{den} of \mathfrak{S}_n onto SE_n have so been constructed with the following properties:

$$\text{inv } \sigma = \text{tot } \phi^{\text{inv}}(\sigma), \quad \text{maj } \sigma = \text{tot } \phi^{\text{maj}}(\sigma), \quad \text{den } \sigma = \text{tot } \phi^{\text{den}}(\sigma).$$

By taking the composition products of ϕ^{inv} , ϕ^{maj} and ϕ^{den} , together with their inverse bijections, we can explicitly construct one-to-one correspondences of \mathfrak{S}_n onto itself, that map “inv” onto “maj”, “inv” onto “den” and “maj” onto “den.”

3. The algebra of the q -binomial coefficients

Consider the product $c = a \cdot b$ of two formal series in the variable u written in the form

$$a = \sum_{i \geq 0} \frac{u^i}{i!} a(i) \quad \text{and} \quad b = \sum_{j \geq 0} \frac{u^j}{j!} b(j).$$

If we want to express the product c in the form

$$c = \sum_{n \geq 0} \frac{u^n}{n!} c(n),$$

we are led to the identity

$$c(n) = \sum_{\substack{i \geq 0, j \geq 0 \\ i+j=n}} \binom{n}{i} a(i) b(j) \quad (n \geq 0),$$

where $\binom{n}{i}$ is the binomial coefficient $\frac{n!}{i!(n-i)!}$.

Now if we replace the factorials $i!$, $j!$, $n!$ occurring in the denominators by their q -analogs $(q; q)_i$, $(q; q)_j$, $(q; q)_n$, as they were defined in (0.1) and if the coefficients $a(i)$, $b(j)$, $c(n)$ are replaced by formal series $a(i, q)$, $b(j, q)$, $c(n, q)$ in the variable q , we obtain the identity

$$c(n, q) = \sum_{\substack{i \geq 0, j \geq 0 \\ i+j=n}} \begin{bmatrix} n \\ i \end{bmatrix} a(i, q) b(j, q),$$

where

$$(3.1) \quad \begin{bmatrix} n \\ i \end{bmatrix} := \frac{(q; q)_n}{(q; q)_i (q; q)_{n-i}} \quad (0 \leq i \leq n).$$

We can also write :

$$(3.2) \quad \begin{bmatrix} n \\ i \end{bmatrix} = \frac{(q^{i+1}; q)_{n-i}}{(q; q)_{n-i}} = \frac{(q^{n-i+1}; q)_i}{(q; q)_i}.$$

The expression $\begin{bmatrix} n \\ i \end{bmatrix}$ is called q -binomial coefficient or *Gaussian polynomial*. It is a remarkable fact that this coefficient is a polynomial in q , with nonnegative integral coefficient. This can be derived in an algebraic manner.

3. THE ALGEBRA OF THE q -BIMOMIAL COEFFICIENTS

In definition (3.1) make the convention that the q -binomial coefficient $\begin{bmatrix} n \\ i \end{bmatrix}$ is zero when condition $0 \leq i \leq n$ does *not* hold. First, we have

$$(3.3) \quad \begin{bmatrix} n \\ 0 \end{bmatrix} = \begin{bmatrix} n \\ n \end{bmatrix} = 1;$$

$$(3.4) \quad \begin{bmatrix} n \\ i \end{bmatrix} = \begin{bmatrix} n \\ n-i \end{bmatrix}.$$

We also have two Pascal Triangle formulas

$$(3.5) \quad \begin{bmatrix} n \\ i \end{bmatrix} = \begin{bmatrix} n-1 \\ i \end{bmatrix} + q^{n-i} \begin{bmatrix} n-1 \\ i-1 \end{bmatrix};$$

$$(3.6) \quad \begin{bmatrix} n \\ i \end{bmatrix} = \begin{bmatrix} n-1 \\ i-1 \end{bmatrix} + q^i \begin{bmatrix} n-1 \\ i \end{bmatrix};$$

that can be derived by mimicking the traditional calculus of the binomial coefficients:

$$\begin{aligned} \begin{bmatrix} n \\ i \end{bmatrix} - \begin{bmatrix} n-1 \\ i \end{bmatrix} &= \frac{(q; q)_{n-1}}{(q; q)_i (q; q)_{n-i}} ((1 - q^n) - (1 - q^{n-i})) \\ &= \frac{(q; q)_{n-1} q^{n-i} (1 - q^i)}{(q; q)_i (q; q)_{n-i}} \\ &= \frac{q^{n-i} (q; q)_{n-1}}{(q; q)_{i-1} (q; q)_{n-i}} = q^{n-i} \begin{bmatrix} n-1 \\ i-1 \end{bmatrix}. \end{aligned}$$

In the same manner,

$$\begin{aligned} \begin{bmatrix} n \\ i \end{bmatrix} - \begin{bmatrix} n \\ n-i \end{bmatrix} &= \begin{bmatrix} n-1 \\ n-i \end{bmatrix} + q^{n-(n-i)} \begin{bmatrix} n-1 \\ n-i-1 \end{bmatrix} \\ &= \begin{bmatrix} n-1 \\ i-1 \end{bmatrix} + q^i \begin{bmatrix} n-1 \\ i \end{bmatrix}. \end{aligned}$$

Finally, the limit $\lim_{q \rightarrow 1} \begin{bmatrix} n \\ i \end{bmatrix} = \binom{n}{i}$ is straightforward.

Relation (3.3) and one of the relations (3.5), (3.6) show that the q -binomial coefficient $\begin{bmatrix} n \\ i \end{bmatrix}$ is a *polynomial* in q , with nonnegative coefficients, of degree $i(n-i)$. The first values of the q -binomial coefficients $\begin{bmatrix} n \\ i \end{bmatrix}$ are shown in Table 3.1.

$$\begin{aligned} \begin{bmatrix} 1 \\ 0 \end{bmatrix} &= \begin{bmatrix} 1 \\ 1 \end{bmatrix} = 1; & \begin{bmatrix} 2 \\ 0 \end{bmatrix} &= \begin{bmatrix} 2 \\ 2 \end{bmatrix} = 1; & \begin{bmatrix} 2 \\ 1 \end{bmatrix} &= 1 + q; & \begin{bmatrix} 3 \\ 0 \end{bmatrix} &= \begin{bmatrix} 3 \\ 3 \end{bmatrix} = 1; \\ \begin{bmatrix} 3 \\ 1 \end{bmatrix} &= \begin{bmatrix} 3 \\ 2 \end{bmatrix} = 1 + q + q^2; & \begin{bmatrix} 4 \\ 0 \end{bmatrix} &= \begin{bmatrix} 4 \\ 4 \end{bmatrix} = 1; & \begin{bmatrix} 4 \\ 1 \end{bmatrix} &= \begin{bmatrix} 4 \\ 3 \end{bmatrix} = 1 + q + q^2 + q^3; \\ \begin{bmatrix} 4 \\ 2 \end{bmatrix} &= 1 + q + 2q^2 + q^3 + q^4; & \begin{bmatrix} 5 \\ 0 \end{bmatrix} &= \begin{bmatrix} 5 \\ 5 \end{bmatrix} = 1; & \begin{bmatrix} 5 \\ 1 \end{bmatrix} &= \begin{bmatrix} 5 \\ 4 \end{bmatrix} = 1 + q + q^2 + q^3 + q^4; \\ \begin{bmatrix} 5 \\ 2 \end{bmatrix} &= \begin{bmatrix} 5 \\ 3 \end{bmatrix} = 1 + q + 2q^2 + 2q^3 + 2q^4 + q^5 + q^6. \end{aligned}$$

Table 3.1

In (1.16) and (1.17) we have obtained two expressions for each one of the two q -exponentials $e_q(u)$ and $E_q(u)$, first as infinite products, then as q -series :

$$(3.7) \quad e_q(u) = \frac{1}{(u; q)_\infty} = \sum_{n \geq 0} \frac{u^n}{(q; q)_n};$$

$$(3.8) \quad E_q(u) = (-u; q)_\infty = \sum_{n \geq 0} q^{n(n-1)/2} \frac{u^n}{(q; q)_n}.$$

By means of the q -binomial coefficients we can obtain the expansions of the *finite* products $1/(u; q)_N$ and $(-u; q)_N$, where N is a nonnegative integer. Those two products can be regarded as *finite versions* of the two q -exponentials $e_q(u)$ and $E_q(u)$.

Proposition 3.1. *We have the identities:*

$$(3.9) \quad \frac{1}{(u; q)_N} = \sum_{n \geq 0} \begin{bmatrix} N+n-1 \\ n \end{bmatrix} u^n;$$

$$(3.10) \quad (-u; q)_N = \sum_{0 \leq n \leq N} \begin{bmatrix} N \\ n \end{bmatrix} q^{n(n-1)/2} u^n.$$

Proof. To derive (3.9) go back to (1.12). We have:

$$\begin{aligned} \frac{1}{(u; q)_N} &= \frac{(q^N u; q)_\infty}{(u; q)_\infty} = \sum_{n \geq 0} (q^N; q)_n \frac{u^n}{(q; q)_n} \\ &= 1 + \sum_{n \geq 1} \frac{(q; q)_{N+n-1}}{(q; q)_{N-1} (q; q)_n} u^n = \sum_{n \geq 0} \begin{bmatrix} N+n-1 \\ n \end{bmatrix} u^n. \end{aligned}$$

To derive (3.10) make use of the q -binomial theorem:

$$\begin{aligned} (-u; q)_N &= \frac{(-u; q)_\infty}{(-uq^N; q)_\infty} = \frac{(q^{-N}(-uq^N); q)_\infty}{(-uq^N; q)_\infty} \\ &= \sum_{n \geq 0} (q^{-N}; q)_n \frac{(-uq^N)^n}{(q; q)_n} = \sum_{0 \leq n \leq N} (q^{-N}; q)_n \frac{(-uq^N)^n}{(q; q)_n}. \end{aligned}$$

The summation is finite, since $(q^{-N}; q)_n$ is zero for every $n \geq N+1$. Now use (1.10) for $\alpha = q^{-N}$, that is,

$$(q^{N+1-n}; q)_n = (q^{-N}; q)_n (-q^N)^n q^{-n(n-1)/2}.$$

We get

$$\begin{aligned} (-u; q)_N &= \sum_{0 \leq n \leq N} (q^{N+1-n}; q)_n q^{n(n-1)/2} \frac{u^n}{(q; q)_n} \\ &= \sum_{0 \leq n \leq N} \begin{bmatrix} N \\ n \end{bmatrix} q^{n(n-1)/2} u^n, \end{aligned}$$

because of (3.2) with the substitutions $n \leftarrow N$ and $i \leftarrow n$. \square

We can also prove (3.9) and (3.10) by induction on N , using the Pascal Triangle formulas (3.5) and (3.6).

4. q -Binomial Combinatorial Structures

For an easy handling of the q -binomial coefficients in Combinatorics it is essential to be familiar with the basic combinatorial structures that admit those coefficients as generating polynomials. For each pair of integers (N, n) we give the description of several pairs (A, f) , where A is a finite set of cardinality $\binom{N}{n}$ and f is a statistic defined on A having the property that

$$(4.1) \quad \sum_{a \in A} q^{f(a)} = \begin{bmatrix} N+n \\ n \end{bmatrix}.$$

Four of those structures are introduced below: the *partitions of integers*, the *nondecreasing sequences of integers*, the *binary words*, the *ordered partitions in two blocks*. Each of these structures has its own geometry and its specific underlying statistic.

4.1. *Partitions of integers.* Formula (3.9) reads

$$(4.2) \quad \frac{1}{(1-u)(1-uq) \cdots (1-uq^N)} = \sum_{n \geq 0} \begin{bmatrix} N+n \\ n \end{bmatrix} u^n.$$

The left-hand side of (4.2) can be expressed as a formal series in the two variables q and u

$$\sum_{n \geq 0} u^n \sum_{m \geq 0} p(m, n, N) q^m,$$

where $p(m, n, N)$ is equal to the number of sequences $(m_0, m_1, m_2, \dots, m_N)$ of nonnegative integers such that

$$(4.3) \quad m_0 + m_1 + \cdots + m_N = n \quad \text{and} \quad 1.m_1 + 2.m_2 + \cdots + N.m_N = m,$$

or, in an equivalent manner, to the number of partitions $1^{m_1} 2^{m_2} \dots N^{m_N}$ of the integer m whose number of parts is *at most equal* to n (because of the occurrence of the coefficient m_0). Hence,

$p(m, n, N)$ is equal to the number of partitions of m in at most n parts, all the parts being at most equal to N .

Notice that $p(m, n, N) = 0$ for $m \geq nN + 1$. Let $\mathcal{P}(n, N)$ be the set of partitions in at most n parts, all of them being at most equal to N (their Ferrers diagrams are then contained in a rectangle of basis N and height n). Let $\|\pi\|$ denote the *weight* of a partition $\pi \in \mathcal{P}(n, N)$, that is, $\|\pi\| = m$ if π is a partition of m . Accordingly,

$$(4.4) \quad \begin{bmatrix} N+n \\ n \end{bmatrix} = \sum_{0 \leq m \leq nN} p(m, n, N) q^m = \sum_{\pi \in \mathcal{P}(n, N)} q^{\|\pi\|}.$$

The first q -binomial model is then the pair $(\mathcal{P}(n, N), \|\cdot\|)$.




Fig. 4.1

There is another way to derive identity (4.4), by using induction on $N + n$, the formula being trivial for $N + n = 1$. We use (3.6), that can be rewritten as

$$\begin{bmatrix} N+n \\ n \end{bmatrix} = \begin{bmatrix} N+(n-1) \\ n-1 \end{bmatrix} + q^n \begin{bmatrix} (N-1)+n \\ n \end{bmatrix}.$$

But the factor $\begin{bmatrix} N+(n-1) \\ n-1 \end{bmatrix}$ is the generating polynomial for the partitions in at most $(n-1)$ parts, all of them being at most equal to N . Call them *of the first kind*. The factor $\begin{bmatrix} (N-1)+n \\ n \end{bmatrix}$ is the generating polynomial for the partitions π whose Ferrers diagram is contained in the rectangle $(N-1) \times n$. Add a column of height n to the left of the Ferrers diagram of each π . We obtain the Ferrers diagram of a partition π' having n parts exactly, all of them being at most equal to N . Say that those partitions π' are *of*

4. q -BINOMIAL COMBINATORIAL STRUCTURES

the second kind. Their generating polynomial is equal to $q^n \begin{bmatrix} (N-1)+n \\ n \end{bmatrix}$ by induction. But, every partition in at most n parts, all at most equal to N , is either of the first kind, or of the second kind (see Fig. 4.1). \square

4.2. Nondecreasing sequences of integers. This model will appear to be extremely convenient, for a great many of combinatorial objects can be easily coded by sequences of integers. For each pair (N, n) of integers let $\text{NDS}(N, n)$ (resp. $\text{IS}(N, n)$) be the set of the nondecreasing (resp. increasing) sequences of nonnegative integers $b = (b_1, b_2, \dots, b_N)$ such that $0 \leq b_1 \leq b_2 \leq \dots \leq b_N \leq n$ (resp. $0 \leq b_1 < b_2 < \dots < b_N \leq n$). As above, let $\text{tot } b := b_1 + b_2 + \dots + b_N$.

Proposition 4.1. *For each pair of integers (N, n) we have:*

$$(4.5) \quad \begin{bmatrix} N+n \\ n \end{bmatrix} = \sum_{b \in \text{NDS}(N, n)} q^{\text{tot } b} = \sum_{b \in \text{NDS}(n, N)} q^{\text{tot } b};$$

$$(4.6) \quad q^{N(N-1)/2} \begin{bmatrix} n+1 \\ N \end{bmatrix} = \sum_{B \in \text{IS}(N, n)} q^{\text{tot } B}.$$

Proof. Notice the symmetry of formula (4.5) in N and n . To derive (4.5) we construct a bijection $\pi \mapsto b$ of $\mathcal{P}(n, N)$ onto $\text{NDS}(n, N)$ that has the property that $\|\pi\| = \text{tot } b$. Let $\pi = (\pi_1 \geq \pi_2 \geq \dots \geq \pi_n \geq 0)$ be a partition in at most n parts, all of them at most equal to N . The bijection is simply given by

$$\pi \mapsto (\pi_n, \dots, \pi_2, \pi_1) = b.$$

Suppose $n \geq N - 1$. To prove (4.6) we use the traditional bijection $B \mapsto b$ that maps each increasing sequence $B \in \text{IS}(N, n)$ onto a nondecreasing sequence $b \in \text{NDS}(N, n - N + 1)$, defined by

$$(0 \leq B_1 < B_2 < \dots < B_N \leq n) \mapsto (0 \leq b_1 \leq b_2 \leq \dots \leq b_N \leq n - N + 1),$$

where $b_1 := B_1$, $b_2 := B_2 - 1$, $b_3 := B_3 - 2$, \dots , $b_N := B_N - N + 1$. It follows that

$$\text{tot } B = \frac{N(N-1)}{2} + \text{tot } b$$

and

$$\begin{aligned} \sum_{B \in \text{IS}(N, n)} q^{\text{tot } B} &= q^{N(N-1)/2} \sum_{b \in \text{NDS}(N, n-N+1)} q^{\text{tot } b} \\ &= q^{N(N-1)/2} \begin{bmatrix} N + (n - N + 1) \\ N \end{bmatrix} = q^{N(N-1)/2} \begin{bmatrix} n+1 \\ N \end{bmatrix}. \quad \square \end{aligned}$$

The second q -binomial model is $(\text{NDS}(N, n), \text{tot})$.

4.3. *Binary words.* Let $\text{BW}(N, n)$ denote the set of all words of length $(N + n)$ having exactly N letters equal to 1 and n letters equal to 0. If $x = x_1x_2 \dots x_{N+n}$ is such a word, the *inversion number*, $\text{inv } x$, of the word x is defined as the number of *subwords* (not simply factors) 10 of the word x .

Example. We can also write the number of 1's that appear to the left of each letter equal to 0, as shown below for the word x .

$$x = 1\ 0\ 0\ 1\ 0\ 1\ 0\ 0\ 1 \\ \cdot\ 1\ 1\ \cdot\ 2\ \cdot\ 3\ 3\ \cdot$$

Hence $\text{inv } x = 1 + 1 + 2 + 3 + 3 = 10$.

Proposition 4.2. *For each pair of integers (N, n) we have:*

$$(4.17) \quad \begin{bmatrix} N+n \\ n \end{bmatrix} = \sum_{x \in \text{BW}(N, n)} q^{\text{inv } x}.$$

Proof. Again, we construct a bijection $\pi \mapsto x$ of $\mathcal{P}(N, n)$ onto $\text{BW}(N, n)$, such that $\|\pi\| = \text{inv } x$. Every partition $\pi \in \mathcal{P}(N, n)$ can be, in its multiplicative version, described as a monomial $i_1^{n_1} i_2^{n_2} \dots i_r^{n_r}$, where $0 \leq i_1 < i_2 < \dots < i_r \leq N$, $n_1 \geq 1$, $n_2 \geq 1$, \dots , $n_r \geq 1$ and $n_1 + n_2 + \dots + n_r = n$. If the number of parts $l(\pi)$ of π is strictly less than n , let $n_1 := n - l(\pi)$ and $i_1 := 0$. Then the partition π has its parts only equal to i_2, \dots, i_r , repeated n_2, \dots, n_r times, respectively. If $l(\pi) = n$, then $1 \leq i_1$ and π has its parts equal to i_1, \dots, i_r , repeated n_1, \dots, n_r times, respectively.

With the partition π associate the word x

$$x := 1^{i_1} 0^{n_1} 1^{i_2 - i_1} 0^{n_2} 1^{i_3 - i_2} 0^{n_3} \dots 0^{n_{r-1}} 1^{i_r - i_{r-1}} 0^{n_r} 1^{N - i_r}.$$

The word x has $i_1 + (i_2 - i_1) + (i_3 - i_2) + \dots + (i_r - i_{r-1}) + (N - i_r) = N$ letters equal to 1 and $n_1 + n_2 + n_3 + \dots + n_{r-1} + n_r = n$ letters equal to 0. Moreover, $\|\pi\| = i_1 \cdot n_1 + i_2 \cdot n_2 + \dots + i_r \cdot n_r = i_1 \cdot n_1 + (i_1 + (i_2 - i_1)) \cdot n_2 + \dots + (i_1 + (i_2 - i_1) + \dots + (i_r - i_{r-1})) \cdot n_r = \text{inv } x$. Finally, the mapping $\pi \mapsto x$ is obviously injective, and then bijective. \square

Remark. There is a geometric manner to see the bijection $\pi \mapsto x$ described in the previous proof. Put the Ferrers diagram of the partition π inside a triangle of basis N and height n . The rim of the Ferrers diagram is a polygonal line, made of vertical and horizontal steps of length 1, starting from the point whose coordinates are $(0, n)$ down to the point with coordinates $(N, 0)$. The rim has exactly n vertical steps and N horizontal

4. q -BINOMIAL COMBINATORIAL STRUCTURES




Fig. 4.2

steps. Now read the rim of π from top to bottom and from left to right and give label 0 (resp. label 1) to each vertical (resp. horizontal) step. The word x thereby obtained is the binary word described in the previous bijection (see Fig. 4.2).

Example. Consider the partition $\pi = (6, 5, 4, 4, 2, 2)$ belonging to $\mathcal{P}(6, 7)$. In its multiplicative version it can be expressed as the monomial $0^1 2^2 4^2 5^1 6^1$. The word x corresponding to that monomial is the word $x = 1^0 0^1 1^{2-0} 0^2 1^{4-2} 0^2 1^{5-4} 0^1 1^{6-5} 0^1 1^{6-6} = 0, 1, 1, 0, 0, 1, 1, 0, 0, 1, 0, 1, 0$, which is the word we can read on the rim of its Ferrers diagram using the previous labelling.

The third q -binomial model is then $(\text{BW}(N, n), \text{inv})$.

4.4. *Ordered Partitions into two blocks.* The word “partition” used in this subsection will refer to (set) partitions. The (set) partition of a finite set S is a collection of subsets (called *blocks*) of S , two by two disjoint, whose union is S . By *ordered partition* of a set it is meant a (set) partition into blocks, together with a linear ordering of those blocks. For convenience, we may assume that some of those blocks are empty.

Now consider the set $[N + n]$ of the $(N + n)$ integers $1, 2, \dots, N + n$. If (A, B) is an ordered partition of $[N + n]$ into two blocks, $\gamma(A)$ (resp. $\gamma(B)$) will designate the *increasing* word whose letters are the elements of A (resp. of B) written in increasing order. There is no inversion of letters in each of the words $\gamma(A), \gamma(B)$, so that the number of inversions $\text{inv}(\gamma(A)\gamma(B))$ in the juxtaposition product $\gamma(A)\gamma(B)$ is equal to the number of pairs (a, b) such that $a \in A, b \in B$ and $a > b$.

Proposition 4.3. *For each pair of integers (N, n) we have*

$$\begin{bmatrix} N + n \\ n \end{bmatrix} = \sum_{(A, B)} q^{\text{inv}(\gamma(A)\gamma(B))},$$

where the sum is over the set $\text{OP}(N, n)$ of all the ordered partitions (A, B) of the set $[N + n]$ into two blocks such that $|A| = N$ and $|B| = n$.

Proof. It follows from Proposition 4.2 that the q -binomial coefficient $\begin{bmatrix} N+n \\ n \end{bmatrix}$ is the generating function for the binary words $x = x_1x_2 \dots x_{N+n}$ having N letters equal to 1 and n letters equal to 0, by the inversion number. With such a binary word x associate the ordered partition (A, B) of $[N+n]$ defined by: $i \in A$ or $i \in B$ depending on whether $x_i = 0$ or $x_i = 1$. The inversion $x_j = 1, x_{j'} = 0, j < j'$ in the word x will then correspond to the inversion $j' > j$ between the element $j' \in A$ and the element $j \in B$. \square

The fourth q -binomial model is then $(\text{OP}(N, n), \text{inv})$.

5. The q -multinomial coefficients

They form a natural q -extension of the multinomial coefficients and can be introduced as follows. For each integer $r \geq 1$ and each sequence of nonnegative integers (m_1, m_2, \dots, m_r) let

$$(5.1) \quad \begin{bmatrix} m_1 + m_2 + \dots + m_r \\ m_1, m_2, \dots, m_r \end{bmatrix}_q := \frac{(q; q)_{m_1+m_2+\dots+m_r}}{(q; q)_{m_1}(q; q)_{m_2} \dots (q; q)_{m_r}}.$$

If there is no ambiguity the subscript q is suppressed. In case $r = 2$ we recover the expression of the Gaussian polynomial studied in the previous section. The fact that the q -multinomial coefficient is a polynomial in q with positive integral coefficients follows from the combinatorial interpretations given in the sequel

When q tends to 1, the q -multinomial coefficient tends to the multinomial coefficient $\binom{m_1+m_2+\dots+m_r}{m_1, m_2, \dots, m_r}$, as is readily verified. We can then expect that the q -multinomial coefficient is the generating polynomial for a set of cardinality $\binom{m_1+m_2+\dots+m_r}{m_1, m_2, \dots, m_r}$ by a certain statistic.

The two combinatorial interpretations of the q -binomial coefficient in terms of classes of partitions and nondecreasing sequences are difficult to be extended to the multinomial case. However, when going from the *binary* words, studied in the previous subsection, to the words whose letters belong to an alphabet of cardinality r ($r \geq 2$) and when the statistic “inv” is extended to those words, the q -multinomial coefficient can easily be interpreted in a combinatorial way. In the sequel the word “rearrangement” of a word w , with or without repeated letters, means any word derived from w by permuting its letters in some order.

For $r \geq 1$ and for each sequence $\mathbf{m} = (m_1, m_2, \dots, m_r)$ of nonnegative integers let $R(\mathbf{m})$ denote the class of all the words of length $m = m_1 + m_2 + \dots + m_r$ which are *rearrangements* of the nondecreasing word $1^{m_1}2^{m_2} \dots r^{m_r}$. The number of such rearrangements is equal to the multinomial coefficient $\binom{m_1+m_2+\dots+m_r}{m_1, m_2, \dots, m_r}$.

5. THE q -MULTINOMIAL COEFFICIENTS

Let $w = x_1 x_2 \dots x_m$ be a word belonging to the class $R(\mathbf{m})$. The *number of inversions*, $\text{inv } w$, of w is defined to be the number of pairs (i, j) such that $1 \leq i < j \leq m$ and $x_i > x_j$. For each word $w = x_1 x_2 \dots x_m$ it is convenient, for each $j = 1, \dots, m$, to determine the number z_j of letters x_i lying *to the left of* x_j such that $x_i > x_j$. Then $\text{inv } w = z_1 + \dots + z_m$.

In the following example the number of inversions of w is determined from the sequence of the z_j 's:

$$\begin{aligned} w &= 3\ 1\ 3\ 4\ 1\ 2\ 5\ 4\ 3 \\ z &= 0\ 1\ 0\ 0\ 3\ 3\ 0\ 1\ 3 \end{aligned}$$

so that $\text{inv } w = \text{tot } z = 1 + 3 + 3 + 1 + 3 = 11$.

Theorem 5.1. *The q -multinomial coefficient $\begin{bmatrix} m_1 + m_2 + \dots + m_r \\ m_1, m_2, \dots, m_r \end{bmatrix}$ is the generating polynomial for the set $R(\mathbf{m})$ by the number of inversions. In other words,*

$$(5.2) \quad \begin{bmatrix} m_1 + m_2 + \dots + m_r \\ m_1, m_2, \dots, m_r \end{bmatrix} = \sum_{w \in R(\mathbf{m})} q^{\text{inv } w}.$$

Proof. Relation (5.2) is banal for $r = 1$ and holds for $r = 2$ by Proposition 4.2. Consider the factorization

$$\begin{aligned} \begin{bmatrix} m_1 + m_2 + \dots + m_{r+1} \\ m_1, m_2, \dots, m_{r+1} \end{bmatrix} &= \frac{(q; q)_{m_1 + m_2 + \dots + m_{r+1}}}{(q; q)_{m_1} (q; q)_{m_2} \dots (q; q)_{m_{r+1}}} \\ &= \frac{(q; q)_{m_1 + m_2 + \dots + m_{r+1}}}{(q; q)_{m_1 + m_2 + \dots + m_r} (q; q)_{m_{r+1}}} \frac{(q; q)_{m_1 + m_2 + \dots + m_r}}{(q; q)_{m_1} (q; q)_{m_2} \dots (q; q)_{m_r}} \\ &= \begin{bmatrix} m_1 + m_2 + \dots + m_{r+1} \\ m_1 + m_2 + \dots + m_r, m_{r+1} \end{bmatrix} \begin{bmatrix} m_1 + m_2 + \dots + m_r \\ m_1, m_2, \dots, m_r \end{bmatrix} \end{aligned}$$

and take a word $w = x_1 x_2 \dots x_{m'}$ from the set $R(m_1, m_2, \dots, m_{r+1})$, so that its length is $m' = m_1 + m_2 + \dots + m_{r+1}$. The inversions $x_i > x_j$ ($i < j$) of w fall into two classes: (i) the inversions of the form $x_i = r + 1 > s = x_j$, where s is one of the integers $1, 2, \dots, r$; (ii) the inversions of the form $x_i = s > t = x_j$, where $r \geq s > t \geq 1$.

Let w_1 denote the word of length m' derived from w by replacing all the letters less than or equal to r by 1 and all the letters equal to $(r + 1)$ by 2. Likewise, let w_2 denote the *subword* of length $m = m_1 + m_2 + \dots + m_r$ obtained from w by deleting *all* the letters equal to $(r + 1)$.

The map $w \mapsto (w_1, w_2)$ is obviously a bijection of $R(m_1, m_2, \dots, m_{r+1})$ onto $R(m_1 + m_2 + \dots + m_r, m_{r+1}) \times R(m_1, m_2, \dots, m_r)$. Moreover,

$$(5.3) \quad \text{inv } w = \text{inv } w_1 + \text{inv } w_2.$$

But by induction on r we have:

$$\begin{aligned} \sum_{w_1} q^{\text{inv } w_1} &= \left[\begin{array}{c} m_1 + m_2 + \cdots + m_{r+1} \\ m_1 + m_2 + \cdots + m_r, m_{r+1} \end{array} \right]; \\ \sum_{w_2} q^{\text{inv } w_2} &= \left[\begin{array}{c} m_1 + m_2 + \cdots + m_r \\ m_1, m_2, \dots, m_r \end{array} \right]. \end{aligned}$$

The identity (5.2) is then a consequence of those identities and of (5.3). \square

6. The MacMahon Verfahren

The German word *Verfahren* means “procedure,” “way of doing,” ... This term applies to the rearrangement method of sequences of numbers, imagined by MacMahon when he was dealing with q -series in a combinatorial context.

Obviously, the fraction $1/(q; q)_m$ is the generating function for the partitions of integers in at most m parts. If we write such a partition in its classical form $\pi = (\pi_1 \geq \pi_2 \geq \cdots \geq \pi_m \geq 0)$, then the sequence $b = (b_1, \dots, b_{m-1}, b_m) := (\pi_m, \dots, \pi_2, \pi_1)$ is a nondecreasing sequence of m nonnegative integers. Extending our previous notation we write $b \in \text{NDS}(m)$, so that

$$(6.1) \quad \frac{1}{(q; q)_m} = \sum_{b \in \text{NDS}(m)} q^{\text{tot } b},$$

where $\text{tot } b = b_1 + \cdots + b_m$.

In section 2 we have introduced the Mahonian statistic on a set of cardinality $n!$ and more essentially on the symmetric group \mathfrak{S}_n . We now extend the definition of that statistic to arbitrary rearrangement classes. A statistic “stat” is said to be *Mahonian*, if for every class $R(\mathbf{m})$ the following identity holds:

$$(6.2) \quad \frac{1}{(q; q)_{m_1 + \cdots + m_r}} \sum_{w \in R(\mathbf{m})} q^{\text{stat } w} = \frac{1}{(q; q)_{m_1} \cdots (q; q)_{m_r}}.$$

Theorem 5.1 says nothing but that the inversion number “inv” is a Mahonian statistic. By using (6.1), identity (6.2) can be rewritten in the form

$$(6.3) \quad \sum_{b \in \text{NDS}(m), w \in R(\mathbf{m})} q^{\text{tot } b + \text{stat } w} = \sum_{b^{(1)}, \dots, b^{(r)}} q^{\text{tot } b^{(1)} + \cdots + \text{tot } b^{(r)}},$$

6. THE MACMAHON VERFAHREN

where $b^{(1)} \in \text{NDS}(m_1), \dots, b^{(r)} \in \text{NDS}(m_r)$. Hence, we can also say that a statistic “stat” is *Mahonian*, if to every pair $(b, w) \in \text{NDS}(m) \times R(\mathbf{m})$ there corresponds a unique sequence $(b^{(1)}, \dots, b^{(r)}) \in \text{NDS}(m_1) \times \dots \times \text{NDS}(m_r)$ such that

$$(6.4) \quad \text{tot } b + \text{stat } w = \text{tot } b^{(1)} + \dots + \text{tot } b^{(r)}.$$

The purpose of this section is to use that definition for making the Major Index appear as a Mahonian statistic, not only for permutations, but for rearrangements of arbitrary words.

Each sequence $(b^{(1)}, \dots, b^{(r)})$ can be mapped, in a bijective way, onto a two-row matrix

$$(6.5) \quad \begin{pmatrix} b_{m_1}^{(1)} & \dots & b_1^{(1)} & b_{m_2}^{(2)} & \dots & b_1^{(2)} & \dots & b_{m_r}^{(r)} & \dots & b_1^{(r)} \\ 1 & \dots & 1 & 2 & \dots & 2 & \dots & r & \dots & r \end{pmatrix},$$

where, on the first row, the *nonincreasing rearrangements* $b_{m_1}^{(1)} \dots b_1^{(1)}, b_{m_2}^{(2)} \dots b_1^{(2)}, \dots, b_{m_r}^{(r)} \dots b_1^{(r)}$ of the sequences $b^{(1)}, b^{(2)}, \dots, b^{(r)}$ have been in that order.

The idea of the MacMahon *Verfahren* is to rearrange the *columns* of the latter matrix in such a way that the elements on the top row will be in *nonincreasing* order (when read from left to right), this being made in a one-to-one manner. The bottom row will then go from $1^{m_1} 2^{m_2} \dots r^{m_r}$ to a rearrangement of that word. To realize the rearrangement of the columns we make use of the following *commutation rule*:

$$(6.6) \quad \text{two columns } \begin{pmatrix} c \\ d \end{pmatrix} \text{ and } \begin{pmatrix} c' \\ d' \end{pmatrix} \text{ commute if and only if } c \neq c'.$$

The commutation rule being given, to each matrix of type (6.5) there corresponds, *in a bijective manner*, a matrix

$$(6.7) \quad \begin{pmatrix} y_1 & y_2 & \dots & y_m \\ x_1 & x_2 & \dots & x_m \end{pmatrix},$$

whose top row is *nonincreasing* and if $y_k = y_{k+1}$, then $x_k \leq x_{k+1}$, or, in an equivalent way,

$$(6.8) \quad x_k > x_{k+1} \implies y_k > y_{k+1}.$$

In other words, if there is a *descent* on the bottom row, there is also a descent on the top row, the converse being not necessarily true.

For example, let $r = 3$, $m_1 = 6$, $m_2 = 2$, $m_3 = 4$, $b^{(1)} = (0, 0, 1, 1, 5, 6)$, $b^{(2)} = (1, 3)$, $b^{(3)} = (1, 1, 4, 5)$, so that $m = m_1 + m_2 + m_3 = 12$. The matrix of type (6.5) reads:

$$\begin{pmatrix} 6 & 5 & 1 & 1 & 0 & 0 & 3 & 1 & 5 & 4 & 1 & 1 \\ 1 & 1 & 1 & 1 & 1 & 1 & 2 & 2 & 3 & 3 & 3 & 3 \end{pmatrix}.$$

Using the commutation rule (6.6) the matrix is transformed into a matrix of type (6.8):

$$\begin{pmatrix} 6 & 5 & 5 & 4 & 3 & 1 & 1 & 1 & 1 & 1 & 0 & 0 \\ 1 & 1 & 3 & \mathbf{3} & \mathbf{2} & 1 & 1 & 2 & 3 & \mathbf{3} & 1 & 1 \end{pmatrix}.$$

The coefficients on the bottom row that are greater than their successors are written in bold-face. We see that the corresponding coefficients on the top row are greater than their successors (property (6.8)).

Go back to the general case and let $v := y_1 y_2 \dots y_m$ denote the word appearing on the top row of the matrix (6.7). It is the *unique nonincreasing* rearrangement of the juxtaposition product $b^{(1)} \dots b^{(r)}$. Next, let $w := x_1 x_2 \dots x_m$ be the word appearing on the bottom row of (6.7). It is a well-defined word belonging to $R(\mathbf{m})$.

For $k = 1, 2, \dots, m$ let z_k be the number of descents in the right factor $x_k x_{k+1} \dots x_m$ of w , that is, the number of subscripts j such that $k \leq j \leq m - 1$ and $x_j > x_{j+1}$; next let $b_k := y_k - z_k$ ($1 \leq k \leq m$). If $x_k > x_{k+1}$, then $z_k = z_{k+1} + 1$ by definition of z_k and also $y_k \geq y_{k+1} + 1$ by (6.8). It follows that $b_k = y_k - z_k \geq y_{k+1} + 1 - (z_{k+1} + 1) = b_{k+1}$. However, if $x_k \leq x_{k+1}$, we always have $y_k \geq y_{k+1}$, since v is nonincreasing and also $z_k = z_{k+1}$. Hence $b_k = y_k - z_k \geq y_{k+1} - z_{k+1} = b_{k+1}$.

We conclude that the sequence b defined by $b := (b_m, \dots, b_2, b_1)$ satisfies the relations $0 \leq b_m \leq \dots \leq b_2 \leq b_1$, so that $b \in \text{NDS}(m)$. Finally, if $z(w)$ designates the sequence (z_1, z_2, \dots, z_m) , we have:

$$\begin{aligned} \text{tot } b^{(1)} + \dots + \text{tot } b^{(r)} &= y_1 + y_2 + \dots + y_m \\ &= (b_1 + z_1) + (b_2 + z_2) + \dots + (b_m + z_m) \\ (6.9) \qquad \qquad \qquad &= \text{tot } b + \text{tot } z(w). \end{aligned}$$

Comparing the last identity with (6.4) we see that $\text{tot } z(w)$ is a new *Mahonian statistic*, if it can be verified that the mapping $(b^{(1)}, \dots, b^{(r)}) \mapsto (b, w)$ is bijective. But the construction that has just been made is perfectly reversible: starting with a word $w \in R(\mathbf{m})$ and a sequence $b = (b_m, \dots, b_2, b_1) \in \text{NDS}(m)$, we first determine the sequence $z(w) = (z_1, \dots, z_m)$. We know that the word $v = y_1 \dots y_{m-1} y_m$ defined by $y_i := b_i + z_i$ ($1 \leq i \leq m$) is nonincreasing. We next form the two-row

6. THE MACMAHON VERFAHREN

matrix $\binom{v}{w}$ and by applying the commutation rule (6.6) we define the sequences $b^{(1)}, \dots, b^{(r)}$ by (6.5). Relation (6.9) obviously holds.

Let us take again the previous example. We had obtained:

$$\binom{v}{w} = \begin{pmatrix} 6 & 5 & 5 & 4 & 3 & 1 & 1 & 1 & 1 & 1 & 0 & 0 \\ 1 & 1 & 3 & 3 & 2 & 1 & 1 & 2 & 3 & 3 & 1 & 1 \end{pmatrix}.$$

We now get

$$\begin{aligned} v &= 6 & 5 & 5 & 4 & 3 & 1 & 1 & 1 & 1 & 1 & 0 & 0 \\ w &= 1 & 1 & 3 & 3 & 2 & 1 & 1 & 2 & 3 & 3 & 1 & 1 \\ z(w) &= 3 & 3 & 3 & 3 & 2 & 1 & 1 & 1 & 1 & 1 & 0 & 0 \\ & 3 & 2 & 2 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 = b \text{ (read from right to left)} \end{aligned}$$

and can verify that

$$\begin{aligned} \text{tot } b^{(1)} + \dots + \text{tot } b^{(r)} &= (6 + 5 + 1 + 1) + (3 + 1) + (5 + 4 + 1 + 1) = 28 \\ &= \text{tot } b + \text{tot } z(w) = (3 + 2 + 2 + 1 + 1) + (12 + 2 + 5) = 28. \end{aligned}$$

Going back to the general case the problem is to characterize the new Mahonian statistic “tot $z(w)$ ” in a more direct way. It is, indeed, the *Major Index*, “maj”, already introduced in section 2 in the case of permutations. Its definition can be extended to the case of arbitrary words.

Definition. Let $w = x_1x_2\dots x_m$ be a word whose letter are taken from the alphabet $\{1, 2, \dots, r\}$. The *Major Index*, “maj w ”, of the word w is defined by

$$(6.10) \quad \text{maj } w := \sum_{1 \leq i \leq m-1} i \chi(x_i > x_{i+1}).$$

Thus, for calculating the Major Index of a word, we determine its descents and their positions. The Major Index is the sum of the positions of its descents.

Proposition 6.2. *For every word w we have: $\text{maj } w = \text{tot } z(w)$.*

Proof. If $w = x_1x_2\dots x_m$ and $1 \leq i \leq m$, we have defined z_i as being the number of descents in the right factor $x_ix_{i+1}\dots x_m$, and $z(w)$ as being the sequence of the z_i 's. If w is of length 1, obviously $\text{maj } w = \text{tot } z(w) = 0$. If w is of length greater than 1, define $w' := x_1x_2\dots x_{m-1}$ and let $z(w') = (z'_1, \dots, z'_{m-1})$. If $x_{m-1} \leq x_m$, then $\text{maj } w = \text{maj } w'x_m = \text{maj } w' = \text{tot } z(w') = \text{tot } z(w'x_m) = \text{tot } z(w)$. If $x_{m-1} > x_m$, then $\text{maj } w = \text{maj } w' + (m - 1)$, since there is a descent in position $(m - 1)$. On the other hand, $z(w) = ((z'_1 + 1), \dots, (z'_{m-1} + 1))$; hence $\text{tot } z(w) = \text{tot } z(w') + (m - 1)$. \square

In the next theorem we state the results derived in this section about the Major Index.

Theorem 6.3. *The Major Index is a q -multinomial statistic; that is, for each rearrangement class $R(\mathbf{m})$ the identity*

$$(6.11) \quad \left[\begin{matrix} m_1 + m_2 + \cdots + m_r \\ m_1, m_2, \dots, m_r \end{matrix} \right] = \sum_{w \in R(\mathbf{m})} q^{\text{maj } w}$$

holds.

The right-hand side of (6.11) is the generating polynomial for the class $R(\mathbf{m})$ by the Major Index, a polynomial that will be denoted by $A_{\mathbf{m}}(q)$. Relation (6.11) can also be rewritten as

$$(6.12) \quad \frac{1}{(q; q)_m} A_{\mathbf{m}}(q) = \frac{1}{(q; q)_{m_1} \cdots (q; q)_{m_r}},$$

which is (6.2) with “maj” replacing “stat.”

Now introduce the algebra of power formal series in the variables u_1, u_2, \dots, u_r . Let $\mathbf{u}^{\mathbf{m}} := u_1^{m_1} u_2^{m_2} \cdots u_r^{m_r}$, also $(\mathbf{u}; q)_{\infty} := (u_1; q)_{\infty} \cdots (u_r; q)_{\infty}$ and remember that $m = m_1 + \cdots + m_r$. Next, multiply (6.12) by $\mathbf{u}^{\mathbf{m}}$ and sum the two sides of the equation with respect to all the sequences $\mathbf{m} = (m_1, \dots, m_r)$ of r nonnegative integers. By using the q -Binomial Theorem (Theorem 1.1) we get the equivalent identity

$$(6.13) \quad \sum_{\mathbf{m}} A_{\mathbf{m}}(q) \frac{\mathbf{u}^{\mathbf{m}}}{(q; q)_m} = \frac{1}{(\mathbf{u}; q)_{\infty}}.$$

7. A refinement of the MacMahon Verfahren

Re-examine the *inverse* mapping of the bijection

$$(7.1) \quad (b^{(1)}, \dots, b^{(r)}) \mapsto (b, w)$$

of $\text{NDS}(m_1) \times \cdots \times \text{NDS}(m_r)$ onto $\text{NDS}(m) \times R(\mathbf{m})$, described in the previous section. Each term z_i in the sequence $z(w) = (z_1, z_2, \dots, z_m)$ counts the number of descents in the right factor $x_i x_{i+1} \cdots x_m$ of the word w . Let $\text{des } w := \sum_{1 \leq i \leq m-1} \chi(x_i > x_{i+1})$ denote the *number of descents* of the word w , so that

$$(7.2) \quad z_1 = \text{des } w.$$

As $y_1 = b_m + z_1$ and since y_1 is the maximum term in the sequence $b^{(1)}, \dots, b^{(r)}$, we also have:

$$(7.3) \quad b_{m_1}^{(1)} \leq b_1 + \text{des } w, \dots, b_{m_r}^{(r)} \leq b_1 + \text{des } w.$$

7. A REFINEMENT OF THE MACMAHON VERFAHREN

Take a nonnegative integer s' and a nondecreasing sequence $b = b_m \dots b_2 b_1$ such that $0 \leq b_m \leq \dots \leq b_1 \leq s'$, i.e., $b \in \text{NDS}(m, s')$; further take a word $w \in R(\mathbf{m})$ and let

$$(7.4) \quad s := s' + \text{des } w.$$

It follows from the inequalities (7.3) that

$$(7.5) \quad b^{(1)} \in \text{NDS}(m_1, s), \dots, b^{(r)} \in \text{NDS}(m_r, s).$$

The bijection constructed in the previous section also has the property stated in the next proposition.

Proposition 7.1. *To each triple (s', b, w) such that $s' \geq 0$, $b \in \text{NDS}(m, s')$ and $w \in R(\mathbf{m})$ there corresponds, in a bijective manner, a sequence $(s, b^{(1)}, \dots, b^{(r)})$, where $s = s' + \text{des } w$ and where $b^{(1)} \in \text{NDS}(m_1, s)$, \dots , $b^{(r)} \in \text{NDS}(m_r, s)$, having the property:*

$$\text{tot } b^{(1)} + \dots + \text{tot } b^{(r)} = \text{tot } b + \text{maj } w.$$

Let $A_{\mathbf{m}}(t, q)$ denote the generating polynomial for $R(\mathbf{m})$ by the bi-statistic (des, maj) :

$$(7.6) \quad A_{\mathbf{m}}(t, q) = \sum_{w \in R(\mathbf{m})} t^{\text{des } w} q^{\text{maj } w}.$$

Then

$$\begin{aligned} \frac{1}{(t; q)_{m+1}} A_{\mathbf{m}}(t, q) &= \sum_{s' \geq 0} t^{s'} \begin{bmatrix} m + s' \\ s' \end{bmatrix} A_{\mathbf{m}}(t, q) && \text{[by (3.9)]} \\ &= \sum_{s' \geq 0} t^{s'} \sum_{b \in \text{NDS}(m, s')} q^{\text{tot } b} A_{\mathbf{m}}(t, q) && \text{[by (4.5)]} \\ &= \sum_{\substack{s' \geq 0, b \in \text{NDS}(m, s'), \\ w \in R(\mathbf{m})}} t^{s' + \text{des } w} q^{\text{tot } b + \text{maj } w} \\ &= \sum_{s \geq 0} t^s \sum_{\substack{s' \geq 0, b \in \text{NDS}(m, s'), \\ w \in R(\mathbf{m}), s' + \text{des } w = s}} q^{\text{tot } b + \text{maj } w} \\ &= \sum_{s \geq 0} t^s \sum_{\substack{b^{(1)} \in \text{NDS}(m_1, s), \dots, \\ b^{(r)} \in \text{NDS}(m_r, s)}} q^{\text{tot } b^{(1)} + \dots + \text{tot } b^{(r)}}, \\ &&& \text{[in view of Proposition 7.1]} \end{aligned}$$

so that

$$(7.7) \quad \frac{1}{(t; q)_{m+1}} A_{\mathbf{m}}(t, q) = \sum_{s \geq 0} t^s \begin{bmatrix} m_1 + s \\ s \end{bmatrix} \dots \begin{bmatrix} m_r + s \\ s \end{bmatrix} \quad \text{[by (4.5)].}$$

As at the end of the previous section, we can express (7.7) as an identity between formal power series in r variables u_1, u_2, \dots, u_r . However, those series will be normalized by denominators of the form $(t; q)_{m+1}$. Again, let $\mathbf{u}^{\mathbf{m}} := u_1^{m_1} u_2^{m_2} \dots u_r^{m_r}$ and let $(\mathbf{u}; q)_{s+1} := (u_1; q)_{s+1} \dots (u_r; q)_{s+1}$. Next, multiply (7.7) by $\mathbf{u}^{\mathbf{m}}$ and sum over all sequences $\mathbf{m} = (m_1, \dots, m_r)$ of r nonnegative integers. Then,

$$\begin{aligned} \sum_{\mathbf{m}} A_{\mathbf{m}}(t, q) \frac{\mathbf{u}^{\mathbf{m}}}{(t; q)_{1+m}} &= \sum_{s \geq 0} t^s \sum_{\mathbf{m}} \mathbf{u}^{\mathbf{m}} \begin{bmatrix} m_1 + s \\ s \end{bmatrix} \dots \begin{bmatrix} m_r + s \\ s \end{bmatrix} \\ &= \sum_{s \geq 0} t^s \left(\sum_{m_1} u_1^{m_1} \begin{bmatrix} m_1 + s \\ s \end{bmatrix} \right) \dots \left(\sum_{m_r} u_r^{m_r} \begin{bmatrix} m_r + s \\ s \end{bmatrix} \right), \end{aligned}$$

so that by using (3.9),

$$(7.8) \quad \sum_{\mathbf{m}} A_{\mathbf{m}}(t, q) \frac{\mathbf{u}^{\mathbf{m}}}{(t; q)_{1+m}} = \sum_{s \geq 0} \frac{t^s}{(\mathbf{u}; q)_{s+1}}.$$

Remark. Identities (7.7) and (7.8) appear to be “ t -extensions” of identities (6.12) and (6.13) derived in the previous section.

8. The Euler-Mahonian polynomials

In the previous section the polynomials $A_{\mathbf{m}}(t, q)$ have been introduced as generating polynomials for the class $R(\mathbf{m})$ by the bi-statistic (des, maj) . In formula (7.7) they appear as *numerators* of rational fractions whose series expansion in t has an explicit form. In fact, formula (7.7) is only another way of looking at their (t, q) -generating function, as obtained in (7.8). However there are other ways of expressing the polynomials *without any reference to any combinatorial interpretation*, as shown in this section. First, we state a definition that will be made valid, once we prove that the four items (1)–(4) are equivalent. For proving the equivalence of those statements we use two methods: a *finite difference q -calculus* and a *q -iteration* that are developed afterwards.

For each multi-index $\mathbf{m} = (m_1, m_2, \dots, m_{r-1}, m_r)$ let $|\mathbf{m}| := m_1 + m_2 + \dots + m_{r-1} + m_r$ (a quantity that has been denoted by m in the previous section) and $\mathbf{m} + 1_r := (m_1, m_2, \dots, m_{r-1}, m_r + 1)$. Also keep the notations \mathbf{u} , $\mathbf{u}^{\mathbf{m}}$, $(\mathbf{u}; q)_{s+1}$; also $[0]_q := 0$ and $[m]_q := 1 + q + \dots + q^{m-1}$ ($m \geq 1$).

Definition 8.1. Let r be a fixed positive integer. A sequence

$$\left(A_{\mathbf{m}}(t, q) = \sum_{s \geq 0} t^s A_{\mathbf{m}, s}(q) \right)$$

8. THE EULER-MAHONIAN POLYNOMIALS

of polynomials in two variables t and q , indexed by a multi-index $\mathbf{m} = (m_1, \dots, m_r)$ of r nonnegative integers, is said to be *Euler-Mahonian*, if one of the following *equivalent* four conditions holds:

(1) For all \mathbf{m} we have:

$$(8.1) \quad \frac{1}{(t; q)_{m+1}} A_{\mathbf{m}}(t, q) = \sum_{s \geq 0} t^s \begin{bmatrix} m_1 + s \\ s \end{bmatrix} \cdots \begin{bmatrix} m_r + s \\ s \end{bmatrix}.$$

(2) The (t, q) -generating function for those polynomials $A_{\mathbf{m}}(t, q)$ is given by:

$$(8.2) \quad \sum_{\mathbf{m}} A_{\mathbf{m}}(t, q) \frac{\mathbf{u}^{\mathbf{m}}}{(t; q)_{1+|\mathbf{m}|}} = \sum_{s \geq 0} \frac{t^s}{(\mathbf{u}; q)_{s+1}}.$$

(3) The recurrence relation holds:

$$(8.3) \quad (1 - q^{m_r+1})A_{\mathbf{m}+1_r}(t, q) = (1 - tq^{1+|\mathbf{m}|})A_{\mathbf{m}}(t, q) - q^{m_r+1}(1 - t)A_{\mathbf{m}}(tq, q).$$

(4) The recurrence relation holds for the coefficients $A_{\mathbf{m},s}(q)$:

$$(8.4) \quad [m_r + 1]_q A_{\mathbf{m}+1_r,s}(q) = [m_r + 1 + s]_q A_{\mathbf{m},s}(q) + q^{s+m_r} [1 + |\mathbf{m}| - s - m_r]_q A_{\mathbf{m},s-1}(q).$$

For the equivalence (1) \Leftrightarrow (2) see the previous section. To verify (3) \Leftrightarrow (4) notice that (8.3) is equivalent, for each $s \geq 0$, to

$$(1 - q^{m_r+1})A_{\mathbf{m}+1_r,s}(q) = A_{\mathbf{m},s}(q) - q^{1+|\mathbf{m}|}A_{\mathbf{m},s-1}(q) - q^{m_r+1+s}A_{\mathbf{m},s}(q) + q^{m_r+1+(s-1)}A_{\mathbf{m},s-1}(q),$$

which, in turn, is equivalent to (8.4) by dividing by $(1 - q)$. We next prove (2) \Rightarrow (3) (resp. (3) \Rightarrow (2)) by means of the *finite difference q -calculus* (resp. the *q -iteration*) given next.

8.1. *A finite difference q -calculus.* Let

$$(8.5) \quad A(t, q; \mathbf{u}) = A(t, q; u_1, \dots, u_r) := \sum_{\mathbf{m}} A_{\mathbf{m}}(t, q) \frac{\mathbf{u}^{\mathbf{m}}}{(t; q)_{1+|\mathbf{m}|}}$$

denote the left-hand side of (8.2) and form the *q -finite difference* applied to the sole variable u_r :

$$D_{u_r} := A(t, q; u_1, \dots, u_r) - A(t, q; u_1, \dots, u_{r-1}, u_r q).$$

We get:

$$\begin{aligned}
 D_{u_r} &= \sum_{\mathbf{m}}_{m_r \geq 1} A_{\mathbf{m}}(t, q) \frac{\mathbf{u}^{\mathbf{m}}}{(t; q)_{1+|\mathbf{m}|}} - \sum_{\mathbf{m}}_{m_r \geq 1} A_{\mathbf{m}}(t, q) \frac{u_1^{m_1} \dots u_{r-1}^{m_{r-1}} (u_r q)^{m_r}}{(t; q)_{1+|\mathbf{m}|}} \\
 &= \sum_{\mathbf{m}} A_{\mathbf{m}+1_r}(t, q) \frac{\mathbf{u}^{\mathbf{m}+1_r}}{(t; q)_{2+|\mathbf{m}|}} \\
 &\quad - \sum_{\mathbf{m}} A_{\mathbf{m}+1_r}(t, q) \frac{u_1^{m_1} \dots u_{r-1}^{m_{r-1}} (u_r q)^{m_r+1}}{(t; q)_{2+|\mathbf{m}|}},
 \end{aligned}$$

so that

$$(8.2) \quad D_{u_r} = \sum_{\mathbf{m}} (1 - q^{m_r+1}) A_{\mathbf{m}+1_r}(t, q) \frac{\mathbf{u}^{\mathbf{m}+1_r}}{(t; q)_{2+|\mathbf{m}|}}.$$

Now use the *right-hand side* of (8.2):

$$\begin{aligned}
 D_{u_r} &= \sum_{s \geq 0} \frac{t^s}{(u_1; q)_{s+1} \dots (u_r; q)_{s+1}} - \sum_{s \geq 0} \frac{t^s}{(u_1; q)_{s+1} \dots (u_r q; q)_{s+1}} \\
 &= \sum_{s \geq 0} \frac{t^s}{(\mathbf{u}; q)_{s+1}} \left[1 - \frac{1 - u_r}{1 - u_r q^{s+1}} \right] \\
 &= u_r \sum_{s \geq 0} \frac{t^s}{(\mathbf{u}; q)_{s+1}} \left[1 - q^{s+1} \frac{1 - u_r}{1 - u_r q^{s+1}} \right] \\
 &= u_r (A(t, q; u_1, \dots, u_r) - qA(tq, q; u_1, \dots, u_{r-1}, u_r q)).
 \end{aligned}$$

This yields:

$$(8.6) \quad \begin{aligned} &A(t, q; u_1, \dots, u_r) - A(t, q; u_1, \dots, u_{r-1}, u_r q) \\ &= u_r (A(t, q; u_1, \dots, u_r) - qA(tq, q; u_1, \dots, u_{r-1}, u_r q)). \end{aligned}$$

Re-write every term on the right-hand side by means of the polynomials $A_{\mathbf{m}}(t, q)$. We get:

$$(8.7) \quad \begin{aligned} u_r A(t, q; \mathbf{u}) &= \sum_{\mathbf{m}} A_{\mathbf{m}}(t, q) \frac{\mathbf{u}^{\mathbf{m}+1_r}}{(t; q)_{1+|\mathbf{m}|}} \\ &= \sum_{\mathbf{m}} (1 - tq^{1+|\mathbf{m}|}) A_{\mathbf{m}}(t, q) \frac{\mathbf{u}^{\mathbf{m}+1_r}}{(t; q)_{2+|\mathbf{m}|}}. \end{aligned}$$

In the same manner,

$$(8.5) \quad \begin{aligned} u_r q A(tq, q; u_1, \dots, u_r q) &= \sum_{\mathbf{m}} A_{\mathbf{m}}(tq, q) \frac{u_1^{m_1} \dots u_{r-1}^{m_{r-1}} (u_r q)^{m_r+1}}{(tq; q)_{1+|\mathbf{m}|}} \\ &= \sum_{\mathbf{m}} q^{m_r+1} (1 - t) A_{\mathbf{m}}(tq, q) \frac{\mathbf{u}^{\mathbf{m}+1_r}}{(t; q)_{2+|\mathbf{m}|}}. \end{aligned}$$

Taking (8.5)—(8.7) into account we deduce the recurrence relation (8.4).

8. THE EULER-MAHONIAN POLYNOMIALS

8.2. *A q-iteration method.* By using the notation (8.5) we see that the recurrence relation (8.3) can be rewritten:

$$(8.8) \quad A(t, q; \mathbf{u}) - A(t, q; u_1, \dots, u_i q \dots, u_r) \\ = u_i A(t, q; \mathbf{u}) - u_i q A(tq, q; u_1, \dots, u_i q, \dots, u_r).$$

Let $A(t, q; \mathbf{u}) := \sum_{s \geq 0} t^s G_s(\mathbf{u}, q)$. We deduce:

$$\sum_{s \geq 0} t^s (1 - u_i) G_s(\mathbf{u}, q) = \sum_{s \geq 0} t^s (1 - u_i q^{s+1}) G_s(u_1, \dots, u_i q, \dots, u_r, q).$$

Now take the coefficient of t^s on each side. We obtain:

$$(8.9) \quad G_s(\mathbf{u}, q) = \frac{1 - u_i q^{s+1}}{1 - u_i} G_s(u_1, \dots, u_i q, \dots, u_r, q),$$

for $i = 1, \dots, r$. Then, let $F_s(\mathbf{u}, q) := G_s(\mathbf{u}, q)(\mathbf{u}; q)_{s+1}$ and use (8.9). For $i = 1, \dots, r$ we obtain the equation

$$(8.10) \quad F_s(\mathbf{u}, q) = F_s(u_1, \dots, u_i q, \dots, u_r, q).$$

But we can write $F_s(\mathbf{u}, q) = \sum_{\mathbf{m}} \mathbf{u}^{\mathbf{m}} F_{s, \mathbf{m}}(q)$, where $F_{s, \mathbf{m}}(q)$ is a (positive) power series in q . Fix the multi-index \mathbf{m} and let m_i be a nonzero component of \mathbf{m} . Relation (8.10) implies: $F_{s, \mathbf{m}}(q) = q^{m_i} F_{s, \mathbf{m}}(q)$. Hence, $F_{s, \mathbf{m}}(q) = 0$ and $F_s(\mathbf{u}, q) = F_{s, 0}(q)$, a quantity that remains to be evaluated. But, by definition of $F_s(u, q)$, we have:

$$F_{s, 0}(q) = F_s(\mathbf{u}, q) \Big|_{\mathbf{u} = 0} = G_s(\mathbf{u}, q)(\mathbf{u}; q)_{s+1} \Big|_{\mathbf{u} = 0} = G_s(0, q) = 1,$$

since $\sum_{s \geq 0} t^s G_s(0, q) = A(t, q; 0) = \frac{1}{(t; q)_1} = \sum_{s \geq 0} t^s$. Thus $G_s(\mathbf{u}, q) = \frac{1}{(\mathbf{u}; q)_{s+1}}$. \square

Remark 8.2. If σ is a permutation of the set $\{1, \dots, r\}$, denote by $\sigma \mathbf{m}$ the sequence $(m_{\sigma(1)}, \dots, m_{\sigma(r)})$. In particular, $R(\sigma \mathbf{m})$ is the class of all the rearrangements of the word $1^{m_{\sigma(1)}} \dots r^{m_{\sigma(r)}}$. As the product of the binomial coefficient on the right-hand side of (8.1) is *symmetric* in m_1, \dots, m_r , we conclude that for every permutation σ we have:

$$(8.11) \quad A_{\sigma \mathbf{m}}(t, q) = A_{\mathbf{m}}(t, q).$$

Remark 8.3. The relations (8.3) and (8.4) provide the evaluations of the first values of the polynomials $A_{\mathbf{m}}(t, q)$. Because of the previous remark it suffices to make the calculations for the nonincreasing sequences \mathbf{m} .

Table of the polynomials $A_{\mathbf{m}}(t, q)$:

$$\begin{aligned} A_{(1)} &= 1; & A_{(1,1)} &= 1 + tq; & A_{(2)} &= 1; & A_{(1,1,1)} &= 1 + t(2q + 2q^2) + t^2q^3; \\ A_{(2,1)} &= 1 + t(q + q^2); & A_{(3)} &= 1; \\ A_{(1,1,1,1)} &= 1 + t(3q + 5q^2 + 3q^3) + t^2(3q^3 + 5q^4 + 3q^5) + t^3q^6; \\ A_{(2,1,1)} &= 1 + t(2q + 3q^2 + 2q^3) + t^2(q^2 + 2q^4 + q^5); \\ A_{(2,2)} &= 1 + t(q + 2q^2 + q^3) + t^2q^4; & A_{(3,1)} &= 1 + t(q + q^2 + q^3); & A_{(4)} &= 1. \end{aligned}$$

9. The insertion technique

In section 7 we have shown that the Euler-Mahonian polynomial $A_{\mathbf{m}}(t, q)$ was the generating polynomial for the class $R(\mathbf{m})$ by the bi-statistic (des, maj) . To derive the result we made use of the MacMahon *Verfahren* and obtained identity (8.1). The natural question is whether we can prove the same result by using one of the two *recurrence relations* (8.3), (8.4). If we dealt with the symmetric group (i.e., with all the m_i 's equal to 1), we would try the traditional *insertion technique*: start with a permutation of order r and study the modification brought to the underlying statistic when the letter $(r + 1)$ is *inserted* into the $(r + 1)$ slots of the permutation. The technique can be applied without any difficulty. Identity (8.4)—with all the m_i 's equal to 1—is then easily derived.

With words with repeated letters the derivation is not straightforward. A transformation called *word marking* must be made on the initial word. The word marking goes as follows. This time we consider the polynomial

$$(9.1) \quad A_{\mathbf{m}}(t, q) = \sum_{s \geq 0} A_{\mathbf{m},s}(q)t^s$$

as being the generating polynomial for $R(\mathbf{m})$ by the bi-statistic (des, maj) , so that $A_{\mathbf{m},s}(q)$ is the generating polynomial for the words $w \in R(\mathbf{m})$ such that $\text{des } w = s$ by the Major Index. Again use the notations $[s]_q := 1 + q + q^2 + \dots + q^{s-1}$, $|\mathbf{m}| := m_1 + \dots + m_r$ and $\mathbf{m} + 1_j := (m_1, \dots, m_j + 1, \dots, m_r)$ for each $j = 1, 2, \dots, r$ and each sequence $\mathbf{m} = (m_1, m_2, \dots, m_r)$.

Proposition 9.1. *Let $1 \leq j \leq r$ and let $A_{\mathbf{m}}(t, q)$ be the generating polynomial for $R(\mathbf{m})$ by the bi-statistic (des, maj) . Then the following*

9. THE INSERTION TECHNIQUE

relations hold:

$$(9.2) \quad (1 - q^{m_j+1})A_{\mathbf{m}+1_j}(t, q) \\ = (1 - tq^{1+|\mathbf{m}|})A_{\mathbf{m}}(t, q) - q^{m_j+1}(1 - t)A_{\mathbf{m}}(tq, q);$$

$$(9.3) \quad [m_j + 1]_q A_{\mathbf{m}+1_j, s}(q) \\ = [m_j + 1 + s]_q A_{\mathbf{m}, s}(q) + q^{s+m_j}[1 + |\mathbf{m}| - s - m_j]_q A_{\mathbf{m}, s-1}(q).$$

Proof. The latter identity is equivalent to the former one, so that only (9.3) is to be proved. From Remark 8.3 this relation is equivalent to the relation formed when j is replaced by any integer in $\{1, \dots, r\}$. It is convenient to prove the relation for $j = 1$ which reads

$$(9.4) \quad (1 + q + \dots + q^{m_1})A_{\mathbf{c}+1_1, s}(q) \\ = (1 + q + \dots + q^{m_1+s})A_{\mathbf{m}, s}(q) + (q^{m_1+s} + \dots + q^{|\mathbf{m}|})A_{\mathbf{m}, s-1}(q).$$

Consider the set $R^*(\mathbf{m} + 1_1, s)$ of 1-*marked* words, i.e., rearrangements w^* of $1^{m_1+1} \dots r^{m_r}$ with s descents such that exactly one letter equal to 1 has been marked. Each word $w \in R(\mathbf{m} + 1_1)$ that has s descents gives rise to $m_1 + 1$ marked words $w^{(0)}, \dots, w^{(m_1)}$. Define

$$\text{maj}^* w^{(i)} := \text{maj} w + n_1,$$

where n_1 is the number of letters equal to 1 to the *right* of the marked 1. Then clearly

$$\sum_{i=0}^{m_1} \text{maj}^* w^{(i)} = (1 + q + \dots + q^{m_1}) \text{maj} w.$$

Hence

$$(1 + q + \dots + q^{m_1})A_{\mathbf{m}+1_1, s}(q) = \sum_{w \in R^*(\mathbf{m}+1_1, s)} q^{\text{maj}^* w}.$$

Let $m = |\mathbf{m}|$ and let the word $w = x_1 x_2 \dots x_m \in R(\mathbf{m})$ have s descents. Say that w has $m + 1$ slots $x_i x_{i+1}$, $i = 0, \dots, m$ (where $x_0 = 0$ and $x_{m+1} = \infty$ by convention). Call the slot $x_i x_{i+1}$ *green* if either $x_i x_{i+1}$ is a descent, $x_i = 1$, or $i = 0$. Call the other slots *red*. Then there are $1 + s + m_1$ green slots and $m - s - m_1$ red slots. Label the green slots $0, 1, \dots, m_1 + s$ from right to left, and label the red slots $m_1 + s + 1, \dots, m$ from left to right.

For example, with $r = 3$, the word $w = 2, 2, 1, 3, 2, 1, 2, 3, 3$ has three descents and ten slots. As $m_1 = 2$, there are eight green slots and two red slots, labelled as follows

slot	0		2		2		1		3		2		1		2		3		3		∞
label	5		6		4		3		2		1		0		7		8		9		

i	$w^{(i)}$	$\text{des } w^{(i)}$	$\text{maj}^* w^{(i)}$
0	2 2 \frown 1 3 \frown 2 \frown 1 1 2 3 3	3	11
1	2 2 \frown 1 3 \frown 2 \frown 1 1 2 3 3	3	12
2	2 2 \frown 1 3 \frown 1 2 \frown 1 2 3 3	3	13
3	2 2 \frown 1 1 3 \frown 2 \frown 1 2 3 3	3	14
4	2 2 \frown 1 1 3 \frown 2 \frown 1 2 3 3	3	15
5	1 2 2 \frown 1 3 \frown 2 \frown 1 2 3 3	3	16
6	2 \frown 1 2 \frown 1 3 \frown 2 \frown 1 2 3 3	4	17
7	2 2 \frown 1 3 \frown 2 \frown 1 2 \frown 1 3 3	4	18
8	2 2 \frown 1 3 \frown 2 \frown 1 2 3 \frown 1 3	4	19
9	2 2 \frown 1 3 \frown 2 \frown 1 2 3 3 \frown 1	4	20

Table 9.1

Denote by $w^{(i)}$ the word obtained from w by inserting a marked 1 into the i -th slot. Then it may be verified that

$$(9.5) \quad \text{des } w^{(i)} = \begin{cases} \text{des } w, & \text{if } i \leq m_1 + s; \\ \text{des } w + 1, & \text{otherwise.} \end{cases}$$

$$(9.6) \quad \text{maj}^* w^{(i)} = \text{maj } w + i.$$

Example. Consider the above word w . In Table 9.1 the values of “des” and “maj*” on $w^{(i)}$. Descents are indicated by \frown and the marked **1** is written in boldface.

So each word $w \in R(\mathbf{m})$ with s descents and $\text{maj } w = n$ gives rise to $m_1 + s + 1$ marked words in $R^*(\mathbf{m} + 1_1, s)$ with maj^* equal to $n, n+1, \dots, n+m_1+s$; and to $m-s-m_1$ marked words in $R^*(\mathbf{m}+1_1, s+1)$ with maj^* equal to $n+m_1+s+1, \dots, n+m$. Hence a word w in $R(\mathbf{m})$ with $s-1$ descents gives rise to $m-s+1-m_1$ marked words in $R^*(\mathbf{m}+1_1, s)$ with maj^* equal to $\text{maj } w+m_1+s, \dots, \text{maj } w+m$. This now proves relation (9.4). \square

10. The two forms of the q -Eulerian polynomials

When the multi-index \mathbf{m} is of the form $(1^r) = (1, 1, \dots, 1)$, the Euler-Mahonian polynomial $A_{\mathbf{m}}(t, q)$ will be denoted by ${}^{\text{maj}}A_r(t, q)$ and referred to as the q -maj-Eulerian polynomial. It also follows from the previous two sections that ${}^{\text{maj}}A_r(t, q)$ is the generating polynomial for the symmetric group \mathfrak{S}_r by the bi-statistic (des, maj) . As for the polynomial $A_{\mathbf{m}}(t, q)$ for an arbitrary \mathbf{m} , the new polynomial ${}^{\text{maj}}A_r(t, q)$ can be characterized in four different ways, as shown in the next definition.

10. THE TWO FORMS OF THE Q-EULERIAN POLYNOMIALS

Definition 10.1. A sequence $({}^{\text{maj}}A_r(t, q))$ of polynomials in two variables t and q , indexed by the integers $r \geq 0$, is said to be *q-maj-Eulerian*, if one of the following *equivalent* four conditions holds:

(1) For every integer $r \geq 0$ we have:

$$(10.1) \quad \frac{1}{(t; q)_{r+1}} {}^{\text{maj}}A_r(t, q) = \sum_{s \geq 0} t^s ([s+1]_q)^r.$$

(2) The ordinary (resp. exponential) generating function for the ratios $\frac{{}^{\text{maj}}A_r(t, q)}{(t; q)_{r+1}}$ is given by:

$$(10.2a) \quad \sum_{r \geq 0} u^r \frac{{}^{\text{maj}}A_r(t, q)}{(t; q)_{r+1}} = \sum_{s \geq 0} t^s \frac{1}{1 - u [s+1]_q};$$

$$(10.2b) \quad \sum_{r \geq 0} \frac{u^r}{r!} \frac{{}^{\text{maj}}A_r(t, q)}{(t; q)_{r+1}} = \sum_{s \geq 0} t^s \exp(u [s+1]_q).$$

(3) The recurrence relation holds:

$$(10.3) \quad (1 - q) {}^{\text{maj}}A_r(t, q) = (1 - tq^r) {}^{\text{maj}}A_{r-1}(t, q) - q(1 - t) {}^{\text{maj}}A_{r-1}(tq, q).$$

(4) With ${}^{\text{maj}}A_r(t, q) := \sum_{s \geq 0} t^s {}^{\text{maj}}A_{r,s}(q)$ the coefficients ${}^{\text{maj}}A_{r,s}(q)$ satisfy the recurrence:

$$(10.4) \quad {}^{\text{maj}}A_{r,s}(q) = [s+1]_q {}^{\text{maj}}A_{r-1,s}(q) + q^s [r-s]_q {}^{\text{maj}}A_{r-1,s-1}(q).$$

In the above definition (10.1), is the specialization of (8.1). However (8.2a) and (8.2b) have no immediate counterparts, but the exponential generating function for the ratios $\frac{1}{(t; q)_{r+1}} {}^{\text{maj}}A_r(t, q)$ has an interesting closed form, as written in (10.2). Finally, (10.3) and (10.4) are straightforward specializations of (8.3) and (8.4) when $\mathbf{m} = 1^{r-1}$.

When $q = 1$ in (10.1), (10.2) and (10.4), we recognize some familiar definitions for the so-called *Eulerian polynomials*. Let us introduce them following the same pattern as above.

Definition 10.2. A sequence $(A_r(t))$ of polynomials in one variable t , indexed by the integers $r \geq 0$, is said to be *Eulerian*, if one of the following *equivalent* five conditions holds:

(1) For every integer $r \geq 0$ we have:

$$(10.5) \quad \frac{1}{(1-t)^{r+1}} A_r(t) = \sum_{s \geq 0} t^s (s+1)^r.$$

(2) The exponential generating function for the ratios $\frac{A_r(t)}{(1-t)^{r+1}}$ is given by:

$$(10.6) \quad \sum_{r \geq 0} \frac{u^r}{r!} \frac{A_r(t)}{(1-t)^{r+1}} = \sum_{s \geq 0} t^s \exp(u(s+1)) = \frac{e^u}{1-te^u}.$$

(3) The following recurrence relation holds:

$$(10.7) \quad A_r(t) = (1 + (r-1)t) A_{r-1}(t) + t(1-t) A'_{r-1}(t),$$

where $A'_{r-1}(t)$ denotes the derivative of the polynomial $A_{r-1}(t)$.

(4) With $A_r(t) := \sum_{s \geq 0} t^s A_{r,s}$ the coefficients $A_{r,s}$ satisfy the recurrence:

$$(10.8) \quad A_{r,s} = (s+1) A_{r-1,s} + (r-s) A_{r-1,s-1}.$$

(5) The exponential generating function for the polynomials reads:

$$(10.9) \quad \sum_{r \geq 0} \frac{u^r}{r!} A_r(t) = \frac{1-t}{-t + \exp(u(t-1))}.$$

Notice that (10.1), (10.2) and (10.4) that *define* the q -maj Eulerian polynomials are reduced to their counterparts (10.5), (10.6) and (10.8) that *define* the Eulerian polynomials, when q is given the value 1. On the other hand, we go from recurrence (10.8) to the q -recurrence (10.4) by replacing the integers $(s+1)$ and $(r-s)$ occurring in the relation by their q -counterparts $[s+1]_q = 1 + q + \dots + q^s$ and $q^s[r-s]_q = q^{r-s} + q^{r-s+1} + \dots + q^{r-1}$, respectively. We then say that the q -maj Eulerian polynomial $A_r(t, q)$ is a q -*analog* of the Eulerian polynomial $A_r(t)$.

Also notice that there is no specialization of (10.3) for $q = 1$. To obtain a recurrence for the polynomials $A_r(t)$ themselves, as shown in (10.7), we start from (10.8) and make the appropriate identifications. Finally, observe that (10.9) is simply derived from (10.6) with the substitution $u \leftarrow u/(1-t)$. The Eulerian polynomial $A_r(t)$ is the generating polynomial for \mathfrak{S}_r by the number of descents “des.”

Another q -extension of the Eulerian polynomials can be achieved by using the defining relation (10.9) or the exponential generating function for the polynomials $t A_n(t)$, that can be directly derived from (10.9) and reads

$$1 + \sum_{n \geq 1} \frac{u^n}{n!} t A_n(t) = \frac{1-t}{1-t \exp((1-t)u)}.$$

10. THE TWO FORMS OF THE Q-EULERIAN POLYNOMIALS

In the above fraction make the substitution $\exp(u) \leftarrow e_q(u)$ and express the fraction so derived as a q -series

$$(10.10) \quad \sum_{n \geq 0} \frac{u^n}{(q; q)_n} \text{inv}A_n(t, q) = \frac{1-t}{1-t e_q((1-t)u)},$$

where the $\text{inv}A_n(t, q)$'s are coefficients to be determined. Identity (10.10) can be rewritten as

$$\sum_{n \geq 0} \text{inv}A_n(t, q) \frac{u^n}{(q; q)_n} = \left(1 - t \sum_{n \geq 1} (1-t)^{n-1} \frac{u^n}{(q; q)_n} \right)^{-1},$$

so that the identity

$$\sum_{n \geq 0} \text{inv}A_n(t, q) \frac{u^n}{(q; q)_n} \cdot \left(1 - t \sum_{n \geq 1} (1-t)^{n-1} \frac{u^n}{(q; q)_n} \right) = 1$$

provides the recurrence: $\text{inv}A_0(t, q) = 1$ and

$$(10.11) \quad \text{inv}A_n(t, q) = \sum_{0 \leq k \leq n-1} \begin{bmatrix} n \\ k \end{bmatrix} \text{inv}A_k(t, q) t (1-t)^{n-1-k} \quad (n \geq 1),$$

so that the coefficients $\text{inv}A_n(t, q)$ are *polynomials* in the two variables t and q with *integral coefficients*. Let q tend to 1 in (10.11) and let ${}^t B_n(t) := \text{inv}A_n(t, q) \big|_{q=1}$ ($n \geq 1$). This yields

$${}^t B_n(t) = \sum_{0 \leq k \leq n-1} \binom{n}{k} {}^t B_k(t) t (1-t)^{n-1-k} \quad (n \geq 1),$$

which, in turn, is equivalent to

$$1 + \sum_{n \geq 1} {}^t B_n(t) \frac{u^n}{n!} = \frac{1-t}{1-t \exp(u(1-t))}.$$

Hence

$$1 + \sum_{n \geq 1} B_n(t) \frac{u^n}{n!} = \frac{1-t}{-t + \exp(u(t-1))},$$

which is the right-hand side of (10.9). Hence $B_n(t) = A_n(t)$ and ${}^t A_n(t)$ is the generating polynomial for \mathfrak{S}_n by the statistic $1 + \text{des}$. Now what can be said about the polynomial $\text{inv}A_n(t, q)$?

Theorem 10.1. For $n \geq 1$ the polynomial $\text{inv}A_n(t, q)$ in the expansion

$$(10.12) \quad \frac{1-t}{1-te_q((1-t)u)} = \sum_{n \geq 0} \text{inv}A_n(t, q) \frac{u^n}{(q; q)_n}$$

is the generating polynomial for \mathfrak{S}_n by the bi-statistic $(1 + \text{des}, \text{inv})$, i.e.,

$$(10.13) \quad \text{inv}A_n(t, q) = \sum_{\sigma \in \mathfrak{S}_n} t^{1+\text{des } \sigma} q^{\text{inv } \sigma}.$$

To prove Theorem 10.1 we will show that the induction formula (10.11) holds for the polynomial defined by (10.13). For $k = 0, 1, \dots, (n-1)$ let $u_k := \begin{bmatrix} n \\ k \end{bmatrix} \text{inv}A_k(t, q) t$; then, by iteration on $k = 0, 1, 2, \dots, (n-1)$, define

$$(10.14) \quad G_{-1} := 0; \quad G_k := u_k + (1-t)G_{k-1}.$$

We see that for proving the theorem it suffices to show that:

$$(10.15) \quad \text{inv}A_n(t, q) = G_{n-1}.$$

But (10.15) follows immediately from the following lemma.

Lemma 10.2. For each $k = 0, 1, \dots, (n-1)$ the polynomial G_k defined by the recurrence (10.14) is the generating polynomial, by the bi-statistic $(1 + \text{des}, \text{inv})$, for the set of the permutations of order n , whose longest increasing right factor is of length at least equal to $(n-k)$, i.e., permutations $\sigma = \sigma(1) \dots \sigma(n)$ such that $\sigma(k+1) < \sigma(k+2) < \dots < \sigma(n)$.

Proof. By Proposition 4.3 we have

$$\begin{bmatrix} n \\ k \end{bmatrix} = \sum_{(A, B)} q^{\text{inv}(\gamma(A)\gamma(B))},$$

where the sum is over all ordered partitions (A, B) of $[n]$ into two blocks such that $|A| = k$ and $|B| = n-k$. Recall that $\gamma(A)$ and $\gamma(B)$ designate the *increasing* words whose letters are the elements of A and of B , respectively. With \mathfrak{S}_A denoting the group of the permutations of A we have

$$\begin{aligned} \begin{bmatrix} n \\ k \end{bmatrix} \text{inv}A_k(t, q) &= \sum_{(A, B)} \sum_{\tau \in \mathfrak{S}_k} q^{\text{inv}(\gamma(A)\gamma(B)) + \text{inv } \tau} t^{1+\text{des } \tau} \\ &= \sum_{(A, B)} \sum_{\tau \in \mathfrak{S}_A} q^{\text{inv}(\gamma(A)\gamma(B)) + \text{inv } \tau} t^{1+\text{des } \tau}, \end{aligned}$$

11. MAJOR INDEX AND INVERSION NUMBER

since A is of cardinality k . As the $(n - k)$ terms of $\gamma(B)$ are in increasing order, the mapping $(\gamma(A)\gamma(B), \tau) \mapsto \sigma$ defined by $\sigma := \tau\gamma(B)$ is a bijection onto the set of the permutations σ , whose longest increasing right factor is of length at least equal to $(n - k)$. Moreover $\text{inv } \sigma = \text{inv } \gamma(A)\gamma(B) + \text{inv } \tau$ and $\text{des } \sigma = \text{des } \tau + \chi(\sigma(k) > \sigma(k+1))$. Let $\mathfrak{S}_{n,k}$ be the set of the permutations whose longest increasing right factor is *exactly* of length $(n - k)$ and F_k be the generating polynomial for $\mathfrak{S}_{n,k}$ by the bi-statistic $(1 + \text{des}, \text{inv})$. Then

$$\begin{aligned} \begin{bmatrix} n \\ k \end{bmatrix} \text{inv}A_k(t, q) &= \sum_{\sigma \in \mathfrak{S}_{n,0} \cup \dots \cup \mathfrak{S}_{n,k}} q^{\text{inv } \sigma} t^{1 + \text{des } \sigma - \chi(\sigma(k) > \sigma(k+1))} \\ &= \sum_{\sigma \in \mathfrak{S}_{n,0} \cup \dots \cup \mathfrak{S}_{n,k-1}} q^{\text{inv } \sigma} t^{1 + \text{des } \sigma} + t^{-1} \sum_{\sigma \in \mathfrak{S}_{n,k}} q^{\text{inv } \sigma} t^{1 + \text{des } \sigma} \\ &= F_0 + \dots + F_{k-1} + t^{-1} F_k. \end{aligned}$$

Hence, by letting $G_k := F_0 + \dots + F_k$ and by multiplying the identity by t ,

$$u_k = tG_{k-1} + (G_k - G_{k-1}) = G_k + (t - 1)G_{k-1}$$

and then

$$G_k = u_k + (1 - t)G_{k-1},$$

which is precisely the induction relation (10.14). \square

The polynomials $\text{maj}A_n(t, q)$ and $\text{inv}A_n(t, q)$ form two q -analogs of the Eulerian polynomial $A_n(t)$. The polynomials $t^{\text{maj}A_n}(t, q)$ and $\text{inv}A_n(t, q)$ already differ for $n = 4$. Notice that Theorem 10.1 implies that $\text{inv}A_n(t, q)$ is a polynomial with *nonnegative* integral coefficients.

Table of the polynomials $\text{maj}A_n(t, q)$ and $\text{inv}A_n(t, q)$.

$$\begin{aligned} \text{maj}A_1(t, q) &= 1; \text{maj}A_2(t, q) = 1 + tq; \text{maj}A_3(t, q) = 1 + 2tq(q + 1) + t^2q^3; \\ \text{maj}A_4(t, q) &= 1 + tq(3q^2 + 5q + 3) + t^2q^3(3q^2 + 5q + 3) + t^3q^6. \\ \text{inv}A_1(t, q) &= t; \text{inv}A_2(t, q) = t + t^2q; \text{inv}A_3(t, q) = t + 2t^2q(q + 1) + t^3q^3; \\ \text{inv}A_4(t, q) &= t + t^2(q^4 + 3q^3 + 4q^2 + 3q) + t^3(3q^5 + 4q^4 + 3q^3 + q^2) + t^3q^6. \end{aligned}$$

11. Major Index and Inversion Number

Again, let $R(\mathbf{m})$ be the set of all rearrangements of $1^{m_1}2^{m_2} \dots r^{m_r}$. In theorems 6.1 and 6.3 it was proved that the generating polynomial for $R(\mathbf{m})$ by the Inversion Number “inv”, on the one hand, and by the Major Index “maj”, on the other hand, was equal to the q -multinomial

coefficient $\left[\begin{smallmatrix} m \\ m_1, m_2, \dots, m_r \end{smallmatrix} \right]$, where $m = m_1 + m_2 + \dots + m_r$. If we write those two generating polynomials under the form

$$\sum_{k \geq 0} q^k |\{w \in R(\mathbf{m}) : \text{inv } w = k\}| \quad \text{and} \quad \sum_{k \geq 0} q^k |\{w \in R(\mathbf{m}) : \text{maj } w = k\}|,$$

we see that for every $k \geq 0$ we have:

$$(11.1) \quad |\{w \in R(\mathbf{m}) : \text{inv } w = k\}| = |\{w \in R(\mathbf{m}) : \text{maj } w = k\}|.$$

Hence, for every $k \geq 0$ there *exists* a bijection of the set $\{w \in R(\mathbf{m}) : \text{inv } w = k\}$ onto the set $\{w \in R(\mathbf{m}) : \text{maj } w = k\}$; this is equivalent to saying that there *exists* a bijection Φ de $R(\mathbf{m})$ onto $R(\mathbf{m})$ with the property that

$$(11.2) \quad \text{maj } w = \text{inv } \Phi(w),$$

for every $w \in R(\mathbf{m})$. This brings up the problem of *constructing* such a bijection, that is to say, of inventing an explicit algorithm that transforms a word $w \in R(\mathbf{m})$ into a word $w' \in R(\mathbf{m})$ such that $\text{maj } w = \text{inv } w'$ in a one-to-one manner.

Of course, when all the m_i 's are equal to 1 and the rearrangement class $R(\mathbf{m})$ is simply the symmetric group \mathfrak{S}_r , the construction of such a bijection can be made by means of the maj- and inv-codings introduced in section 2. For arbitrary rearrangement classes we have to follow another route, but the route will be richer, as further properties will be given for free, in particular when we restrict ourselves to the symmetric group.

11.1. *How to construct a bijection.* The proofs of Theorems 6.1 and 6.3 were so different in nature that there was no hint for imagining any immediate construction. However, if we make up a table of the first values of the Major Index and Inversion Number for small classes $R(\mathbf{m})$, we *observe* a further property. Let $L(w)$ (“*L*” for “*last*”) be the *last* letter of the word w . Then, for every $k \geq 0$ and $x \in X = \{1, 2, \dots, r\}$ we have:

$$(11.3) \quad |\{w \in R(\mathbf{m}) : \text{inv } w = k, L(w) = x\}| \\ = |\{w \in R(\mathbf{m}) : \text{maj } w = k, L(w) = x\}|.$$

In Fig. 11.1 the values of those two statistics have been calculated for the words of the set $R(2, 1, 1)$, the rearrangements of the word 1, 1, 2, 3. In the first, second, third table the words ending by 1, 2, 3 are respectively listed. We can observe that the distribution of “maj” and “inv” in each table is the same.

11. MAJOR INDEX AND INVERSION NUMBER

w	maj	inv
1, 2, 3, 1	3	2
1, 3, 2, 1	5	3
2, 1, 3, 1	4	3
2, 3, 1, 1	2	4
3, 1, 2, 1	4	4
3, 2, 1, 1	3	5

w	maj	inv
1, 1, 3, 2	3	1
1, 3, 1, 2	2	2
3, 1, 1, 2	1	3

w	maj	inv
1, 1, 2, 3	0	0
1, 2, 1, 3	2	1
2, 1, 1, 3	1	2

Fig. 11.1

For each letter x belonging to a linearly ordered alphabet X and each word w let $\text{bot}_x(w)$ (resp. $\text{top}_x(w)$), resp. $|w|_x$ denote the number of letters in w which are less than or equal to (resp. greater than, resp. equal to) x . In particular,

$$(11.4) \quad \text{bot}_x(w) + \text{top}_x(w) = |w| \quad (\text{length of } w).$$

Further, let $R(\mathbf{m})x$ be the set of words wx , where $w \in R(\mathbf{m})$. If w' is a rearrangement of w , the following properties hold:

$$(11.5) \quad \text{inv } wx = \text{inv } w + \text{top}_x(w);$$

$$(11.6) \quad \text{maj } wx = \begin{cases} \text{maj } w, & \text{if } L(w) \leq x; \\ \text{maj } w + \text{bot}_x(w) + \text{top}_x(w), & \text{if } L(w) > x. \end{cases}$$

Suppose that property (11.3) holds for every class $R(\mathbf{m})$. Then there exists a bijection $w \mapsto w'$ of $R(\mathbf{m})$ onto itself such that $\text{maj } w = \text{inv } w'$ and $L(w) = L(w')$. In the same manner, the letter x being given, there also exists a bijection $w \mapsto w''$ such that $\text{maj } wx = \text{inv } w''x$.

If $L(w) \leq x$, we then have:

$$\begin{aligned} L(w') &= L(w); \\ \text{inv } w' &= \text{maj } w \\ &= \text{maj } wx && [\text{by (11.6)}] \\ &= \text{inv } w''x \\ &= \text{inv } w'' + \text{top}_x(w') && [\text{by (11.5)}]. \end{aligned}$$

If $L(w) > x$, we also have:

$$\begin{aligned} L(w') &= L(w); \\ \text{inv } w' &= \text{maj } w \\ &= \text{maj } wx - \text{bot}_x(w) - \text{top}_x(w) && [\text{by (11.6)}] \\ &= \text{inv } w''x - \text{bot}_x(w) - \text{top}_x(w) \\ &= \text{inv } w'' - \text{bot}_x(w') && [\text{by (11.5)}]. \end{aligned}$$

Consequently, if property (11.3) holds, there exists a bijection γ_x of $R(\mathbf{m})$ onto itself, namely $w' \mapsto w''$, having the property

$$(11.7) \quad \text{inv } \gamma_x(w') = \begin{cases} \text{inv } w' - \text{top}_x(w'), & \text{if } L(w') \leq x ; \\ \text{inv } w' + \text{bot}_x(w'), & \text{if } L(w') > x. \end{cases}$$

Conversely, if there exists a bijection $\gamma_x : w' \mapsto w''$ that satisfies (11.7), we can define a bijection Φ of each class of rearrangements of words onto itself by letting

$$(11.8) \quad \Phi(w) := w,$$

if w is of length 1, and for each nonempty word w and each letter x by letting

$$(11.9) \quad \Phi(wx) := \gamma_x(\Phi(w))x.$$

Thus by induction we determine the image $\Phi(w)$ of w , then apply the bijection γ_x to $\Phi(w)$, finally the letter x is juxtaposed at the end of the word.

Theorem 11.1. *The following two statements are equivalent:*

- (a) *Property (11.3) holds for every class.*
- (b) *For every letter x there exists a bijection $\gamma_x : w' \mapsto w''$ such that property (11.7) holds; moreover, the bijection Φ defined by (11.8) and (11.9) has the properties:*

$$(11.10) \quad \text{maj } w = \text{inv } \Phi(w) \quad \text{and} \quad L(w) = L(\Phi(w)).$$

Proof. The implication (b) \Rightarrow (a) is straightforward. To prove the converse it suffices to verify (11.10). First, $L(w) = L(\Phi(w))$ by the definition of Φ given in (11.9). Then, for each nonempty word w and each letter x such that $L(w) \leq x$ we have:

$$\begin{aligned} \text{inv } \Phi(wx) &= \text{inv } \gamma_x(\Phi(w))x \\ &= \text{inv } \gamma_x(\Phi(w)) + \text{top}_x(w) && \text{by (11.5)]} \\ &= (\text{inv } \Phi(w) - \text{top}_x(\Phi(w))) + \text{top}_x(w) && \text{[by (11.7)]} \\ &= \text{maj } w \\ &= \text{maj } wx. \end{aligned}$$

If $L(w) > x$, we have:

$$\begin{aligned} \text{inv } \Phi(wx) &= \text{inv } \gamma_x(\Phi(w)) + \text{top}_x(w) \\ &= (\text{inv } \Phi(w) + \text{bot}_x(w)) + \text{top}_x(w) \\ &= \text{maj } w + |w| \\ &= \text{maj } wx. \quad \square \end{aligned}$$

11. MAJOR INDEX AND INVERSION NUMBER

Thus, the construction of a bijection Φ boils down to the constructions of bijections γ_x having property (11.7).

11.2. *The binary case.* Let a, b be two elements of the alphabet such that $a < b$ and consider a (binary) word w' of the class $R(m_a, m_b)$ with $m_a, m_b \geq 1$. We have

$$\begin{aligned} \text{bot}_x(w') &= \begin{cases} 0, & \text{if } x < a; \\ |w'|_a, & \text{if } a \leq x < b; \\ |w'|, & \text{if } b \leq x; \end{cases} \\ \text{top}_x(w') &= \begin{cases} |w'|, & \text{if } x < a; \\ |w'|_b, & \text{if } a \leq x < b; \\ 0, & \text{if } b \leq x. \end{cases} \end{aligned}$$

Condition (11.7) can be rewritten as

$$(11.11) \quad \text{inv } \gamma_x(w') = \begin{cases} \text{inv } w', & \text{if } x < a; \\ \text{inv } w' - |w'|_b, & \text{if } L(w') = a \leq x < b; \\ \text{inv } w' + |w'|_a, & \text{if } a \leq x < L(w') = b; \\ \text{inv } w', & \text{if } b = L(w') \leq x. \end{cases}$$

We can take the identity map for γ_x when $x < a$ or $b = L(w') \leq x$. Let vy belong to $R(m_a, m_b)$ with y the last letter equal to a or b . The *most straightforward* transformation we can think of for γ_x in the remaining two cases is

$$(11.12) \quad \gamma_x(vy) := yv,$$

which is obviously bijective and satisfies (11.11). The next theorem is then a consequence of Theorem 11.1.

Theorem 11.2. *Let $\{a, b\}$ be a two-letter alphabet ($a < b$) and for each $x = a, b$ let γ_x be defined, for every binary word v in the letters a, b , by*

$$(11.13) \quad \gamma_a(va) = av \quad \text{and} \quad \gamma_b(vb) = vb.$$

Then, the transformation Φ , as defined in (11.8) and (11.9), is a bijection of every rearrangement class onto itself having the property:

$$\text{maj } w = \text{inv } \Phi(w) \quad \text{and} \quad L(w) = L(\Phi(w)).$$

Example. Starting with $w = 0, 0, 1, 0, 1, 1, 0$, so that $\text{maj } w = 3+6 = 9$, we successively have:

v		$\Phi(v)$
0		0
0, 0	$\gamma_0(0) = 0$	0, 0
0, 0, 1	$\gamma_1(0, 0) = 0, 0$	0, 0, 1
0, 0, 1, 0	$\gamma_0(0, 0, 1) = 1, 0, 0$	1, 0, 0, 0
0, 0, 1, 0, 1	$\gamma_1(1, 0, 0, 0) = 1, 0, 0, 0$	1, 0, 0, 0, 1
0, 0, 1, 0, 1, 1	$\gamma_1(1, 0, 0, 0, 1) = 1, 0, 0, 0, 1$	1, 0, 0, 0, 1, 1
0, 0, 1, 0, 1, 1, 0	$\gamma_0(1, 0, 0, 0, 1, 1) = 1, 1, 0, 0, 0, 1$	1, 1, 0, 0, 0, 1, 0

Thus $\Phi(w) = 1, 1, 0, 0, 0, 1, 0$ and $\text{inv } \Phi(w) = 9 = \text{maj } w$.

11.3. *From the binary to the general case.* The next step is to start with the *natural* bijection γ_x , introduced in (11.13) and see how it can be extended to arbitrary words while keeping property (11.7).

For each $n \geq 1$ let $E_n := \{0, 1, 2, \dots, n\}$ and let E_n^* denote the set of all finite words whose letters belong to E_n . For $x, x_i \in E_n$ define

$$(11.14) \quad \beta_x(x_i) := \begin{cases} 0, & \text{if } x_i \leq x; \\ 1, & \text{if } x_i > x; \end{cases}$$

and for each word $w = x_1x_2 \dots x_m \in E_n^*$ let

$$u := \beta_x(x_1)\beta_x(x_2) \dots \beta_x(x_m).$$

If $i_1 < i_2 < \dots < i_a$ (resp. $j_1 < j_2 < \dots < j_b$) is the sequence of the subscripts i such that $x_i \leq x$ (resp. subscripts j such that $x_j > x$), let w_0, w_1 be the subwords: $w_0 = x_{i_1}x_{i_2} \dots x_{i_a}$ and $w_1 = x_{j_1}x_{j_2} \dots x_{j_b}$. Also let

$$(11.15) \quad B_x(w) := (u, w_0, w_1).$$

Proposition 11.3. *For each $x \in E_n$ the mapping B_x defined in (11.15) is a bijection of E_n^* onto the set of triples (u, w_0, w_1) such that $|w| = |u|$, $|u|_0 = |w_0|$ and $|u|_1 = |w_1|$ having the further property that*

$$(11.16) \quad \text{inv } w = \text{inv } u + \text{inv } w_0 + \text{inv } w_1.$$

Proof. Relation (11.16) simply indicates a sorting of the inversions within the word w . The bijective property is obvious. \square

Now, let w be a word in E_n^* and let x be a letter. Form the chain

$$(11.17) \quad w \xrightarrow{B_x} (u, w_0, w_1) \mapsto (\gamma_x(u), w_0, w_1) \xrightarrow{B_x^{-1}} w',$$

where B_x^{-1} designates the inverse of B_x , and define $\gamma_x(w) := w'$.

11. MAJOR INDEX AND INVERSION NUMBER

When w is a binary word, *i. e.* $n = 1$, the above chain gives back the value of $\gamma_x(w)$ defined in the previous subsection. Assume $n \geq 2$ and let $w = x_1x_2 \dots x_m$ be an arbitrary word. If all its letters are greater than (resp. less than or equal to) x , then $B_w(w) = (1^m, e, w)$ (resp. $B_x(w) = (0^m, w, e)$), so that $\gamma_x(w) = w$. Condition (11.7) holds.

If $L(w) \leq x$ and there is at least one letter of w greater than x , then $L(u) = 0$ and $\text{inv } \gamma_x(w) = \text{inv } \gamma_x u + \text{inv } w_0 + \text{inv } w_1 = \text{inv } u - |u|_1 + \text{inv } w_0 + \text{inv } w_1 = \text{inv } u - \text{top}_x(w) + \text{inv } w_0 + \text{inv } w_1 = \text{inv } w - \text{top}_x(w)$.

If $L(w) > x$ and there is at least one letter of w less than or equal to x , then $L(u) = 1$ and $\text{inv } \gamma_x(w) = \text{inv } \gamma_x u + \text{inv } w_0 + \text{inv } w_1 = \text{inv } u + |u|_0 + \text{inv } w_0 + \text{inv } w_1 = \text{inv } u + \text{bot}_x(w) + \text{inv } w_0 + \text{inv } w_1 = \text{inv } w + \text{bot}_x(w)$. Thus, condition (11.7) always holds.

Accordingly, our program is fulfilled: by means of the bijections defined in (11.13) for binary words, then, the bijections B_x defined in (11.15) and finally the chain (11.17), we hold a family of bijections γ_x that, when incorporated in the definition of Φ given in (11.9), provide a bijection having properties (11.10).

The three steps (11.13), (11.15), (11.17) can be combined and described more quickly by making use of the *x-factorisation* defined as follows: let x be a letter and w a word. If the last letter $L(w)$ of w is less than or equal to (resp. greater than) x , the word w admits the unique factorization:

$$(v_1y_1, v_2y_2, \dots, v_py_p),$$

called its *x-factorisation* having the following properties:

- (i) each y_i ($1 \leq i \leq p$) is a *letter* verifying $y_i \leq x$ (resp. $y_i > x$);
- (ii) each v_i ($1 \leq i \leq p$) is a factor which is either empty or has all its letters greater than (resp. smaller than or equal to) x .

We then let:

$$(11.18) \quad \gamma_x(w) := y_1v_1y_2v_2 \dots y_pv_p.$$

We see again that γ_x is a bijection of each class $R(\mathbf{m})$ onto itself, since it maps each *x-factorisation* $(v_1y_1, v_2y_2, \dots, v_py_p)$, obtained by cutting the word *after* every letter less than or equal to (resp. greater than) x onto the factorisation $(y_1v_1, y_2v_2, \dots, y_pv_p)$ obtained by cutting the word *before* every letter less than or equal to (resp. greater than) x .

The transformation Φ itself, as defined in (11.9), can also be described in the following algorithmic manner:

Algorithm for Φ . Let $w = x_1x_2 \dots x_m$;

1. Put $i := 1$, $w'_i := x_1$;
2. If $i = m$, let $\Phi(w) := w'_i$ and stop; else continue;
3. If the last letter of w'_i is less than or equal to (resp. greater than) x_{i+1} , cut w'_i after every letter less than or equal to (resp. greater than) x_{i+1} ;
4. In each compartment of w'_i determined by the previous cuttings, move the last letter in the compartment to the beginning of it; let v' be the word obtained after all those moves; put $w'_{i+1} := v'x_{i+1}$; replace i by $i + 1$ and go to 2.

For example, the image of $w = 4, 3, 5, 1, 1, 3, 4, 2, 3$ under Φ is obtained as follows:

$$\begin{aligned}
 w'_1 &= 4 \mid \\
 w'_2 &= 4 \mid 3 \mid \\
 w'_3 &= 4 \mid 3 \mid 5 \mid \\
 w'_4 &= 4351 \mid \\
 w'_5 &= 1 \mid 43 \mid 51 \mid \\
 w'_6 &= 1 \mid 3 \mid 4 \mid 1 \mid 53 \mid \\
 w'_7 &= 13 \mid 4 \mid 13 \mid 5 \mid 4 \mid \\
 w'_8 &= 3 \mid 1 \mid 43 \mid 1 \mid 542 \mid \\
 \Phi(w) = w'_9 &= 3, 1, 3, 4, 1, 2, 5, 4, 3.
 \end{aligned}$$

The descents in the word $w = 4, 3, 5, 1, 1, 3, 4, 2, 3$ are in position 1, 3 and 7, so that $\text{maj } w = 11$. But $\Phi(w) = 3, 1, 3, 4, 1, 2, 5, 4, 3$ has $\text{inv } \Phi(w) = 11$ inversions. Finally, w and $\Phi(w)$ end with the same letter, namely 3.

11.4. *Further properties of the transformation.* As we now see, the transformation Φ just defined preserves other statistics, first, the set-valued statistic *inverse ligne of route* and also the subword-valued statistics *right-to-left minimum letter subword* and *right-to-left maximum letter subword* we now define.

If a multiplicity \mathbf{m} reads $\mathbf{m} = (m_1, \dots, m_i, 0, \dots, 0, m_j, \dots, m_r)$ with $1 \leq i < j \leq r$ and $m_i, m_j \geq 1$, we say that j is the *successor* of i in \mathbf{m} and we write $j = \text{succ } i$. Of course, if all the components of \mathbf{m} are positive, the successor of each i ($1 \leq i \leq r - 1$) is $(i + 1)$. Let w be a word in the class $R(\mathbf{m})$.

Definition. The *inverse ligne of route* of w is defined to be the set, denoted by $\text{Iligne } w$, of all the letters i such that the rightmost occurrence of $\text{succ}(i)$ lies *to the left* of the rightmost occurrence of i . In an equivalent manner, the letter i is said to belong to the *inverse ligne of route* of w ,

presence of the “s” in our notation for “Rmals.” Finally, let $y := x_{i_1}$, then $\text{Rmil } w = y^{m_y} v$, where v has no letter equal to y .

Four other analogous definitions “Lmip,” “Lmap,” “Lmil” and “Lmals” can be introduced by considering the subwords *from left to right*, instead of right to left. We will do it later on for permutations only.

Remark. For a permutation w the numerical statistics $\# \text{Lmil } w$ and $\# \text{Lmals } w$ have been considered long ago by probabilists under the names of *lower records* and *upper records*.

Theorem 11.3. *Let Φ be the transformation constructed in subsection 11.3. Then, the following properties hold for every word w :*

- (a) $\text{maj } w = \text{inv } \Phi(w)$;
- (b) $\Phi(w)$ is a rearrangement of w and the restriction of Φ to each rearrangement class $R(\mathbf{m})$ is a bijection of $R(\mathbf{m})$ onto itself;
- (c) $\text{Iligne } w = \text{Iligne } \Phi(w)$;
- (d) $\text{Rmil } w = \text{Rmil } \Phi(w)$;
- (e) $\text{Rmals } w = \text{Rmals } \Phi(w)$.

The next corollary is an immediate consequence of Theorem 1.2.

Corollary 11.4. *Let A be a finite set of integers, u, v be two words and $R(\mathbf{m})$ be a rearrangement class. Then, the statistics “maj” and “inv” are equidistributed on the subclass*

$$\{w \in R(\mathbf{m}) : (\text{Iligne}, \text{Rmil}, \text{Rmals}) w = (A, u, v)\}.$$

In Fig. 11.2 the values of these statistics have been calculated for the words belonging to the class $R(2, 1, 1)$, the rearrangements of the word 1, 1, 2, 3. There are five tables corresponding to five subclasses characterized by a value (A, u, v) of the triple $(\text{Iligne}, \text{Rmil}, \text{Rmals})$. Notice that within each subclass the statistics “inv” and “maj” are equidistributed.

When Φ is restricted to \mathfrak{S}_n , the group of permutations of $1, 2, \dots, n$, i.e., the rearrangement class $R(1^n)$, several properties can also be derived. For each permutation w let

$$\text{imaj } w := \sum_i i \chi(i \in \text{Iligne } w).$$

It is immediate to verify

$$\text{imaj } w = \text{maj } w^{-1},$$

where w^{-1} denotes the inverse permutation of w . We then deduce the following corollary that is proved in section 5.

11. MAJOR INDEX AND INVERSION NUMBER

w	maj	inv	lligne	Rmil	Rmals
1, 2, 3, 1	3	2	1	1, 1	3, 1
2, 1, 3, 1	4	3	1	1, 1	3, 1
2, 3, 1, 1	2	4	1	1, 1	3, 1

w	maj	inv	lligne	Rmil	Rmals
1, 1, 3, 2	3	1	2	1, 1, 2	3, 2
1, 3, 1, 2	2	2	2	1, 1, 2	3, 2
3, 1, 1, 2	1	3	2	1, 1, 2	3, 2

w	maj	inv	lligne	Rmil	Rmals
1, 3, 2, 1	5	3	1, 2	1, 1	3, 2, 1
3, 1, 2, 1	4	4	1, 2	1, 1	3, 2, 1
3, 2, 1, 1	3	5	1, 2	1, 1	3, 2, 1

w	maj	inv	lligne	Rmil	Rmals
1, 1, 2, 3	0	0	\emptyset	1, 1, 2, 3	3

w	maj	inv	lligne	Rmil	Rmals
1, 2, 1, 3	2	1	1	1, 1, 3	3
2, 1, 1, 3	1	2	1	1, 1, 3	3

Fig. 11.2

Corollary 11.4. *The two statistics “inv” and “maj” are equally distributed on each set of permutations having a given ligne of route A , a given left-to-right maximum place subword C and a given right-to-left maximum place subword D . In other words, let*

$$S := \{w \in \mathfrak{S}_n : \text{Ligne } w = A, \text{Lmap } w = C, \text{Rmap } w = D\};$$

then

$$\sum_{w \in S} q^{\text{inv } w} = \sum_{w \in S} q^{\text{maj } w}.$$

The preceding corollary is an extension of the classical result (see, e.g. [Lo02, Theorem 11.4.4]), where no restriction is made neither on “Lmap,” nor on “Rmap”, but only on “Ligne.”

Proof of Theorem 11.3. Properties (a) and (b) have been proved with Theorem 11.1. For the remaining ones we proceed as follows:

Proof of Property (c).

Consider the mapping $wx \mapsto \Phi(wx)$ defined by (11.9), where wx belongs to the rearrangement class $R(\mathbf{m})$. Let z a letter of wx having a successor $z' := \text{succ}(z)$ in \mathbf{m} . Suppose $L(w) \leq x$ (resp. $L(w) > x$). There are four cases to be considered:

(i) $z < z' < x$; let \mathcal{P} be the property “the rightmost occurrence of z' is to the left of the rightmost occurrence of z .” Then, all the following properties are equivalent: (1) \mathcal{P} holds for wx ; (2) \mathcal{P} holds for w ; (3) \mathcal{P} holds for $\Phi(w)$ [by induction]; (4) \mathcal{P} holds for $y_1 y_2 \dots y_p$; (resp. (4) \mathcal{P} holds for $v_1 v_2 \dots v_p$;) (5) \mathcal{P} holds for $y_1 v_1 y_2 v_2 \dots y_p v_p = \gamma_x(\Phi(w))$; (6) \mathcal{P} holds for $\Phi(wx) = \gamma_x(\Phi(w))x$.

(ii) $x < z < z'$; same proof, only property (4) is to reverse: (4) \mathcal{P} holds for $v_1 v_2 \dots v_p$; (resp. (4) \mathcal{P} holds for $y_1 y_2 \dots y_p$;))

(iii) $x = z < z'$; then $x \in \text{Iigne } wx$ and $x \in \text{Iigne } \Phi(wx)$, since $L(\Phi(w)) = L(w) = x$.

(iv) $z < z' = x$; then $x \notin \text{Iigne } wx$ and $x \notin \text{Iigne } \Phi(wx)$. \square

Proof of Property (d).

Assume that (d) holds for a nonempty word w and let x be a letter. If all the letters of w are equal, the result is banal. Otherwise, let $\text{Rmil}(w) := x_{i_1}x_{i_2}\dots x_{i_a}$, so that, by induction $\text{Rmil } \Phi(w) = x_{i_1}x_{i_2}\dots x_{i_a}$. Notice that $L(w) = L(\Phi(w)) = x_m = x_{i_a}$ and the smallest letter of w is equal to x_{i_1} .

If all the letters of w are less than x , then $\text{Rmil}(wx) = x = \text{Rmil } \Phi(wx)$. If it is not the case, but still if $L(w) \leq x$, then

$$\begin{aligned} x_{i_1}x_{i_2}\dots x_{i_a}x &= \text{Rmil}(wx) \\ &= \text{Rmil}(\Phi(w)x) && \text{[by induction]} \\ &= \text{Rmil}(y_1y_2\dots y_px) && \text{[by definition of the } x\text{-factorization]} \\ &= \text{Rmil}(y_1v_1y_2v_2\dots y_pv_px) \\ &= \text{Rmil } \Phi(wx). && \text{[by definition of } \Phi \end{aligned}$$

If $L(w) > x$, there is a unique integer k such that $1 \leq k \leq a - 1$ and $x_{i_1} \leq \dots \leq x_{i_k} \leq x < x_{i_{k+1}} \leq \dots \leq x_{i_a}$. Then

$$\begin{aligned} x_{i_1}\dots x_{i_k}x &= \text{Rmil}(wx) \\ &= \text{Rmil}(\Phi(w)x) && \text{[by induction]} \\ &= \text{Rmil}(v_1v_2\dots v_px) && \text{[by definition of the } x\text{-factorization]} \\ &= \text{Rmil}(y_1v_1y_2v_2\dots y_pv_px) \\ &= \text{Rmil } \Phi(wx). && \square \text{ [by definition of } \Phi \end{aligned}$$

Proof of Property (e).

Again, the result is banal if all the letters of w are identical. Otherwise, let $\text{Rmals}(w) := x_{j_1}x_{j_2}\dots x_{j_b}$, so that $\text{Rmals } \Phi(w) = x_{j_1}x_{j_2}\dots x_{j_b}$. If all the letters of w are less than or equal to x , then $\text{Rmals } wx = x = \text{Rmals } \Phi(wx)$. If it is not the case, but still $L(w) \leq x$, there is a unique integer k such that $1 \leq k \leq b - 1$ and $x_{j_1} > \dots > x_{j_k} > x \geq x_{j_{k+1}} > \dots > x_{j_b}$. Then

$$\begin{aligned} x_{j_1}\dots x_{j_k}x &= \text{Rmals}(wx) \\ &= \text{Rmals}(\Phi(w)x) && \text{[by induction]} \\ &= \text{Rmals}(v_1v_2\dots v_px) && \text{[by definition of the } x\text{-factorization]} \\ &= \text{Rmals}(y_1v_1y_2v_2\dots y_pv_px) \\ &= \text{Rmals } \Phi(wx). && \text{[by definition of } \Phi \end{aligned}$$

If $L(w) > x$, then

$$\begin{aligned}
 x_{j_1} \dots x_{j_k} x &= \text{Rmals}(wx) \\
 &= \text{Rmals}(\Phi(w)x) && \text{[by induction]} \\
 &= \text{Rmals}(v_1 v_2 \dots v_p x) && \text{[by definition of the } x\text{-factorization]} \\
 &= \text{Rmals}(y_1 v_1 y_2 v_2 \dots y_p v_p x) \\
 &= \text{Rmals } \Phi(wx). \quad \square && \text{[by definition of } \Phi]
 \end{aligned}$$

This achieves the proof of Theorem 11.3.

11.5. *Application to permutations.* Consider the classes $R(\mathbf{m})$ of words without repeated letters (all the m_i 's are equal to 1), i.e., the *permutations*. The *inverse ligne of route* of a permutation $w = x_1 x_2 \dots x_r$ (of the word $12 \dots r$) is then the set of all the integers j such that $1 \leq j \leq r - 1$ and $(j + 1)$ is *to the left of* j within the word $x_1 x_2 \dots x_r$.

The *ligne of route* of a permutation w is the set, $\text{Ligne } w$, of the integers j such that $x_j > x_{j+1}$, so that

$$\text{des } w = |\text{Ligne } w| \quad \text{and} \quad \text{maj } w = \sum_j j \quad (j \in \text{Ligne } w).$$

It is readily seen that

$$(11.19) \quad \text{Iligne } w = \text{Ligne } w^{-1},$$

where w^{-1} designates the inverse of the permutation w . Also let

$$\text{ides } w := |\text{Iligne } w| \quad \text{and} \quad \text{imaj } w := \sum_j j \quad (j \in \text{Iligne } w).$$

The statistic ‘‘imaj’’ is called the *Inverse Major Index* of w . Also let $\mathbf{i}(w)$ denote the inverse w^{-1} of the permutation w . As $\text{inv } \mathbf{i}(w) = \text{inv } w$, it follows from the property (c) that the chain

$$w \xrightarrow{\mathbf{i}} w_1 \xrightarrow{\Phi^{-1}} w_2 \xrightarrow{\mathbf{i}} w_3 \xrightarrow{\Phi} w_4 \xrightarrow{\mathbf{i}} w_5$$

has the properties:

$$\begin{aligned}
 \text{Ligne } w &= \text{Iligne } w_1 = \text{Iligne } w_2 = \text{Ligne } w_3; \\
 \text{des } w &= \text{ides } w_1 = \text{ides } w_2 = \text{des } w_3; \\
 \text{maj } w &= \text{imaj } w_1 = \text{imaj } w_2 = \text{maj } w_3 = \text{inv } w_4 = \text{inv } w_5; \\
 \text{inv } w &= \text{inv } w_1 = \text{maj } w_2 = \text{imaj } w_3 = \text{imaj } w_4 = \text{maj } w_5.
 \end{aligned}$$

This result has several consequences that are now stated.

Corollary 11.5. *The six pairs (maj, inv), (imaj, inv), (imaj, maj), (maj, imaj), (inv, imaj), (inv, maj) are equally distributed on each symmetric group \mathfrak{S}_n , i.e., with w running over \mathfrak{S}_n we have:*

$$(11.20) \quad \sum_w q_1^{\text{maj } w} q_2^{\text{inv } w} = \sum_w q_1^{\text{maj } w} q_2^{\text{inv } w} = \sum_w q_1^{\text{imaj } w} q_2^{\text{maj } w} \\ = \sum_w q_1^{\text{maj } w} q_2^{\text{imaj } w} = \sum_w q_1^{\text{inv } w} q_2^{\text{imaj } w} = \sum_w q_1^{\text{inv } w} q_2^{\text{maj } w}.$$

Corollary 11.6. *With w running over \mathfrak{S}_n we have:*

$$(11.21) \quad \text{inv}A_n(t, q) = \sum_w t^{\text{des } w} q^{\text{inv } w} = \sum_w t^{\text{des } w} q^{\text{imaj } w}.$$

Proof. Consider the bijection $w \mapsto w_3$. \square

Corollary 11.7. *The two statistics “inv” and “imaj” have the same distribution on each set of permutations having a given ligne of route. In other words, for every subset $A \subset [n - 1]$ the following identity holds:*

$$\sum_w q^{\text{inv } w} = \sum_w q^{\text{imaj } w} \quad (w \in \mathfrak{S}_n, \text{Ligne } w = A).$$

Proof. Again consider the bijection $w \mapsto w_3$. \square

We make a further use of the transformations, \mathbf{i} , \mathbf{c} , \mathbf{r} of the dihedral group. Recall that \mathbf{c} is the *complement to $(n + 1)$* and \mathbf{r} the *reverse image* that map the permutation w , written as a linear word $w = x_1 \dots x_n$, onto

$$\mathbf{c} w := (n + 1 - x_1)(n + 1 - x_2) \dots (n + 1 - x_n); \\ \mathbf{r} w := x_n \dots x_2 x_1.$$

As $\text{inv } \mathbf{i} w = \text{inv } w$ and also $\text{Rmap } \mathbf{i} w = \text{Rmals } w$, $\text{Lmap } \mathbf{i} w = \text{Rmil } w$ (easy to verify), it follows from Theorem 11.3 that the chain

$$w \xrightarrow{\mathbf{i}} w_1 \xrightarrow{\Phi} w_2 \xrightarrow{\mathbf{i}} w_3$$

has the properties:

$$\begin{aligned} \text{Ligne } w &= \text{Iligne } w_1 = \text{Iligne } w_2 = \text{Ligne } w_3; \\ \text{Lmap } w &= \text{Rmil } w_1 = \text{Rmil } w_2 = \text{Lmap } w_3; \\ \text{Rmap } w &= \text{Rmals } w_1 = \text{Rmals } w_2 = \text{Rmap } w_3; \\ \text{imaj } w &= \text{maj } w_1 = \text{inv } w_2 = \text{inv } w_3. \end{aligned}$$

Hence, for each triple (A, B, C) the bijection $w \mapsto w_3$ maps each set of permutations w such that $(\text{Ligne}, \text{Lmap}, \text{Rmap}) w = (A, B, C)$ onto itself with the property that $\text{imaj } w = \text{inv } w_3$. This proves the following corollary.

Corollary 11.8. *The two statistics “inv” and “imaj” are equally distributed on each set of permutations having a given ligne of route A , a given left-to-right maximum place subword C and a given right-to-left maximum place subword D . In other words, let*

$$S := \{w \in \mathfrak{S}_n : \text{Ligne } w = A, \text{Rmap } w = C, \text{Lmap } w = D\};$$

then

$$\sum_{w \in S} q^{\text{inv } w} = \sum_{w \in S} q^{\text{imaj } w}.$$

There are other consequences we can deduce from Theorem 11.3 by taking the composition product of Φ with other operations of the dihedral group. First, we can verify that $\text{inv } \mathbf{rc} w = \text{inv } w$ and $\text{Ligne } \mathbf{rc} w = n - \text{Ligne } w := \{n - i : i \in \text{Ligne } w\}$, so that

$$\text{maj } \mathbf{rc} w = \sum_i (n - i) \chi(x_i > x_{i+1}),$$

a statistic that will be denoted by $\text{comaj } w$. Let $\text{Lmil } w$ (resp. $\text{Lmals } w$) denote the *left-to-right* minimum (resp. maximum) letter subword of the permutation w . Again, it is easy to verify that

$$\text{Lmil } \mathbf{rc} w = n + 1 - \text{Rmals } w, \quad \text{Lmals } \mathbf{rc} w = n + 1 - \text{Rmil } w.$$

Consider the sequence:

$$w \xrightarrow{\mathbf{rc}} w_1 \xrightarrow{\Phi} w_2 \xrightarrow{\mathbf{rc}} w_3.$$

Then

$$\begin{aligned} \text{Iligne } w &= n - \text{Iligne } w_1 = n - \text{Iligne } w_2 = \text{Iligne } w_3; \\ \text{Lmil } w &= n + 1 - \text{Rmals } w_1 = n + 1 - \text{Rmals } w_2 = \text{Lmil } w_3; \\ \text{Lmals } w &= n + 1 - \text{Rmil } w_1 = n + 1 - \text{Rmil } w_2 = \text{Lmals } w_3; \\ \text{comaj } w &= \text{maj } w_1 = \text{inv } w_2 = \text{inv } w_3. \end{aligned}$$

This implies the following corollary.

Corollary 11.9. *The two statistics “inv” and “imaj” are equally distributed on each set of permutations having a given inverse ligne of route A , a given left-to-right minimum letter subword C and a given left-to-right maximum letter subword D . In other words, let*

$$S := \{w \in \mathfrak{S}_n : \text{Iligne } w = A, \text{Lmil } w = C, \text{Lmals } w = D\}.$$

then

$$\sum_{w \in S} q^{\text{inv } w} = \sum_{w \in S} q^{\text{comaj } w};$$

that can also be expressed as:

$$\sum_{w \in S'} q^{\text{inv } w} = \sum_{w \in S'} q^{\text{comaj } w},$$

where $S' := \{w \in \mathfrak{S}_n : \text{Ligne } w^{-1} = A, \text{Lmip } w^{-1} = C, \text{Lmap } w^{-1} = D\}$.

12. Major Index and Inverse Major Index

For each $n \geq 0$ let $A_n(q_1, q_2)$ be the generating polynomial for \mathfrak{S}_n by the pair (maj, imaj) :

$$(12.1) \quad A_n(q_1, q_2) := \sum_{\sigma \in \mathfrak{S}_n} q_1^{\text{maj } \sigma} q_2^{\text{imaj } \sigma}.$$

Corollary 11.3 shows that there are five other ways of expressing such a polynomial. By analogy with the one-basis q -ascending factorials, introduce the following notations:

$$(u; q_1, q_2)_{r,s} := \begin{cases} 1, & \text{if } r \text{ or } s \text{ is zero;} \\ \prod_{0 \leq i \leq r-1} \prod_{0 \leq j \leq s-1} (1 - uq_1^i q_2^j), & \text{if } r, s \geq 1, \end{cases}$$

$$(12.2) \quad (u; q_1, q_2)_{\infty, \infty} := \lim_{r,s} (u; q_1, q_2)_{r,s} = \prod_{i \geq 0} \prod_{j \geq 0} (1 - uq_1^i q_2^j).$$

The purpose of this section is to prove the following theorem.

Theorem 12.1. *The bibasic generating function for the polynomials $A_n(q_1, q_2)$ is given by:*

$$(12.3) \quad \sum_{n \geq 0} A_n(q_1, q_2) \frac{u^n}{(q_1; q_1)_n (q_2; q_2)_n} = \frac{1}{(u; q_1, q_2)_{\infty, \infty}}.$$

The term ‘‘bibasic’’ refers to the normalization $(q_1; q_1)_n (q_2; q_2)_n$ of the denominator as a product of two q -ascending factorials. *A priori*, there was no evidence that such a normalization was to be introduced. In fact, the infinite product on the right-hand side preexisted in the literature, ready to be unearthed for combinatorial purposes. It could also be regarded as a specialization of the celebrated Cauchy infinite product

$$\prod_{i \geq 1, j \geq 1} \frac{1}{1 - ux_i y_j}$$

with the substitutions $x_i \leftarrow q_1^{i-1}$, $y_j \leftarrow q_j^{j-1}$, for which several expansions are known, in particular in terms of Schur functions. That method will be exploited more in detail in subsequent chapters.

In the proof of Theorem 12.1 the starting point is the infinite product $1/(u; q_1, q_2)_{\infty, \infty}$ that is first expanded as an infinite series and shown to be the generating function for pairs of finite sequences, called *biwords*. This is the “manipulatorics” part of the proof. The next step, the “combinatorics” part, consists of mapping each such a biword onto a triple (σ, b', c') , where σ is a permutation and where (b', c') is another biword that is precisely counted by the product $1/(q_1; q_1)_n (q_2; q_2)_n$. The construction of that mapping may be regarded as another application of the MacMahon *Verfahren*. Accordingly, let us decompose the proof into those two parts.

12.1. *The biword expansion.* On the right-hand side of (12.3) each fraction $1/(1 - uq_1^i q_2^j)$ expands into a geometric series $\sum_{a_{ij} \geq 0} (uq_1^i q_2^j)^{a_{ij}}$, whose first nonconstant term is the monomial $uq_1^i q_2^j$. Hence, the coefficient of the monomial $u^\alpha q_1^\beta q_2^\gamma$ in the expansion of the infinite product $1/(u; q_1, q_2)_{\infty, \infty}$ is equal to the coefficient of the same monomial in the *finite* product $\prod_{i \leq \alpha, j \leq \beta} 1/(1 - uq_1^i q_2^j)$. We can then write $1/(u; q_1, q_2)_{\infty, \infty}$ as the series

$$(12.4) \quad \sum_A \prod_{i,j} (uq_1^i q_2^j)^{a_{ij}} = \sum_A u^{\sum a_{ij}} q_1^{\sum i a_{ij}} q_2^{\sum j a_{ij}},$$

where A runs over the set of all matrices of the form $A = (a_{ij})$ ($i \geq 0, j \geq 0$), whose entries a_{ij} are integers which are all zero except finitely many of them. Beside the null matrix we can then express each such matrix as a bounded matrix having at least one nonzero entry on its rightmost column and its lowest row.

For example,

$$A = \begin{matrix} & \begin{matrix} 0 & 1 & 2 & 3 & 4 \end{matrix} \\ \begin{matrix} 0 \\ 1 \\ 2 \\ 3 \end{matrix} & \begin{pmatrix} 0 & 0 & 1 & 0 & 2 \\ 1 & 0 & 0 & 3 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 2 & 0 & 2 \end{pmatrix} \end{matrix}$$

is such a matrix.

Now with each matrix A we associate a two-row matrix or *biword* $\mathbf{b} = \begin{pmatrix} b \\ c \end{pmatrix} = \begin{pmatrix} b_1 \cdots b_n \\ c_1 \cdots c_n \end{pmatrix}$, with integral entries such that

$$(12.5) \quad \sum_{i,j} a_{ij} = n, \quad \sum_{i,j} i a_{ij} = b_1 + \cdots + b_n, \quad \sum_{i,j} j a_{ij} = c_1 + \cdots + c_n;$$

$$(12.6) \quad (b_1, c_1) \leq (b_2, c_2) \leq \cdots \leq (b_n, c_n)$$

with respect to the lexicographic order. We then say that the biword \mathbf{b} is *nondecreasing*.

The sum in (12.4) is then replaced by:

$$(12.7) \quad \frac{1}{(u; q_1, q_2)_{\infty, \infty}} = \sum_{n \geq 0} u^n \sum_{\mathbf{b}} q_1^{\text{tot } b} q_2^{\text{tot } c},$$

where \mathbf{b} runs over all nondecreasing biwords (whose biletters are pairs of nonnegative integers) of length n . To fulfill the relations (12.5) start with matrix A , read its rows from left to right and top to bottom and, for each *positive* entry a_{ij} write down a_{ij} biletters $\begin{pmatrix} i \\ j \end{pmatrix}$ one after another. The number of biletters written in this way is then equal to $\sum_{i,j} a_{ij}$. Moreover, on the top (resp. bottom) row of \mathbf{b} each number i (resp. j) is repeated a_{ij} times. Consequently, the last two conditions of (12.5) hold.

Finally, as the biletters $\begin{pmatrix} b_i \\ c_i \end{pmatrix}$ were written starting with the first row from left to right, then the second row, \dots those biletters are in *increasing lexicographic order* when the biword \mathbf{b} is read from left to right, so that the relations (12.6) hold. Conversely, when starting with a nondecreasing biword, we can reconstruct the matrix A in a unique manner.

For example, to the matrix A above there corresponds the nondecreasing biword

$$\mathbf{b} = \begin{pmatrix} b \\ c \end{pmatrix} = \begin{pmatrix} 0 & 0 & 0 & 1 & 1 & 1 & 1 & 3 & 3 & 3 & 3 & 3 \\ 2 & 4 & 4 & 0 & 3 & 3 & 3 & 1 & 2 & 2 & 4 & 4 \end{pmatrix},$$

a biword of length $n = 12$ and such that $\text{tot } b = 19$, $\text{tot } c = 32$.

12.2. *Another application of the MacMahon Verfahren.* For convenience define the *Comajor Index* of a permutation $\sigma \in \mathfrak{S}_n$ by

$$(12.8) \quad \text{comaj } \sigma := \sum_{1 \leq i \leq n-1} (n-i) \chi(\sigma(i) > \sigma(i+1)).$$

We now construct a bijection $\mathbf{b} \mapsto (\sigma, b', c')$ that maps each nondecreasing biword $\mathbf{b} = \begin{pmatrix} b \\ c \end{pmatrix}$ of length n onto a triple (σ, b', c') , where $\sigma \in \mathfrak{S}_n$ and $b', c' \in \text{NDS}(n)$, that is, σ is a permutation of order n and b', c' are both nondecreasing words of length n whose letters are nonnegative integers. Moreover, the bijection has the properties:

$$(12.9) \quad \text{tot } b = \text{comaj } \sigma + \text{tot } b', \quad \text{tot } c = \text{comaj } \sigma^{-1} + \text{tot } c'.$$

To do so start with a nondecreasing biword $\mathbf{b} = \begin{pmatrix} b \\ c \end{pmatrix}$. Since the biletters $\begin{pmatrix} b_i \\ c_i \end{pmatrix}$ are written in nondecreasing order with respect to the lexicographic order, the following property holds:

$$(12.10) \quad c_i > c_{i+1} \Rightarrow b_i < b_{i+1}.$$

For each $i = 1, 2, \dots, n$ let

$$(12.11) \quad \sigma(i) = |\{j : 1 \leq j \leq n, c_j < c_i\}| + |\{j : 1 \leq j \leq i, c_j = c_i\}|.$$

In other words, $\sigma(i)$ is equal to the number of letters in c which are less than c_i , plus the number of letters equal to c_i , but lie to the left of c_i , with c_i included. This defines a permutation σ of order n .

By construction $c_i > c_{i+1}$ if and only if $\sigma(i) > \sigma(i+1)$. For $1 \leq i \leq n$ let y_i be the number of integers j such that $1 \leq j \leq i-1$ and $\sigma(j) > \sigma(j+1)$, which is also the number of integers j such that $1 \leq j \leq i-1$ and $c_j > c_{j+1}$. The word $y = y_1 \dots y_n$ is nondecreasing. Moreover,

$$(12.12) \quad \text{comaj } \sigma = \text{tot } y = y_1 + \dots + y_n$$

and (12.10) implies

$$(12.13) \quad y_i < y_{i+1} \Leftrightarrow \sigma(i) > \sigma(i+1) \Leftrightarrow c_i > c_{i+1} \Rightarrow b_i < b_{i+1}.$$

As $y_1 = 0$, the relations (12.13) imply that

$$(12.14) \quad y_i \leq b_i \quad (1 \leq i \leq n).$$

We then define a *nondecreasing* word $b' = b'_1 \dots b'_n$ by

$$(12.15) \quad b'_i := b_i - y_i \quad (1 \leq i \leq n).$$

Finally, because of (12.12) the first of the relations (12.9) holds.

Again use the previous example. Under the nondecreasing word $\begin{pmatrix} b \\ c \end{pmatrix}$ we have written the values of σ , of y and of $b' = b - y$.

$$\begin{array}{r} b = 0 \ 0 \ 0 \ 1 \ 1 \ 1 \ 1 \ 3 \ 3 \ 3 \ 3 \ 3 \\ c = 2 \ 4 \ 4 \ 0 \ 3 \ 3 \ 3 \ 1 \ 2 \ 2 \ 4 \ 4 \\ \sigma = 3 \ 9 \ 10 \ 1 \ 6 \ 7 \ 8 \ 2 \ 4 \ 5 \ 11 \ 12 \\ y = 0 \ 0 \ 0 \ 1 \ 1 \ 1 \ 1 \ 2 \ 2 \ 2 \ 2 \ 2 \\ b' = 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 1 \ 1 \ 1 \ 1 \ 1 \end{array}$$

We can also start with the biword $\begin{pmatrix} c \\ b \end{pmatrix}$ (and not $\mathbf{b} = \begin{pmatrix} b \\ c \end{pmatrix}$) and rearrange its biletters $\begin{pmatrix} c_i \\ b_i \end{pmatrix}$ in increasing order. We obtain a biword $\tilde{\mathbf{b}}$ whose top row is the nondecreasing rearrangement of the word c .

With the running example we get:

$$\tilde{\mathbf{b}} = \begin{pmatrix} 0 \ 1 \ 2 \ 2 \ 2 \ 3 \ 3 \ 3 \ 4 \ 4 \ 4 \ 4 \\ 1 \ 3 \ 0 \ 3 \ 3 \ 1 \ 1 \ 1 \ 0 \ 0 \ 3 \ 3 \end{pmatrix}.$$

Let us go back to the general case. By definition of σ given in (12.11), there are exactly $\sigma(i)$ biletters $\begin{pmatrix} b_j \\ c_j \end{pmatrix}$ such that $1 \leq j \leq n$ and $c_j < c_i$ or such that $1 \leq j \leq i$ and $c_j = c_i$. As in the biword $\tilde{\mathbf{b}}$ we have sorted the biletter $\begin{pmatrix} c_j \\ b_j \end{pmatrix}$ in increasing order, the biletter $\begin{pmatrix} c_i \\ b_i \end{pmatrix}$ will be found in $\tilde{\mathbf{b}}$ in the $\sigma(i)$ -th position. Hence $\tilde{\mathbf{b}} = \begin{pmatrix} c_{\sigma^{-1}(1)} \cdots c_{\sigma^{-1}(n)} \\ b_{\sigma^{-1}(1)} \cdots b_{\sigma^{-1}(n)} \end{pmatrix}$. Let τ be the permutation defined as in (12.11) when the counting is applied to the nondecreasing biword $\tilde{\mathbf{b}}$, i.e.,

$$\begin{aligned} \tau(i) &= |\{j : 1 \leq j \leq n, b_{\sigma^{-1}(j)} < b_{\sigma^{-1}(i)}\}| \\ &\quad + |\{j : 1 \leq j \leq i, b_{\sigma^{-1}(j)} = b_{\sigma^{-1}(i)}\}|. \end{aligned}$$

We also have:

$$\tau\sigma(i) = |\{j : 1 \leq j \leq n, b_j < b_i\}| + |\{j : 1 \leq j \leq \sigma(i), b_{\sigma^{-1}(j)} = b_i\}|.$$

But by definition of σ , if $b_k = b_l$, we have $\sigma(k) < \sigma(l) \Leftrightarrow k < l$. Therefore, if $b_{\sigma^{-1}(j)} = b_i$, we have $j < \sigma(i) \Leftrightarrow \sigma^{-1}(j) < i$. Since b is nondecreasing, we obtain:

$$\begin{aligned} \tau\sigma(i) &= |\{j : 1 \leq j \leq n, b_j < b_i\}| + |\{j : 1 \leq \sigma^{-1}(j) \leq i, b_{\sigma^{-1}(j)} = b_i\}| \\ &= |\{j : 1 \leq j \leq i, b_j < b_i\}| + |\{j : 1 \leq j \leq i, b_j = b_i\}| \\ &= i. \end{aligned}$$

Therefore, $\tau = \sigma^{-1}$.

Under the same procedure, to the nondecreasing biword $\tilde{\mathbf{b}}$ there corresponds a pair (τ, c') , in a one-to-one manner, where $\tau = \sigma^{-1}$ and where c' is a nondecreasing word. Moreover, the second relation of (12.9) holds.

Keeping the same example we determine σ^{-1} , y and c' , where this time the word y serves to calculate $\text{comaj } \sigma^{-1}$:

$$\begin{aligned} \tilde{c} &= 0 \ 1 \ 2 \ 2 \ 2 \ 3 \ 3 \ 3 \ 4 \ 4 \ 4 \ 4 \\ \tilde{b} &= 1 \ 3 \ 0 \ 3 \ 3 \ 1 \ 1 \ 1 \ 0 \ 0 \ 3 \ 3 \\ \sigma^{-1} &= 4 \ 8 \ 1 \ 9 \ 10 \ 5 \ 6 \ 7 \ 2 \ 3 \ 11 \ 12 \\ y &= 0 \ 0 \ 1 \ 1 \ 1 \ 2 \ 2 \ 2 \ 3 \ 3 \ 3 \ 3 \\ c' &= 0 \ 0 \ 1 \ 1 \ 1 \ 1 \ 1 \ 1 \ 1 \ 1 \ 1 \ 1 \end{aligned}$$

Now by using the bijection $\mathbf{b} \mapsto (\sigma, b', c')$ just obtained we can derive:

$$\sum_{n \geq 0} u^n \sum_{\mathbf{b}} q_1^{\text{tot } b} q_2^{\text{tot } c} = \sum_{n \geq 0} u^n \sum_{\substack{\sigma \in \mathfrak{S}_n \\ b', c' \in \text{NDS}(n)}} q_1^{\text{comaj } \sigma + \text{tot } b'} q_2^{\text{comaj } \sigma^{-1} + \text{tot } c'};$$

and then by (6.1):

13. A FOUR-VARIABLE DISTRIBUTION

$$\sum_{n \geq 0} u^n \sum_{\mathbf{b}} q_1^{\text{tot } b} q_2^{\text{tot } c} = \sum_{n \geq 0} u^n \frac{1}{(q_1; q_1)_n (q_2; q_2)_n} \sum_{\sigma \in \mathfrak{S}_n} q_1^{\text{comaj } \sigma} q_2^{\text{comaj } \sigma^{-1}}.$$

There remains to show that the polynomial $A_n(q_1, q_2)$, defined in (12.1), is also equal to $\sum_{\sigma \in \mathfrak{S}_n} q_1^{\text{comaj } \sigma} q_2^{\text{comaj } \sigma^{-1}}$. This can be proved by means of the bijection \mathbf{rc} that maps the permutation σ onto the permutation $\mathbf{rc} \sigma$ defined by

$$\mathbf{rc} \sigma(i) := n + 1 - \sigma(n + 1 - i) \quad (1 \leq i \leq n).$$

It is easily seen that

$$\text{comaj } \mathbf{rc} \sigma = \text{maj } \sigma.$$

This completes the proof of Theorem 12.1.

13. A four-variable distribution

In section 7 we have introduced the Euler-Mahonian polynomial $A_{\mathbf{m}}(t, q)$ as a *t-extension* of the polynomial $A_{\mathbf{m}}(q)$ by noticing that the combinatorial correspondence used in the calculation of the generating function for the $A_{\mathbf{m}}(q)$'s had a further property. We will do the same for the bijection constructed in the previous section and derive what could be called a *t₁, t₂-extension* of formula (12.3).

Consider the inverse bijection $(\sigma, b', c') \mapsto \mathbf{b}$ described in 12.2. The nondecreasing word y defined just before (12.12), such that $\text{tot } y = \text{comaj } \sigma$, has the further property

$$(13.1) \quad y_n = \text{des } \sigma.$$

Now consider the *finite* product

$$\frac{1}{(u; q_1, q_2)_{r+1, s+1}} = \prod_{0 \leq i \leq r} \prod_{0 \leq j \leq s} \frac{1}{1 - u q_1^i q_2^j}.$$

It can be expanded into the series

$$\sum_A \prod_{i, j} (u q_1^i q_2^j)^{a_{ij}} = \sum_A u^{\sum a_{ij}} q_1^{\sum i a_{ij}} q_2^{\sum j a_{ij}},$$

but this time the matrices A are $(r+1) \times (s+1)$ -matrices. The nondecreasing biword $\mathbf{b} = \begin{pmatrix} b \\ c \end{pmatrix} = \begin{pmatrix} b_1 & \dots & b_n \\ c_1 & \dots & c_n \end{pmatrix}$ that corresponds to such a matrix has the further property:

$$\max b_i = b_n \leq r \quad \text{and} \quad \max c_i \leq s.$$

Properties (12.15) and (13.1) imply: $b'_n = b_n - y_n \leq r - \text{des } \sigma$, with an analogous property for the c'_i 's, that is $c'_n \leq s - \text{des } \sigma^{-1}$.

Start with a quintuple (σ, r', s', b', c') , where

$$(13.2) \quad \sigma \in \mathfrak{S}_n, \quad r', s' \geq 0, \quad b' \in \text{NDS}(n, r'), \quad c' \in \text{NDS}(n, s').$$

Under the inverse bijection $(\sigma, b', c') \mapsto \mathbf{b}$ the quintuple is mapped onto a triple (r, s, \mathbf{b}) such that $r = r' + \text{des } \sigma$, $s = s' + \text{des } \sigma^{-1}$, $\mathbf{b} = \begin{pmatrix} b \\ c \end{pmatrix}$ with $b \in \text{NDS}(n, r)$ and \tilde{c} (the nondecreasing rearrangement of c) in $\text{NDS}(n, s)$. We can then write

$$\begin{aligned} \sum_{r, s \geq 0} \frac{t_1^r t_2^s}{(u; q_1, q_2)_{r+1, s+1}} &= \sum_{r, s \geq 0} t_1^r t_2^s \sum_{n \geq 0} u^n \sum_{\substack{b \in \text{NDS}(n, r), \\ c \in \text{NDS}(n, s)}} q_1^{\text{tot } b} q_2^{\text{tot } c} \\ &= \sum_{r, s \geq 0} \sum_{n \geq 0} u^n \sum_{(\sigma, r', s', b', c')} t_1^{r'+\text{des } \sigma} t_2^{s'+\text{des } \sigma^{-1}} q_1^{\text{comaj } \sigma + \text{tot } b'} q_2^{\text{comaj } \sigma^{-1} + \text{tot } c'}, \end{aligned}$$

where the relations (13.2) hold and also $r = r' + \text{des } \sigma$, $s = s' + \text{des } \sigma^{-1}$. Let

$$(13.3) \quad A_n(t_1, t_2, q_1, q_2) := \sum_{\sigma \in \mathfrak{S}_n} t_1^{\text{des } \sigma} t_2^{\text{des } \sigma^{-1}} q_1^{\text{comaj } \sigma} q_2^{\text{comaj } \sigma^{-1}}.$$

It follows that

$$\begin{aligned} \sum_{r, s \geq 0} \frac{t_1^r t_2^s}{(u; q_1, q_2)_{r+1, s+1}} &= \sum_{n \geq 0} u^n A_n(t_1, t_2, q_1, q_2) \sum_{\substack{r' \geq 0 \\ b' \in \text{NDS}(n, r')}} t_1^{r'} q_1^{\text{tot } b'} \sum_{\substack{s' \geq 0 \\ c' \in \text{NDS}(n, s')}} t_2^{s'} q_2^{\text{tot } c'} \\ (13.4) \quad &= \sum_{n \geq 0} u^n A_n(t_1, t_2, q_1, q_2) \frac{1}{(t_1; q_1)_{n+1} (t_2; q_2)_{n+1}}, \end{aligned}$$

by (3.9) and (4.5).

Formula (13.4) provides an expression for the generating function for the polynomials $A_n(t_1, t_2, q_1, q_2)$. Observe the nature of the denominators. They are products of a q_1 -ascending factorial by q_2 -ascending factorial. There remains to verify that

$$(13.5) \quad A_n(t_1, t_2, q_1, q_2) = \sum_{\sigma \in \mathfrak{S}_n} t_1^{\text{des } \sigma} t_2^{\text{ides } \sigma} q_1^{\text{maj } \sigma} q_2^{\text{imaj } \sigma},$$

where, by analogy with “imaj”, the symbol “ides” means

$$(13.6) \quad \text{ides } \sigma := \text{des } \sigma^{-1}.$$

14. SYMMETRIC FUNCTIONS

Again this is proved by means of the bijection \mathbf{rc} that maps each permutation σ onto the permutation $\mathbf{rc} \sigma$ defined by

$$\mathbf{rc} \sigma(i) := n + 1 - \sigma(n + 1 - i) \quad (1 \leq i \leq n).$$

Clearly, we have

$$\text{comaj } \mathbf{rc} \sigma = \text{maj } \sigma, \quad \text{des } \mathbf{rc} \sigma = \text{des } \sigma.$$

We have then proved the following result.

Theorem 13.1. *The bibasic generating function for the polynomials $A_n(t_1, t_2, q_1, q_2)$, as defined in (13.5), is given by*

$$(13.7) \quad \sum_{n \geq 0} A_n(t_1, t_2, q_1, q_2) \frac{u^n}{(t_1; q_1)_{n+1} (t_2; q_2)_{n+1}} = \sum_{r, s \geq 0} \frac{t_1^r t_2^s}{(u; q_1, q_2)_{r+1, s+1}}.$$

Specializations. The right-hand side of (13.7) is symmetric in the pairs (t_1, q_1) , (t_2, q_2) , so that the polynomial $A_n(t_1, q_1, t_2, q_2)$ is also symmetric in (t_1, q_1) , (t_2, q_2) . This can also be seen by using the bijection $\sigma \mapsto \sigma^{-1}$ of \mathfrak{S}_n on itself. In particular, both specializations $A_n(t, 1, q, 1)$ and $A_n(1, t, 1, q)$ are equal. Moreover,

$$(13.8) \quad A_n(t, 1, q, 1) = A_n(1, t, 1, q) = \text{maj}A(t, q),$$

where $\text{maj}A(t, q)$ is the q -maj-Eulerian polynomial defined in section 10. On the other hand, it follows from Corollary 11.4 that

$$(13.9) \quad t A_n(t, 1, 1, q) = t A_n(1, t, 1, q) = \text{inv}A(t, q),$$

where $\text{inv}A(t, q)$ is the q -inv-Eulerian polynomial, also defined in section 10.

Tables of the polynomials $A_n(t_1, q_1, t_2, q_2)$ for $n = 3, 4, 5$ are shown in Fig. 13.1. Keeping in mind (13.8), the following notations have been used: $\text{maj}A(t, q) = \sum_k A_{n,k}(q)$ and $A_{n,k} := A_{n,k}(1)$ (the Eulerian coefficient). Notice the numerous symmetries within the tables.

	ides \rightarrow	0	1	2	$n = 3$	
	imaj \rightarrow	0	1 2	3	$A_{3,k}(q)$	$A_{3,k}$
des	maj					
\downarrow	\downarrow					
0	0	1			1	1
1	1		1 1		2	
	2		1 1		2	4
2	3			1	1	1

	ides \rightarrow	0	1	2	3	$n = 4$	
	imaj \rightarrow	0	1 2 3	3 4 5	6	$A_{4,k}(q)$	$A_{4,k}$
des	maj						
\downarrow	\downarrow						
0	0	1				1	1
1	1		1 1 1			3	
	2		1 2 1	1		5	11
	3		1 1 1			3	
2	3			1 1 1		3	
	4		1	1 2 1		5	11
	5			1 1 1		3	
3	6				1	1	1

	ides \rightarrow	0	1	2	3	4	$n = 5$	
	imaj \rightarrow	0	1 2 3 4	3 4 5 6 7	6 7 8 9	10	$A_{5,k}(q)$	$A_{5,k}$
des	maj							
\downarrow	\downarrow							
0	0	1					1	1
1	1		1 1 1 1				4	26
	2		1 2 2 1	1 1 1			9	
	3		1 2 2 1	1 1 1			9	
	4		1 1 1 1				4	
2	3			1 1 2 1 1			6	66
	4		1 1	1 3 4 3 1	1 1		16	
	5		1 1	2 4 6 4 2	1 1		22	
	6		1 1	1 3 4 3 1	1 1		16	
	7			1 1 2 1 1			6	
3	6				1 1 1 1		4	26
	7			1 1 1	1 2 2 1		9	
	8			1 1 1	1 2 2 1		9	
	9				1 1 1 1		4	
4	10					1	1	1

Fig. 13.1

14. Symmetric Functions

The Cauchy identity for Schur functions will be an essential tool for deriving several combinatorial formulas for symmetric group statistics. It matters to have a brief account for the algebra of symmetric function and a complete description of the combinatorial properties of the Schur functions. This is the content of the next three sections.

14. SYMMETRIC FUNCTIONS

14.1. *Partitions of integers.* Those structures remain the privileged objects in the theory of symmetric functions. As already discussed in section 4, by *partition* of an integer $n \geq 1$ it is meant a nondecreasing sequence $\lambda = (\lambda_1, \lambda_2, \dots)$ of nonnegative integers, where finitely many of them are nonzero. The nonzero elements of λ are the *parts* of the partition. The *number of parts* is denoted by $l(\lambda)$ and the *weight*, denoted by $|\lambda|$, of the partition λ is defined by

$$|\lambda| = \lambda_1 + \lambda_2 + \dots$$

When $|\lambda| = n$, we say that λ is a *partition* of the integer n . The symbol \mathcal{P}_n will designate the set of partitions of n .

The *multiplicative notation* of the partition λ is reads as $\lambda = 1^{m_1} 2^{m_2} \dots$, where for each $i = 1, 2, \dots$ the exponent m_i is equal to the number of parts of λ equal to i . The integer $m_i = m_i(\lambda)$ is called the *multiplicity* of i in λ .

For example, $\lambda = (5, 4, 4, 2, 1, 1)$ is a partition of $n = 17$, whose multiplicative notation reads $1^2 2^1 3^0 4^2 5^1$.

The *shape* of a partition $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_r)$ ($\lambda_r \geq 1$) is the set of all the $|\lambda|$ points $(1, 1), (1, 2), \dots, (1, \lambda_1), (2, 1), (2, 2), \dots, (2, \lambda_2), \dots, (r, 1), (r, 2), \dots, (r, \lambda_r)$ located in the north-eastern quadrant \mathbb{N}^2 of \mathbb{Z}^2 . Each shape if also represented by a set of squared boxes left justified, where every point of the previous sequence is the center of a box. For example, the partition $\lambda = (5, 4, 4, 2, 1, 1)$ is represented by the shape drawn in Fig. 14.1




Fig. 14.1

This geometric representation is also called *Ferrers diagram* and denoted by the same symbol λ .

The *conjugate* partition of $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_r)$ is the partition $\lambda' = 1^{\lambda_1 - \lambda_2} 2^{\lambda_2 - \lambda_3} \dots$ (written in multiplicative notation) or $\lambda' = (\lambda'_1, \lambda'_2, \dots, \lambda'_{\lambda_1})$, where $\lambda'_i := |\{j : \lambda_j \geq i\}|$ ($i = 1, \dots, \lambda_1$). In particular, $\lambda'_1 = l(\lambda)$. The Ferrers diagram of λ' is obtained by taking the symmetry of the Ferrers diagram of λ with respect of the line $y = x$ of the plane. With the previous example we have $\lambda' = 1^1 2^0 3^2 4^1 5^0 6^1 = (6, 4, 3, 3, 1)$.

14.2. *The algebra of symmetric functions.* Let $\mathbb{Z}[x_1, \dots, x_n]$ be the ring of polynomials in n variables with integral coefficients. A polynomial in that ring is said to be *symmetric*, if it is invariant by permutation of its variables. Let Λ_n be the subring of $\mathbb{Z}[x_1, \dots, x_n]$ of all symmetric polynomials. For every $k \geq 0$ let Λ_n^k be the set of all homogeneous symmetric polynomials of degree k , the zero polynomial included. Then,

$$\Lambda_n = \bigoplus_{k \geq 0} \Lambda_n^k.$$

For each $\alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{N}^n$ let x^α be the monomial $x^\alpha := x_1^{\alpha_1} \dots x_n^{\alpha_n}$ and for each partition λ of length $l(\lambda) \leq n$ let $m_\lambda(x_1, \dots, x_n) := \sum x^\alpha$ denote the sum of all (distinct) monomials x^α , where α is a permutation of $\lambda = (\lambda_1, \dots, \lambda_n)$. The polynomial m_λ is called *monomial symmetric polynomial*. We also use the notation $\sum x_1^{\lambda_1} x_2^{\lambda_2} \dots$ in place of m_λ .

For example, with $n = 4$ variables,

$$\begin{aligned} m_{(1)} &= \sum x_i = x_1 + x_2 + x_3 + x_4; \\ m_{(1,1)} &= \sum x_i x_j = x_1 x_2 + x_1 x_3 + x_1 x_4 + x_2 x_3 + x_2 x_4 + x_3 x_4; \\ m_{(2,1)} &= \sum x_i^2 x_j = x_1^2 x_2 + x_1^2 x_3 + x_1^2 x_4 + x_2^2 x_1 + x_2^2 x_3 + x_2^2 x_4 + x_3^2 x_1 + x_3^2 x_2 + x_3^2 x_4 + x_4^2 x_1 + x_4^2 x_2 + x_4^2 x_3. \end{aligned}$$

The monomial symmetric polynomials $m_\lambda(x_1, \dots, x_n)$ form a \mathbb{Z} -basis for Λ_n , when λ is restricted to the set of all partitions of length $l(\lambda) \leq n$. On the other hand, the polynomials $m_\lambda(x_1, \dots, x_n)$ ($l(\lambda) \leq n$; $|\lambda| = k$) form a \mathbb{Z} -basis for Λ_n^k . When $n \geq k$, that is to say, when the number of variables is *large*, the set of *all* the polynomials m_λ such that $|\lambda| = k$ form a \mathbb{Z} -basis for Λ_n^k . Hence, $\dim \Lambda_n^k = p(k)$, the number of partitions of k .

In the theory of symmetric functions we assume that the number of variables is finite, but *large*; some authors prefer to deal with infinitely many of them, but the statements of certain properties are less intuitive. To make the notion of largeness more precise, we deal, not with polynomials, but with sequences of polynomials $f = (f_n)$ ($n \geq 0$), where each term f_n belongs to Λ_n^k and where for each pair $m \geq n$ the polynomials f_m and f_n satisfy the compatibility property:

$$f_m(x_1, \dots, x_n, 0, \dots, 0) = f_n(x_1, \dots, x_n).$$

Let Λ^k be the set of the sequences $f = (f_n)$, where each term f_n is a symmetric polynomial of degree k . We can show that for $n \geq k$ the mapping ρ_n^k of Λ^k into Λ_n^k that sends $f = (f_n)$ onto f_n is an *isomorphism*. Consequently, Λ^k is of dimension $p(k)$. The set of the sequences $m_\lambda = (m_\lambda(x_1, \dots, x_n))$ ($n \geq 0$) such that $|\lambda| = k$ is a \mathbb{Z} -basis for Λ^k . We then let

$$\Lambda = \bigoplus_{\lambda \geq 0} \Lambda^k.$$

14. SYMMETRIC FUNCTIONS

The elements of Λ are formal series with integral coefficients in the functions m_λ . The ring Λ is called the *ring of the symmetric functions*. For every $n \geq 0$ the mapping of Λ into Λ_n that sends the formal series $\sum_\lambda a_\lambda m_\lambda$ onto the symmetric function $\sum_\lambda a_\lambda m_\lambda(x_1, \dots, x_n)$ is a surjective homomorphism for $n \geq 0$ and maps the formal series $\sum_\lambda a_\lambda m_\lambda$ such that $|\lambda| \leq n$ onto the corresponding series, in an injective manner.

14.3. *The classical bases.* For each $r \geq 1$ the polynomial

$$e_r = m_{1^r} = \sum x_1 x_2 \dots x_r = \sum_{1 \leq i_1 < i_2 < \dots < i_r} x_{i_1} x_{i_2} \dots x_{i_r}$$

is called the r -th *elementary symmetric function*. By convention, $e_0 = 1$. The generating function for the e_r 's is obviously:

$$(14.1) \quad E(u) = \sum_{r \geq 0} e_r u^r = \prod_{i \geq 1} (1 + x_i u).$$

For each partition $\lambda = (\lambda_1, \lambda_2, \dots)$ we define:

$$e_\lambda = e_{\lambda_1} e_{\lambda_2} \dots$$

Notice that if λ is written $\lambda = 1^{m_1} 2^{m_2} \dots$ (multiplicative notation), then $e_\lambda = e_1^{m_1} e_2^{m_2} \dots$.

For each $r \geq 0$ define the *homogeneous symmetric function* of degree r by

$$h_r := \sum_{|\lambda| = r} m_\lambda$$

In particular, $h_0 = 1$, $h_1 = e_1$, $h_2 = m_{(2)} + m_{(1,1)}$, $h_3 = m_{(3)} + m_{(2,1)} + m_{(1,1,1)}$. Now

$$\begin{aligned} \prod_{i \geq 1} (1 - x_i u)^{-1} &= \prod_{i \geq 1} \sum_{k_i \geq 0} (x_i u)^{k_i} \\ &= \sum_{r \geq 0} u^r \sum_{1 \leq i_1 < \dots < i_m} \sum_{\substack{(k_1, \dots, k_m) \\ \sum k_i = r}} x_{i_1}^{k_1} \dots x_{i_m}^{k_m} \\ &= \sum_{r \geq 0} u^r \sum_{|\lambda| = r} \sum_{(k_1, \dots, k_m)} \sum_{1 \leq i_1 < \dots < i_m} x_{i_1}^{k_1} \dots x_{i_m}^{k_m} \end{aligned}$$

[where (k_1, \dots, k_m) is a rearrangement of $(\lambda_1, \dots, \lambda_m)$]

$$= \sum_{r \geq 0} u^r \sum_{|\lambda| = r} m_\lambda = \sum_{r \geq 0} u^r h_r.$$

Thus, the generating function for the h_r 's is

$$(14.2) \quad H(u) = \sum_{r \geq 0} h_r u^r = \prod_{i \geq 1} (1 - x_i u)^{-1}.$$

For every partition $\lambda = (\lambda_1, \lambda_2, \dots)$ define

$$h_\lambda := h_{\lambda_1} h_{\lambda_2} \dots$$

The *power sums* p_r are defined by

$$p_r := m_{(r)} = \sum_i x_i^r;$$

furthermore, let

$$p_\lambda := p_{\lambda_1} p_{\lambda_2} \dots$$

The generating function for the p_r 's, defined by $P(u) := \sum_{r \geq 1} p_r u^{r-1}$, can also be expressed as

$$\begin{aligned} P(u) &= \sum_{i \geq 1} \sum_{r \geq 1} x_i^r u^{r-1} = \sum_{i \geq 1} \frac{x_i}{1 - x_i u} = \sum_{i \geq 1} \frac{d}{du} \log \frac{1}{1 - x_i u} \\ &= \frac{d}{du} \log \prod_i \frac{1}{1 - x_i u} = \frac{d}{du} \log H(u) = \frac{H'(u)}{H(u)}. \end{aligned}$$

The proof of the following theorem can be found in Macdonald [Ma95].

Theorem 14.1. *The functions e_λ (resp. h_λ) form a \mathbb{Z} -basis for Λ . The functions p_λ form a \mathbb{Q} -basis for Λ .*

Theorem 14.2. *We have the relations:*

- (i)
$$\prod_{i \geq 1} (1 + x_i u) = \sum_{r \geq 0} e_r u^r = E(u);$$
- (ii)
$$\prod_{i \geq 1} (1 - x_i u)^{-1} = \sum_{r \geq 0} h_r u^r = H(u);$$
- (iii)
$$H(u)E(-u) = 1;$$
- (iv)
$$\sum_{0 \leq r \leq n} (-1)^r e_r h_{n-r} = 0 \quad (n \geq 1)$$
- (v)
$$\frac{H'(u)}{H(u)} = \sum_{r \geq 1} p_r u^{r-1} = P(u);$$

14. SYMMETRIC FUNCTIONS

$$\begin{aligned}
 \text{(vi)} \quad & \frac{E'(u)}{E(u)} = P(-u); \\
 \text{(vii)} \quad & nh_n = \sum_{1 \leq r \leq n} p_r h_{n-r}; \\
 \text{(viii)} \quad & \prod_{i,j} \frac{1}{1-x_i y_j} = \sum_{n \geq 0} \sum_{|\lambda|=n} m_\lambda(x) h_\lambda(y); \\
 \text{(ix)} \quad & \prod_i \frac{1}{1-x_i u} = \exp \sum_{n \geq 1} p_n(x) \frac{u^n}{n}; \\
 \text{(x)} \quad & \prod_{i,j} \frac{1}{1-x_i y_j} = \exp \sum_{n \geq 1} \frac{p_n(x) p_n(y)}{n};
 \end{aligned}$$

Proof. Relations (i), (ii) have already been proved; (iii) is a straightforward consequence of (i) and (ii). Identity (iv) is derived by considering the coefficient of t^r on the two sides of (iii). Identity (v) has been proved and (vi) follows from (iii) and (v). When (v) is written in the form $H'(u) = H(u)P(u)$, we obtain (vii).

For the proof of (viii) write

$$\begin{aligned}
 \prod_i \prod_j \frac{1}{1-x_i y_j} &= \prod_i \sum_{r \geq 0} x_i^r h_r(y) && \text{[by (ii)]} \\
 &= \sum_{n \geq 0} \sum_{(r_1, \dots, r_n)} h_{r_1}(y) \dots h_{r_n}(y) \sum_{1 \leq i_1 < \dots < i_n} x_{i_1}^{r_1} \dots x_{i_n}^{r_n} \\
 &= \sum_{n \geq 0} \sum_{|y|=n} h_\lambda(y) m_\lambda(x) = \sum_{n \geq 0} \sum_{|\lambda|=n} h_\lambda(x) m_\lambda(y).
 \end{aligned}$$

For (ix) write:

$$\begin{aligned}
 \log \prod_i \frac{1}{1-x_i u} &= \sum_i \log \frac{1}{1-x_i u} = \sum_i \sum_{n \geq 1} \frac{x_i^n}{n} u^n \\
 &= \sum_{n \geq 1} \frac{u^n}{n} \sum_i x_i^n = \sum_{n \geq 1} \frac{u^n}{n} p_n(x).
 \end{aligned}$$

Finally,

$$\log \prod_i \prod_j \frac{1}{1-x_i y_j} = \sum_i \sum_{n \geq 1} \frac{x_i^n}{n} p_n(y) = \sum_n p_n(x) p_n(y) \frac{1}{n}. \quad \square$$

Let us give further relations between h_n , e_n and p_λ . If $\lambda = 1^{m_1} 2^{m_2} \dots$ is a partition, define

$$z_\lambda := 1^{m_1} m_1! 2^{m_2} m_2! \dots$$

Notice that if C_λ designates the set of the permutations of order n whose cycle structure is given by $1^{m_1}2^{m_2}\dots n^{m_n}$, in other words, if C_λ is the set of all the permutations having m_1 cycles of length 1, m_2 cycles of length 2, \dots , then $z_\lambda = n!/|C_\lambda|$.

Theorem 14.3. *We have:*

$$\begin{aligned} H(u) &= \sum_{\lambda} \frac{1}{z_{\lambda}} p_{\lambda} u^{|\lambda|}; & E(u) &= \sum_{\lambda} (-1)^{|\lambda|-l(\lambda)} \frac{1}{z_{\lambda}} p_{\lambda} u^{|\lambda|}; \\ h_n &= \sum_{|\lambda|=n} \frac{1}{z_{\lambda}} p_{\lambda}; & e_n &= \sum_{|\lambda|=n} (-1)^{|\lambda|-l(\lambda)} \frac{1}{z_{\lambda}} p_{\lambda}. \end{aligned}$$

Proof. It suffices to prove the first identity. But $P(u) = (d/du) \log H(u)$ implies

$$\begin{aligned} H(u) &= \exp \sum_{r \geq 1} p_r \frac{u^r}{r} = \prod_{r \geq 1} \exp \left(p_r \frac{u^r}{r} \right) = \prod_{r \geq 1} \sum_{m_r \geq 0} \frac{(p_r u^r)^{m_r}}{r^{m_r}} \frac{1}{m_r!} \\ &= \sum_{\lambda} u^{\sum r m_r} \frac{\prod p_r^{m_r}}{\prod r^{m_r} m_r!} = \sum_{\lambda} u^{|\lambda|} p_{\lambda} \frac{1}{z_{\lambda}}. \quad \square \end{aligned}$$

15. Schur Functions

Let $\varepsilon(\sigma)$ designate the *signature* of the permutation σ . A polynomial $P(x) = P(x_1, \dots, x_n)$ in n variables is said to be *alternant* or *antisymmetric*, if for every permutation σ of $(1, 2, \dots, n)$ the following relation holds:

$$P(x_{\sigma 1}, x_{\sigma 2}, \dots, x_{\sigma n}) = \varepsilon(\sigma) P(x_1, x_2, \dots, x_n).$$

Let A_n (resp. A_n^k) be the space of all the alternants in n variables (resp. in n variables and homogeneous of degree k , the zero polynomial included). Then every alternant $P(x)$ that belongs to A_n^k can be written

$$P(x) = \sum_{\substack{\alpha_1 > \alpha_2 > \dots > \alpha_n \\ |\alpha| = \alpha_1 + \dots + \alpha_n = k}} c(\alpha) \det(x_i^{\alpha_j})_{(1 \leq i, j \leq n)},$$

since, if $P(x)$ contains the term $x_1^{\alpha_1} \dots x_n^{\alpha_n}$ with coefficient $c(\alpha)$, it also contains the term $x_{\sigma 1}^{\alpha_1} \dots x_{\sigma n}^{\alpha_n}$ with the coefficient $\varepsilon(\alpha)c(\alpha)$. On the other hand, the α_i 's are all distinct, for if it were not the case, every determinant $\det(x_i^{\alpha_j})$ would be zero. Also notice that $c(\alpha)$ is the coefficient of x^α in $P(x)$.

15. SCHUR FUNCTIONS

As $\alpha_1 > \alpha_2 > \dots > \alpha_n$, we can write $\alpha_i = \lambda_i + n - i$ ($i = 1, 2, \dots$), so that $\lambda_1 \geq \lambda_2 \geq \dots > \lambda_n$ and $k = |\alpha| = \lambda_1 + \lambda_2 + \dots + n(n-1)/2$. Thus,

$$P(x) = \sum_{|x|=k=n(n-1)/2} c(\lambda) \det(x_i^{\lambda_j+n-j})_{(1 \leq i, j \leq n)}.$$

Consequently, there exists no alternant in n variables of degree (strictly) less than $n(n-1)/2$.

The determinant $a_\delta = \det(x_i^{n-j})$ ($1 \leq i, j \leq n$), where $\delta = (n-1, n-2, \dots, 1, 0)$ is the Vandermonde determinant, equal to $\prod_{i < j} (x_i - x_j)$. The determinant $a_\alpha = \det(x_i^{\alpha_j})$ can be written $a_\alpha = a_{\lambda+\delta} = \det(x_i^{\lambda_j+n-j})$. But if two α_i are equal, the determinant is zero. Therefore, it is *divisible* by $(x_i - x_j)$ ($i \neq j$) and then by the product $a_\delta = \prod_{i < j} (x_i - x_j)$, that is, by a_δ .

Definition. The quotient $s_\lambda(x_1, \dots, x_n) := a_{\lambda+\delta}/a_\delta$ is called the *Schur function* in the variables x_1, \dots, x_n associated with the partition λ .

The Schur function $a_{\lambda+\delta}$ is *symmetric* and *homogeneous* of degree k . This follows from the fact that it is the ratio of two alternants. On the other hand, the alternants $a_{\lambda+\delta}$ ($|\lambda| = k, l(\lambda) \leq n$) form a basis for $A_n^{k+n(n-1)/2}$. The mapping $A \mapsto a_\delta Q$ is an isomorphism of Λ_n^k onto $A_n^{k+n(n-1)/2}$, the kernel being zero, since $a_\delta Q = 0 \Rightarrow Q = 0$. The following theorem has then be proved.

Theorem 15.1. *The Schur functions $s_\lambda(x_1, \dots, x_n)$ ($|\lambda| = k, l(\lambda) \leq n$) form a \mathbb{Z} -basis for Λ_n^k and the Schur functions $s_\lambda(x_1, \dots, x_n)$ ($l(\lambda) \leq n$) form a \mathbb{Z} -basis for Λ_n .*

There is a compatibility relation that holds for the Schur functions, as shown in the next proposition.

Proposition 15.2. *Let $l(\lambda) = l$ and p, q be two integers such that $l \leq p < q$. Then,*

$$s_\lambda(x_1, \dots, x_p) = s_\lambda(x_1, \dots, x_p, x_{p+1}, \dots, x_q) \Big|_{x_{p+1} = \dots = x_q = 0}.$$

Proof. It suffices to verify the proposition for $q = p + 1$. First, $a_{\lambda+\delta}(x_1, \dots, x_{p+1})$ is equal to

$$\begin{vmatrix} x_1^{\lambda_1+p} & \dots & x_1^{\lambda_p+1} & x_1^{\lambda_{p+1}} \\ \dots & \dots & \dots & \dots \\ x_p^{\lambda_1+p} & \dots & x_p^{\lambda_p+1} & x_p^{\lambda_{p+1}} \\ x_{p+1}^{\lambda_1+p} & \dots & x_{p+1}^{\lambda_p+1} & x_{p+1}^{\lambda_{p+1}} \end{vmatrix} = \begin{vmatrix} x_1^{\lambda_1+p} & \dots & x_1^{\lambda_p+1} & 1 \\ \dots & \dots & \dots & \dots \\ x_p^{\lambda_1+p} & \dots & x_p^{\lambda_p+1} & 1 \\ x_{p+1}^{\lambda_1+p} & \dots & x_{p+1}^{\lambda_p+1} & 1 \end{vmatrix},$$

since $\lambda_{p+1} = 0$. Hence, $a_{\lambda+\delta}(x_1, \dots, x_p, 0)$ is equal to

$$\begin{aligned} \begin{vmatrix} x_1^{\lambda_1+p} & \dots & x_1^{\lambda_p+1} & 1 \\ \dots & \dots & \dots & \dots \\ x_p^{\lambda_1+p} & \dots & x_p^{\lambda_p+1} & 1 \\ 0 & \dots & 0 & 1 \end{vmatrix} &= x_1 \dots x_p \begin{vmatrix} x_1^{\lambda_1+p-1} & \dots & x_1^{\lambda_p} \\ \dots & \dots & \dots \\ x_p^{\lambda_1+p-1} & \dots & x_p^{\lambda_p} \end{vmatrix} \\ &= x_1 \dots x_p a_{\lambda+\delta}(x_1, \dots, x_p). \end{aligned}$$

Thus,

$$s_{\lambda}(x_1, \dots, x_p, 0) = \frac{x_1 \dots x_p a_{\lambda+\delta}(x_1, \dots, x_p)}{x_1 \dots x_p a_{\delta}(x_1, \dots, x_p)} = s_{\lambda}(x_1, \dots, x_p). \quad \square$$

We can then define the Schur functions as infinite sequences $s_{\lambda} = (s_{\lambda}(x_1, \dots, x_n))$ ($n \geq 0$). Thus the s_{λ} 's form a \mathbb{Z} -basis for Λ and the s_{λ} 's ($|\lambda| = k$) form a \mathbb{Z} -basis for Λ^k .

The Schur functions can also be expressed as plain determinants in the h_k 's and also the e_k 's, as shown in the next proposition. Remember that λ' designates the conjugate of the partition λ .

Proposition 15.3. *We have*

$$\begin{aligned} s_{\lambda} &= \det(h_{\lambda_i-i+j})_{(1 \leq i, j \leq n)} \quad (n \geq l(\lambda)); \\ s_{\lambda} &= \det(e_{\lambda'_i-i+j})_{(1 \leq i, j \leq m)} \quad (m \geq l(\lambda')); \end{aligned}$$

where, by convention, the coefficients are zero when the subscripts of the h_k 's or the e_k 's are strictly negative.

Proof. We only give the proof of the first identity. Start with the formula (iii) of Theorem 14.2, that is, $H(u)E(-u) = 1$ and let $e_i^{(k)}$ be the elementary symmetric function of $x_1, \dots, x_{k-1}, x_{k+1}, \dots, x_n$, with $E^{(k)}(u)$ being the generating function $\sum_{r=0}^{n-1} e_r^{(k)} u^r$. Obviously,

$$\sum_{p \geq 0} h_p u^p \sum_{r=0}^{n-1} e_r^{(k)} (-u)^r = (1 - x_k u)^{-1}.$$

Now consider a sequence (b_1, \dots, b_n) of nonnegative integers and determine the coefficient of u^{b_m} on the two sides of the previous equation. We get

$$\sum_{r=0}^{n-1} h_{b_m-r} (-1)^r e_r^{(k)} = x_k^{b_m},$$

where with $r + j = n$

15. SCHUR FUNCTIONS

$$(15.1) \quad \sum_{r=1}^n h_{b_m-n+r} (-1)^{n-j} e_{n-j}^{(k)} = x_k^{b_m}.$$

Let \mathcal{E} be the $n \times n$ -matrix whose k -th row is given by

$$(15.2) \quad ((-1)^{n-1} e_{n-1}^{(k)}, \dots, (-1) e_1^{(k)}, e_0^{(k)})$$

and let \mathcal{H} be the matrix whose m -th column $\mathcal{H}_{\cdot, m}$ reads

$$(15.3) \quad \mathcal{H}_{\cdot, m}^t = (h_{b_m-n+1}, h_{b_m-n+2}, \dots, h_{b_m}).$$

The left-hand side of identity (15.1) can be rewritten as a matrix product, so that

$$\mathcal{H}\mathcal{E} = (x_k^{b_m}).$$

Now take the determinant of each side: $\det \mathcal{H} \cdot \det \mathcal{E} = \det(x_k^{b_m})$. When $b = \delta$, we get $1 \times \det \mathcal{E} = \det(x_k^{n-j}) = a_\delta$, so that $\det(h_{b_j-n+i}) a_\delta = \det(x_i^{b_j})$. With $b_i := \lambda_i + n - i$, this can be rewritten as

$$\det(h_{\lambda_j+n-j-n+i}) a_\delta = \det(x_i^{\lambda_j+n-j}),$$

i.e.,

$$\det(h_{\lambda_j-j+i}) = \det(h_{\lambda_i-i+j}) = \det(x_i^{\lambda_j+n-j}) / \det(x_i^{n-j}) = s_\lambda. \quad \square$$

Let ν, θ be two Ferrers diagrams such that $\nu \supset \theta$. The set difference $\nu \setminus \theta$, usually denoted by ν/θ , is called a *skew diagram*. When $n \geq l(\nu)$, we can define the *skew Schur function* $s_{\nu/\theta}(x) = s_{\nu/\theta}(x_1, \dots, x_n)$ by the determinantal expression

$$(15.4) \quad s_{\nu/\theta}(x) = \det(h_{\nu_i-\theta_j-i+j}) \quad (1 \leq i, j, \leq n).$$

As above, it can be shown that also

$$(15.5) \quad s_{\nu/\theta}(x) = \det(e_{\nu'_i-\theta'_j-i+j}) \quad (1 \leq i, j, \leq n).$$

when $n \geq l(\nu')$. The skew Schur Function $s_{\nu/\theta}(x)$ is a symmetric function that reduces to $s_\nu(x)$ when θ is the zero partition.

16. The Cauchy identity

We have already found an expression for the expansion of the product $\prod(1 - x_i y_j)^{-1}$, namely

$$(16.1) \quad \prod_{i,j} (1 - x_i y_j)^{-1} = \sum_{\lambda} m_{\lambda}(x) h_{\lambda}(y).$$

Two other expressions can be obtained:

$$(16.2) \quad \prod_{i,j} (1 - x_i y_j)^{-1} = \sum_{\lambda} z_{\lambda}^{-1} p_{\lambda}(x) p_{\lambda}(y);$$

$$(16.3) \quad \prod_{i,j} (1 - x_i y_j)^{-1} = \sum_{\lambda} s_{\lambda}(x) s_{\lambda}(y).$$

In those expansions λ runs over all partitions of integers. First, (16.2) follows from

$$\prod_{i,j} (1 - x_i y_j)^{-1} = \exp \sum_{n \geq 1} p_n(x) p_n(y) \frac{1}{n},$$

since the right-hand side can be written

$$\begin{aligned} \prod_{n \geq 1} \sum_{k_n \geq 0} p_n(x)^{k_n} p_n(y)^{k_n} \frac{1}{n^{k_n} k_n!} &= \sum_{(n_i)} \sum_{(k_{n_i})} \prod_i \frac{p_{n_i}(x)^{k_{n_i}} p_{n_i}(y)^{k_{n_i}}}{n_i^{k_{n_i}} k_{n_i}!} \\ &= \sum_{\lambda} \frac{1}{z_{\lambda}} p_{\lambda}(x) p_{\lambda}(y). \end{aligned}$$

This proves (16.2). The third identity follows from the Binet-Cauchy identity that reads:

$$\det((1 - x_i y_j)^{-1})_{(1 \leq i, j \leq n)} = a_{\delta}(x) a_{\delta}(y) \prod_{i,j=1}^n (1 - x_i y_j)^{-1}.$$

The proof of the Binet-Cauchy identity can be made as follows. Multiply the i -th row of the determinant $\det((1 - x_i y_j)^{-1})$ by the product $\prod_{k=1}^n (1 - x_i y_k)$ and do it for each $i = 1, \dots, n$. The entry in (i, j) becomes

$$\begin{aligned} \prod_{k \neq j} (1 - x_i y_k) &= \sum_{r=0}^{n-1} x_i^r (-1)^r e_r^{(j)}(y) \quad [e_r^{(j)}(y) = e_r(y_1, \dots, \check{y}_j, \dots, y_n)] \\ &= \sum_{r=1}^n x_i^{n-r} (-1)^{n-r} e_{n-r}^{(j)}(y). \end{aligned}$$

As product of matrices this can be read as

$$\left(\prod_{k \neq j} (1 - x_i y_k) \right)_{(i,j)} = (x_i^{n-j})_{(i,j)} \left((-1)^{n-i} e_{n-i}^{(j)}(y) \right)_{(i,j)};$$

and as product of determinants as

$$\begin{aligned} \det \left(\prod_{k \neq j} (1 - x_i y_k) \right) &= \prod_{i,j} (1 - x_i y_j) \det((1 - x_i y_j)^{-1}) \\ &= \det(x_i^{n-j}) \det((-1)^{n-i} e_{n-i}^{(j)}(y)) \\ &= a_{\delta}(x) a_{\delta}(y), \end{aligned}$$

15. SCHUR FUNCTIONS

remembering that $a_\delta(x) = \det(x_i^{n-j})$ and $\det((-1)^{n-i}e_{n-i}^{(j)}(y)) = \det \mathcal{E} = a_\delta(y)$, using the notations of the previous section.

Now to derive (16.3) we expand each term $(1 - x_i y_j)^{-1}$ in the determinant:

$$\begin{aligned} \det((1 - x_i y_j)^{-1}) &= \det\left(\left(\sum_{m \geq 0} x_i^m y_j^m\right)\right) \\ &= \det\left(\sum_{\alpha_1} \begin{pmatrix} x_1^{\alpha_1} y_1^{\alpha_1} \\ \vdots \\ x_n^{\alpha_1} y_1^{\alpha_1} \end{pmatrix}, \dots, \sum_{\alpha_n} \begin{pmatrix} x_1^{\alpha_n} y_n^{\alpha_n} \\ \vdots \\ x_n^{\alpha_n} y_n^{\alpha_n} \end{pmatrix}\right) \\ &= \sum_{\alpha} \det(x_i^{\alpha_j} y_j^{\alpha_j}) \quad [\alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{N}^n] \\ &= \sum_{\alpha} \det\left(y_1^{\alpha_1} \begin{pmatrix} x_1^{\alpha_1} \\ \vdots \\ x_n^{\alpha_1} \end{pmatrix}, \dots, y_n^{\alpha_n} \begin{pmatrix} x_1^{\alpha_n} \\ \vdots \\ x_n^{\alpha_n} \end{pmatrix}\right) \\ &= \sum_{\alpha} y^\alpha \det(x_i^{\alpha_j}) = \sum_{\alpha} y^\alpha a_\alpha(x). \end{aligned}$$

As $a_\alpha(x) = 0$ if the α_i 's are not all distinct, this yields:

$$\begin{aligned} \det((1 - x_i y_j)^{-1}) &= \sum_{\beta_1 > \dots > \beta_n \geq 0} \sum_{\sigma \in \mathfrak{S}_n} y^{\sigma\beta} a_{\sigma\beta}(x) \\ &= \sum_{\beta} \sum_{\sigma \in S_n} y^{\sigma\beta} \varepsilon(\sigma) a_\beta(x) \\ &= \sum_{\beta} a_\beta(x) a_\beta(y) = \sum_{\lambda} a_{\lambda+\delta}(x) a_{\lambda+\delta}(y), \end{aligned}$$

where $l(\lambda) \leq n$. Hence,

$$\prod_{i,j=1}^n (1 - x_i y_j)^{-1} = \sum_{\lambda} s_\lambda(x_1, \dots, x_n) s_\lambda(y_1, \dots, y_n).$$

The identity also holds for infinitely many variables (x_i, y_j) , for the coefficient of $x_{i_1}^{r_1} \dots x_{i_n}^{r_n} y_{j_1}^{r_1} \dots y_{j_n}^{r_n}$ ($i_1 < \dots < i_n$; $j_1 < \dots < j_n$) in the product $\prod_{i,j \geq 1} (1 - x_i y_j)^{-1}$ is equal to the coefficient of the same monomial in the *finite* product $\prod_{1 \leq i,j \leq N} (1 - x_i y_j)^{-1}$, where $i_n, j_n \leq N$. The identity holding in the finite case, the previous coefficient is equal to the coefficient of the same monomial in $s_\lambda(x_1, \dots, x_N) s_\lambda(y_1, \dots, y_N)$, and also in $s_\lambda(x_1, \dots, x_M) s_\lambda(y_1, \dots, y_M)$ for every $M \geq N$, because of the compatibility property of the s_λ 's: $s_\lambda(x_1, \dots, x_n, 0, \dots, 0) = s_\lambda(x_1, \dots, x_n)$. \square

Identity (16.3) is referred to as the *Cauchy identity* and is used in many combinatorial contexts, especially under the form

$$(16.4) \quad \sum_{n \geq 0} u^n \sum_{|\lambda|=n} s_\lambda(x) s_\lambda(y) = \prod_{i,j} \frac{1}{1 - ux_i y_j}.$$

17. The combinatorial definition of the Schur functions

The theory of symmetric functions would not have such an impact in Combinatorics, if there was no interpretation of the Schur functions in terms of sums of tableau evaluations. The tableaux in questions are called semi-standard and can be introduced as follows.

First, recall that the *Ferrers diagram* associated with a partition $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_r)$ is the set of ordered pairs (i, j) of the Euclidean plane with the property that $1 \leq i \leq \lambda_j$, $1 \leq j \leq r$. Each Ferrers diagram is usually identified with its corresponding partition. For instance, the following Ferrers diagram

$$\begin{array}{cccc} \times & \times & & \\ \times & \times & \times & \\ \times & \times & \times & \\ \times & \times & \times & \times \end{array}$$

corresponds to the partition $\lambda = (5, 3, 3, 2)$.

Let λ be a Ferrers diagram with n points and let $i_1 i_2 \dots i_n$ be a nonincreasing word of length n whose letters are integers. Suppose that those n letters are written on the n points of λ in such a way that every column (resp. every row) is (strictly increasing) from bottom to top (resp. nondecreasing from left to right). The configuration τ thereby obtained is called a *semi-standard tableau*, of *content* $\{i_1, i_2, \dots, i_n\}$ and of *shape* λ . If $i_1 i_2 \dots i_n$ is the word $1, 2, \dots, n$ the semi-standard tableau is called *standard* of order n .

For instance,

$$\tau = \begin{array}{cccc} & 6 & 7 & \\ & 4 & 4 & 8 \\ & 3 & 3 & 5 \\ 1 & 2 & 2 & 4 & 5 \end{array}$$

is a semi-standard tableau of shape $(5, 3, 3, 2)$ and of content $12^2 3^2 4^3 5^2 6^7 8$.

Let $x = \{x_1, x_2, \dots\}$ be a set of variables (finite or infinite) and τ be a semi-standard tableau of shape λ and of content $\{i_1 \leq i_2 \leq \dots \leq i_n\}$. If the cardinality of x is greater than or equal to i_n , the *x-evaluation* of τ is defined to be the monomial $x(\tau) := x_{i_1} x_{i_2} \dots x_{i_n}$, i.e., the product

$$\prod x_{\tau(i,j)},$$

where (i, j) runs over all points (i, j) of the Ferrers diagram λ .

17. THE COMBINATORIAL DEFINITION OF THE SCHUR FUNCTIONS

For each alphabet $x = \{x_1, x_2, \dots, x_n\}$ with $n \geq 7$ the semi-standard tableau of the running example has the x -evaluation $x_1 x_2^2 x_3^2 x_4^3 x_5^2 x_6 x_7 x_8$.

Theorem 17.1. *The Schur function $s_\lambda(x)$ associated with the partition λ in the alphabet x is equal to*

$$(17.1) \quad s_\lambda(x) = \sum_{\tau} x(\tau),$$

where the summation is over all semi-standard tableaux τ of shape λ .

Thus Theorem 17.1 provides a *combinatorial definition* of the Schur functions. Notice that with that definition it is not at all obvious that each Schur function is symmetric in the x_i 's. This follows from the algebraic definition given in section 15, although there are several ways of proving the symmetry property using (17.1) only. To obtain the expansion of $s_\lambda(x)$ it suffices to list the semi-standard tableaux of contents $\{1^{c_1}, 2^{c_2}, \dots\}$ with $c_1 \geq c_2 \geq \dots \geq 1$. As the Schur function is symmetric, the expansion will also include the semi-standard tableaux of contents $\{1^{c_{\sigma 1}}, 2^{c_{\sigma 2}}, \dots\}$ for each permutation σ of the subscripts.

For example, consider the partition $\lambda = (3, 1)$ and the alphabet $x = \{x_1, x_2, x_3, x_4\}$. The semi-standard tableaux τ of shape λ whose contents are of the form $\{1^{c_1}, 2^{c_2}, \dots\}$ with $c_1 \geq c_2 \geq \dots \geq 1$ are the following

$$\begin{array}{ccccccc} 2 & 2 & 3 & 2 & 4 & 3 & 2 \\ 1\ 1\ 1; & 1\ 1\ 2; & 1\ 1\ 2; & 1\ 1\ 3; & 1\ 2\ 3; & 1\ 2\ 4; & 1\ 3\ 4. \end{array}$$

According to Theorem 17.1 we have:

$$s_{(3,1)}(x_1, x_2, x_3, x_4) = \sum x_1^3 x_2 + \sum x_1^2 x_2^2 + 2 \sum x_1^2 x_2 x_3 + 3 x_1 x_2 x_3 x_4.$$

Notice that the first summation involves twelve monomials. For $1 \leq i < j \leq 4$, the monomial $x_i^3 x_j$ (resp. $x_i x_j^3$) corresponds to the semi-standard

tableau $\begin{array}{c} j \\ i\ i\ i \end{array}$ (resp. $\begin{array}{c} j \\ i\ j\ j \end{array}$).

To prove Theorem 17.1 we start with the very first definition of the Schur function as a ratio of two determinants, as given in Section 15. An alternate proof consists of using Proposition 15.3, but this involves other combinatorial techniques

Let λ and μ be two partitions (or Ferrers diagrams) such that μ is contained in λ . The set difference $\lambda \setminus \mu$ is called a *skew tableau* and denoted by λ/μ . A skew tableau is called a *horizontal strip*, if each column contains at most one box (or one cross) of the skew tableau. Let λ_i and μ_i denote




Fig. 17.1

the parts of the partitions λ and μ , respectively. As shown in Fig. 17.1, the skew tableau λ/μ is a horizontal strip if and only if for all i we have $\lambda_{i+1} \leq \mu_i$, i.e.,

$$\lambda_1 \geq \mu_1 \geq \lambda_2 \geq \mu_2 \geq \lambda_3 \geq \dots$$

Let τ be a semi-standard tableau of shape λ and of content $1^{c_1} 2^{c_2} \dots n^{c_n}$. Let $\lambda_0 := \emptyset$ and $\lambda_n := \lambda$. For each $i = 1, 2, \dots, n$ the subset of the boxes of τ containing an integer at most equal to i (resp. an integer equal to i) is a semi-standard tableau (resp. a horizontal strip) we denote by $\lambda^{(i)}$ (resp. $\lambda^{(i)}/\lambda^{(i-1)}$). Let $\lambda_j^{(i)}$ denote the parts of $\lambda^{(i)}$. Hence, for each $i = 1, 2, \dots, n$ the following inequalities holds

$$\lambda_1^{(i)} \geq \lambda_1^{(i-1)} \geq \lambda_2^{(i)} \geq \lambda_2^{(i-1)} \geq \lambda_3^{(i)} \geq \lambda_3^{(i-1)} \geq \dots$$

With those notations the x -evaluation of τ is given by:

$$x(\tau) = x_1^{|\lambda^{(1)}| - |\lambda^{(0)}|} x_2^{|\lambda^{(2)}| - |\lambda^{(1)}|} \dots x_n^{|\lambda^{(n)}| - |\lambda^{(n-1)}|}.$$

Theorem 17.1 can then be rephrased as follows.

Theorem 17.1'. *We have:*

$$s_\lambda(x_1, \dots, x_n) = \sum_{\substack{\emptyset = \lambda^{(0)} \subset \lambda^{(1)} \subset \dots \subset \lambda^{(n)} = \lambda \\ \lambda^{(i)}/\lambda^{(i-1)} \text{ horizontal strip}}} x_1^{|\lambda^{(1)}| - |\lambda^{(0)}|} x_2^{|\lambda^{(2)}| - |\lambda^{(1)}|} \dots x_n^{|\lambda^{(n)}| - |\lambda^{(n-1)}|}.$$

The formula holds for $n = 1$, since the only Schur functions $s_\lambda(x_1)$ in one variable x_1 are the monomials $x_1^{|\lambda|}$, where $\lambda = \lambda^{(1)}$ is reduced to a single part. The formula can be rewritten

$$s_\lambda(x_1, \dots, x_n) = \sum_{\lambda^{(n-1)} \subset \lambda} x_n^{|\lambda| - |\lambda^{(n-1)}|} \sum_{\lambda^{(0)} \subset \lambda^{(1)} \subset \dots \subset \lambda^{(n-1)}} x_1^{|\lambda^{(1)}| - |\lambda^{(0)}|} \dots x_{n-1}^{|\lambda^{(n-1)}| - |\lambda^{(n-2)}|},$$

keeping in mind that each skew tableau $\lambda^{(i)}/\lambda^{(i-1)}$ is a horizontal strip. The second summation is nothing but the formula of the theorem for

17. THE COMBINATORIAL DEFINITION OF THE SCHUR FUNCTIONS

$(n - 1)$ variables. Let $\mu := \lambda^{(n-1)}$ and use the induction hypothesis. We are left to prove the identity

$$(\star) \quad s_\lambda(x_1, \dots, x_n) = \sum_{\substack{\mu \\ \lambda/\mu \text{ horizontal strip}}} x_n^{|\lambda|-|\mu|} s_\mu(x_1, x_2, \dots, x_{n-1}).$$

To prove (\star) we proceed as follows. Consider the determinant

$$a_\lambda(x_1, \dots, x_n) := \det(x_i^{\lambda_j+n-j})_{(1 \leq i, j \leq n)}$$

and evaluate

$$a_\lambda(x_1, \dots, x_{n-1}, 1) = \begin{vmatrix} x_1^{\lambda_1+n-1} & \dots & x_1^{\lambda_n+n-n} \\ \vdots & \ddots & \vdots \\ x_{n-1}^{\lambda_1+n-1} & \dots & x_{n-1}^{\lambda_n+n-n} \\ 1 & \dots & 1 \end{vmatrix}.$$

Subtract the $(j + 1)$ -st column from the j -th column for each $j = 1, 2, \dots, n - 1$. We get:

$$\begin{vmatrix} & & & x_1^{\lambda_n} \\ & E & & \\ & & & x_{n-1}^{\lambda_n} \\ 0 & \dots & 0 & 1 \end{vmatrix},$$

where E is the matrix $(x_i^{\lambda_j+n-j} - x_i^{\lambda_{j+1}+n-j-1})_{(1 \leq i, j \leq n-1)}$. But

$$\begin{aligned} x_i^{\lambda_j+n-j} - x_i^{\lambda_{j+1}+n-j-1} &= (x_i - 1) \sum_{\lambda_{j+1}+n-j-1 \leq t \leq \lambda_j+n-j-1} x_i^t \\ &= (x_i - 1) \sum_{\lambda_{j+1} \leq \mu_j \leq \lambda_j} x_i^{\mu_j+n-j-1}. \end{aligned}$$

Hence

$$\begin{aligned} \det E &= \det \left((x_i - 1) \sum_{\lambda_{j+1} \leq \mu_j \leq \lambda_j} x_i^{\mu_j+n-j-1} \right)_{(1 \leq i, j \leq n-1)} \\ &= \prod_{1 \leq i \leq n} (x_i - 1) \times \sum_{\substack{(\lambda_{j+1} \leq \mu_j \leq \lambda_j) \\ (1 \leq j \leq n-1)}} \det(x_i^{\mu_j+n-j-1})_{(1 \leq i, j \leq n-1)}. \end{aligned}$$

Now if we divide $\det E$ by

$$\Delta(x_1, \dots, x_{n-1}, 1) = \prod_{1 \leq i \leq n} (x_i - 1) \cdot \Delta(x_1, \dots, x_{n-1}),$$

we obtain the Schur function $s_\lambda(x_1, \dots, x_{n-1}, 1)$. Consequently,

$$\begin{aligned} s_\lambda(x_1, \dots, x_{n-1}, 1) &= \sum_{\substack{(\lambda_{j+1} \leq \mu_j \leq \lambda_j) \\ (1 \leq j \leq n-1)}} \frac{\det(x_i^{\mu_j + n - 1 - j})_{(1 \leq i, j \leq n-1)}}{\Delta(x_1, \dots, x_{n-1})} \\ &= \sum_{\substack{\mu \\ \lambda/\mu \text{ horizontal strip}}} s_\mu(x_1, \dots, x_{n-1}). \end{aligned}$$

As $s_\lambda(x_1, \dots, x_{n-1}, x_n)$ is of degree $|\lambda|$ with respect to the set of all variables and as $s_\mu(x_1, \dots, x_{n-1})$ is of degree $|\mu|$, we get:

$$s_\lambda(x_1, \dots, x_{n-1}, x_n) = \sum_{\substack{\mu \\ \lambda/\mu \text{ horizontal strip}}} x_n^{|\lambda| - |\mu|} s_\mu(x_1, \dots, x_{n-1}),$$

which is formula (\star) . \square

There is also a combinatorial definition for the skew Schur function that reads

$$(17.2) \quad s_{\nu/\theta}(x) = \sum_{\tau} x(\tau),$$

where the summation is over all semi-standard tableaux τ of shape ν/θ . Those semi-standard tableaux obey the same rules as the semi-standard tableaux occurring in (17.1): every column (resp. every row) is strictly increasing from bottom to top (resp. nondecreasing from left to right). However their shapes are skew diagrams.

For instance,

$$\tau = \begin{array}{cc} 6 & 8 \\ 1 & 5 & 5 \\ & & 2 \\ & & & 1 & 2 & 2 \end{array}$$

is such a semi-standard tableau of shape ν/θ with $\nu = (6, 4, 3, 2)$ and $\theta = (3, 3)$.

To derive (17.2) we have to use another combinatorial technique, since we have to start from a determinantal definition such as (15.4) and not a ratio of two determinants. The proof of (17.2) is not given in this memoir.

18. The inverse ligne of route of a standard tableau

In this section we consider standard tableaux whose shapes can be skew diagrams ν/θ , as introduced in section 15. If the skew diagram has n

18. THE INVERSE LIGNE OF ROUTE OF A STANDARD TABLEAU

boxes (or crosses), the entries written on those crosses are the n integers $1, 2, \dots, n$, the rows (resp. columns) being increasing when read from left to right (resp. from bottom to top). For instance,

$$T = \begin{array}{cccc} & 6 & 8 & \\ & 4 & 5 & 9 \\ & & & 3 \\ & & & 1 & 2 & 7 \end{array}$$

is a standard tableau of shape ν/θ with $\nu = 6, 4, 3, 2$ and $\theta = 3, 3$.

By analogy with the permutations, we can define the *inverse ligne of route* of the standard tableau T as the set, $\text{Iligne } T$, of all k such that $(k+1)$ is *above* k in T (or, equivalently, to its left). With the above example we have

$$\text{Iligne } T = \{2, 3, 5, 7\}.$$

Notice that for a standard tableau no ligne of route is introduced. The word “inverse” has been here added for convenience. We further define:

$$(18.1) \quad \text{ides } T := |\text{Iligne } T| \quad \text{and} \quad \text{imaj } T := \sum_j j \quad (j \in \text{Iligne } T).$$

A more refined statistic is the *y-vector* of T defined as the word $y(T) = y(1)y(2)\dots y(n)$, where

$$(18.2) \quad y(i) := \#\{j \geq i : j \in \text{Iligne } T\}.$$

Using the same example, $\text{ides } T = 4$, $\text{imaj } T = 2 + 3 + 5 + 7 = 17$ and

$$y(T) = \begin{array}{cccccccccc} & 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 \\ & 4 & 4 & 3 & 2 & 2 & 1 & 1 & 0 & 0 \end{array}$$

The next proposition is straightforward and given without proof.

Proposition 18.1. *For each standard tableau T whose y -vector reads $y(T) = y(1)y(2)\dots y(n)$ the following holds:*

$$\text{ides } T = y(1), \quad \text{imaj } T = \text{tot } y(T) = y(1) + y(2) + \dots + y(n).$$

Now let $\{x_1, x_2, \dots\}$ and $\{q_1, q_2, \dots\}$ be two sets of independent variables. If $x_{c(1)}x_{c(2)}\dots x_{c(n)}$ is a monomial written in such a way that $c(1) \geq c(2) \geq \dots \geq c(n)$, define

$$\phi_{\mathbf{q}}(x_{c(1)}x_{c(2)}\dots x_{c(n)}) := q_1^{c(1)-1} q_2^{c(2)-1} \dots q_n^{c(n)-1}$$

and let $\phi_{\mathbf{q}}$ act linearly on the algebra of the polynomials in the x_i 's. If the variables q_i 's are distinct, then $\phi_{\mathbf{q}}$ cannot be a ring homomorphism. For a standard tableau T of order n it will be convenient to use the notation: $\mathbf{q}^{y(T)} := q_1^{y(1)} q_2^{y(2)} \dots q_n^{y(n)}$. Also, for each variable t introduce

$$[t; \mathbf{q}]_n := \begin{cases} 1 - t, & \text{if } n = 0; \\ (1 - t)(1 - tq_1)(1 - tq_1q_2) \dots (1 - tq_1 \dots q_n), & \text{if } n \geq 1. \end{cases}$$

Theorem 18.2. *Let ν/θ be a skew diagram with n points. Then, the following identities hold:*

$$(18.3) \quad \phi_{\mathbf{q}}(s_{\nu/\theta}(x_1, x_2, \dots)) = \frac{1}{\prod_{i=1}^n (1 - q_1 \dots q_i)} \sum_{T, T \text{ of shape } \nu/\theta} \mathbf{q}^{y(T)};$$

$$(18.4) \quad \sum_{r \geq 0} t^r \phi_{\mathbf{q}}(s_{\nu/\theta}(x_1, \dots, x_{r+1})) = \frac{1}{[t; \mathbf{q}]_{n+1}} \sum_{T, T \text{ of shape } \nu/\theta} t^{\text{idex } T} \mathbf{q}^{y(T)};$$

the summations on the two right-hand sides being over all standard tableaux of shape ν/θ .

When all the variables q_i are equal to a single variable q , the homomorphism $\phi_{\mathbf{q}}$, that will then be denoted by ϕ_q , becomes a *ring homomorphism*. We then have

$$(18.5) \quad \phi_q(s_{\nu/\theta}(x_1, x_2, \dots)) = s_{\nu/\theta}(1, q, q^2, \dots);$$

$$(18.6) \quad \phi_q(s_{\nu/\theta}(x_1, \dots, x_{r+1})) = s_{\nu/\theta}(1, q, \dots, q^r).$$

In view of Proposition 18.1 this implies the subsequent corollary.

Corollary 18.3. *Let ν/θ be a skew diagram with n points. Then, the following identities hold:*

$$(18.7) \quad s_{\nu/\theta}(1, q, q^2, \dots) = \frac{1}{(q; q)_n} \sum_{T, T \text{ of shape } \nu/\theta} q^{\text{imaj } T};$$

$$(18.8) \quad \sum_{r \geq 0} t^r s_{\nu/\theta}(1, q, \dots, q^r) = \frac{1}{(t; q)_{n+1}} \sum_{T, T \text{ of shape } \nu/\theta} t^{\text{idex } T} q^{\text{imaj } T};$$

the summations on the two right-hand sides being over all standard tableaux of shape ν/θ .

The proof of Theorem 18.2 is based upon the *combinatorial* definition of Schur functions (or skew Schur functions), as discussed in the previous section, and a bijection

$$\tau \mapsto (T, d)$$

18. THE INVERSE LIGNE OF ROUTE OF A STANDARD TABLEAU

of the set of semi-standard tableaux onto pairs (T, d) , where T is a standard tableau having the same shape as τ and d a nonincreasing sequence of integers.

The *semi-standard* tableaux we will consider this time are *non-increasing* along the rows (from left to right) and (strictly) decreasing (from bottom to top) along the columns, as opposed to the definition given in the previous section. This convention will facilitate the calculations. For instance

$$\tau = \begin{array}{cccc} 2 & 1 & & \\ 3 & 3 & 1 & \\ & & & 5 \\ & & & 8 & 6 & 2 \end{array}$$

is such a semi-standard tableau, whose shape is ν/θ with $\nu = (6, 4, 3, 2)$ and $\theta = (3, 3)$.

Each such semi-standard tableau τ determines a total ordering on the points (i, j) of its diagram ν/θ in the following manner: (i, j) is said to be less than (i', j') , if the integer $\tau(i, j)$ written on (i, j) is greater than $\tau(i', j')$, or if $\tau(i, j) = \tau(i', j')$ and (i, j) is to the left of (i', j') , that is, $i < i'$.

Suppose that the diagram ν/θ has n elements; write k on the point (i, j) if (i, j) is the smallest k -th element using the previous ordering. This produces a *standard tableau* T , of order n , of shape ν/θ . Reading the elements $\tau(i, j)$ from smallest to greatest yields a non-increasing sequence $c = c(1)c(2) \dots c(n)$, called the *content* of τ . Moreover, we have

$$(18.9) \quad k \in \text{Iligne } T \Rightarrow c(k) > c(k + 1).$$

As $i \in \text{Iligne } T \Rightarrow y(i) = y(i + 1) + 1$ and $i \notin \text{Iligne } T \Rightarrow y(i) = y(i + 1)$, we see that the sequence $d := d(1)d(2) \dots d(n)$, defined by $d(i) := c(i) - 1 - y(i)$ for each $i = 1, 2, \dots, n$, satisfies $d(1) \geq d(2) \geq \dots \geq d(n) \geq 0$. Moreover, the mapping $\tau \mapsto (T, d)$ is *bijective*.

With the previous semi-standard tableau τ the standard tableau T is the tableau given in the beginning of this section. We further have:

$$\begin{array}{cccccccccc} & 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 \\ c = & 8 & 6 & 5 & 3 & 3 & 2 & 2 & 1 & 1 \\ y(T) = & 4 & 4 & 3 & 2 & 2 & 1 & 1 & 0 & 0 \\ d = & 3 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \end{array}$$

Next, for each semi-standard tableau τ of content $c = c(1) \dots c(n)$ define

$$x(\tau) := x_{c(1)} \cdots x_{c(n)},$$

so that

$$\phi_{\mathbf{q}}(x(\tau)) := q_1^{c(1)-1} q_2^{c(2)-1} \dots q_n^{c(n)-1}.$$

Using the above bijection $\tau \mapsto (T, d)$, we get

$$\begin{aligned}\phi_{\mathbf{q}}(x(\tau)) &= q_1^{y(1)} q_2^{y(2)} \cdots q_n^{y(n)} q_1^{d(1)} q_2^{d(2)} \cdots q_n^{d(n)} \\ &= \mathbf{q}^{y(T)} \mathbf{q}^d.\end{aligned}$$

Accordingly,

$$\sum_{\tau, \tau \text{ of shape } \nu/\theta} \phi_{\mathbf{q}}(x(\tau)) = \sum_{T, T \text{ of shape } \nu/\theta} \mathbf{q}^{y(T)} \sum_d \mathbf{q}^d.$$

As the last summation is over all non-increasing sequences $d = d(1) \geq \cdots \geq d(n) \geq 0$, we obtain

$$(18.10) \quad \sum_{\tau, \tau \text{ of shape } \nu/\theta} \phi_{\mathbf{q}}(x(\tau)) = \frac{1}{\prod_{i=1}^n (1 - q_1 \cdots q_i)} \sum_{T, T \text{ of shape } \nu/\theta} \mathbf{q}^{y(T)}.$$

Now make use of the combinatorial definition for the skew Schur function

$$s_{\nu/\theta}(x_1, x_2, \dots) = \sum_{\tau, \tau \text{ of shape } \nu/\theta} x(\tau),$$

and take the homomorphic image of identity (18.10) under $\phi_{\mathbf{q}}$. This yields (18.3).

For the proof of (18.4) proceed as follows. In the expansion of the skew Schur function $s_{\nu/\theta}(x_1, x_2, \dots, x_{r+1})$ in the finite alphabet $\{x_1, x_2, \dots, x_{r+1}\}$ the semi-standard tableaux τ of content c such that $c(1) \geq r + 2$ bring no contribution. Consequently,

$$\sum_{c(1)=r+1} x(c) = s_{\nu/\theta}(x_1, x_2, \dots, x_{r+1}) - s_{\nu/\theta}(x_1, x_2, \dots, x_r);$$

hence,

$$\begin{aligned}\sum_r t^r \sum_{c(1)=r+1} x(c) &= \sum_r t^r (s_{\nu/\theta}(x_1, x_2, \dots, x_{r+1}) - s_{\nu/\theta}(x_1, x_2, \dots, x_r)) \\ &= (1 - t) \sum_r t^r s_{\nu/\theta}(x_1, x_2, \dots, x_{r+1}).\end{aligned}$$

On the other hand, by using (18.10) we get

$$\begin{aligned}\phi_{\mathbf{q}}\left(\sum_r t^r \sum_{c(1)=r+1} x(c)\right) &= \sum_{\tau, \tau \text{ of shape } \nu/\theta} (tq_1)^{c(1)-1} q_1^{c(2)-1} \cdots q_n^{c(n)-1} \\ &= \frac{1}{\prod_{i=1}^n (1 - tq_1 \cdots q_i)} \sum_{T, T \text{ of shape } \nu/\theta} t^{\text{ides } T} \mathbf{q}^{y(T)},\end{aligned}$$

since $y(1) = \text{ides } T$. Thus identity (18.4) is proved.

19. The Robinson-Schensted Correspondence

This is one of the most celebrated correspondences. Its main interest in Combinatorics lies in the fact that geometric properties held by permutations can be transferred onto tableaux. Remember that we have introduced inverse lignes of route for both permutations and standard tableaux. One of our problems will be to relate those two lignes of routes through the Robinson-Schensted correspondence.

19.1. *The Schensted-Knuth algorithm.* In section 12.2 each nondecreasing biword \mathbf{b} has been mapped onto a triple (σ, b', c') , where σ is a permutation and b', c' two nondecreasing words. The Schensted-Knuth algorithm will map such a biword onto a pair of two semi-standard tableaux *of the same shape*, that shape being a (right) Ferrers diagram. Again take up the example of section 12.1

$$\mathbf{b} = \begin{pmatrix} b \\ c \end{pmatrix} = \begin{pmatrix} 0 & 0 & 0 & 1 & 1 & 1 & 1 & 3 & 3 & 3 & 3 & 3 \\ 2 & 4 & 4 & 0 & 3 & 3 & 3 & 1 & 2 & 2 & 4 & 4 \end{pmatrix},$$

a biword of length $n = 12$, which is nondecreasing in the sense that the billetters $(0, 2), (0, 4), \dots$, are in nondreasing order with respect to the lexicographic order. The Schensted-Knuth algorithm applied to \mathbf{b} can be described as follows:

start with two empty tableaux $P = \begin{array}{|} \hline \\ \hline \end{array}, Q = \begin{array}{|} \hline \\ \hline \end{array}$. Next, insert the first entry 0 on the top row b into Q and the first entry 2 on the bottom row c into P , to obtain $P = \begin{array}{|} \hline 2 \\ \hline \end{array}, Q = \begin{array}{|} \hline 0 \\ \hline \end{array}$. As $2 \leq 4 \leq 4$ (on the bottom row c), insert 44 next to 2 onto the same row in P and record the positions by inserting the corresponding entries 00 (on the top row b) onto analogous places in Q :

$$P = \begin{array}{|} \hline 2 & 4 & 4 \\ \hline \end{array}, Q = \begin{array}{|} \hline 0 & 0 & 0 \\ \hline \end{array}.$$

Next, the fourth letter 0 of c will *bump up the smallest entry in P that is strictly larger than itself*. This is 2. We also record the new position occupied by 2 on the second row in P by writing 1 (the fourth entry on the top row) into Q :

$$P = \begin{array}{|} \hline 2 \\ \hline 0 & 4 & 4 \\ \hline \end{array}, Q = \begin{array}{|} \hline 1 \\ \hline 0 & 0 & 0 \\ \hline \end{array}.$$

Then, the fifth letter 3 of c will *bump up the smallest leftmost entry on the first row in P that is strictly larger than itself*. This is the leftmost 4. There is no entry on the second row of P smaller than 4, so that 4 is

inserted next to the right of 2. We then record the new position occupied by 4 on the second row in P by writing 1 (the fifth letter of b) into Q :

$$P = \begin{array}{|c|c|} \hline 2 & 4 \\ \hline 0 & 3 & 4 \\ \hline \end{array}, \quad Q = \begin{array}{|c|c|} \hline 1 & 1 \\ \hline 0 & 0 & 0 \\ \hline \end{array}.$$

Again, the sixth letter 3 of c will *bump up the smallest leftmost entry on the first row in P that is strictly larger than itself*. This is 4. There is no entry on the second row of P smaller than 4, so that 4 is inserted to the right of the second row. We then record the new position occupied by 4 on the second row in P by writing 1 (the sixth letter of b) into Q :

$$P = \begin{array}{|c|c|c|} \hline 2 & 4 & 4 \\ \hline 0 & 3 & 3 \\ \hline \end{array}, \quad Q = \begin{array}{|c|c|c|} \hline 1 & 1 & 1 \\ \hline 0 & 0 & 0 \\ \hline \end{array}.$$

The seventh letter of c is 3. There is no entry on the first row in P that is greater than 3, so that 3 is inserted to the right of the first row and the seventh letter 1 of b is written to the right of the first row in Q :

$$P = \begin{array}{|c|c|c|} \hline 2 & 4 & 4 \\ \hline 0 & 3 & 3 & 3 \\ \hline \end{array}, \quad Q = \begin{array}{|c|c|c|c|} \hline 1 & 1 & 1 \\ \hline 0 & 0 & 0 & 1 \\ \hline \end{array}.$$

The eighth letter of c is 1. It will *bump up the smallest leftmost entry on the first row in P that is strictly larger than itself*. This is the leftmost 3. In its turn that entry 3 will *bump up the smallest leftmost entry on the second row in P that is strictly larger than itself*. This is the leftmost 4. The third row is empty, so that 4 will form a new one-entry row for P . We then record the new position occupied by 4 on the third row in P by writing 3 (the eighth letter of b) into Q :

$$P = \begin{array}{|c|c|c|} \hline 4 \\ \hline 2 & 3 & 4 \\ \hline 0 & 1 & 3 & 3 \\ \hline \end{array}, \quad Q = \begin{array}{|c|c|c|} \hline 3 \\ \hline 1 & 1 & 1 \\ \hline 0 & 0 & 0 & 1 \\ \hline \end{array}.$$

Using the same insertion technique we successively get

$$P = \begin{array}{|c|c|} \hline 4 & 4 \\ \hline 2 & 3 & 3 \\ \hline 0 & 1 & 2 & 3 \\ \hline \end{array}, \quad Q = \begin{array}{|c|c|} \hline 3 & 3 \\ \hline 1 & 1 & 1 \\ \hline 0 & 0 & 0 & 1 \\ \hline \end{array}; \quad P = \begin{array}{|c|c|} \hline 4 & 4 \\ \hline 2 & 3 & 3 & 3 \\ \hline 0 & 1 & 2 & 2 \\ \hline \end{array}, \quad Q = \begin{array}{|c|c|} \hline 3 & 3 \\ \hline 1 & 1 & 1 & 3 \\ \hline 0 & 0 & 0 & 1 \\ \hline \end{array};$$

and finally

$$P = \begin{array}{|c|c|} \hline 4 & 4 \\ \hline 2 & 3 & 3 & 3 \\ \hline 0 & 1 & 2 & 3 & 4 & 4 \\ \hline \end{array}, \quad Q = \begin{array}{|c|c|} \hline 3 & 3 \\ \hline 1 & 1 & 1 & 3 \\ \hline 0 & 0 & 0 & 1 & 3 & 3 \\ \hline \end{array}.$$

Theorem 19.1. *The insertion algorithm*

$$\mathbf{b} = \begin{pmatrix} b \\ c \end{pmatrix} = \begin{pmatrix} b_1 & b_2 & \dots & b_n \\ c_1 & c_2 & \dots & c_n \end{pmatrix} \mapsto PQ$$

provides a bijection of the set of all nondecreasing biwords onto the set of ordered pairs PQ of semi-standard tableaux, of the same shape, the content of P (resp. of Q) being $\{c_1, c_2, \dots, c_n\}$ (resp. $\{b_1, b_2, \dots, b_n\}$).

It is not our intention to give a formal proof of the theorem. The only difficult part is to see that the algorithm provides two *tableaux* which are both *semi-standard*. The fact that both P and Q have the same shape is then evident. Also, the algorithm for the reverse construction is easy to formulate.

In the sequel, *standard Young tableau* of order n means standard tableau whose shape is a (right) Ferrers diagram corresponding to a partition of n . Let \mathfrak{T}_n (resp. $\mathfrak{T}_n^{(2)}$) be the set of all standard Young tableaux of order n (resp. of all ordered pairs PQ of standard Young tableaux of order n of the same shape).

Corollary 19.2. *When restricted to permutations σ of order n written as increasing biwords, the insertion algorithm*

$$\sigma \mapsto PQ$$

establishes a bijection of the group \mathfrak{S}_n of $n!$ elements onto $\mathfrak{T}_n^{(2)}$. The bijection is called the Robinson-Schensted correspondence.

19.2. *A combinatorial proof of the Cauchy identity.* Let

$$A = (a_{ij})_{(i,j \geq 0)} \mapsto \mathbf{b} = \begin{pmatrix} b \\ c \end{pmatrix} = \begin{pmatrix} b_1 & b_2 & \dots & b_n \\ c_1 & c_2 & \dots & c_n \end{pmatrix}$$

be the bijection constructed in section 12.1 that maps each matrix $A = (a_{ij})$ ($i, j \geq 0$) with integral entries such that $\sum_{i,j} a_{ij} = n$ onto a nondecreasing biword of length n . By construction the billetter $\binom{i}{j}$ occurs a_{ij} times in \mathbf{b} . Accordingly, if $x = (x_0, x_1, x_2, \dots)$ and $y = (y_0, y_1, y_2, \dots)$ are two alphabets of variables, we have

$$(19.1) \quad x_{b_1} x_{b_2} \dots x_{b_n} y_{c_1} y_{c_2} \dots y_{c_n} = \prod_{i,j} (x_i y_j)^{a_{ij}}.$$

Now consider the Schensted-Knuth bijection $\mathbf{b} \mapsto PQ$. The product (19.1) is equal to the product of the x -evaluation $x(Q)$ by the y -evaluation $y(P)$.

When the matrix A runs over all matrices with integral entries whose finitely many of them are nonzero, the pair PQ runs over all pairs of semi-standard tableaux, of the same shape. For short, designate the shape of a tableau P by $|P|$. We can then write:

$$\begin{aligned}
 \prod_{i,j}(1 - x_i y_j)^{-1} &= \prod_{i,j} \sum (x_i y_j)^{a_{ij}} \\
 &= \sum_A \prod_{i,j} (x_i y_j)^{a_{ij}} \\
 &= \sum_{PQ} x(Q)y(P) \\
 &= \sum_{\lambda} \sum_{|P|=|Q|=\lambda} x(Q)y(P) \\
 &= \sum_{\lambda} \left(\sum_{|Q|=\lambda} x(Q) \right) \left(\sum_{|P|=\lambda} y(P) \right) \\
 &= \sum_{\lambda} s_{\lambda}(y) s_{\lambda}(x),
 \end{aligned}$$

by the very combinatorial definition for the Schur functions given in section 17.

In the same vein we can prove the dual Cauchy identity, i.e.,

$$(19.2) \quad \prod_{i,j}(1 + x_i y_j) = \sum_{\lambda} s_{\lambda'}(x) s_{\lambda}(y),$$

where λ' still designates the conjugate partition. The product on the left-hand side is equal to

$$\sum_A \prod_{i,j} (x_i y_j)^{a_{ij}},$$

but this time the entries of the matrices are only 0 or 1. The corresponding biwords \mathbf{b} have *distinct* biletters, so that we can derive an inserting process in which each inserted element bumps up the entry that is *larger than or equal to* instead of *the smallest leftmost entry strictly larger than* itself. Recording the tableau Q as before, we obtain a pair PQ , of tableaux of the same shape, such that the transpose P^T of P and Q itself are semi-standard. An analogous calculation leads to identity (19.2).

19.3. *Geometric properties of correspondence.* The ligne of route “Ligne” and the inverse ligne of route “Iligne” of a permutation have been defined in section 11 and the inverse ligne of route “Iligne” of a tableau in section 18.

Theorem 19.3. *Let $\sigma \mapsto PQ$ be the Robinson-Schensted correspondence. Then*

$$(19.3) \quad \text{Ligne } \sigma = \text{Iligne } Q; \quad \text{Iligne } \sigma = \text{Iligne } P.$$

This property can be verified by a careful study of the various steps in the construction of the Schensted-Knuth algorithm. We omit the detailed proof. It is essential to notice that the Q -tableau retains the geometric information on the permutation σ , on the one hand, and the P -tableau on the inverse permutation σ^{-1} , on the other hand.

For instance, with

$$\sigma = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 \\ 3 & 1 & 4 & 2 & 5 \end{pmatrix} \mapsto PQ = \begin{array}{cc} 3 & 4 & & 2 & 4 \\ 1 & 2 & 5 & , & 1 & 3 & 5 \end{array}$$

we observe the equalities: $\text{Ligne } \sigma = \{1, 3\} = \text{Iligne } Q$ and $\text{Iligne } \sigma = \{2\} = \text{Iligne } P$.

The Robinson-Schensted correspondence is a useful ingredient in the construction of bijections of the symmetric group onto itself. For instance, suppose that we have a bijection $Q \mapsto Q^J$ of \mathfrak{S}_n onto itself that preserves the shape and has the property: $\text{Iligne } Q^J = n - \text{Iligne } Q = \{n - i : i \in \text{Iligne } Q\}$. Then the chain

$$\sigma \mapsto PQ \mapsto PQ^J \mapsto \mathbf{j}\sigma,$$

where the rightmost arrow stands for the inverse of the Robinson correspondence, is a bijection of \mathfrak{S}_n onto itself such that

$$\text{Iligne } \mathbf{j}\sigma = \text{Iligne } \sigma, \quad \text{Ligne } \mathbf{j}\sigma = n - \text{Ligne } \sigma.$$

The other important property of the correspondence (for which there exist many proofs) can be stated as follows:

$$(19.4) \quad [\sigma \mapsto PQ] \Leftrightarrow [\sigma^{-1} \mapsto QP].$$

19.4. *A permutation statistic distribution.* As was done in section 18 for standard tableaux, we can introduce the y -vector of a permutation $\sigma = \sigma(1) \dots \sigma(n)$ and define it as the word $y(\sigma) = y_\sigma(1) \dots y_\sigma(n)$, where

$$y_\sigma(i) := \#\{j \geq i : j \in \text{Ligne } \sigma\}.$$

Hence, $\text{des } \sigma = y_\sigma(1)$, $\text{maj } \sigma = \text{tot } y(\sigma) = y_\sigma(1) + \cdots + y_\sigma(n)$. We can also consider the y -vector for the inverse σ^{-1} of σ . Taking two sequences (p_1, p_2, \dots) and (q_1, q_2, \dots) of variables define the monomials

$$\mathbf{q}^{y(\sigma)} := q_1^{y_\sigma(1)} \cdots q_n^{y_\sigma(n)}, \quad \mathbf{p}^{y(\sigma^{-1})} := p_1^{y_{\sigma^{-1}}(1)} \cdots p_n^{y_{\sigma^{-1}}(n)}$$

and consider the polynomial

$$A_n(t, s, \mathbf{q}, \mathbf{p}) := \sum_{\sigma \in \mathfrak{S}_n} t^{\text{des } \sigma} s^{\text{des } \sigma^{-1}} \mathbf{q}^{y(\sigma)} \mathbf{p}^{y(\sigma^{-1})}.$$

The homomorphism $\phi_{\mathbf{q}}$ that has been introduced just after Proposition 18.1, keeps the same meaning. Also let $\phi_{\mathbf{p}}$ be the analogous homomorphism defined by means of the variables p_i 's.

Theorem 19.4. *The bi-factorial generating function for the polynomials $A_n(t, s, \mathbf{q}, \mathbf{p})$ is given by:*

$$\sum_{n \geq 0} \frac{u^n}{[t; \mathbf{q}]_{n+1} [s; \mathbf{p}]_{n+1}} A_n(t, s, \mathbf{q}, \mathbf{p}) = \phi_{\mathbf{q}} \phi_{\mathbf{p}} \sum_{l, k} t^l s^k \prod_{\substack{1 \leq j \leq l+1 \\ 1 \leq i \leq k+1}} \frac{1}{1 - u x_j y_i}.$$

Again, we emphasize the fact that $\phi_{\mathbf{q}}$ and $\phi_{\mathbf{p}}$ are (non-ring) homomorphisms acting on the x_j 's and the y_i 's, respectively. The infinite product on the right-hand is to be extended first in monomials in the x_j 's and the y_i 's and the mappings $\phi_{\mathbf{p}}$, $\phi_{\mathbf{q}}$ are to be applied next to each monomial.

The proof of Theorem 19.4 is a consequence of the geometric properties of the Robinson-Schensted correspondence, as stated in Theorem 19.3, and of the Cauchy identity. From (19.3) we can write:

$$A_n(t, s, \mathbf{q}, \mathbf{p}) = \sum_{|\lambda|=n} \sum_{|P|=|Q|=\lambda} s^{\text{idex } Q} t^{\text{idex } P} \mathbf{p}^{y(Q)} \mathbf{q}^{y(P)}$$

Hence,

$$\begin{aligned} & \frac{1}{[t; \mathbf{q}]_{n+1} [s; \mathbf{p}]_{n+1}} A_n(t, s, \mathbf{q}, \mathbf{p}) \\ &= \sum_{|\lambda|=n} \left(\frac{1}{[t; \mathbf{q}]_{n+1}} \sum_{|Q|=\lambda} t^{\text{idex } Q} \mathbf{q}^{y(Q)} \right) \left(\frac{1}{[s; \mathbf{p}]_{n+1}} \sum_{|P|=\lambda} s^{\text{idex } P} \mathbf{p}^{y(P)} \right) \\ &= \sum_{|\lambda|=n} \left(\sum_{l \geq 0} t^l \phi_{\mathbf{q}}(s_\lambda(x_1, \dots, x_{l+1})) \right) \left(\sum_{k \geq 0} s^k \phi_{\mathbf{p}}(s_\lambda(y_1, \dots, y_{k+1})) \right) \end{aligned}$$

[by (18.4).]

Thus

$$\begin{aligned}
 & \sum_{n \geq 0} \frac{u^n}{[t; \mathbf{q}]_{n+1} [s; \mathbf{p}]_{n+1}} A_n(t, s, \mathbf{q}, \mathbf{p}) \\
 &= \phi_{\mathbf{q}} \phi_{\mathbf{p}} \sum_{l, k} t^l s^k \sum_{n \geq 0} u^n \sum_{|\lambda|=n} s_{\lambda}(x_1, \dots, x_{l+1}) s_{\lambda}(y_1, \dots, y_{k+1}) \\
 &= \phi_{\mathbf{q}} \phi_{\mathbf{p}} \sum_{l, k} t^l s^k \prod_{\substack{1 \leq j \leq l+1 \\ 1 \leq i \leq k+1}} \frac{1}{1 - ux_j y_i} \quad [\text{by the Cauchy formula (16.4).}]
 \end{aligned}$$

When all the variables q_j (resp. the variables p_i) are equal to a single variable q (resp. p), the homomorphisms become ring homomorphisms. We recover formula (13.7) that gives the bi-basic generating functions by the four-statistic (des, ides, maj, imaj).

20. Eulerian Calculus; the first extensions

The next three sections will be devoted to deriving the various extensions of the fundamental identity for the Eulerian polynomials ($A_n(t)$) ($n \geq 0$) (see (10.9)) that reads

$$(20.1) \quad \frac{1-t}{-t + \exp(u(t-1))} = \sum_{n \geq 0} \frac{u^n}{n!} A_n(t),$$

when the exponential occurring in the denominator of the rational fraction is replaced by the q -exponential (as was done in section 10), then by a Bessel function J or by a q -Bessel function \mathbf{J} , in its infinite or finite form, and also when the fraction itself is replaced by

$$(20.2) \quad \frac{(1-t) \exp(u(t-1)X)}{-t + \exp(u(t-1)(X+Y))} = \sum_{n \geq 0} \frac{u^n}{n!} B_n(X, Y, t),$$

that stands for the natural extension of the fraction, when going from the group A_n of the *permutations* to the group B_n of the *signed permutations*.

Those various extensions are symbolized by the diagram of Fig. 20.1, where the identities (20.1) and (20.2) sit on the vertices (\cdot, \cdot, \cdot) and $(\text{sgn}, \cdot, \cdot)$, respectively, and where the horizontal arrow \rightarrow stands for the passage from the permutations to the signed permutations, the oblique arrow \nearrow stands for the q -extension, the uparrow \uparrow for the extension by means of a Bessel function, in its infinite form (J) for the first level and in its finite form (J_k) for the second one. All those terms will be fully explained in the sequel.




Fig. 20.1

Of course, we could start from the most general extension that sits on (sgn, q, J_k) and get the other results by successive specializations. However proceeding in such a way would overlook the local techniques that have their own interest. We then proceed from the particular to the general, at least in the beginning. When we have enough material, we will attack (sgn, q, J_k) itself and obtain the remaining identities with their combinatorial interpretations.

20.1. *The signed permutations.* A signed permutation of order n can be defined as a pair (σ, ε) , where σ is a permutation $\sigma = \sigma(1)\sigma(2) \dots \sigma(n)$ and $\varepsilon = \varepsilon(1)\varepsilon(2) \dots \varepsilon(n)$ is a word of length n in the alphabet $\{x, y\}$. The word ε is called a *xy-word*; the number of letters in ε equal to x (resp. equal to y) is denoted by $\ell(\varepsilon|x)$ (resp. $\ell(\varepsilon|y)$). Also $\sigma_{\varepsilon|x}$ (resp. $\sigma_{\varepsilon|y}$) designates the *subword* of σ made of all letters $\sigma(i)$ such that $\varepsilon(i) = x$ (resp. $\varepsilon(i) = y$).

Definition. The integer i is said to be a *descent* of the signed permutation (σ, ε) , if one of the following conditions is verified

- (i) $i = n$ and $\varepsilon(n) = x$;
- (ii) $1 \leq i \leq n - 1$, $\varepsilon(i) = x$, $\varepsilon(i + 1) = y$;
- (iii) $1 \leq i \leq n - 1$, $\varepsilon(i) = \varepsilon(i + 1)$ and $\sigma(i) > \sigma(i + 1)$.

The *number of descents* of (σ, ε) is denoted by $\text{des}(\sigma, \varepsilon)$.

Example. Let

$$\begin{array}{cccccccccc} & 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 \\ \sigma = & 6 & 7 & 2 & 4 & 3 & 1 & 8 & 5 & 9 \\ \varepsilon = & x & y & y & y & x & x & y & x & x \end{array}$$

be a signed permutation. Then $\sigma_{\varepsilon|x} = 63159$, $\sigma_{\varepsilon|y} = 7248$ and $\text{des}(\sigma, \varepsilon) = \text{Card}\{1, 2, 5, 6, 9\} = 5$.

Another way of introducing the descents is to start with the linear order $(n, x) > \cdots > (1, x) > (n, y) > \cdots > (1, y)$, keep item (i) and replace conditions (ii) and (iii) above by

(ii') $1 \leq i \leq n - 1$ and $(\sigma(i), \varepsilon(i)) > (\sigma(i + 1), \varepsilon(i + 1))$.

Now let

$$(20.3) \quad B_n(X, Y, t) = \sum_{(\sigma, \varepsilon)} X^{\ell(\varepsilon|x)} Y^{\ell(\varepsilon|y)} t^{\text{des}(\sigma, \varepsilon)}$$

be the generating polynomial for the set of signed permutations of order n by the number of descents.

Theorem 20.1. *The following identity holds:*

$$(20.4) \quad \frac{(1 - t) \exp(u(t - 1)X)}{-t + \exp(u(t - 1)(X + Y))} = \sum_{n \geq 0} \frac{u^n}{n!} B_n(X, Y, t).$$

Notice that when $X = 0$ and $Yu = u$ in (20.4) we recover identity (20.1) with their combinatorial interpretation.

For the proof of Theorem 20.1 we proceed as follows. Define the statistic $\text{des}_k \sigma$ of a permutation (not a signed permutation) $\sigma = \sigma(1) \dots \sigma(n)$ to be

$$\text{des}_k \sigma := \begin{cases} \text{des } \sigma, & \text{if } 1 \leq \sigma(n) \leq n - k; \\ 1 + \text{des } \sigma, & \text{if } n - k + 1 \leq \sigma(n) \leq n; \end{cases}$$

and let

$$A_n^k(t) := \sum_{\sigma} t^{\text{des}_k \sigma} \quad (\sigma \in \mathfrak{S}_n).$$

Because of item (i) in the definition of a descent in a signed permutation we can write:

$$(20.5) \quad B_n(X, Y, t) = \sum_{k=0}^n \binom{n}{k} X^k Y^{n-k} A_n^k(t).$$

Now the “ des_k ” interpretation given for $A_n^k(t)$ provides the following recurrence relation

$$(20.6) \quad A_n^k(t) = A_n^{k-1}(t) + (t - 1)A_{n-1}^{k-1}(t) \quad (1 \leq k \leq n),$$

since $tA_{n-1}^{k-1}(t)$ is the generating polynomial for the permutations ending with $(n-k+1)$ by “ des_k ,” while $A_n^{k-1}(t) - A_{n-1}^{k-1}(t)$ is the generating polynomials for the other permutations. By iteration,

$$(20.7) \quad A_n^k(t) = A_n^0(t) + \binom{k}{1}(t-1)A_{n-1}^0(t) + \binom{k}{2}(t-1)^2A_{n-2}^0(t) \\ + \cdots + \binom{k}{k}(t-1)^kA_{n-k}^0(t).$$

Using Umbral Calculus (dear to the late professor Rota, who had made an extensive usage of that approach) $A^n \equiv A_n \equiv A_n^0(t)$ and $A^0 \equiv 1$, formula (20.7) may be rewritten as

$$(20.8) \quad A_n^k(t) = A^{n-k}((t-1) + A)^k.$$

In view of (20.5) and (20.8) we then have

$$(20.9) \quad B_n(X, Y, t) = (X(t-1) + (X+Y)A)^n, \quad A^n \equiv A_n.$$

Now exponential generating function (20.1) for the Eulerian polynomials may be rewritten as

$$\exp(uA) = \frac{1-t}{-t + \exp((t-1)u)},$$

so that

$$\sum_{n \geq 0} \frac{u^n}{n!} B_n(X, Y, t) = \sum_{n \geq 0} \frac{u^n}{n!} (X(t-1) + (X+Y)A)^n \\ = \exp(uX(t-1)) \exp(u(X+Y)A) \\ = \frac{(1-t) \exp(uX(t-1))}{-t + \exp(u(X+Y)(t-1))}. \quad \square$$

20.2. Pairs of permutations. Let $\pi = \pi(1)\pi(2)\dots\pi(n)$ and $\rho = \rho(1)\rho(2)\dots\rho(n)$ be two permutations of order n . The *number of common descents* of π and ρ , denoted by $\text{ddes}(\pi, \rho)$, is defined to be the number of integers k such that $1 \leq k \leq n-1$ and $\pi(k) > \pi(k+1)$, $\rho(k) > \rho(k+1)$. The generating polynomial for the Cartesian product $\mathfrak{S}_n \times \mathfrak{S}_n$ by “ ddes ” is denoted by

$$(20.10) \quad A_n^{\text{ddes}}(t) := \sum_{(\pi, \rho)} t^{\text{ddes}(\pi, \rho)} \quad ((\pi, \rho) \in \mathfrak{S}_n \times \mathfrak{S}_n).$$

Now, let

$$J_0(u) = \sum_{n \geq 0} (-1)^n \frac{(u/2)^{2n}}{n! n!}$$

be the *Bessel function of the first kind, of order zero* and let

$$(20.10) \quad J(u) := J_0(2\sqrt{u}) = \sum_{n \geq 0} (-1)^n \frac{u^n}{n! n!}.$$

Theorem 20.3. *The bi-exponential generating function for the polynomials $A_n^{\text{ddes}}(t)$ ($n \geq 0$) is given by:*

$$(20.11) \quad \frac{1-t}{-t + J(u(1-t))} = 1 + \sum_{n \geq 1} \frac{u^n}{n! n!} A_n^{\text{ddes}}(t).$$

Notice that we go from the fraction in (20.1) to the fraction in (20.11) by merely replacing “exp” by “ J .” The proof of Theorem 20.3 is quite similar to the proof of Theorem 10.1 that gives the generating function for the polynomials ${}^{\text{inv}}A_n(t, q)$. The fraction in identity (20.11) can be rewritten as

$$\left(1 + \sum_{n \geq 1} (-1)^n (1-t)^{n-1} \frac{u^n}{n! n!} \right)^{-1},$$

so that (20.11) is equivalent to

$$\left(1 - \sum_{n \geq 1} (-1)^{n-1} (1-t)^{n-1} \frac{u^n}{n! n!} \right) \cdot \sum_{n \geq 0} A_n^{\text{ddes}}(t) \frac{u^n}{n! n!} = 1$$

and provides the recurrence $A_0^{\text{ddes}}(t) = 1$ and

$$(20.12) \quad A_n^{\text{ddes}}(t) = \sum_{1 \leq k \leq n} \binom{n}{k} \binom{n}{k} (t-1)^{k-1} A_{n-k}^{\text{ddes}}(t) \quad (n \geq 1).$$

Let F_k be the generating polynomial for the pairs of permutations (π, ρ) of order n , such that both π and ρ have their *longest decreasing rightmost factors (l.d.r.f.)* equal to k by “ddes.” Also, let $G_k := F_k + F_{k+1} + \cdots + F_n$. Under those conditions we have

$$\binom{n}{k} \binom{n}{k} t^k A_{n-k}^{\text{ddes}}(t) = tF_k + G_{k+1},$$

so that

$$\begin{aligned}
 \sum_{1 \leq k \leq n} \binom{n}{k} \binom{n}{k} (t-1)^{k-1} A_{n-k}^{\text{ddes}}(t) &= \sum_{k=1}^n \left(\frac{F_k}{t^{k-1}} + \frac{G_{k+1}}{t^k} \right) (t-1)^{k-1} \\
 &= \sum_{k=1}^n \frac{1}{t^k} \left(t(G_k - G_{k+1}) + G_{k+1} \right) (t-1)^{k-1} \\
 &= \sum_{k=1}^n \frac{1}{t^k} \left(tG_k - (t-1)G_{k+1} \right) (t-1)^{k-1} \\
 &= \sum_{k=1}^n G_k \frac{(t-1)^{k-1}}{t^{k-1}} - G_{k+1} \frac{(t-1)^k}{t^k} \\
 &= G_1 = A_n^{\text{ddes}}(t),
 \end{aligned}$$

by definition of the G_k 's. \square

Remark. The proof of the previous theorem shows that if instead of a pair of permutations we consider a finite sequence of permutations, we can also calculate the generating function for those finite sequences by their common descent. This is one of the extensions that will be proposed in the next section.

Let $A_{n,k}^{\text{ddes}} := |\{(\pi, \rho) \in \mathfrak{S}_n \times \mathfrak{S}_n : \text{ddes}(\pi, \rho) = k\}|$. The first values of the coefficients $A_{n,k}^{\text{ddes}}$, calculated by means of the recurrence (20.12), are shown in Table 20.2.

$k =$	0	1	2	3	4
$n = 1$	1				
2	3	1			
3	19	16	1		
4	211	299	65	1	
5	3651	7346	3156	246	1

Table 20.2 (distribution of ddes on $\mathfrak{S}_n \times \mathfrak{S}_n$)

20.3. *The q -extension.* The q -extension of (20.1) was already done in Theorem 10.1. Recall that when the exponential is replaced by the q -exponential E_q , the expansion of the fraction involves the generating polynomials for \mathfrak{S}_n by the bi-statistic $(1 + \text{des}, \text{inv})$. We will also consider the replacement of the exponential by the other q -exponential e_q . The formulas obtained together with their combinatorial interpretations will be discussed in the next section as specializations of more general results.

Referring to the diagram represented in Fig. 20.1 the three extensions $(\text{sgn}, \cdot, \cdot)$ [signed permutations], (\cdot, \cdot, J) [pairs of permutations] and (\cdot, q, \cdot) [the q -extension] have been described.

20.4. *The t, q -maj extension for signed permutations.* In § 20.1 the notion of *descent* for a signed permutation has been introduced, so that the *major index* can also be defined by adding the *positions* at which a descent occurs. In the example shown in § 20.1 the descents occur at positions 1, 2, 5, 6, 9 so that $\text{maj}(\sigma, \varepsilon) = 1 + 2 + 5 + 6 + 9 = 23$.

Now form the generating polynomial for the set of signed permutations of order n by the pair (des, maj) , that is,

$$(20.13) \quad B_n(X, Y, t, q) = \sum_{(\sigma, \varepsilon)} X^{\ell(\varepsilon|x)} Y^{\ell(\varepsilon|y)} t^{\text{des}(\sigma, \varepsilon)} q^{\text{maj}(\sigma, \varepsilon)}.$$

As we now see, the generating function for those polynomials $B_n(X, Y, t, q)$ ($n \geq 0$) can be derived, by *q-analogizing* the calculation made in § 20.1 for the polynomial $B_n(X, Y, t)$. For $0 \leq k \leq n$ and each (ordinary) permutation σ let

$$(20.14) \quad \begin{aligned} \text{des}_k \sigma &:= \begin{cases} \text{des } \sigma, & \text{if } 1 \leq \sigma(n) \leq n - k; \\ 1 + \text{des } \sigma, & \text{if } n - k + 1 \leq \sigma(n) \leq n; \end{cases} \\ \text{maj}_k \sigma &:= \text{maj } \sigma + n \chi(n - k + 1 \leq \sigma(n) \leq n); \end{aligned}$$

and let

$$A_n^k(t, q) := \sum_{\sigma} t^{\text{des}_k \sigma} q^{\text{maj}_k \sigma} \quad (\sigma \in \mathfrak{S}_n).$$

If I is a subset of $[n] = \{1, 2, \dots, n\}$ let $B_I(t, q)$ denote the generating polynomial for the signed permutations $(\sigma, \varepsilon) = (\sigma(1), \varepsilon(1)) \dots (\sigma(n), \varepsilon(n))$ containing all the letters (i, x) when $i \in I$ and all the letters (j, y) when $j \in [n] \setminus I$. Because of the definition of a descent for a signed permutation, we have

$$(20.15) \quad B_I(t, q) = A_n^k(t, q) \quad \text{if } \#I = k.$$

Hence

$$B_n(X, Y, t, q) = \sum_{k=0}^n \binom{n}{k} X^k Y^{n-k} A_n^k(t, q).$$

Mimicking (20.6) we also have:

$$A_n^k(t, q) = A_n^{k-1}(t, q) + (tq^n - 1)A_{n-1}^{k-1}(t, q) \quad (1 \leq k \leq n).$$

By iteration,

$$A_n^k(t, q) = \sum_{s=0}^k \binom{k}{s} (tq^n - 1)(tq^{n-1} - 1) \dots (tq^{n-s+1} - 1) B_{n-s}^0(t, q).$$

Hence

$$\begin{aligned} A_n(X, Y; t, q) &= \sum_{0 \leq s \leq k \leq n} \frac{n!}{(n-k)! s! (k-s)!} X^k Y^{n-k} \\ &\quad \times (tq^n - 1)(tq^{n-1} - 1) \dots (tq^{n-s+1} - 1) B_{n-s}^0(t, q) \end{aligned}$$

and

$$\begin{aligned} \sum_{n \geq 0} \frac{u^n}{n!} \frac{B_n(X, Y; t, q)}{(t; q)_{n+1}} &= \sum_{s, n, m \geq 0} \frac{u^{s+n+m} X^{s+n} Y^m}{s! n! m!} \frac{(-1)^s}{(t; q)_{n+m+1}} B_{n+m}^0(t, q) \\ &= \exp(-uX) \sum_{r \geq 0} \frac{(u(X+Y))^r}{r!} \frac{A_r^0(t, q)}{(t; q)_{r+1}}. \end{aligned}$$

But $A_s^0(t, q)$ is the generating polynomial by $(\text{des}_0, \text{maj}_0)$, that is, the q -maj Eulerian polynomial ${}^{\text{maj}}A_s(t, q)$, introduced in Definition 10.1. As, by (10.2b)

$$\sum_{r \geq 0} \frac{u^r}{r!} \frac{A_r^0(t, q)}{(t; q)_{r+1}} = \sum_{s \geq 0} t^s \exp(u[s+1]_q),$$

we conclude that

$$\begin{aligned} \sum_{n \geq 0} \frac{u^n}{n!} \frac{B_n(X, Y; t, q)}{(t; q)_{n+1}} &= \exp(-uX) \sum_{s \geq 0} t^s \exp(u(X+Y)[s+1]_q) \\ &= \sum_{n \geq 0} t^s \exp(u(Xq[s]_q + Y[s+1]_q)). \end{aligned}$$

Proposition 20.4. *Let $B_n(X, Y; t, q)$ be the generating polynomial for the signed permutations by (des, maj) , as defined in (20.13). Then*

$$(20.16) \quad \frac{B_n(X, Y; t, q)}{(t; q)_{n+1}} = \sum_{s \geq 0} t^s (Xq[s]_q + Y[s+1]_q)^n;$$

$$(20.17) \quad \sum_{n \geq 0} u^n \frac{B_n(X, Y; t, q)}{(t; q)_{n+1}} = \sum_{n \geq 0} t^s \frac{1}{1 - u(Xq[s]_q + Y[s+1]_q)};$$

$$(20.18) \quad \sum_{n \geq 0} \frac{u^n}{n!} \frac{B_n(X, Y; t, q)}{(t; q)_{n+1}} = \sum_{n \geq 0} t^s \exp(u(Xq[s]_q + Y[s+1]_q)).$$

For each subset $I \subset [n]$ let $B_I(t, q)$ be the generating polynomial for the signed permutations containing all the letters (i, x) with $i \in I$ and all the letters (j, y) with $j \in [n] \setminus I$ by (des, maj) . If $\#I = k$, we have:

$$(20.19) \quad \frac{B_I(t, q)}{(t; q)_{n+1}} = \sum_{s \geq 0} t^s q^k [s]_q^k [s+1]_q^{n-k}.$$

Proof. Identity (20.18) has just been proved. The other formulas can then be easily deduced. \square

20.5. *A first inversion number for signed permutations.* As explained in the next section, the inversion number for signed permutations that will be further studied will be motivated by analytical reasons. However, we can also look for an inversion number, that reduces to the usual “inv” for the ordinary permutations, that has the further property of being equidistributed with “maj” on the set of signed permutations.

Keep the notations of Proposition 20.4 and let SP_I be the set of signed permutations of order n containing all the letters (i, x) with $i \in I$ and all the letters (j, y) with $j \in [n] \setminus I$. By letting $t = 1$ in (20.19) we get

$$B_I(1, q) = q^k \frac{(q; q)_n}{(1 - q)^n}.$$

The most (?) natural inversion number, say, “Inv”, whose distribution on the set SP_I is given by $B_I(1, q)$, can be defined by mapping each $(\sigma, \varepsilon) \in SP_I$ onto

$$\text{Inv}(\sigma, \varepsilon) := \text{inv } \sigma + \# I.$$

In the notations of § 20.1

$$(20.20) \quad \text{Inv}(\sigma, \varepsilon) := \text{inv } \sigma + \ell(\varepsilon|x).$$

To avoid any confusion, denote the major index of a signed permutation w by $\text{Maj } w$. To construct a bijection Ψ having the property that

$$(20.21) \quad \text{Maj}(\sigma, \varepsilon) = \text{Inv } \Psi(\sigma, \varepsilon),$$

proceed as follows. It suffices to do so for each subset SP_I such that $I = \{n - k + 1, n - k + 2, \dots, n\}$ ($0 \leq k \leq n$). Each signed permutation in such an SP_I is then a permutation of the *increasing* word

$$(1, y)(2, y) \dots (n - k, y)(n - k + 1, x) \dots (n, x),$$

Introduce a new letter $*$ and impose the ordering

$$(n - k, y) < * < (n - k + 1, x).$$

Consider the alphabet $\{(1, y), \dots, (n - k, y), *, (n - k + 1, x), \dots, (n, x)\}$ and let w belong to SP_I . The transformation Φ , introduced in Theorem 11.3, maps $w*$ (the signed permutation w , linearly written as a permutation, with the juxtaposition of the letter $*$ at the end) onto

$$(20.22) \quad \Phi(w*) = w'*,$$

where w' is a rearrangement of w . Furthermore,

$$(20.23) \quad \text{maj}(w*) = \text{inv } \Phi(w*) = \text{inv}(w'*).$$

The bijection Ψ of SP_I onto itself is then defined by

$$(20.24) \quad \Psi(w) * := \Phi(w *) = w' * .$$

Theorem 20.5. *The mapping Ψ defined in (20.24) is a bijection of each class SP_I onto itself such that $\text{Maj } w = \text{Inv } \Psi(w)$ holds for every $w \in SP_I$.*

Proof. As Φ is a bijection of each rearrangement class onto itself and because of (20.22), the mapping Ψ is itself a bijection. Furthermore, let $w = x_1 x_2 \dots x_n \in SP_I$. Then $\text{Maj } w = \text{maj } w + n\chi(x_n \in I) = \text{maj}(w *) = \text{inv } \Phi(w *) = \text{inv}(\Psi(w) *) = \text{Inv } \Psi(w)$. \square

Example. Let $n = 6$, $I = \{4, 5, 6\}$ and consider the signed permutation $w = (3, y)(1, y)(6, x)(5, x)(2, y)(4, x)$, that will be rewritten as $w = 3, 1, \mathbf{6}, \mathbf{5}, 2, \mathbf{4}$. With $3 < * < \mathbf{4}$ the image of $w *$ under Φ (defined in § 11.3) is $\Phi(w *) = \mathbf{6}, 3, \mathbf{5}, 1, \mathbf{4}, 2, *$, so that $\Psi(w) = \mathbf{6}, 3, \mathbf{5}, 1, \mathbf{4}, 2$. We verify that $\text{Maj } w = 1 + 3 + 4 + 6 = 14$. Moreover $\Psi(w)$ has 11 inversions and $\#I = 3$, so that $\text{Inv } \Psi(w) = 11 + 3 = 14$.

21. Eulerian Calculus; the analytic choice

When the symmetric group \mathfrak{S}_n is regarded as a Coxeter group generated by the transpositions $(i, i + 1)$ ($1 \leq i \leq n - 1$), the length $l_{\text{Cox}}(\sigma)$ of a permutation σ is simply the number of inversions $\text{inv } \sigma$. As recalled in the previous subsection 20.3, the generating function for the bi-statistic $(1 + \text{des}, l_{\text{Cox}}) = (1 + \text{des}, \text{inv})$ on the symmetric groups $A_n = \mathfrak{S}_n$ has been derived by taking a suitable q -analog of the generating function for the single statistic “des.”

As we already have an expression for the generating function for “des” on the groups B_n , namely (20.4), the natural question arises: can we find an analogous approach to obtain the generating function for $(\text{des}, l_{\text{Cox}})$ on the groups B_n ? The answer is yes (see Exercise 30), but the formula we can derive does not fit any longer in the set-up of the q -series; its analytic manipulation is cumbersome.

Accordingly, if we wish to remain in the algebra of the q -series and obtain the desired extension for the signed permutations, identity (20.2) must play the role played by identity (20.1) for the plain permutations. In the first place it is necessary to find a q -analog for (20.2), with an interesting combinatorial interpretation.

We definitely adopt that approach of *analytical* nature. The statistic “inv” for the signed permutations will directly come out of that interpretation.

21.1. *Inversions for signed permutations.* A good choice for a q -analog of formula (20.2) is to replace the product of the two exponentials

in the denominator by a product of two q -exponentials (and not by $e_q((t-1)(X+Y))$ that would give a q -expansion with some negative terms.) In the q -series expansion

$$(21.1) \quad \frac{(1-t)e_q((t-1)X)}{-t + e_q((t-1)X)e_q((t-1)Y)} = \sum_{n \geq 0} \frac{1}{(q; q)_n} B_n(X, Y, t, q)$$

the coefficients $B_n(X, Y, t, q)$ are polynomials with *nonnegative integral* coefficients. One of the consequences of the main result proved in this section is to show that each $B_n(X, Y, t, q)$ is the generating polynomial for the signed permutations of order n by a bi-statistic “(des, coinv)”:

$$(21.2) \quad B_n(X, Y, t, q) = \sum_{(\sigma, \varepsilon)} X^{\ell(\varepsilon|x)} Y^{\ell(\varepsilon|y)} t^{\text{des}(\sigma, \varepsilon)} q^{\text{coinv}(\sigma, \varepsilon)}.$$

Of course, “des(σ, ε)” is the number of descents, as was defined in subsection 20.2; and “coinv(σ, ε)” is the *number of co-inversions* in the signed permutation, defined as follows.

A pair of integers (i, j) is said to be an *inversion* (resp. a *co-inversion*) of the signed permutation (σ, ε) of order n , if one of the following conditions is verified

- (i) $\varepsilon(i) = \varepsilon(j)$, $i < j$; and $\sigma(i) > \sigma(j)$ (resp. $\sigma(i) < \sigma(j)$);
- (ii) $\varepsilon(i) = y$, $\varepsilon(j) = x$ and $\sigma(i) > \sigma(j)$.

Let $\text{inv}(\sigma, \varepsilon)$ (resp. $\text{coinv}(\sigma, \varepsilon)$) denote the *number of inversions* (resp. the *number of co-inversions*) of (σ, ε) .

Take up again the example of section 20.2. We have:

$$\begin{aligned} \text{inv} \begin{pmatrix} 6 & 7 & 2 & 4 & 3 & 1 & 8 & 5 & 9 \\ x & y & y & y & x & x & y & x & x \end{pmatrix} &= 4 + 2 + 11 = 17; \\ \text{coinv} \begin{pmatrix} 6 & 7 & 2 & 4 & 3 & 1 & 8 & 5 & 9 \\ x & y & y & y & x & x & y & x & x \end{pmatrix} &= 6 + 4 + 11 = 21. \end{aligned}$$

The first values of the polynomials $B_n(X, Y, t, q)$, i.e., the generating polynomials for the signed permutations by the pair (des, coinv), are shown in Table 21.1.

$$\begin{aligned} B_1 &= tX + Y; & B_2 &= t(t+q)X^2 + 2t(1+q)XY + (t+q)Y^2; \\ B_3 &= (2tq + 2tq^2 + t^2 + q^3)Y^3 + t(1+q+q^2)(3q+2t+1)XY^2 \\ &\quad + t(1+q+q^2)(2q+ tq + 3t)X^2Y + t(2tq + 2tq^2 + t^2 + q^3)X^3. \end{aligned}$$

Table 21.1 (distribution of (des, coinv))

Another consequence of our main result is to show that if in (21.1) each q -exponential $e_q(u)$ is replaced by the *second* Q -exponential $E_Q(u)$, the polynomials $B_n(X, Y, t, Q)$ on the right-hand side are the generating polynomials for the signed permutations by the pair (des, inv). The first values of those polynomials are shown in Table 21.2.

$$\begin{aligned} B_1 &= Y + tX; & B_2 &= (1+tQ)Y^2 + 2t(1+Q)XY + t(1+tQ)X^2; \\ B_3 &= (1+2tQ+2tQ^2+t^2Q^3)Y^3 + t(1+Q+Q^2)(3+Q+2tQ)XY^2 \\ &\quad + t(1+Q+Q^2)(3+Q+2tQ)X^2Y + t(1+2tQ+2tQ^2+t^2Q^3)X^3. \end{aligned}$$

Table 21.2 (distribution of (des, inv))

When restricted to the elements of \mathfrak{S}_n , the statistic “inv” for B_n is nothing but the usual number of inversions for the permutations. Likewise, formula (21.1) with $X = 0$ specializes into the analog of (10.12) for e_q .

Thus, (21.1) with its combinatorial interpretation (21.2) is our q -extension of (20.4). In our diagram of Fig. 20.1 that extension sits on vertex (sgn, q, \cdot).

21.2. Basic Bessel Functions. The next step is to study the expansion of (21.1), when the q -exponentials are replaced by q -analogs of Bessel functions. Formally, the left-hand side of (21.1) is to be replaced by the fraction

$$(21.3) \quad \frac{(1-t) \mathbf{J}((1-t)X; \mathbf{Q}, \mathbf{q})}{-t + \mathbf{J}((1-t)X; \mathbf{Q}, \mathbf{q}) \mathbf{J}((1-t)Y; \mathbf{Q}, \mathbf{q})},$$

where \mathbf{J} is a basic Bessel function defined below. Our problem will be to study the expansion of that fraction in the algebra of series in one or several bases.

Let L, l be two fixed nonnegative integers and $\mathbf{Q} = (Q_1, Q_2, \dots, Q_L)$, $\mathbf{q} = (q_1, q_2, \dots, q_l)$ two sequences of variables. Define

$$\begin{aligned} \mathbf{Q}^{\binom{n}{2}} &:= Q_1^{\binom{n}{2}} \dots Q_L^{\binom{n}{2}}, \\ (\mathbf{Q}; \mathbf{Q})_n &:= (Q_1; Q_1)_n \dots (Q_L; Q_L)_n, \\ (\mathbf{q}; \mathbf{q})_n &:= (q_1; q_1)_n \dots (q_l; q_l)_n. \end{aligned}$$

Using the H-notation for the *Hadamard product* of two series, that is,

$$\left(\sum_{n \geq 0} \alpha_n u^n \right) \text{H} \left(\sum_{n \geq 0} \beta_n u^n \right) = \sum_{n \geq 0} (\alpha_n \beta_n) u^n,$$

we define *the several-basis Bessel function* by

$$(21.4) \quad \mathbf{J}(u; \mathbf{Q}, \mathbf{q}) := \left(\sum_{n \geq 0} (-u)^n \right) \text{H} E_{Q_1}(u) \text{H} E_{Q_2}(u) \text{H} \cdots \\ \text{H} e_{q_1}(u) \text{H} e_{q_2}(u) \text{H} \cdots$$

We can also write:

$$(21.5) \quad \mathbf{J}(u; \mathbf{Q}, \mathbf{q}) := \sum_{n \geq 0} (-1)^n \frac{Q \binom{n}{2}}{(Q; Q)_n} \frac{1}{(q; q)_n} u^n.$$

If $L = l = 1$, then

$$\mathbf{J}(u; Q, q) = \sum_{n \geq 0} (-1)^n \frac{Q \binom{n}{2}}{(Q; Q)_n} \frac{1}{(q; q)_n} u^n.$$

Finally, for $L = 0$ and $l = 1$, we regain the q -exponential

$$\mathbf{J}(u; -, q) = \sum_{n \geq 0} (-1)^n \frac{1}{(q; q)_n} u^n = e_q(-u),$$

and for $L = 1$ and $l = 0$, the Q -exponential

$$\mathbf{J}(u; Q, -) = \sum_{n \geq 0} (-1)^n \frac{Q \binom{n}{2}}{(Q; Q)_n} u^n = E_Q(-u).$$

The coefficients in the series expansion of the fraction (21.3) will be shown to be polynomials with positive integral coefficients. They actually are generating polynomials for combinatorial objects, called *signed multipermutations* by a multivariable statistic that includes a single statistic referred to as the *number of descents*.

Definition. Each triple $(\underline{\Sigma}, \underline{\sigma}, \varepsilon)$ is called a *signed multipermutation* of order n if $\underline{\Sigma} = (\Sigma_1, \dots, \Sigma_L)$ and $\underline{\sigma} = (\sigma_1, \dots, \sigma_l)$ are two sequences of L and l permutations of order n , respectively, and if ε is an xy -word of length n .

Definition. Each integer i is said to be a *descent* of the signed multipermutation $(\underline{\Sigma}, \underline{\sigma}, \varepsilon)$, if one of the following three conditions is satisfied:

- (i) $i = n$ and $\varepsilon(n) = x$;
- (ii) $1 \leq i \leq n - 1$, $\varepsilon(i) = x$, $\varepsilon(i + 1) = y$;
- (iii) $1 \leq i \leq n - 1$, $\varepsilon(i) = \varepsilon(i + 1)$ and $\Sigma_1(i) > \Sigma_1(i + 1), \dots, \Sigma_L(i) > \Sigma_L(i + 1)$, and also $\sigma_1(i) > \sigma_1(i + 1), \dots, \sigma_l(i) > \sigma_l(i + 1)$;

Let $\text{ddes}(\underline{\Sigma}, \underline{\sigma}, \varepsilon)$ denote the *number of descents* of $(\underline{\Sigma}, \underline{\sigma}, \varepsilon)$.

The inversion and co-inversion numbers “inv” and “coinv” of a signed permutation (σ, ε) have been defined in § 21.1. With a signed multipermutation $(\underline{\Sigma}, \underline{\sigma}, \varepsilon)$ we further use the notations:

$$\begin{aligned} \mathbf{Q}^{\text{inv}(\underline{\Sigma}, \varepsilon)} &= Q_1^{\text{inv}(\Sigma_1, \varepsilon)} \dots Q_L^{\text{inv}(\Sigma_L, \varepsilon)}; \\ \mathbf{q}^{\text{coinv}(\underline{\sigma}, \varepsilon)} &= q_1^{\text{coinv}(\sigma_1, \varepsilon)} \dots q_l^{\text{coinv}(\sigma_l, \varepsilon)}. \end{aligned}$$

Let (σ, ε) be a signed permutation, where $\sigma \in \mathcal{S}_n$ and $\varepsilon = \varepsilon(1) \dots \varepsilon(n)$ is an xy -word. Recall that the word $\sigma_{\varepsilon|x}$ (resp. $\sigma_{\varepsilon|y}$) is defined to be the subword of $\sigma(1) \dots \sigma(n)$ made of all letters $\sigma(i)$ such that $\varepsilon(i) = x$ (resp. $\varepsilon(i) = y$). Introduce the number of inversions “inv($\sigma_{\varepsilon|y}, \sigma_{\varepsilon|x}$)” between the words $\sigma_{\varepsilon|y}$ and $\sigma_{\varepsilon|x}$ by

$$\text{inv}(\sigma_{\varepsilon|y}, \sigma_{\varepsilon|x}) = \#\{(i, j) : \varepsilon(i) = y, \varepsilon(j) = x, \sigma(i) > \sigma(j)\}.$$

Another way of defining the statistics “inv” and “coinv” is then to let

$$\begin{aligned} \text{inv}(\sigma, \varepsilon) &= \text{inv} \sigma_{\varepsilon|x} + \text{inv} \sigma_{\varepsilon|y} + \text{inv}(\sigma_{\varepsilon|y}, \sigma_{\varepsilon|x}); \\ \text{coinv}(\sigma, \varepsilon) &= \text{coinv} \sigma_{\varepsilon|x} + \text{coinv} \sigma_{\varepsilon|y} + \text{inv}(\sigma_{\varepsilon|y}, \sigma_{\varepsilon|x}). \end{aligned}$$

Definition. Suppose that ε has α letters equal to x and β letters equal to y ($\alpha + \beta = n$). We say that the permutation σ is *compatible with ε* (also that the signed permutation (σ, ε) is *compatible*), if $\text{inv}(\sigma_{\varepsilon|y}, \sigma_{\varepsilon|x}) = 0$, or, in an equivalent manner, if the subword $\sigma_{\varepsilon|y}$ is a rearrangement of the word $12 \dots \beta$ (and then $\sigma_{\varepsilon|x}$ is a rearrangement of $(\beta + 1)(\beta + 2) \dots n$ (of length α)).

A signed multipermutation $(\underline{\Sigma}, \underline{\sigma}, \varepsilon)$ such that $\underline{\Sigma} = (\Sigma_1, \dots, \Sigma_L)$ and $\underline{\sigma} = (\sigma_1, \dots, \sigma_l)$ is said to be *compatible*, if the permutations $\Sigma_1, \dots, \Sigma_L, \sigma_1, \dots, \sigma_l$ are all compatible with ε .

Let

$$(21.6) \quad B_{\alpha, \beta}(t, \mathbf{Q}, \mathbf{q}) := \sum_{\substack{(\underline{\Sigma}, \underline{\sigma}, \varepsilon) \text{ (comp.)} \\ \ell(\varepsilon|x) = \alpha, \ell(\varepsilon|y) = \beta}} t^{\text{ddes}(\underline{\Sigma}, \underline{\sigma}, \varepsilon)} \mathbf{Q}^{\text{inv}(\underline{\Sigma}, \varepsilon)} \mathbf{q}^{\text{coinv}(\underline{\sigma}, \varepsilon)}$$

be the generating polynomial for the *compatible* signed multipermutations $(\underline{\Sigma}, \underline{\sigma}, \varepsilon)$ such that ε contains α letters equal to x and β letters equal to y , by the statistic (ddes, inv, coinv). Also let

$$(21.7) \quad \begin{aligned} B_n(X, Y, t, \mathbf{Q}, \mathbf{q}) \\ := \sum_{(\underline{\Sigma}, \underline{\sigma}, \varepsilon)} X^{\ell(\varepsilon|x)} Y^{\ell(\varepsilon|y)} t^{\text{ddes}(\underline{\Sigma}, \underline{\sigma}, \varepsilon)} \mathbf{Q}^{\text{inv}(\underline{\Sigma}, \varepsilon)} \mathbf{q}^{\text{coinv}(\underline{\sigma}, \varepsilon)}, \end{aligned}$$

the sum being over all signed multipermutations of length n .

Proposition 21.1. *We have the identity*

$$(21.8) \quad B_n(X, Y, t, \mathbf{Q}, \mathbf{q}) = \sum_{\alpha+\beta=n} \begin{bmatrix} n \\ \alpha \end{bmatrix}_{\mathbf{Q}} \begin{bmatrix} n \\ \alpha \end{bmatrix}_{\mathbf{q}} X^\alpha Y^\beta B_{\alpha,\beta}(t, \mathbf{Q}, \mathbf{q})$$

Proof. To avoid cumbersome notations assume that $L = l = 1$. Let $(\Sigma, \sigma, \varepsilon)$ be a signed bipermutation of length n such that $\ell(\varepsilon|x) = \alpha$. Let I denote the set of the α letters of the word $\sigma_{\varepsilon|x}$. There exists a unique increasing bijection f_Σ (resp. f_σ) mapping the set $\Sigma(I)$ (resp. $\sigma(I)$) onto the interval $[n - \alpha + 1, n]$ and the set $\Sigma([n] \setminus I)$ (resp. $\sigma([n] \setminus I)$) onto the interval $[1, n - \alpha]$. The triple $(f_\Sigma \circ \Sigma, f_\sigma \circ \sigma, \varepsilon)$ is then a compatible signed permutation. The mapping $(\Sigma, \sigma, \varepsilon) \mapsto (f_\Sigma \circ \Sigma, f_\sigma \circ \sigma, \varepsilon)$ is *surjective*. Moreover, if $(\Sigma_0, \sigma_0, \varepsilon)$ is compatible, the number of preimages $(\Sigma, \sigma, \varepsilon)$ mapped over $(\Sigma_0, \sigma_0, \varepsilon)$ by that surjection is equal to $\binom{n}{\alpha} \binom{n}{\alpha}$.

If $(\Sigma_0, \sigma_0, \varepsilon)$ is compatible and if $(f_\Sigma \circ \Sigma, f_\sigma \circ \sigma, \varepsilon) = (\Sigma_0, \sigma_0, \varepsilon)$, then

$$\begin{aligned} \text{ddes}(\Sigma, \sigma, \varepsilon) &= \text{ddes}(\Sigma_0, \sigma_0, \varepsilon); \\ \text{inv}(\sigma, \varepsilon) &= \text{inv}(\sigma_0, \varepsilon) + \text{inv}(\sigma_{\varepsilon|y}, \sigma_{\varepsilon|x}); \\ \text{coinv}(\sigma, \varepsilon) &= \text{coinv}(\sigma_0, \varepsilon) + \text{inv}(\sigma_{\varepsilon|y}, \sigma_{\varepsilon|x}), \end{aligned}$$

since f_Σ and f_σ are increasing mappings.

Therefore, the sum $\sum_{(\Sigma, \sigma, \varepsilon)} Q^{\text{inv}(\Sigma, \varepsilon)} q^{\text{coinv}(\sigma, \varepsilon)}$ over all triples $(\Sigma, \sigma, \varepsilon)$ such that $(f_\Sigma \circ \Sigma, f_\sigma \circ \sigma, \varepsilon) = (\Sigma_0, \sigma_0, \varepsilon)$, is equal to

$$\begin{aligned} Q^{\text{inv}(\Sigma_0, \varepsilon)} q^{\text{coinv}(\sigma_0, \varepsilon)} \sum_{(\Sigma, \sigma, \varepsilon)} Q^{\text{inv}(\Sigma_{\varepsilon|y}, \Sigma_{\varepsilon|x})} q^{\text{inv}(\sigma_{\varepsilon|y}, \sigma_{\varepsilon|x})} \\ = Q^{\text{inv}(\Sigma_0, \varepsilon)} q^{\text{coinv}(\sigma_0, \varepsilon)} \begin{bmatrix} n \\ \alpha \end{bmatrix}_{\mathbf{Q}} \begin{bmatrix} n \\ \alpha \end{bmatrix}_{\mathbf{q}}. \end{aligned}$$

Hence,

$$\begin{aligned} B_n(X, Y, t, \mathbf{Q}, \mathbf{q}) &= \sum_{\alpha+\beta=n} X^\alpha Y^\beta \sum_{\substack{(\Sigma_0, \sigma_0, \varepsilon) \text{ (comp.)} \\ \ell(\varepsilon|x)=\alpha}} t^{\text{ddes}(\Sigma_0, \sigma_0, \varepsilon)} \sum_{\substack{(\Sigma, \sigma, \varepsilon) \\ (f_\Sigma \circ \Sigma, f_\sigma \circ \sigma, \varepsilon) = (\Sigma_0, \sigma_0, \varepsilon)}} Q^{\text{inv}(\Sigma, \varepsilon)} q^{\text{coinv}(\sigma, \varepsilon)} \\ &= \sum_{0 \leq \alpha \leq n} X^\alpha Y^{n-\alpha} \begin{bmatrix} n \\ \alpha \end{bmatrix}_{\mathbf{Q}} \begin{bmatrix} n \\ \alpha \end{bmatrix}_{\mathbf{q}} B_{\alpha,\beta}(t, \mathbf{Q}, \mathbf{q}). \quad \square \end{aligned}$$

The main result of this section that will be proved in the next subsection is the following theorem.

Theorem 21.2. *The following identities hold:*

$$\begin{aligned}
 (21.9) \quad & \frac{(1-t) \mathbf{J}((1-t)X; \mathbf{Q}, \mathbf{q})}{-t + \mathbf{J}((1-t)X; \mathbf{Q}, \mathbf{q}) \mathbf{J}((1-t)Y; \mathbf{Q}, \mathbf{q})} \\
 &= \sum_{n \geq 0} \frac{1}{(\mathbf{Q}; \mathbf{Q})_n (\mathbf{q}; \mathbf{q})_n} B_n(X, Y, t, \mathbf{Q}, \mathbf{q}) \\
 &= \sum_{\alpha, \beta \geq 0} \frac{X^\alpha Y^\beta}{(\mathbf{Q}; \mathbf{Q})_\alpha (\mathbf{q}; \mathbf{q})_\alpha (\mathbf{Q}; \mathbf{Q})_\beta (\mathbf{q}; \mathbf{q})_\beta} B_{\alpha, \beta}(t, \mathbf{Q}, \mathbf{q}).
 \end{aligned}$$

The list of the first values of the polynomials $B_{\alpha, \beta}(t, Q, q)$ is shown in Table 21.3. Remember that to obtain the polynomials $B_n(X, Y, t, Q, q)$ (for $L = l = 1$) identity (21.8) is to be used.

$$\begin{aligned}
 B_{0,1} &= 1; & B_{1,0} &= t; \\
 B_{0,2} &= Qq + q + tQ + 1; & B_{1,1} &= 2t; & B_{2,0} &= t(Qq + q + tQ + 1); \\
 B_{0,3} &= 2Qq + 2Q^2q + 2q + 2Q^3tq + 2Q^3tq^2 + 2Q^2q^3 + 2q^2 \\
 &\quad + q^3 + 1 + 2Qq^3 + 2tQq + 2tQ^2q + 2q^2tQ + 2q^2tQ^2 \\
 &\quad + 2Qq^2 + Q^3t^2 + 2tQ + 2tQ^2 + 2Q^2q^2 + Q^3q^3 \\
 B_{1,2} &= t(3Qq + 3q + 2tQ + Q + 3) \\
 B_{2,1} &= t(2Qq + tQq + 2q + tq + 3tQ + 2 + t) \\
 B_{3,0} &= tB_{0,3}
 \end{aligned}$$

Table 21.3 (distribution of (ddes, inv, coinu) over the compatible signed bipermutations)

The list of the first values of the polynomials $B_n(X, Y, t, Q, q)$ (for $L = l = 1$) is shown in Table 21.3. If in Table 21.3 we let $q = 0$ (resp. $Q = 0$), we recover Table 21.1 (resp. Table 21.2). On the other hand, the table contains the first values of the Gaussian polynomials $1+Q$, $1+Q+Q^2$ and $1+q$, $1+q+Q^2$. That occurrence will be fully explained in the next subsection.

Back to the diagram in Fig. 20.1 we see that Theorem 21.2 refers to extension (sgn, q, J) and the distribution of (ddes, inv, coinu) on signed multipermutations. When $X = 0$ in (21.9), we go from signed multipermutations to the multipermutations themselves. This corresponds to vertex (\cdot, q, J) . For $L = 0, l = 1$ we have the specialization (sgn, q, \cdot) , corresponding to identity (21.1) with a combinatorial interpretation in terms of number of descents and co-inversions. Finally, vertex (sgn, \cdot, J) may be regarded as the specialization with all the variables Q_i and q_i equal to 1.

Theorem 21.2 provides a combinatorial interpretation in terms of signed multipermutations counted only by their numbers of descents.

21.3. *The iterative method.* To prove Theorem 21.2 we proceed as follows. Let $(Q)_n := (Q; Q)_n$, $(q)_n := (q; q)_n$ and again assume that $L = l = 1$. Identity (21.9) (in its second form) may be rewritten

$$\begin{aligned} \left(1 - \sum_{i+j \geq 1} \frac{(t-1)^{i+j-1} Q^{\binom{i}{2} + \binom{j}{2}} X^i Y^j}{(Q)_i (q)_i (Q)_j (q)_j}\right)^{-1} \mathbf{J}((1-t)X; Q, q) \\ = \sum_{\alpha, \beta \geq 0} \frac{X^\alpha Y^\beta}{(Q)_\alpha (q)_\alpha (Q)_\beta (q)_\beta} B_{\alpha, \beta}(t, Q, q), \end{aligned}$$

an identity of the form $(1-R)^{-1}S = T$. Express it in the form $S = (1-R)T$ and look for the coefficients of $X^\alpha Y^\beta$ on both sides. Then the identity is equivalent to the set of the two recurrence formulas

$$(21.10) \quad B_{n,0} = Q^{\binom{n}{2}} (t-1)^n + \sum_{1 \leq i \leq n} \begin{bmatrix} n \\ i \end{bmatrix}_Q \begin{bmatrix} n \\ i \end{bmatrix}_q (t-1)^{i-1} Q^{\binom{i}{2}} B_{n-i,0};$$

$$(21.11) \quad B_{\alpha, \beta} = \sum_{\substack{0 \leq i \leq \alpha \\ 0 \leq j \leq \beta \\ i+j \geq 1}} \begin{bmatrix} \alpha \\ i \end{bmatrix}_Q \begin{bmatrix} \alpha \\ i \end{bmatrix}_q \begin{bmatrix} \beta \\ j \end{bmatrix}_Q \begin{bmatrix} \beta \\ j \end{bmatrix}_q (t-1)^{i+j-1} Q^{\binom{i}{2} + \binom{j}{2}} B_{\alpha-i, \beta-j};$$

with $n \geq 1$, $\alpha + \beta \geq 1$ and $B_{0,0} = 1$.

First, if $(\Sigma, \sigma, \varepsilon)$ is a signed bipermutation such that ε contains α letters x and β letters y , both signed permutations (Σ, ε) , (σ, ε) are permutations of the billetters $(1, y) \dots (\beta, y)(1 + \beta, x) \dots (\alpha + \beta, x)$. Consequently, the left factor $(\Sigma(1), \sigma(1), \varepsilon(1)) \dots (\Sigma(k), \sigma(k), \varepsilon(k))$ ($1 \leq k \leq \alpha + \beta$) of $(\Sigma, \sigma, \varepsilon)$ is *decreasing*, if we have $\Sigma(1) > \dots > \Sigma(k)$ and $\sigma(1) > \dots > \sigma(k)$. Each compatible signed bipermutation [in short, *c.s.b.*] has a *longest decreasing left factor* [in short, *l.d.l.f.*] of length at least equal to 1. Each letter $(\Sigma(i), \sigma(i), \varepsilon(i))$ of the signed bipermutation $(\Sigma, \sigma, \varepsilon)$ is said to be *large* (resp. *small*), if $\varepsilon(i) = x$ (resp. $\varepsilon(i) = y$).

In the sequel the *c.s.b.* are supposed to have α large letters and β small letters. Let $F(i, j)$ denote the generating function for those *c.s.b.*, whose *l.d.l.f.* is of length $(i + j)$ and start with i large letters followed by j small letters ($0 \leq i \leq \alpha$, $0 \leq j \leq \beta$, $i + j \geq 1$). Let $G(\geq i, \geq 0, \geq k)$ (resp. $G(= i, \geq j, \geq k)$) designate the generating function for those *c.s.b.*, whose *l.d.l.f.* is of length at least equal to k and start with at least i large letters (resp. and start with i large letters followed by at least j small letters). Let

$$u_{i,j} = \begin{bmatrix} \alpha \\ i \end{bmatrix}_Q \begin{bmatrix} \alpha \\ i \end{bmatrix}_q \begin{bmatrix} \beta \\ j \end{bmatrix}_Q \begin{bmatrix} \beta \\ j \end{bmatrix}_q Q^{\binom{i}{2} + \binom{j}{2}} B_{\alpha-i, \beta-j},$$

so that (21.10) and (21.11) can be rewritten as

$$(21.12) \quad B_{\alpha,0} = Q^{\binom{\alpha}{2}}(t-1)^\alpha + \sum_{1 \leq i \leq \alpha} u_{i,0}(t-1)^{i-1} \quad (\beta = 0);$$

$$(21.13) \quad B_{\alpha,\beta} = \sum_{\substack{0 \leq i \leq \alpha \\ 0 \leq j \leq \beta \\ i+j \geq 1}} u_{i,j}(t-1)^{i+j-1} \quad (\beta \geq 1).$$

Lemma 21.3. *The following identities hold:*

$$(21.14) \quad t^i u_{i,0} = tF(i, 0) + G(\geq i, \geq 0, \geq i+1);$$

$$(21.15) \quad t^{i+j} u_{i,j} = tF(i, j) + G(= i, \geq j+1, \geq i+j+1), \quad \text{if } j \geq 1.$$

Proof. The product $t^i u_{i,0}$ is the generating function for the *c.s.b.* starting with a decreasing factor having i large letters, with the restriction that the initial number of descents is counted i (instead of $i-1$) when the *l.d.l.f.* is exactly of length i . In that generating function are included, besides the *c.s.b.* just described, the *c.s.b.* whose *l.d.l.f.* contains i large letters and at least a small letter, plus all the *c.s.b.* whose *l.d.l.f.* contains at least $i+1$ large letters and no small letter. This proves (21.14).

When $j \geq 1$, the product $t^{i+j} u_{i,j}$ is the generating function for the *c.s.b.* starting with a decreasing factor having i large letters and at least j small letters, still with the restriction that the initial number of descents is counted $i+j$ (instead of $i+j-1$) when the *l.d.l.f.* is of length $i+j$. Beside those *c.s.b.* the generating function involves all the *c.s.b.* whose *l.d.l.f.* contains i large letters and at least $j+1$ small letters. \square

As $F(i, 0) = G(\geq i, \geq 0, \geq i) - G(\geq i, \geq 0, \geq i+1)$, it follows from (21.14) that

$$(21.16) \quad G(\geq i, \geq 0, \geq i) = t^{i-1} u_{i,0} + \frac{t-1}{t} G(\geq i, \geq 0, \geq i+1);$$

it also follows from (21.15), when $j \geq 1$, that

$$(21.17) \quad G(=i, \geq j, \geq i+j) = t^{i+j-1} u_{i,j} + \frac{t-1}{t} G(=i, \geq j+1, \geq i+j+1).$$

Moreover, we have the identity:

$$(21.18) \quad G(\geq i, \geq 0, \geq i+1) = G(=i, \geq 1, \geq i+1) \\ + G(\geq i+1, \geq 0, \geq i+1).$$

22. EULERIAN CALCULUS; FINITE ANALOGS OF BESSEL FUNCTIONS

Since $B_{\alpha,\beta}$ is equal to $G(\geq 1, \geq 0, \geq 1) + G(= 0, \geq 1, \geq 1)$, those three formulas furnish the calculation for $B_{\alpha,\beta}$ as we now see. By iteration,

$$\begin{aligned}
 (21.19) \quad B_{\alpha,\beta} &= u_{1,0} + (t-1)u_{2,0} + \cdots + (t-1)^{\alpha-2}u_{\alpha-1,0} \\
 &\quad + \frac{(t-1)^{\alpha-1}}{t^{\alpha-1}}G(\geq \alpha, \geq 0, \geq \alpha) \\
 &\quad + \frac{(t-1)^{\alpha-1}}{t^{\alpha-1}}G(= \alpha-1, \geq 1, \geq \alpha) \\
 &\quad + \cdots + \frac{(t-1)}{t}G(= 1, \geq 1, \geq 2) + G(= 0, \geq 1, \geq 1).
 \end{aligned}$$

If $\beta = 0$, we have $G(\geq \alpha, \geq 0, \geq \alpha) = G(= \alpha, = 0, = \alpha) = F(\alpha, 0) = t^\alpha Q^{\binom{\alpha}{2}}$ and all quantities of the form $G(= i, \geq 1, \geq i+1)$ are null. As $u_{\alpha,0} = Q^{\binom{\alpha}{2}}$, we get

$$\begin{aligned}
 B_{\alpha,0} &= u_{1,0} + (t-1)u_{2,0} + \cdots + (t-1)^{\alpha-2}u_{\alpha-1,0} + t(t-1)^{\alpha-1}Q^{\binom{\alpha}{2}} \\
 &= u_{1,0} + (t-1)u_{2,0} + \cdots + (t-1)^{\alpha-2}u_{\alpha-1,0} + (t-1)^{\alpha-1}u_{\alpha,0} \\
 &\quad + (t-1)^\alpha Q^{\binom{\alpha}{2}},
 \end{aligned}$$

which is nothing but formula (21.12) to be proved.

When $\beta \geq 1$, we can pursue the iteration in (21.19) and because $G(\geq \alpha+1, \geq 0, \geq \alpha+1) = 0$ obtain

$$\begin{aligned}
 B_{\alpha,\beta} &= u_{1,0} + (t-1)u_{2,0} + \cdots + (t-1)^{\alpha-1}u_{\alpha,0} \\
 &\quad + \frac{(t-1)^\alpha}{t^\alpha}G(= \alpha, \geq 1, = \alpha+1) \\
 &\quad + \cdots + \frac{(t-1)}{t}G(= 1, \geq 1, \geq 2) + G(= 0, \geq 1, \geq 1).
 \end{aligned}$$

Then each expression $G(= i, \geq 1, \geq i+1)$ is iterated by making use of the relations $G(= i, \geq \beta+1, \geq i+\beta+1) = 0$ ($0 \leq i \leq \alpha$). We then obtain (21.13). \square

22. Eulerian Calculus; finite analogs of Bessel functions

Looking back on the diagram in Fig. 20.1 there remains to show how a further extension of Theorem 21.1 can be made and what meaning be given to the top level of the diagram, especially to vertex (sgn, q, J_k) . As was shown in the previous section, the letter “ J ” without subscript refers to the basic Bessel function. The subscript “ k ” will refer to a finite analog of that function, that is now introduced.

Rewrite the two identities of Proposition 4.1 as:

$$(22.1) \quad \begin{bmatrix} n+k \\ n \end{bmatrix}_q = \sum_{0 \leq b_1 \leq \dots \leq b_n < k+1} q^{\text{tot } b},$$

$$(22.2) \quad Q^{\binom{n}{2}} \begin{bmatrix} K+1 \\ n \end{bmatrix}_Q = \sum_{0 \leq B_1 < \dots < B_n < K+1} Q^{\text{tot } B}.$$

They hold for all integers $k \geq 0$ and $K \geq 0$. On the other hand, as we have (see (6.1))

$$(22.3) \quad \frac{1}{(q; q)_n} = \sum_{0 \leq b_1 \leq \dots \leq b_n} q^{\text{tot } b},$$

$$(22.4) \quad \frac{Q^{\binom{n}{2}}}{(Q; Q)_n} = \sum_{0 \leq B_1 < \dots < B_n} Q^{\text{tot } B},$$

we can let

$$(22.5) \quad \begin{bmatrix} n+\infty \\ n \end{bmatrix}_q := \frac{1}{(q; q)_n},$$

$$(22.6) \quad Q^{\binom{n}{2}} \begin{bmatrix} \infty+1 \\ n \end{bmatrix}_Q := \frac{Q^{\binom{n}{2}}}{(Q; Q)_n},$$

so that (22.1) and (22.2) are valid for all k and K finite or infinite.

Let $e_q^k(u)$ and $E_Q^K(u)$ denote the generating functions for (22.1) and (22.2), i.e.,

$$(22.7) \quad e_q^k(u) = \sum_{n \geq 0} \begin{bmatrix} n+k \\ n \end{bmatrix}_q u^n, \quad \text{still equal to } \frac{1}{(u; q)_{k+1}},$$

$$(22.8) \quad E_Q^K(u) = \sum_{n \geq 0} Q^{\binom{n}{2}} \begin{bmatrix} K+1 \\ n \end{bmatrix}_Q u^n.$$

In particular, the two q -exponentials may rewrite as

$$e_q(u) = e_q^\infty(u) = \sum_{n \geq 0} \frac{1}{(q; q)_n} u^n,$$

$$E_Q(u) = E_Q^\infty(u) = \sum_{n \geq 0} \frac{Q^{\binom{n}{2}}}{(Q; Q)_n} u^n.$$

22. EULERIAN CALCULUS; FINITE ANALOGS OF BESSEL FUNCTIONS

Let K, k be two fixed nonnegative integers. Using the Hadamard product (see (21.4)) we define the K, k -finite analog basic Bessel function $\mathbf{J}_k^K(u; Q, q)$ by

$$(22.9) \quad \mathbf{J}_k^K(u; Q, q) := \left(\sum_{n \geq 0} (-u)^n \right) \mathbb{H} E_Q^K(u) \mathbb{H} e_q^k(u);$$

or, in an equivalent manner, by

$$(22.10) \quad \mathbf{J}_k^K(u; Q, q) := \sum_{n \geq 0} (-1)^n Q^{\binom{n}{2}} \begin{bmatrix} K+1 \\ n \end{bmatrix}_Q \begin{bmatrix} n+k \\ n \end{bmatrix}_q u^n.$$

When K and k are infinite, then

$$(22.11) \quad \mathbf{J}_\infty^\infty(u; Q, q) = \mathbf{J}(u; Q, q) = \sum_{n \geq 0} (-1)^n \frac{Q^{\binom{n}{2}}}{(Q, Q)_n} \frac{1}{(q; q)_n} u^n,$$

which is the basic Bessel function defined and used in the previous section. We could take the Hadamard product of more than two Bessel functions; our results would still be valid, but awkward to state and handle.

Following the same pattern as in the previous two sections consider the fraction

$$(22.12) \quad F = F^{(K, M)}_{(k, m)} := \frac{(1-t) \mathbf{J}_k^K((1-t)X; P, p)}{-t + \mathbf{J}_k^K((1-t)X; P, p) \mathbf{J}_m^M((1-t)Y; Q, q)}$$

and expand it first as a generating function for *signed biwords*, then for *signed bipermutations*. Notice that the above fraction involves *two* Bessel functions $\mathbf{J}_k^K((1-t)X; P, p)$ and $\mathbf{J}_m^M((1-t)Y; Q, q)$.

22.1. *Signed biwords.* The next step is to express F as a homomorphic image of a generating function for the so-called *signed biwords*. We just make use of the combinatorial expressions given in (22.1) and (22.2). Let x, X, y, Y, P, p, Q, q designate independent variables and K, k, M, m four fixed nonnegative integers. Consider the alphabet $\mathfrak{A} = \mathfrak{A}^{(K, M)}_{(k, m)}$ whose

elements are the triletters $\begin{pmatrix} C \\ c \\ z \end{pmatrix}$, where $z = x$ or y , and C, c are integers such that

$$\begin{cases} 0 \leq C \leq K, 0 \leq c \leq k, & \text{if } z = x; \\ 0 \leq C \leq M, 0 \leq c \leq m, & \text{if } z = y. \end{cases}$$

Let $\Phi(\cdot) = \Phi(\cdot; P, p, Q, q, X, Y)$ be the map that sends each triletter $\begin{pmatrix} C \\ c \\ z \end{pmatrix}$ onto

$$\Phi\left(\begin{pmatrix} C \\ c \\ z \end{pmatrix}\right) = \begin{cases} P^C p^c X, & \text{if } z = x; \\ Q^C q^c Y, & \text{if } z = y. \end{cases}$$

Next, form the free monoid \mathfrak{A}^* generated by \mathfrak{A} . It consists of all the three-row matrices, called *signed biwords*,

$$(22.13) \quad w = \begin{pmatrix} B \\ b \\ \varepsilon \end{pmatrix} = \begin{pmatrix} B(1) & B(2) & \dots & B(n) \\ b(1) & b(2) & \dots & b(n) \\ \varepsilon(1) & \varepsilon(2) & \dots & \varepsilon(n) \end{pmatrix}$$

The *length* of w is n and denoted by $\ell(w)$. The signed biword of length 0 is the empty word, denoted by e . When $\varepsilon = x^n$ (resp. $\varepsilon = y^n$) and also when $B(1) < B(2) < \dots < B(n)$ and $b(1) \leq b(2) \leq \dots \leq b(n)$, we say that the signed word w is *nondecreasing*. Let ND_x (resp. ND_y) be the set of all nondecreasing signed biwords, whose bottom row has only x 's (resp. only y 's). The set of the juxtaposition products $w'w''$, where $w' \in \text{ND}_x$ and $w'' \in \text{ND}_y$, is denoted by $\text{ND}_x \text{ND}_y$.

By defining $\Phi(w)$ as the product of the images under Φ of the trileters of w , we extend Φ , by linearity, to a homomorphism of \mathfrak{A}^* into the ring of polynomials in the variables P, p, X, Q, q, Y .

Proposition 22.1. *Let*

$$G := \left(1 - \sum_w (t-1)^{\ell(w)-1} (w)\right)^{-1} \left(\sum_{w'} (t-1)^{\ell(w')} w'\right),$$

where w runs over $\text{ND}_x \text{ND}_y \setminus \{e\}$ and w' over ND_x . Then

$$F = \Phi(G).$$

Proof. It follows from (22.1) and (22.2) that

$$\mathbf{J}(X) := \mathbf{J}_k^K((1-t)X; P, p) = \sum_w (t-1)^{\ell(w)} \Phi(w) \quad (w \in \text{ND}_x);$$

$$\mathbf{J}(Y) := \mathbf{J}_m^M((1-t)Y; Q, q) = \sum_w (t-1)^{\ell(w)} \Phi(w) \quad (w \in \text{ND}_y);$$

$$\mathbf{J}(X)\mathbf{J}(Y) = \sum_w (t-1)^{\ell(w)} \Phi(w) \quad (w \in \text{ND}_x \text{ND}_y).$$

Hence, with w running over $\text{ND}_x \text{ND}_y$,

$$\begin{aligned} \frac{1-t}{-t + \mathbf{J}(X)\mathbf{J}(Y)} &= \frac{1-t}{-t + \sum_w (t-1)^{\ell(w)} \Phi(w)} \\ &= \left(1 - \sum_{w \neq e} (t-1)^{\ell(w)-1} \Phi(w)\right)^{-1}. \end{aligned}$$

We obtain the desired identity by multiplying by $\mathbf{J}(X)$ on the right. \square

Definition. Let w be a signed biword, as in (22.13), and i an integer. We say that i is a *rise* for w , if one of the following three conditions holds:

- (i) $i = n$ and $\varepsilon(n) = x$;
- (ii) $1 \leq i \leq n - 1$, $\varepsilon(i) = x$, $\varepsilon(i + 1) = y$;
- (iii) $1 \leq i \leq n - 1$, $\varepsilon(i) = \varepsilon(i + 1)$, $B(i) < B(i + 1)$ and $b(i) \leq b(i + 1)$.

The *number of rises* of the signed biword w is denoted by $\text{rise } w$.

Proposition 22.2. *The following identity holds:*

$$(22.14) \quad G = \sum_w t^{\text{rise } w} w$$

and consequently

$$(22.15) \quad F = \sum_w t^{\text{rise } w} \Phi(w),$$

where w runs over all signed biwords in \mathfrak{A}^* .

To prove Proposition 22.2 we make use of a classical inversion formula that is stated and proved in various contexts. Let X^* be the free monoid generated by a set X and form the large algebra of X^* over a ring Ω of polynomials. The elements of that algebra are the formal sums $\sum c(w) w$, where w runs over X^* and $c(w)$ belongs to Ω . Let $a : X^2 \rightarrow \Omega$ be a given mapping that we extend to a mapping $a : X^* \rightarrow \Omega$ by sending each word $w = x_1 x_2 \dots x_n \in X^*$ to

$$a(w) = \begin{cases} a(x_1, x_2) \dots a(x_{n-1}, x_n), & \text{if } n \geq 2; \\ 1, & \text{if } n = 0 \text{ or } 1. \end{cases}$$

Next define:

$$\bar{a}(w) := \begin{cases} (a(x_1, x_2) - 1) \dots (a(x_{n-1}, x_n) - 1), & \text{if } n \geq 2; \\ 1, & \text{if } n = 0 \text{ or } 1. \end{cases}$$

Further, let U, V be two nonempty subsets of the alphabet X . The expressions U^+ and U^*V stand for the sets of all *nonempty* words $w = x_1 x_2 \dots x_n$ all letters of which are in U (resp. whose *rightmost* letter x_n is in V and other letters are in U).

Lemma 22.3. *In the large algebra of X^* we have the identity:*

$$(22.16) \quad \sum_{w \in U^*V} a(w) w = \left(1 - \sum_{w \in U^+} \bar{a}(w) w \right)^{-1} \times \sum_{w \in U^*V} \bar{a}(w) w.$$

Proof. Multiply each side by $(1 - \sum_{w \in U^+} \bar{a}(w) w)$ on the left and verify that the coefficients of each word w on both sides are the same. \square

In formula (22.16) take the following ingredients: first, $X := \mathfrak{A} \cup \{\infty\}$, where “ ∞ ” is the triletter (∞, ∞, ∞) (writing the triletter horizontally for typographical reasons). Let $u = (C, c, z)$ and $u' = (C', c', z')$ be two triletters belonging to X .

For “ a ” take the mapping:

$$a(u, u') = \begin{cases} t, & \text{if } z = x \text{ and } z' = \infty; \\ t, & \text{if } z = x \text{ and } z' = y; \\ t, & \text{if } z = z' \text{ and } C < C' \text{ and } c \leq c'; \\ 1, & \text{otherwise.} \end{cases}$$

For V take the singleton $\{\infty\}$ and for U the alphabet \mathfrak{A} . Formula (22.16) is rewritten, after simplification on the right by ∞ , as

$$(22.17) \quad \sum_{w \in \mathfrak{A}^*} a(w \infty) w = \left(1 - \sum_{w \in \mathfrak{A}^* \setminus \{e\}} \bar{a}(w) w\right)^{-1} \times \sum_{w \in \mathfrak{A}^*} \bar{a}(w \infty) (w).$$

Let $w = u^{(1)}u^{(2)} \dots u^{(n)}$ be a signed biword of length n , written as a word with n triletters. Let $u^{(n+1)} := \infty$. Then, the statistic $\text{rise}(w)$ can also be expressed as:

$$(22.18) \quad \text{rise}(w) = \sum_{1 \leq i \leq n} \chi(a(u^{(i)} u^{(i+1)}) = t).$$

For every signed biword w we then have:

$$\begin{aligned} a(w \infty) &= t^{\text{rise } w}, \\ \bar{a}(w) &= \begin{cases} (t-1)^{\ell(w)-1}, & \text{if } w \in \text{ND}_x \text{ND}_y \text{ and } \ell(w) \geq 2; \\ 1, & \text{if } \ell(w) = 0 \text{ or } 1; \\ 0, & \text{otherwise;} \end{cases} \\ \bar{a}(w \infty) &= \begin{cases} (t-1)^{\ell(w)}, & \text{if } w \in \text{ND}_x \text{ and } \ell(w) \geq 1; \\ 1, & \text{if } \ell(w) = 0; \\ 0, & \text{otherwise.} \end{cases} \end{aligned}$$

With those ingredients the left-hand side of identity (22.17) is equal to $\sum_{w \in \mathfrak{A}^*} t^{\text{rise } w} w$ and the right-hand side is the expression found for G in

Proposition 22.1. \square

22.2. Signed bipermutations. Although Proposition 22.2 provides an expression for the fraction F in terms of *biwords* and involves one statistic, namely the *number of rises*, the combinatorial objects are still rudimentary. To obtain an expression in terms of combinatorial objects having a richer geometry, such as permutations or signed permutations, it is necessary to imagine a further correspondence. This is the purpose of this subsection and the next one.

The definition of a signed multipermutation $(\underline{\Sigma}, \underline{\sigma}, \varepsilon)$ was given in § 21.2. When $\underline{\Sigma}$ (resp. $\underline{\sigma}$) consists of a single permutation Σ (resp. σ), we speak of a *signed bipermutation* $(\Sigma, \sigma, \varepsilon)$, as in the present subsection. With such a signed bipermutation we associate the subset $I_{\varepsilon|x} := \{i_1, \dots, i_{\ell(\varepsilon|x)}\}$ (resp. $I_{\varepsilon|y} := \{j_1, \dots, j_{\ell(\varepsilon|y)}\}$) of all the integers i such that $\varepsilon(i) = x$ (resp. $\varepsilon(i) = y$).

Suppose that $(\Sigma, \sigma, \varepsilon)$ is *compatible* in the sense of Proposition 21.2 and consider the restriction of the bijection $i \mapsto \Sigma(i)$ to $I_{\varepsilon|y}$ (resp. to $I_{\varepsilon|x}$); it is a *bijection*, denoted by $\Sigma_{\varepsilon|y}$, of $I_{\varepsilon|y}$ onto the interval $[1, \ell(\varepsilon|y)]$ (resp. a bijection $\Sigma_{\varepsilon|x}$ of $I_{\varepsilon|x}$ onto the interval $[\ell(\varepsilon|y) + 1, n]$). In the same manner, two restrictions of σ are considered: $\sigma_{\varepsilon|y}$ of $I_{\varepsilon|y}$ onto $[1, \ell(\varepsilon|y)]$ and $\sigma_{\varepsilon|x}$ of $I_{\varepsilon|x}$ onto $[\ell(\varepsilon|y) + 1, n]$.

For instance,

$$(22.19) \quad \begin{aligned} \Sigma &= (\mathbf{1\ 2\ 3\ 4\ 5\ 6\ 7\ 8\ 9}) \\ \sigma &= (\mathbf{7\ 5\ 3\ 4\ 2\ 1\ 8\ 9\ 6}) \\ \varepsilon &= (x\ x\ y\ y\ y\ y\ x\ x\ x) \end{aligned}$$

is a compatible signed bipermutation of order $n = 9$. Also $I_{\varepsilon|x} = \{\mathbf{1, 2, 7, 8, 9}\}$, $I_{\varepsilon|y} = \{3, 4, 5, 6\}$.

The restrictions $\Sigma_{\varepsilon|x}$, $\Sigma_{\varepsilon|y}$, $\sigma_{\varepsilon|x}$, $\sigma_{\varepsilon|y}$, written as two-row matrices, together with their inverses read:

$$\begin{aligned} \Sigma_{\varepsilon|x} &= \begin{pmatrix} \mathbf{1\ 2\ 7\ 8\ 9} \\ \mathbf{7\ 5\ 8\ 9\ 6} \end{pmatrix}, \quad \Sigma_{\varepsilon|y} = \begin{pmatrix} \mathbf{3\ 4\ 5\ 6} \\ \mathbf{3\ 4\ 2\ 1} \end{pmatrix}, \quad \sigma_{\varepsilon|x} = \begin{pmatrix} \mathbf{1\ 2\ 7\ 8\ 9} \\ \mathbf{6\ 7\ 5\ 8\ 9} \end{pmatrix}, \quad \sigma_{\varepsilon|y} = \begin{pmatrix} \mathbf{3\ 4\ 5\ 6} \\ \mathbf{2\ 4\ 3\ 1} \end{pmatrix}, \\ \Sigma_{\varepsilon|x}^{-1} &= \begin{pmatrix} \mathbf{5\ 6\ 7\ 8\ 9} \\ \mathbf{2\ 9\ 1\ 7\ 8} \end{pmatrix}, \quad \Sigma_{\varepsilon|y}^{-1} = \begin{pmatrix} \mathbf{1\ 2\ 3\ 4} \\ \mathbf{6\ 5\ 3\ 4} \end{pmatrix}, \quad \sigma_{\varepsilon|x}^{-1} = \begin{pmatrix} \mathbf{5\ 6\ 7\ 8\ 9} \\ \mathbf{7\ 1\ 2\ 8\ 9} \end{pmatrix}, \quad \sigma_{\varepsilon|y}^{-1} = \begin{pmatrix} \mathbf{1\ 2\ 3\ 4} \\ \mathbf{6\ 3\ 5\ 4} \end{pmatrix}. \end{aligned}$$

We do not repeat the definitions of “des” and “maj” for permutations, but introduce the definition of “rise” and “rmaj” (referring to the “rises” and no longer to the “descents”). If ρ is a permutation, written as a linear word $\rho = \rho(1)\rho(2) \dots \rho(n)$. Then, define

$$\begin{aligned} \text{rise } \rho &:= \sum_{1 \leq i \leq n-1} \chi(\rho(i) < \rho(i+1)); \\ \text{rmaj } \rho &:= \sum_{1 \leq i \leq n-1} i \chi(\rho(i) < \rho(i+1)). \end{aligned}$$

The statistics “des” and “maj” will be determined for $\Sigma_{\varepsilon|x}^{-1}$, $\sigma_{\varepsilon|y}^{-1}$ and the statistics “rise” and “rmaj” for $\sigma_{\varepsilon|x}^{-1}$, $\sigma_{\varepsilon|y}^{-1}$, by means of word codes, as it was done in § 12.2 for the application of the MacMahon *Verfahren*. To avoid

cumbersome notations suppose that those four bijections are identified with the bottom rows of their defining two-row matrices, respectively denoted by $S_x := S_x(1) \dots S_x(\ell(\varepsilon|x))$, $S_y := S_y(1) \dots S_y(\ell(\varepsilon|y))$, $s_x := s_x(1) \dots s_x(\ell(\varepsilon|x))$, $s_y := s_y(1) \dots s_y(\ell(\varepsilon|y))$.

For each $i = 1, \dots, \ell(\varepsilon|x)$ let $Z_x(i)$ be the number of j such that $i \leq j \leq \ell(\varepsilon|x) - 1$ and $S_x(j) > S_x(j+1)$, that is, to the *number of descents* to the right of $S_x(i)$ and form the word $Z_x := Z_x(1) \dots Z_x(\ell(\varepsilon|x))$. Define Z_y in the same way, replacing “ x ” by “ y ”.

For s_x and s_y the word “descent” is to be replaced by “rise.” For each $i = 1, \dots, \ell(\varepsilon|x)$ let $z_x(i)$ be the number of j such that $i \leq j \leq \ell(\varepsilon|x) - 1$ and $s_x(j) < s_x(j+1)$ and form the word $z_x := z_x(1) \dots z_x(\ell(\varepsilon|x))$. Define z_y in the same way, replacing “ x ” by “ y ”.

With the above example we get

$$\begin{matrix} S_x = (\mathbf{29178}) & S_y = (\mathbf{6534}) & s_x = (\mathbf{71289}) & s_y = (\mathbf{6354}) \\ Z_x = (\mathbf{11000}) & Z_y = (\mathbf{2100}) & z_x = (\mathbf{33210}) & z_y = (\mathbf{1100}) \end{matrix}.$$

Proposition 22.4. *We have:*

$$\begin{aligned} \text{des } \Sigma_{\varepsilon|x}^{-1} &= Z_x(1); & \text{maj } \Sigma_{\varepsilon|x}^{-1} &= \text{tot } Z_x = Z_x(1) + \dots + Z_x(\ell(\varepsilon|x)); \\ \text{des } \Sigma_{\varepsilon|y}^{-1} &= Z_y(1); & \text{maj } \Sigma_{\varepsilon|y}^{-1} &= \text{tot } Z_y = Z_y(1) + \dots + Z_y(\ell(\varepsilon|y)); \\ \text{rise } \sigma_{\varepsilon|x}^{-1} &= z_x(1); & \text{rmaj } \sigma_{\varepsilon|x}^{-1} &= \text{tot } z_x = z_x(1) + \dots + z_x(\ell(\varepsilon|x)); \\ \text{rise } \sigma_{\varepsilon|y}^{-1} &= z_y(1); & \text{rmaj } \sigma_{\varepsilon|y}^{-1} &= \text{tot } z_y = z_y(1) + \dots + z_y(\ell(\varepsilon|y)). \end{aligned}$$

Proof. With the words Z_x , Z_y , z_x and z_y we have another way of counting descents and rises. \square

To further simplify the notations the following shorthand will be used:

$$\begin{aligned} \text{idex } \Sigma &:= \text{des } \Sigma_{\varepsilon|x}^{-1}; & \text{imaj } \Sigma &:= \text{maj } \Sigma_{\varepsilon|x}^{-1}; \\ \text{idey } \Sigma &:= \text{des } \Sigma_{\varepsilon|y}^{-1}; & \text{imaj } \Sigma &:= \text{maj } \Sigma_{\varepsilon|y}^{-1}; \\ \text{irise } \sigma &:= \text{rise } \sigma_{\varepsilon|x}^{-1}; & \text{irmaj } \sigma &:= \text{rmaj } \sigma_{\varepsilon|x}^{-1}; \\ \text{irise } \sigma &:= \text{rise } \sigma_{\varepsilon|y}^{-1}; & \text{irmaj } \sigma &:= \text{rmaj } \sigma_{\varepsilon|y}^{-1}. \end{aligned}$$

With each compatible signed bipermutation $(\Sigma, \sigma, \varepsilon)$ is associated the eleven-variable monomial

$$(22.20) \quad \Psi(\Sigma, \sigma, \varepsilon) := X^{\ell(\varepsilon|x)} Y^{\ell(\varepsilon|y)} t^{\text{ddes}(\Sigma, \sigma, \varepsilon)} \\ \times U^{\text{idex } \Sigma} V^{\text{idey } \Sigma} P^{\text{imaj } \Sigma} Q^{\text{imaj } \Sigma} u^{\text{irise } \sigma} v^{\text{irise } \sigma} p^{\text{irmaj } \sigma} q^{\text{irmaj } \sigma}.$$

Our purpose is to work out the generating function for the polynomials

$$(22.21) \quad \mathbf{B}_{\alpha,\beta} := \sum \Psi(\Sigma, \sigma, \varepsilon),$$

where the sum is over all compatible signed bipermutations $(\Sigma, \sigma, \varepsilon)$ such that $\ell(\varepsilon|x) = \alpha$ and $\ell(\varepsilon|y) = \beta$.

22.3. Signed biwords and compatible bipermutations. Using the notations of the previous subsection and Proposition 22.2, we prove the following theorem.

Theorem 22.5. *The factorial generating function for the polynomials $\mathbf{B}_{\alpha,\beta}$ defined in (22.21) is given by*

$$\begin{aligned} \sum_{\binom{K,M}{k,m}} U^K V^M u^k v^m \frac{(1-t) \mathbf{J}_k^K((1-t)X; P, p)}{-t + \mathbf{J}_k^K((1-t)X; P, p) \mathbf{J}_m^M((1-t)Y; Q, q)} \\ = \sum_{\alpha \geq 0, \beta \geq 0} e_P^\alpha(U) e_Q^\beta(V) e_p^\alpha(u) e_q^\beta(v) \mathbf{B}_{\alpha,\beta}. \end{aligned}$$

By (22.12) and (22.15) the left-hand side of the previous identity is equal to

$$\sum_{\binom{K,M}{k,m}} R^K S^M r^k s^m F_{k,m}^{(K,M)} = \sum_{\binom{K,M}{k,m}} U^K V^M u^k v^m \sum_{w \in \mathfrak{A}_{k,m}^{(K,M)*}} t^{\text{rise } w} \Phi(w),$$

so that there remains to find an appropriate correspondence between signed biwords and compatible signed bipermutations.

Again use the notation (22.13) for the signed biword w and still let $I_{\ell(\varepsilon|x)}$ (resp. $I_{\ell(\varepsilon|y)}$) be the set of all i such that $\varepsilon(i) = x$ (resp. $\varepsilon(i) = y$). Then let $B|x := B_x(1) \dots B_x(\ell(\varepsilon|x))$ (resp. $B|y := B_y(1) \dots B_y(\ell(\varepsilon|y))$) be the *nonincreasing* rearrangement of all letters $B(i)$ such that $i \in I_{\ell(\varepsilon|x)}$ (resp. $i \in I_{\ell(\varepsilon|y)}$). Let $b|x := b_x(1) \dots b_x(\ell(\varepsilon|x))$ (resp. $b|y := b_y(1) \dots b_y(\ell(\varepsilon|y))$) have analogous definitions when B is replaced by b . Hence (see § 22.1) the image $\Phi(w)$ of the signed biword w reads:

$$(22.22) \quad \Phi(w) = P^{\text{tot } B|x} Q^{\text{tot } B|y} p^{\text{tot } b|x} q^{\text{tot } b|y} X^{\ell(\varepsilon|x)} Y^{\ell(\varepsilon|y)}.$$

For example, let $K = 8$, $M = 6$, $k = 9$, $m = 8$ and consider the signed biword $w \in \mathfrak{A}_{9,8}^{(8,6)}$:

$$w = \begin{pmatrix} B \\ b \\ \varepsilon \end{pmatrix} = \begin{pmatrix} 4 & 7 & 1 & 0 & 4 & 6 & 3 & 3 & 5 \\ 9 & 5 & 4 & 0 & 0 & 6 & 9 & 1 & 0 \\ x & x & y & y & y & y & x & x & x \end{pmatrix}$$

Then $B|x = 75433$, $B|y = 6410$, $b|x = 99510$, $b|y = 6400$, $\ell(\varepsilon|x) = 5$, $\ell(\varepsilon|y) = 4$, so that $\Phi(w) = P^{22}Q^{11}p^{24}q^{10}X^5Y^4$. The rises of w are: 2 (because $\varepsilon(2) = y$, $\varepsilon(3) = x$), 4 and 5 (because $\varepsilon(4) = \varepsilon(5) = \varepsilon(6) = y$, $B(4) = 0 < B(5) = 4 < B(6) = 6$, $b(4) = 0 \leq b(5) = 0 \leq b(6) = 6$) and finally 9 (because $n = 9$ and $\varepsilon(9) = x$). Hence, rise $w = 4$.

Definition of the mapping $w = (B, b, \varepsilon) \mapsto (\Sigma, \sigma, \varepsilon)$. First, the x, y -word ε remains the same. If $\varepsilon(i) = y$ let

$$(22.23) \quad \Sigma(i) := |\{j : 1 \leq j \leq n, B(j) > B(i), \varepsilon(j) = y\}| \\ + |\{j : 1 \leq j \leq i, B(j) = B(i), \varepsilon(j) = y\}|;$$

and if $\varepsilon(i) = x$ let

$$(22.23') \quad \Sigma(i) := \ell(\varepsilon|y) + |\{j : 1 \leq j \leq n, B(j) > B(i), \varepsilon(j) = x\}| \\ + |\{j : 1 \leq i \leq j, B(j) = B(i), \varepsilon(j) = x\}|.$$

Next for each $i = 1, 2, \dots, n$ and $\varepsilon(i) = y$ let

$$(22.24) \quad \sigma(i) := |\{j : 1 \leq j \leq n, b(j) > b(i), \varepsilon(j) = y\}| \\ + |\{j : i \leq j \leq n, b(j) = b(i), \varepsilon(j) = y\}|.$$

When $\varepsilon(i) = x$, let

$$(22.24') \quad \sigma(i) := \ell(\varepsilon|y) + |\{j : 1 \leq j \leq n, b(j) > b(i), \varepsilon(j) = x\}| \\ + |\{j : i \leq j \leq n, b(j) = b(i), \varepsilon(j) = x\}|.$$

Notice that the only difference in the definitions of σ and Σ is the following: if $b(i) = b(j)$ (resp. $B(i) = B(j)$) and $i < j$, then $\sigma(i) > \sigma(j)$ (resp. $\Sigma(i) < \Sigma(j)$). Obviously, $(\Sigma, \sigma, \varepsilon)$ is a compatible *signed bipermutation*.

For instance, to the above signed biword $w = (B, b, \varepsilon)$ there corresponds the compatible signed permutation displayed in (22.19). Let us materialize the correspondence with the following matrix.

$$(22.25) \quad \begin{array}{l} B = \\ b = \\ \varepsilon = \\ \Sigma = \\ \sigma = \end{array} \begin{pmatrix} \mathbf{1} & \mathbf{2} & \mathbf{3} & \mathbf{4} & \mathbf{5} & \mathbf{6} & \mathbf{7} & \mathbf{8} & \mathbf{9} \\ \mathbf{4} & \mathbf{7} & \mathbf{1} & \mathbf{0} & \mathbf{4} & \mathbf{6} & \mathbf{3} & \mathbf{3} & \mathbf{5} \\ \mathbf{9} & \mathbf{5} & \mathbf{4} & \mathbf{0} & \mathbf{0} & \mathbf{6} & \mathbf{9} & \mathbf{1} & \mathbf{0} \\ x & x & y & y & y & y & x & x & x \\ \mathbf{7} & \mathbf{5} & \mathbf{3} & \mathbf{4} & \mathbf{2} & \mathbf{1} & \mathbf{8} & \mathbf{9} & \mathbf{6} \\ \mathbf{6} & \mathbf{7} & \mathbf{2} & \mathbf{4} & \mathbf{3} & \mathbf{1} & \mathbf{5} & \mathbf{8} & \mathbf{9} \end{pmatrix}.$$

Once the signed bipermutation $(\Sigma, \sigma, \varepsilon)$ has been obtained, we determine the words S_x, Z_x, \dots and the statistics $\text{idex}_x, \text{idex}_y, \dots$, as they were introduced in the previous subsection; also, we let

$$(22.26) \quad \begin{aligned} K' &:= K - \text{idex}_x \Sigma; & M' &:= M - \text{idex}_y \Sigma; \\ k' &:= k - \text{irise}_x \sigma; & m' &:= m - \text{idex}_y \sigma; \end{aligned}$$

and define the words:

$$\Lambda := B|x - Z_x, \quad \Theta := B|y - Z_y, \quad \lambda := b|x - z_x, \quad \theta := b|y - z_y,$$

where the difference is defined letter by letter. For instance,

$$\Lambda = \Lambda(1) \cdots \Lambda(\ell(\varepsilon|x)) := (B_x(1) - Z_x(1)) \cdots (B_x(\ell(\varepsilon|x)) - Z_x(\ell(\varepsilon|x))).$$

Proposition 22.6. *The integers K' , M' , k' , m' are nonnegative. Furthermore,*

- (i) $K' \geq \Lambda(1) \geq \cdots \geq \Lambda(\ell(\varepsilon|x)) \geq 0$; $M' \geq \Theta(1) \geq \cdots \geq \Theta(\ell(\varepsilon|y)) \geq 0$;
- (ii) $k' \geq \lambda(1) \geq \cdots \geq \lambda(\ell(\varepsilon|x)) \geq 0$; $m' \geq \theta(1) \geq \cdots \geq \theta(\ell(\varepsilon|y)) \geq 0$;
- (iii) $\text{tot } B|x = \text{imaj } \Sigma_x + \text{tot } \Lambda$; $\text{tot } B|y = \text{imaj } \Sigma_y + \text{tot } \Theta$;
- (iv) $\text{tot } b|x = \text{imaj } \sigma_x + \text{tot } \lambda$; $\text{tot } b|y = \text{imaj } \sigma_y + \text{tot } \theta$.

Also the rises of w and the descents of $(\Sigma, \sigma, \varepsilon)$ coincide. In particular,

- (v) $\text{rise } w = \text{ddes}(\Sigma, \sigma, \varepsilon)$.

Finally, the mapping

$$\left(\binom{KM}{k m}, w \right) \mapsto \left(\binom{K'M'}{k' m'}, \Lambda, \Theta, \lambda, \theta, (\Sigma, \sigma, \varepsilon) \right)$$

is bijective.

With our running example, the correspondence between w and $(\Sigma, \sigma, \varepsilon)$ was shown in (22.25). Also $K = 8$; $M = 6$, $k = 9$, $m = 8$. The other parameters are (see the first calculations just before Proposition 22.4): $K' = K - Z_x(1) = 8 - 1 = 7$, $M' = M - Z_y(1) = 6 - 2 = 4$, $k' = k - z_x(1) = 9 - 3 = 6$, $m' = m - z_y(1) = 8 - 1 = 7$. Finally, $\Lambda = B|x - Z_x = \mathbf{6\ 4\ 4\ 3\ 3}$, $\Theta = 4\ 3\ 1\ 0$, $\lambda = \mathbf{6\ 6\ 3\ 0\ 0}$, $\theta = 5\ 3\ 0\ 0$.

The proof of Proposition 22.6 follows the pattern derived earlier for the MacMahon *Verfahren*.

The final calculation uses all those items, as well as identity (22.3):

$$\begin{aligned} & \sum_{\binom{KM}{k m}, w} U^K V^M u^k v^m t^{\text{rise } w} \Phi(w) \\ &= \sum_{(\Sigma, \sigma, \varepsilon)} \Psi(\Sigma, \sigma, \varepsilon) \sum_{\binom{K'M'}{k' m'}} U^{K'} V^{M'} u^{k'} v^{m'} \sum_{\binom{\Lambda \Theta}{\lambda \theta}} P^{\text{tot } \Lambda} Q^{\text{tot } \Theta} p^{\text{tot } \lambda} q^{\text{tot } \theta} \\ &= \sum_{\alpha \geq 0, \beta \geq 0} \sum_{\substack{\ell(\varepsilon|x) = \alpha \\ \ell(\varepsilon|y) = \beta}} \sum_{(\Sigma, \sigma, \varepsilon)} \Psi(\Sigma, \sigma, \varepsilon) e_P^\alpha(U) e_Q^\beta(V) e_p^\alpha(u) e_q^\beta(v) \\ &= \sum_{\alpha \geq 0, \beta \geq 0} e_P^\alpha(U) e_Q^\beta(V) e_p^\alpha(u) e_q^\beta(v) \mathbf{B}_{\alpha, \beta}. \quad \square \end{aligned}$$

22.4. *The last specializations.* With $X = 0$ in the identity of Theorem 22.5 we get:

$$(22.27) \quad \sum_{\binom{M}{m}} V^M v^m \frac{1-t}{-t + \mathbf{J}_m^M((1-t)Y; Q, q)}$$

$$= \sum_{\beta \geq 0} e_Q^\beta(V) e_q^\beta(v) Y^\beta \sum_{(\Sigma, \sigma)} t^{\text{ddes}(\Sigma, \sigma)} V^{\text{idcs } \Sigma} Q^{\text{imaj } \Sigma} v^{\text{irise } \sigma} q^{\text{irmaj } \sigma}.$$

The sum on the right-hand side is over all pairs of permutations (Σ, σ) . This can be regarded as the specialization sitting on vertex (\cdot, q, J_k) in the diagram of Fig. 20.1.

When the parameters K, M, k, m are not finite, the identity of Theorem 22.5 is derived as follows: multiply both sides successively by $(1-U)$ and let $U = 1$, multiply by $(1-V)$ and let $V = 1$, multiply by $(1-u)$ and let $u = 1$, multiply by $(1-v)$ and let $v = 1$. We obtain the identity:

$$(22.28) \quad \frac{(1-t) \mathbf{J}((1-t)X; P, p)}{-t + \mathbf{J}((1-t)X; P, p) \mathbf{J}((1-t)Y; Q, q)}$$

$$= \sum_{\alpha \geq 0, \beta \geq 0} \frac{1}{(P; P)_\alpha} \frac{1}{(Q; Q)_\beta} \frac{1}{(p; p)_\alpha} \frac{1}{(q; q)_\beta} X^\alpha Y^\beta$$

$$\times \sum_{\substack{(\Sigma, \sigma, \varepsilon) \text{ (comp.)} \\ \ell(\varepsilon|x) = \alpha, \ell(\varepsilon|y) = \beta}} t^{\text{ddes}(\Sigma, \sigma, \varepsilon)} P^{\text{imaj}_x \Sigma} Q^{\text{imaj}_y \Sigma} p^{\text{irmaj}_x \sigma} q^{\text{irmaj}_y \sigma}.$$

The latter underlying generating function is over the *compatible* signed bipermutations only, while identity (21.6), that corresponds to vertex (sgn, q, J) , is over *all* signed bipermutations and even multipermutations. However identity (22.8) involves *two* distinct Bessel functions $\mathbf{J}(\cdot; P, p)$, $\mathbf{J}(\cdot; Q, q)$, instead of only one. When we reduce the number of bases, say, when $(P, p) = (Q, q)$, we can still regain (21.6), as is now explained.

For each signed bipermutation (not necessarily compatible) $(\Sigma, \sigma, \varepsilon)$ let

$$\text{inv}(\sigma_{\varepsilon|y}, \sigma_{\varepsilon|x}) := \#\{(i, j) : \varepsilon(i) = y, \varepsilon(j) = x, \sigma(i) > \sigma(j)\},$$

a quantity equal to zero when the bipermutation is compatible. Also let

$$\text{imaj}(\Sigma, \varepsilon) := \text{imaj}_x(\Sigma, \varepsilon) + \text{imaj}_y(\Sigma, \varepsilon) + \text{inv}(\sigma_{\varepsilon|y}, \sigma_{\varepsilon|x})$$

$$\text{irmaj}(\sigma, \varepsilon) := \text{irmaj}_x(\sigma, \varepsilon) + \text{irmaj}_y(\sigma, \varepsilon) + \text{inv}(\sigma_{\varepsilon|y}, \sigma_{\varepsilon|x}).$$

22. EULERIAN CALCULUS; FINITE ANALOGS OF BESSEL FUNCTIONS

Now in (22.8) let $(P, p) = (Q, q)$; the right-hand side can then be rewritten as:

$$\sum_{n \geq 0} \frac{1}{(Q; Q)_n (q; q)_n} \sum_{\alpha + \beta = n} \begin{bmatrix} n \\ \alpha \end{bmatrix}_Q \begin{bmatrix} n \\ \alpha \end{bmatrix}_q X^\alpha Y^\beta \times \sum_{\substack{(\Sigma, \sigma, \varepsilon) \text{ (comp.)} \\ \ell(\varepsilon|x) = \alpha, \ell(\varepsilon|y) = \beta}} t^{\text{ddes}(\Sigma, \sigma, \varepsilon)} Q^{\text{imaj}(\Sigma, \varepsilon)} q^{\text{irmaj}(\sigma, \varepsilon)}.$$

Because of the definition of “imaj” and “irmaj” for a signed permutation the polynomial $B'_n(X, Y, t, Q, q)$ defined by

$$(22.29) \quad B'_n(X, Y, t, Q, q) := \sum_{(\Sigma, \sigma, \varepsilon)} X^{\ell(\varepsilon|x)} Y^{\ell(\varepsilon|y)} t^{\text{ddes}(\Sigma, \sigma, \varepsilon)} Q^{\text{imaj}(\Sigma, \varepsilon)} q^{\text{irmaj}(\sigma, \varepsilon)},$$

the sum being over all signed bipermutations of length n , is also equal to

$$\sum_{\alpha + \beta = n} \begin{bmatrix} n \\ \alpha \end{bmatrix}_Q \begin{bmatrix} n \\ \alpha \end{bmatrix}_q X^\alpha Y^\beta \sum_{\substack{(\Sigma, \sigma, \varepsilon) \text{ (comp.)} \\ \ell(\varepsilon|x) = \alpha, \ell(\varepsilon|y) = \beta}} t^{\text{ddes}(\Sigma, \sigma, \varepsilon)} Q^{\text{imaj} \Sigma} q^{\text{irmaj} \sigma}.$$

Hence, when $(P, p) = (Q, q)$ identity (22.28) takes the form

$$(22.30) \quad \frac{(1-t) \mathbf{J}((1-t)X; Q, q)}{-t + \mathbf{J}((1-t)X; Q, q) \mathbf{J}((1-t)Y; Q, q)} = \sum_{n \geq 0} \frac{1}{(Q; Q)_n (q; q)_n} B'_n(X, Y, t, Q, q).$$

The comparison with (21.6) shows that the right-hand sides of (21.6) and (22.30) involve different statistics: $\text{inv}(\Sigma, \varepsilon), \text{coinv}(\sigma, \varepsilon)$ for (21.6) (when reduced to bipermutations) and $\text{imaj}(\Sigma, \varepsilon), \text{irmaj}(\sigma, \varepsilon)$ for (22.30). However, those two formulas imply the identity

$$(22.31) \quad B_n(X, Y, t, Q, q) = B'_n(X, Y, t, Q, q).$$

In order to show that Theorem 22.5 that contains the most general extension sitting on (sgn, q, J_k) implies (21.6) sitting on (sgn, q, J) we prove identity (22.31) *combinatorially* by means of a direct bijection.

A bijection for $(\text{inv}, \text{coinv})$ and $(\text{imaj}, \text{irmaj})$. The main ingredient is the bijection Φ of Theorem 11.2 that is used in the following context. Let π be a bijection of a finite set $I = \{i_1 < i_2 < \dots < i_k\}$, of integers onto a finite set of integers $\pi(I) := \{j_1 < j_2 < \dots < j_k\}$. The

permutation π is presented as a two-row matrix $\begin{pmatrix} i_1 & \dots & i_k \\ \pi(i_1) & \dots & \pi(i_k) \end{pmatrix}$. Let \mathbf{i} denote the correspondence that sends each bijection onto its inverse and consider the chain

$$\begin{pmatrix} i_1 & \dots & i_k \\ \pi(i_1) & \dots & \pi(i_k) \end{pmatrix} \xrightarrow{\mathbf{i}} \begin{pmatrix} j_1 & \dots & j_k \\ \pi^{-1}(j_1) & \dots & \pi^{-1}(j_k) \end{pmatrix} \\ \xrightarrow{\Phi} \begin{pmatrix} j_1 & \dots & j_k \\ y_1 & \dots & y_k \end{pmatrix} \xrightarrow{\mathbf{i}} \begin{pmatrix} i_1 & \dots & i_k \\ \pi'(i_1) & \dots & \pi'(i_k) \end{pmatrix},$$

that is rewritten:

$$\pi \xrightarrow{\mathbf{i}} \mathbf{i}\pi \xrightarrow{\Phi} \Phi\mathbf{i}\pi \xrightarrow{\mathbf{i}} \mathbf{i}\Phi\mathbf{i}\pi.$$

As proved in Theorem 11.2, the transformation Φ sends the *word* $\pi^{-1}(j_1) \dots \pi^{-1}(j_k)$ onto a *rearrangement* $y_1 \dots y_k$ of that word, which is the bottom row of the third matrix above. Moreover,

$$\text{maj}(\pi^{-1}(j_1) \dots \pi^{-1}(j_k)) = \text{inv}(y_1 \dots y_k),$$

which implies

$$\text{maj}\mathbf{i}\pi = \text{inv}\mathbf{i}\Phi\mathbf{i}\pi \quad \text{and} \quad \text{rmaj}\mathbf{i}\pi = \text{coinv}\mathbf{i}\Phi\mathbf{i}\pi.$$

Also $\text{ligne}\mathbf{i}\pi = \text{ligne}\Phi\mathbf{i}\pi$. Hence,

$$(22.32). \quad \text{Ligne}\pi = \text{Ligne}\mathbf{i}\Phi\mathbf{i}\pi.$$

Now each signed bipermutation $(\Sigma, \sigma, \varepsilon)$ is characterized by the two subsets $I_{\varepsilon|x}, I_{\varepsilon|y}$ and its restrictions $\Sigma_x, \Sigma_y, \sigma_x, \sigma_y$ to those two subsets, as they were defined in § 22.2. This gives a sense to the notation

$$[I_{\varepsilon|x}, I_{\varepsilon|y}, \Sigma_x, \Sigma_y, \sigma_x, \sigma_y]$$

to designate a signed bipermutation $(\Sigma, \sigma, \varepsilon)$.

Next form the composition product $\mathbf{i}\Phi\mathbf{i}$ described above and consider the mapping $\Delta : (\Sigma, \sigma, \varepsilon) \mapsto (\Sigma', \sigma', \varepsilon)$ defined by

$$\begin{aligned} (\Sigma, \sigma, \varepsilon) &:= [I_{\varepsilon|x}, I_{\varepsilon|y}, \Sigma_x, \Sigma_y, \sigma_x, \sigma_y]; \\ (\Sigma', \sigma', \varepsilon) &:= [I_{\varepsilon|x}, I_{\varepsilon|y}, \mathbf{i}\Phi\mathbf{i}\Sigma_x, \mathbf{i}\Phi\mathbf{i}\Sigma_y, \mathbf{i}\Phi\mathbf{i}\sigma_x, \mathbf{i}\Phi\mathbf{i}\sigma_y]. \end{aligned}$$

Proposition 22.6. *The mapping Δ is a bijection of the set of signed bipermutations of length n onto itself, having the following properties:*

- (i) $\text{ddes}(\Sigma, \sigma, \varepsilon) = \text{ddes}(\Sigma', \sigma', \varepsilon)$;

(ii) $\text{imaj}(\Sigma, \varepsilon) = \text{inv}(\Sigma', \varepsilon)$ and $\text{irmaj}(\sigma, \varepsilon) = \text{coinv}(\sigma', \varepsilon)$.

Proof. Property (i) follows from (22.32), since $\text{Ligne } \sigma_x = \text{Ligne } \mathbf{i}\Phi\mathbf{i} \sigma_x$ and analogous relations for the other restrictions. Finally,

$$\text{inv}(\Sigma_{\varepsilon|y}, \Sigma_{\varepsilon|x}) = \text{inv}(\Sigma'_{\varepsilon|y}, \Sigma'_{\varepsilon|x}) \text{ and } \text{inv}(\sigma_{\varepsilon|y}, \sigma_{\varepsilon|x}) = \text{inv}(\sigma'_{\varepsilon|y}, \sigma'_{\varepsilon|x}).$$

Then,

$$\begin{aligned} \text{imaj}(\Sigma, \varepsilon) &= \text{maj} \Sigma_{\varepsilon|x}^{-1} + \text{maj} \Sigma_{\varepsilon|y}^{-1} + \text{inv}(\Sigma_{\varepsilon|y}, \Sigma_{\varepsilon|x}) \\ &= \text{inv} \Sigma'_{\varepsilon|x} + \text{inv} \Sigma'_{\varepsilon|y} + \text{inv}(\Sigma'_{\varepsilon|y}, \Sigma'_{\varepsilon|x}) = \text{inv}(\Sigma', \varepsilon); \\ \text{irmaj}(\sigma, \varepsilon) &= \text{rmaj} \sigma_{\varepsilon|x}^{-1} + \text{rmaj} \sigma_{\varepsilon|y}^{-1} + \text{inv}(\sigma_{\varepsilon|y}, \sigma_{\varepsilon|x}) \\ &= \text{coinv} \sigma'_{\varepsilon|x} + \text{coinv} \sigma'_{\varepsilon|y} + \text{inv}(\sigma'_{\varepsilon|y}, \sigma'_{\varepsilon|x}) = \text{coinv}(\sigma, \varepsilon). \quad \square \end{aligned}$$

23. Eulerian Calculus; multi-indexed polynomials

As seen in section 10, the classical Eulerian polynomials $A_n(t) = \sum_{\sigma \in \mathfrak{S}_n} t^{\text{des } \sigma}$ ($n \geq 0$) have the following exponential generating function

$$(23.1) \quad 1 + \sum_{n \geq 1} A_n(t) \frac{u^n}{n!} = \frac{1-t}{-t + \exp(u(t-1))},$$

so that

$$(23.2) \quad 1 + \sum_{n \geq 1} t A_n(t) \frac{u^n}{n!} = \frac{1-t}{1-t \exp(u(1-t))}$$

provides the exponential generating function for the polynomials ${}^0A_n(t) := t A_n(t) = \sum_{\sigma \in \mathfrak{S}_n} t^{1+\text{des } \sigma}$ ($n \geq 1$) and ${}^0A_0(t) := 1$. In the previous three

sections we have worked out various extensions of the rational function occurring in (23.1). In the present one we will rather use the second one, because the major results dealing with multi-indexed Eulerian polynomials have been introduced in that way and have become classical. The change, however, is minor.

23.1. The bi-indexed Eulerian polynomials. In the fraction occurring in (23.2) replace $\exp(u(1-t))$ by the product $e_p((1-t)u) e_q((1-t)v)$ where u, v, p, q are independent variables. Then expand the fraction as a series normalized by products of the form $(p; p)_n (q; q)_m$. We obtain:

$$(23.3) \quad \sum_{\substack{n \geq 0, \\ m \geq 0}} \frac{u^n v^m}{(p; p)_n (q; q)_m} A_{n,m}(t; p, q) = \frac{1-t}{1 - t e_p((1-t)u) e_q((1-t)v)}.$$

To see that $A_{n,m}(t; p, q)$ is a *polynomial* with *integral* coefficients we may rewrite (23.3) as $L = (1 - R)^{-1}$. The identity $L(1 - R) = 1$ yields the recurrence relation

$$(23.4) \quad A_{n,m}(t; p, q) = \sum_{i,j} \begin{bmatrix} n \\ i \end{bmatrix}_p \begin{bmatrix} m \\ j \end{bmatrix}_q A_{i,j}(t; p, q) t(1-t)^{n-i+m-j-1},$$

where $0 \leq i \leq n$, $0 \leq j \leq m$ and $(i, j) \neq (n, m)$. As $A_{0,0}(t; p, q) = 1$, relation (23.4) shows by induction that each $A_{n,m}(t; p, q)$ is a polynomial with integral coefficients. To show that the coefficients are indeed positive and of sum $(n + m)!$ requires more analysis and will be a consequence of the theorem below.

Of course, a product of more than two q -exponentials could have been taken and the results below would have been very similar. The advantage of keeping only two allows a much easier reading.

The second class of polynomials under study will be denoted by $A_{n,m}(t; p)$. There is no risk of confusion with the previous ones, as they are in two variables t and p instead of three. No confusion either with the q -Eulerian polynomials $A_n(t, q)$, as the present ones are double indexed. Those polynomials $A_{n,m}(t; p)$ are defined by

$$(23.5) \quad \sum_{\substack{n \geq 0, \\ m \geq 0}} \frac{u^n v^m}{(p; p)_n m!} A_{n,m}(t; p) = \frac{1 - t}{1 - t e_p((1 - t)u) \exp((1 - t)v)}.$$

Let $q = 1$ in (23.4). Then $A_{n,m}(t; p, q)$ is transformed into a polynomial in two variables t and p . Furthermore, the Gaussian polynomial $\begin{bmatrix} m \\ j \end{bmatrix}_q$ becomes the ordinary binomial coefficient $\binom{m}{j}$. Hence (23.3) itself is transformed into identity (23.5). Thus

$$(23.6) \quad A_{n,m}(t; p) = A_{n,m}(t; p, 1).$$

Notice that when $u = 0$ (resp. $v = 0$) identity (23.5) specializes into the exponential (resp. basic) generating function for the Eulerian polynomials $\mathcal{A}_n(t)$ introduced above (resp. for the polynomials ${}^{\text{inv}}A_n(t, q)$).

The ligne and inverse ligne of route “Ligne” and “Iligne,” that have been defined in the previous sections for permutations, are being partitioned into two parts. Let (n, m) be an ordered pair of nonnegative integers and σ be a permutation of order $n + m$. Define:

$$\begin{aligned} \text{Iligne}_1 \sigma &:= \{r : 1 \leq r \leq n - 1, \sigma^{-1}(r) > \sigma^{-1}(r + 1)\}, \\ \text{Iligne}_2 \sigma &:= \{r : n + 1 \leq r \leq n + m - 1, \sigma^{-1}(r) > \sigma^{-1}(r + 1)\}, \end{aligned}$$

$$\begin{aligned} \text{imaj}_1 \sigma &:= \sum \{r : r \in \text{Iligne}_1 \sigma\}, \\ \text{imaj}_2 \sigma &:= \sum \{r - n : r \in \text{Iligne}_2 \sigma\}, \\ \text{inv}_1 \sigma &:= \#\{(r, r') : 1 \leq r < r' \leq n, \sigma^{-1}(r) > \sigma^{-1}(r')\}, \\ \text{inv}_2 \sigma &:= \#\{(r, r') : n + 1 \leq r < r' \leq n + m, \sigma^{-1}(r) > \sigma^{-1}(r')\}. \end{aligned}$$

When $m = 0$ (resp. $n = 0$), the statistics $\text{imaj}_1 \sigma$ and $\text{inv}_1 \sigma$ (resp. $\text{imaj}_2 \sigma$ and $\text{inv}_2 \sigma$) are the familiar inverse major index and inversion number of the permutation σ , respectively.

Theorem 23.1. *Let n, m be two nonnegative integers. The generating polynomial for the group \mathfrak{S}_{n+m} of the permutations of order $n + m$ by the three-variable statistic $(1 + \text{des}, \text{imaj}_1, \text{imaj}_2)$ is equal to $A_{n,m}(t; p, q)$. In other words, if $A_{n,m}(t; p, q)$ is defined by identity (23.3), then*

$$(23.7) \quad A_{n,m}(t; p, q) = \sum_{\sigma \in \mathfrak{S}_{n+m}} t^{1+\text{des } \sigma} p^{\text{imaj}_1 \sigma} q^{\text{imaj}_2 \sigma}.$$

We also have:

$$(23.8) \quad A_{n,m}(t; p, q) = \sum_{\sigma \in \mathfrak{S}_{n+m}} t^{1+\text{des } \sigma} p^{\text{inv}_1 \sigma} q^{\text{inv}_2 \sigma}.$$

For the proof of (23.7) we use the Schur function algebra and the Robinson-Schensted correspondence, as was already done for the proof of Theorem 19.4. Take up again the previous notations with a pair (n, m) of integers and σ designating a permutation of order $n + m$. Let τ_1 (resp. τ_2) be the restriction of σ to the set $\sigma^{-1}([1, n])$ (resp. $\sigma^{-1}([n + 1, n + m])$). Using the Robinson-Schensted correspondence (see § Corollary 19.2) each bijection τ_j ($j = 1, 2$) is mapped onto a pair (P_j, Q_j) of Young tableaux of the same *shape* that we shall denote by λ_j .

It follows from the properties of the correspondence that the entries of P_1 (resp. P_2) are the elements of the interval $[1, n]$ (resp. $[n + 1, n + m]$) and the entries of Q_1 (resp. Q_2) are those of the set $\sigma^{-1}([1, n])$ (resp. $\sigma^{-1}([n + 1, n + m])$). Let $T_1 := P_1$ and T_2 be the Young tableau obtained from P_2 by replacing each entry r by $r - n$ and let U be the product $U = Q_1 \otimes Q_2$. This means that U is the skew tableau obtained by placing Q_2 to the right of Q_1 and just *under* it. Thus T_1 (resp. T_2) is a Young tableau of shape λ_1 (resp. λ_2) whose entries are $1, 2, \dots, n$ (resp. $1, 2, \dots, m$); the entries of U are $1, 2, \dots, n$ and the shape of U is the skew shape $\lambda_1 \otimes \lambda_2$.

We summarize all this by writing :

$$(23.9) \quad \begin{aligned} \text{shape } T_j &= \lambda_j \quad (j = 1, 2); \quad |\lambda_1| = n; \quad |\lambda_2| = m; \\ \text{shape } U &= \lambda_1 \otimes \lambda_2; \quad |\lambda_1| + |\lambda_2| = n + m. \end{aligned}$$

For instance, let $(n, m) = (5, 4)$ and

$$\sigma = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 \\ 8 & 6 & 4 & 5 & 2 & 1 & 9 & 7 & 3 \end{pmatrix}; \quad \sigma^{-1} = \left(\begin{array}{ccccc|ccccc} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 \\ 6 & 5 & 9 & 3 & 4 & 2 & 8 & 1 & 7 \end{array} \right).$$

Then

$$\tau_1 = \begin{pmatrix} 3 & 4 & 5 & 6 & 9 \\ 4 & 5 & 2 & 1 & 3 \end{pmatrix}; \quad \tau_2 = \begin{pmatrix} 1 & 2 & 7 & 8 \\ 8 & 6 & 9 & 7 \end{pmatrix}.$$

Under the Robinson-Schensted correspondence

$$\begin{aligned} \tau_1 \mapsto (P_1, Q_1) &= \begin{pmatrix} 4 & 6 \\ 2 & 5 & 5 & 9 \\ 1 & 3 & 3 & 4 \end{pmatrix}; \quad \lambda_1 = (2, 2, 1); \\ \tau_2 \mapsto (P_2, Q_2) &= \begin{pmatrix} 8 & 9 & 2 & 8 \\ 6 & 7 & 1 & 7 \end{pmatrix}; \quad \lambda_2 = (2, 2); \end{aligned}$$

and

$$\begin{array}{ccccccc} & 4 & & & & & 6 \\ & & 2 & 5 & & & & 5 & 9 \\ T_1 = P_1 = 1 & 3 & ; & T_2 = P_2 - n = 1 & 2 & ; & U = Q_1 \otimes Q_2 = 3 & 4 & \\ & & & & & & & & 2 & 8 \\ & & & & & & & & & 1 & 7 \end{array}.$$

The inverse ligne of route (see section 18) of U is $\text{lligne } U = \{1, 2, 4, 5, 7, 8\}$, so that $\text{imaj } U = 27$ and $\text{ides } U = 6$.

It follows from Theorem 19.3 that the mapping

$$(\sigma, n, m) \mapsto (\lambda_1, \lambda_2; T_1, T_2; U)$$

is a bijection having properties (23.9) and also

$$(23.10) \quad \begin{aligned} \text{lligne}_1 \sigma &= \text{lligne } T_1 & \text{lligne}_2 \sigma - n &= \text{lligne } T_2; \\ \text{Ligne } \sigma &= \text{lligne } U. \end{aligned}$$

With our working example $\text{lligne}_1 \sigma = \{1, 3\} = \text{lligne } T_1$; $\text{lligne}_2 \sigma = \{2\} = \text{lligne } T_2$; $\text{Ligne } \sigma = \{1, 2, 4, 5, 7, 8\} = \text{lligne } U$.

It follows from (23.10) that

$$(23.11) \quad \text{imaj}_j \sigma = \text{imaj } T_j \quad (j = 1, 2); \quad \text{des } \sigma = \text{ides } U.$$

23. EULERIAN CALCULUS: MULTI-INDEXED POLYNOMIALS

Let $t {}^1A_n(t; p, q)$ denote the right-hand side of (23.7). Then by (23.10) and (23.11)

$$(23.12) \quad {}^1A_n(t; p, q) = \sum_{(\lambda_1, \lambda_2)} \sum_{(T_1, T_2, U)} t^{\text{idex } U} p^{\text{imaj } T_1} q^{\text{imaj } T_2},$$

where the first sum is over the ordered pairs (λ_1, λ_2) of partitions satisfying $|\lambda_1| = n$, $|\lambda_2| = m$, and the second over all triples (T_1, T_2, U) satisfying (23.9).

We next use identity (18.7) as well as a specialization of (18.8) obtained by letting $q = 1$, i.e.,

$$(23.13) \quad \sum_{r \geq 0} t^r s_{\nu/\theta}(1^{r+1}) = \frac{1}{(1-t)^{n+1}} \sum_{T, T \text{ of shape } \nu/\theta} t^{\text{idex } T},$$

where $s_{\nu/\theta}(1^{r+1})$ is the skew Schur function obtained by taking an alphabet of $(r+1)$ letters all equal to 1. We have :

$$\begin{aligned} & \frac{{}^1A_n(t; p, q)}{(1-t)^{n+m+1}(p; p)_n(q; q)_m} \\ &= \frac{1}{(1-t)^{n+m+1}(p; p)_n(q; q)_m} \sum_{(T_1, T_2, U)} t^{\text{idex } U} p^{\text{imaj } T_1} q^{\text{imaj } T_2} \\ &= \sum_{(\lambda_1, \lambda_2)} \sum_r t^r s_{\lambda_1 \otimes \lambda_2}(1^{r+1}) s_{\lambda_1}(1, p, p^2, \dots) s_{\lambda_2}(1, q, q^2, \dots) \\ &= \sum_{(\lambda_1, \lambda_2)} \sum_r t^r s_{\lambda_1}(1^{r+1}) s_{\lambda_1}(1, p, p^2, \dots) s_{\lambda_2}(1^{r+1}) s_{\lambda_2}(1, q, q^2, \dots), \end{aligned}$$

as U is of shape $\lambda_1 \otimes \lambda_2$. In the last step we have used the fundamental multiplicative property of the Schur functions : $s_{\lambda \otimes \mu}(x) = s_{\lambda}(x) s_{\mu}(x)$. Now, the Cauchy identity (16.4) for Schur functions yields

$$\begin{aligned} \sum_{\lambda_1} u^{|\lambda_1|} s_{\lambda_1}(1^{r+1}) s_{\lambda_1}(1, p, p^2, \dots) &= \prod_{d \geq 1} \frac{1}{(1 - up^{d-1})^{r+1}} = \frac{1}{(u; p)_{\infty}^{r+1}}; \\ \sum_{\lambda_2} v^{|\lambda_2|} s_{\lambda_2}(1^{r+1}) s_{\lambda_2}(1, q, q^2, \dots) &= \prod_{d \geq 1} \frac{1}{(1 - vq^{d-1})^{r+1}} = \frac{1}{(v; q)_{\infty}^{r+1}}. \end{aligned}$$

As $|\lambda_1| = m$, $|\lambda_2| = n$, we may write

$$\sum_{n, m} \frac{u^n v^m}{(1-t)^{n+m+1}(p; p)_n(q; q)_m} {}^1A_n(t; p, q)$$

$$\begin{aligned}
&= \sum_{n,m} u^n v^m \sum_{(\lambda_1, \lambda_2)} \sum_r t^r s_{\lambda_1}(1^{r+1}) s_{\lambda_1}(1, p, \dots) s_{\lambda_2}(1^{r+1}) s_{\lambda_2}(1, q, \dots) \\
&= \sum_r t^r \sum_{(\lambda_1, \lambda_2)} u^{|\lambda_1|} v^{|\lambda_2|} s_{\lambda_1}(1^{r+1}) s_{\lambda_1}(1, p, \dots) s_{\lambda_2}(1^{r+1}) s_{\lambda_2}(1, q, \dots) \\
&= \sum_r t^r \left(\sum_{\lambda_1} u^{|\lambda_1|} s_{\lambda_1}(1^{r+1}) s_{\lambda_1}(1, p, \dots) \right) \\
&\quad \times \left(\sum_{\lambda_2} v^{|\lambda_2|} s_{\lambda_2}(1^{r+1}) s_{\lambda_2}(1, q, \dots) \right) \\
&= \sum_r t^r \frac{1}{(u; p)_\infty^{r+1}} \frac{1}{(v; q)_\infty^{r+1}} = \frac{1}{-t + (u; p)_\infty (v; q)_\infty}.
\end{aligned}$$

Next replace $u/(1-t)$ by u and $v/(1-t)$ by v . This yields

$$\begin{aligned}
\sum_{n,m} \frac{u^n v^m}{(p; q)_n (q; q)_m} {}^1A_n(t; p, q) &= \frac{1-t}{-t + ((1-t)u; p)_\infty ((1-t)v; q)_\infty} \\
&= \frac{1-t}{-t + E_p((t-1)u) E_q((t-1)v)},
\end{aligned}$$

using the notation of the second q -exponential. Since $E_p(u) e_p(-u) = 1$, we obtain:

$$\begin{aligned}
1 + \sum_{\substack{(n,m) \\ \neq (0,0)}} \frac{u^n v^m}{(p; q)_n (q; q)_m} A_n(t; p, q) &= t \frac{1-t}{-t + E_p((t-1)u) E_q((t-1)v)} - t + 1 \\
&= \frac{1-t}{1 - t e_p((1-t)u) e_q((1-t)v)}.
\end{aligned}$$

This proves identity (23.7).

There are two methods for proving identity (23.8): first, the *iterative method* that has already been used for the proofs of Theorems 10.1, 20.3 and 21.1. The recurrence relation (23.4) is to be considered and the summand to be combinatorially interpreted by introducing the notion of the longest increasing rightmost factor. As one of the previous proofs could be reproduced almost verbatim, the proof is omitted. The second proof consists of constructing a bijection $\sigma \mapsto \sigma'$ of \mathfrak{S}_{n+m} onto itself having the property that

$$\text{Ligne } \sigma = \text{Ligne } \sigma', \quad \text{imaj}_j \sigma = \text{inv}_j \sigma' \quad (j = 1, 2).$$

Again, the bijection Φ of Theorem 11.2 could also be used in the same way as was done in Proposition 22.6. As the similarity is so strong, the construction is omitted. \square

23.2. *The Désarménien Verfahren.* This method, based on the algebra of symmetric functions, is quite effective for deriving congruence properties of certain polynomials, especially the Eulerian polynomials and also the bi-indexed Eulerian polynomials introduced in the previous subsection.

Take up again the notations introduced in § 14.3, especially formulas (ii) and (ix) of Theorem 14.2, that merge into

$$(23.14) \quad H(u; x) = \prod_{j \geq 1} (1 - ux_j)^{-1} = \exp \sum_{r \geq 1} u^r \frac{\mathbf{p}_r(x)}{r},$$

where the “ p ” has been written in boldface as “ \mathbf{p} ” to avoid confusion with a further base denoted by “ p .” Also recall that a partition of an integer n is a non-increasing sequence $\lambda = (\lambda_1, \lambda_2, \dots)$ or a word $\lambda = 1^{m_1} 2^{m_2} \dots$ (the multiplicative notation) with the meaning that λ has m_1 parts λ_i equal to 1, m_2 parts λ_i equal to 2, etc. As usual, to each partition λ is attached the constant

$$z_\lambda = 1^{m_1} 2^{m_2} \dots m_1! m_2! \dots$$

and the *power symmetric function*

$$\mathbf{p}_\lambda(x) = \mathbf{p}_{\lambda_1}(x) \mathbf{p}_{\lambda_2}(x) \dots = \mathbf{p}_1(x)^{m_1} \mathbf{p}_2(x)^{m_2} \dots$$

Finally, $|\lambda| = n$ means that λ is a partition of n and the notation $l(\lambda)$ stands for the number of parts of λ .

Now for each partition $\lambda = 1^{m_1} 2^{m_2} \dots$ of the integer n introduce the polynomial:

$$(23.15) \quad T_\lambda(q) := \frac{(q; q)_n}{\prod_j (1 - q^j)^{m_j}} = (q; q)_n \mathbf{p}_\lambda(1, q, q^2, \dots).$$

By induction on the number of *distinct* parts of λ it is easily verified that $T_\lambda(q)$ is a polynomial of degree $n(n - 1)/2$. In the following lemma needed for deriving congruences properties for the polynomials $T_\lambda(q)$ we denote the k -th cyclotomic polynomial by $\Phi_k(q)$ with the convention that $\Phi_1(q) = 1 - q$.

Lemma 23.2. *Let $n = ka + b$ ($0 \leq b \leq k - 1$). Then*

$$\frac{(q; q)_n}{(1 - q^k)^a} \equiv k^a a! (q; q)_b \pmod{\Phi_k(q)}.$$

Proof. Write $(q; q)_n = \left(\prod_{0 \leq j \leq a-1} (q^{kj+1}; q)_{k-1} \right) (q^k; q^k)_a (q^{ka+1}; q)_b$. As $q^k \equiv 1 \pmod{\Phi_k(q)}$, we have $(q^{ka+1}; q)_b \equiv (q; q)_b$ and $(q^{kj+1}; q)_{k-1} \equiv (q; q)_{k-1}$.

When $q = \zeta$ is a k -th primitive root of unity, $(\zeta; \zeta)_{k-1} = \prod_{1 \leq l \leq k-1} (1 - \zeta^l)$, which is the value in $X = 1$ of the polynomial $(1 - X^k)/(1 - X) = 1 + X + \dots + X^{k-1}$, that is, k . Therefore, $(q; q)_{k-1} \equiv k \pmod{\Phi_k(q)}$. Moreover, $(q^k; q^k)_a$ is divisible by $(1 - q^k)^a$ and $\lim_{q \rightarrow \zeta} \frac{(q; q)_a}{(1 - q^a)} = a!$ \square

Proposition 23.3. *Let $n = ka + b$, $0 \leq b \leq k - 1$ and $\lambda = 1^{m_1} 2^{m_2} \dots$ be a partition of n . Then the following congruences hold :*

- (i) *if $m_k \neq a$, then $T_\lambda(q) \equiv 0 \pmod{\Phi_k(q)}$.*
- (ii) *if $m_k = a$, let $\lambda^* = 1^{m_1} \dots (k - 1)^{m_{k-1}} (k + 1)^{m_{k+1}} \dots$ be the partition of b derived from λ by deleting the $m_k = a$ parts equal to k . Then*

$$(23.16) \quad T_\lambda(q) \equiv k^a a! T_{\lambda^*}(q) \pmod{\Phi_k(q)}.$$

Proof. The proposition is an immediate consequence of the previous lemma and the definition of $T_\lambda(q)$ given in (23.15). \square

Let $F(u)$ be a formal power series in one variable u . As the two series $e_q(u)e^v - 1$ and $e_q(u)e_p(v) - 1$ have no constant term, it makes sense to expand the following two series as

$$(23.17) \quad F(e_q(u)e^v - 1) = \sum_{n \geq 0, m \geq 0} \frac{u^n}{(q; q)_n} \frac{v^m}{m!} K_{n,m}(q);$$

$$(23.18) \quad F(e_q(u)e_p(v) - 1) = \sum_{n \geq 0, m \geq 0} \frac{u^n}{(q; q)_n} \frac{v^m}{(p; p)_m} K_{n,m}(p, q).$$

Theorem 23.4. *Let $n = ka + b$ with $0 \leq b \leq k - 1$ and $m \geq 0$. Then*

$$(23.19) \quad K_{ka+b,m}(q) \equiv K_{b,m+a}(q) \pmod{\Phi_k(q)}.$$

Furthermore, if $0 \leq m \leq k - 1$, then

$$(23.20) \quad K_{ka+b,m}(q, p) \equiv K_{b,m+ka}(q, p) \pmod{\Phi_k(q)} \pmod{\Phi_k(p)},$$

where the double congruence means that the two congruences are to be taken in succession.

Notice that the subscripts in the polynomials K are different in (23.19) and (23.20), so that the first congruence is not a specialization of the second one.

23. EULERIAN CALCULUS: MULTI-INDEXED POLYNOMIALS

Proof. Consider the following expansion in the variables u and v

$$\begin{aligned} \left(\sum_{r \geq 1} u^r \frac{\mathbf{p}_r(x)}{r} + v \right)^N &= \sum_{m=0}^N \binom{N}{m} v^m \left(\sum_{r \geq 1} u^r \frac{\mathbf{p}_r(x)}{r} \right)^{N-m} \\ &= \sum_{m=0}^n \binom{N}{m} v^m \sum_{n \geq 0} u^n \sum_{m_1, m_2, \dots} \frac{(N-m)!}{1!^{m_1} 2!^{m_2} \dots} \frac{\mathbf{p}_1(x)^{m_1}}{1^{m_1}} \frac{\mathbf{p}_2(x)^{m_2}}{2^{m_2}} \dots \\ &\quad [n+m \geq N, m_1 + m_2 + \dots = N-m; 1.m_1 + 2.m_2 + \dots = n.] \\ &= N! \sum_{0 \leq m \leq N \leq n+m} \frac{v^m}{m!} u^n \sum_{\substack{|\lambda|=n \\ l(\lambda)=N-m}} \frac{\mathbf{p}_\lambda(x)}{z_\lambda}. \end{aligned}$$

Replacing the alphabet x by the successive powers of q yields

$$(23.21) \quad \left(\sum_{r \geq 1} u^r \frac{\mathbf{p}_r(1, q, q^2, \dots)}{r} + v \right)^N = \sum_{0 \leq m \leq N \leq n+m} \frac{u^n}{(q; q)_n} \frac{v^m}{m!} K_{n,m}^{(N)}(q),$$

where

$$(23.22) \quad K_{n,m}^{(N)}(q) = N! \sum_{\substack{|\lambda|=n \\ l(\lambda)=N-m}} \frac{T_\lambda(q)}{z_\lambda}.$$

When $m_k = a$, then $l(\lambda) = l(\lambda^*) + a$ and $z_\lambda = z_{\lambda^*} k^a a!$, so that

$$K_{n,m}^{(N)}(q) \equiv N! \sum_{\substack{|\lambda^*=b \\ l(\lambda^*)=N-m-a}} \frac{T_{\lambda^*}(q)}{z_{\lambda^*}} \pmod{\Phi_k(q)}.$$

In particular by (23.22)

$$K_{b,m+a}^{(N)}(q) = N! \sum_{\substack{|\lambda^*=b \\ l(\lambda^*)=N-m-a}} \frac{T_{\lambda^*}(q)}{z_{\lambda^*}}.$$

Therefore

$$(23.23) \quad K_{ka+b,m}^{(N)}(q) \equiv K_{b,m+a}^{(N)}(q) \pmod{\Phi_k(q)}.$$

Now, by (23.14),

$$e_q(u)e^v = \prod_{j \geq 1} (1 - uq^{j-1})^{-1} e^v = \exp \left(\sum_{r \geq 1} u^r \frac{\mathbf{p}_r(1, q, q^2, \dots)}{r} + v \right),$$

so that, if $F(u) = \sum_{i \geq 0} c_i u^i$, we get

$$\begin{aligned}
 F(e_q(u)e^v - 1) &= \sum_{i \geq 0} c_i (e_q(u)e^v - 1)^i \\
 &= \sum_{i \geq 0} c_i \left(\exp \left(\sum_{r \geq 1} u^r \frac{\mathbf{p}_r(1, q, q^2, \dots)}{r} + v \right) - 1 \right)^i \\
 &= \sum_{N \geq 0} \frac{C_N}{N!} \left(\sum_{r \geq 1} u^r \frac{\mathbf{p}_r(1, q, q^2, \dots)}{r} + v \right)^N \\
 &\quad \text{[for some coefficients } C_N \text{ (} N \geq 0 \text{);]} \\
 &= \sum_{N \geq 0} \frac{C_N}{N!} \sum_{0 \leq m \leq N \leq n+m} \frac{u^n}{(q; q)_n} \frac{v^m}{m!} K_{n,m}^{(N)}(q) \\
 &= \sum_{n \geq 0, m \geq 0} \frac{u^n}{(q; q)_n} \frac{v^m}{m!} K_{n,m}(q),
 \end{aligned}$$

where $K_{n,m}(q) = \sum_{N=0}^{n+m} \frac{C_N}{N!} K_{n,m}^{(N)}(q)$, since the formal series (23.21) is of order N in the variables u and v . Hence, the congruence property for $K_{n,m}(q)$ is a consequence of the congruence property (23.23) satisfied by the polynomials $K_{n,m}^{(N)}(q)$ themselves.

The proof of (23.20) is quite similar. This time we start with the expansion of the series $\sum_{r \geq 1} \left(\frac{u^r \mathbf{p}_r(x) + v^r \mathbf{p}_r(y)}{r} \right)^N$, where y stands for a second alphabet. We have

$$\begin{aligned}
 \left(\sum_{r \geq 1} \frac{u^r \mathbf{p}_r(x) + v^r \mathbf{p}_r(y)}{r} \right)^N &= \sum_{A+B=N} \binom{N}{A} \left(\sum_{r \geq 1} u^r \frac{\mathbf{p}_r(x)}{r} \right)^A \left(\sum_{r \geq 1} v^r \frac{\mathbf{p}_r(y)}{r} \right)^B \\
 &= \sum_{A+B=N} \binom{N}{A} \left(\sum_{l(\lambda)=A} A! \frac{u^{|\lambda|} \mathbf{p}_\lambda(x)}{z_\lambda} \right) \left(\sum_{l(\mu)=B} B! \frac{v^{|\mu|} \mathbf{p}_\mu(y)}{z_\mu} \right) \\
 &= N! \sum_{\substack{n \geq 0, m \geq 0 \\ n+m \geq N}} u^n v^m \sum_{\substack{|\lambda|=n, |\mu|=m \\ l(\lambda)+l(\mu)=N}} \frac{\mathbf{p}_\lambda(x)}{z_\lambda} \frac{\mathbf{p}_\mu(y)}{z_\mu}.
 \end{aligned}$$

Again, if we replace the alphabet x (resp. y) by the successive powers of q (resp. of p), we obtain

$$\left(\sum_{r \geq 1} \frac{u^r \mathbf{p}_r(1, q, \dots) + v^r \mathbf{p}_r(1, p, \dots)}{r} \right)^N = \sum_{\substack{n \geq 0, m \geq 0 \\ n+m \geq N}} \frac{u^n}{(q; q)_n} \frac{v^m}{(p; p)_m} K_{n,m}^{(N)}(q, p),$$

where

$$(23.24) \quad K_{n,m}^{(N)}(q, p) = N! \sum_{\substack{|\lambda|=n, |\mu|=m \\ l(\lambda)+l(\mu)=N}} \frac{T_\lambda(q)}{z_\lambda} \frac{T_\mu(p)}{z_\mu}.$$

In the last summation the sum of the lengths of the two partitions is to remain constant, equal to N . We then use the bijection $(\lambda, \mu) \mapsto (\lambda^*, \mu^+)$, where μ^+ is derived from μ by *adding* a parts equal to k . We have

$$\begin{aligned} K_{ka+b,m}^{(N)}(q, p) &\equiv N! \sum_{\substack{|\lambda^*|=b, |\mu|=m+ka \\ l(\lambda^*)+l(\mu^+)=N}} \frac{T_{\lambda^*}(q)}{z_{\lambda^*}} \frac{T_{\mu^+}(p)}{z_{\mu^+}} \pmod{\Phi_k(q)} \pmod{\Phi_k(p)} \\ &\equiv K_{b,m+ka}^{(N)}(q, p). \end{aligned}$$

For the coefficients $K_{n,m}(q, p)$ of the series $F(e_p(u)e_q(v) - 1)$ we also find the expression

$$K_{n,m}(q, p) = \sum_{N=0}^{n+m} \frac{C_N}{N!} K_{n,m}^{(N)}(q, p)$$

for some coefficients C_N ($N \geq 0$). The congruence property for the $K_{n,m}(q, p)$ follows from (23.24). \square

23.3. Congruences for bi-indexed polynomials. We apply the techniques of the previous subsection to the sequence of the bi-indexed Eulerian polynomials (in *two* variables) $A_{n,m}(t, q)$ defined by identity (23.5). Notice that it has two specializations

$$(23.25) \quad A_{n,0}(t, q) = \text{inv}A_n(t, q), \quad A_{0,m}(t, q) = {}^0A_m(t),$$

where $\text{inv}A_n(t, q)$ is the q -inv-Eulerian polynomial introduced in section 10 and ${}^0A_m(t) = 1$ if $m = 0$ and $tA_m(t)$ (with $A_m(t)$ being the traditional Eulerian polynomial) if $m \geq 1$. In particular, $A_{1,0}(t, q) = A_{0,1}(t, q) = t$. The recurrence relation (23.4) with $q = 1$ provides the recurrence relation for the polynomials $A_{n,m}(t, q)$ and shows by induction that

$$(23.26) \quad A_{1,m}(t, q) = \text{inv}A_{0,m+1}(t, q) = {}^0A_{m+1}(t) \quad (m \geq 0).$$

Theorem 23.5. *Let n and k be two positive integers and let $n = ka + b$, $0 \leq b \leq k - 1$ be the Euclidean division of n by k . Then the following congruence holds:*

$$(23.27) \quad A_{ka+b,m}(t, q) \equiv (1-t)^{(k-1)a} A_{b,m+a}(t, q) \pmod{\Phi_k(q)}.$$

Furthermore, if $0 \leq m \leq k - 1$, then

$$(23.28) \quad A_{ka+b,m}(t; q, p) \equiv A_{b,m+ka}(t; q, p) \pmod{\Phi_k(q)} \pmod{\Phi_k(p)},$$

For $k = 1$, $m = 1$:

$$(23.29) \quad A_{a,1}(t, q) = A_{0,a+1}(t, q) = {}^0A_{a+1}(t);$$

for $k = 2$, $b = 0, 1$, $m = 0$:

$$(23.30) \quad \begin{aligned} \text{inv}A_{2a+b}(t, q) &= A_{0,2a+b} \equiv (1-t)^a A_{b,a}(t, q) \pmod{(1+q)} \\ &\equiv (1-t)^a {}^0A_{a+b}(t) \pmod{(1+q)}. \end{aligned}$$

Proof. The generating function for the polynomials $A_{n,m}(t, q)$ may be rewritten as

$$\sum_{n \geq 0, m \geq 0} \frac{A_{n,m}(t, q)}{(1-t)^{n+m}} \frac{u^n}{(q; q)_n} \frac{v^m}{m!} = \sum_{i \geq 0} \left(\frac{t}{1-t} \right)^i (e_q(u) \exp v - 1)^i.$$

By Proposition 23.4 the coefficients $K_{n,m}(q) := A_{n,m}(t, q)/(1-t)^{n+m}$ satisfy the congruence (23.19):

$$\frac{A_{ka+b,m}(t, q)}{(1-t)^{ka+b+m}} \equiv \frac{A_{b,m+a}(t, q)}{(1-t)^{b+m+a}} \pmod{\Phi_k(q)}.$$

In an analogous manner

$$\sum_{n \geq 0, m \geq 0} \frac{A_{n,m}(t; q, p)}{(1-t)^{n+m}} \frac{u^n}{(q; q)_n} \frac{v^m}{(p; p)_m} = \sum_{i \geq 0} \left(\frac{t}{1-t} \right)^i (e_q(u)e_p(u) - 1)^i$$

and congruence (23.20) applies. \square

23.4. *The signed Eulerian numbers.* Consider the q -inv-Eulerian polynomials $\text{inv}A_n(t, q) = \sum_{\sigma \in \mathfrak{S}_n} t^{1+\text{des } \sigma} q^{\text{inv } \sigma}$ introduced in section 10 and define the *signed Eulerian polynomial* to be

$$(23.30) \quad \text{sgn}A_n(t) := \text{inv}A_n(t, -1) := \sum_{k=1}^n \text{sgn}A_{n,k} t^k.$$

The integers $\text{sgn}A_{n,k}$ ($1 \leq k \leq n$) are called the *signed Eulerian numbers*. Relation (23.30) shows that

$$(23.31) \quad \text{sgn}A_{n,k} = \sum_{\sigma} \text{sgn } \sigma,$$

where the sum ranges over all permutations $\sigma \in \mathfrak{S}_n$ such that $1+\text{des } \sigma = k$. Their first values are displayed in Table 23.1. Recall that ${}^0A_n(t)$ is the generating polynomial for \mathfrak{S}_n by the statistic “ $1 + \text{des}$ ” (see section 10).

23. EULERIAN CALCULUS: MULTI-INDEXED POLYNOMIALS

$k =$	1	2	3	4	5	6	7
$n = 1$	1						
2	1	-1					
3	1	0	-1				
4	1	-1	-1	1			
5	1	2	-6	2	1		
6	1	1	-8	8	-1	-1	
7	1	8	-19	0	19	-8	-1

Table 23.1

Theorem 23.10. Let ${}^{\text{sgn}}A_n(t)$ be the signed Eulerian polynomial defined in (23.30). Then

$$(23.32a) \quad {}^{\text{sgn}}A_{2n}(t) = (1-t)^n {}^0A_n(t);$$

$$(23.32b) \quad {}^{\text{sgn}}A_{2n+1}(t) = (1-t)^n {}^0A_{n+1}(t).$$

Proof. This has already been proved in Theorem 23.5 with relation (23.30), since ${}^{\text{sgn}}A_n(t) \equiv {}^{\text{inv}}A_n(t, q) \pmod{1+q}$. Another proof consists in starting with the recurrence relation (10.11) for the polynomials ${}^{\text{inv}}A_n(t, q)$ and let $q \mapsto -1$ in that relation. As

$$\begin{aligned} \lim_{q \rightarrow -1} \begin{bmatrix} 2m \\ 2i \end{bmatrix}_q &= \lim_{q \rightarrow -1} \begin{bmatrix} 2m+1 \\ 2i \end{bmatrix}_q = \lim_{q \rightarrow -1} \begin{bmatrix} 2m+1 \\ 2i+1 \end{bmatrix}_q = \binom{m}{i}, \\ \lim_{q \rightarrow -1} \begin{bmatrix} 2m \\ 2i+1 \end{bmatrix}_q &= 0, \end{aligned}$$

we are led to the recurrences

$$(23.33) \quad {}^{\text{sgn}}A_{2n}(t) = t(1-t)^{2n-1} + \sum_{1 \leq i \leq n-1} \binom{n}{i} {}^{\text{sgn}}A_{2i}(t) t(1-t)^{2n-1-2i};$$

$$(23.34) \quad {}^{\text{sgn}}A_{2n+1}(t) = t(1-t)^{2n} + \sum_{1 \leq i \leq n} \binom{n}{i} {}^{\text{sgn}}A_{2i}(t) t(1-t)^{2n-2i} \\ + \sum_{0 \leq i \leq n-1} \binom{n}{i} {}^{\text{sgn}}A_{2i+1}(t) t(1-t)^{2n-2i-1}.$$

When $q \rightarrow 1$ in (10.11) we get the recurrence relation for the polynomials ${}^0A_n(t)$, i.e.,

$$(23.35) \quad {}^0A_n(t) = t(1-t)^{n-1} + \sum_{1 \leq i \leq n-1} \binom{n}{i} {}^0A_i(t) t(1-t)^{n-1-i}.$$

By comparing (23.33) and (23.35) we see that the polynomials $(1 - t)^n A_n(t)$ satisfy the same recurrence relation as the polynomials ${}^{\text{sgn}}A_{2n}(t)$. Hence (23.32a) holds. To establish (23.32b) an easy induction on n suffices. \square

Having the classical relations for the (unsigned) Eulerian numbers $A_{n,k}$ at our disposal, as derived in (10.5)–(10.9), it is straightforward from Theorem 23.10 to prove the following relations:

$$(23.36a) \quad {}^{\text{sgn}}A_{2n,k} = {}^{\text{sgn}}A_{2n-1,k} - {}^{\text{sgn}}A_{2n-1,k-1};$$

$$(23.36b) \quad {}^{\text{sgn}}A_{2n+1,k} = k {}^{\text{sgn}}A_{2n,k} + (2n - k + 2) {}^{\text{sgn}}A_{2n,k-1};$$

together with the analogs of the Worpitzky formulas

$$\sum_i \binom{2n+i}{i} {}^{\text{sgn}}A_{2n,k-i} = k^n;$$

$$\sum_i \binom{2n-1+i}{i} {}^{\text{sgn}}A_{2n-1,k-i} = k^n;$$

and the polynomial relations

$${}^{\text{sgn}}A_{2n}(t) = (1 - t) {}^{\text{sgn}}A_{2n-1}(t);$$

$${}^{\text{sgn}}A_{2n+1}(t) = (2n + 1) t {}^{\text{sgn}}A_{2n}(t) + t(1 - t) {}^{\text{sgn}}A'_{2n}(t);$$

where ${}^{\text{sgn}}A'_{2n}(t)$ denotes the derivative of ${}^{\text{sgn}}A_{2n}(t)$.

24. The basic and bibasic trigonometric functions

By analogy with classical analysis start with the traditional Euler identity

$$(24.1) \quad e_L = \cos_L + i \sin_L,$$

where e_L is an analog of the traditional complex exponential. When we choose $e_L = e_q(iu)$ (resp. $e_L = e_p(iu)e_q(iv)$), we obtain the definitions of the q -sine, $\sin_q(u)$, and q -cosine, $\cos_q(u)$, (resp. of the p, q -sine, $\sin_{p,q}(u, v)$, and p, q -cosine, $\cos_{p,q}(u, v)$.) Of course, the other q -exponential $E_q(u)$ could be used, but this does not lead to significantly different results.

The goal of this section is to define the other functions \tan_L and \sec_L , to derive the recurrence formulas for the coefficients of their expansions and, finally, to obtain combinatorial interpretations for the coefficients.

24.1. *The basic and bibasic tangent and secant functions.* First, recall some easy results on the p - or q -partial derivative. If $A(u, v) =$

$A(u, v; p, q) = \sum_{n \geq 0, m \geq 0} A_{n,m} u^n v^m / ((p; p)_n (q; q)_m)$ is a (p, q) -bibasic series in the variables u and v , define:

$$(24.2) \quad \partial_p A(u, v) := \frac{A(u, v) - A(pu, v)}{u};$$

$$(24.3) \quad \partial_q A(u, v) := \frac{A(u, v) - A(u, qv)}{u}.$$

The following relations are immediate

$$(24.4) \quad \partial_p (A(u, v)B(u, v)) = A(u, v) \partial_p B(u, v) + A(pu, v) \partial_p B(u, v);$$

$$(24.5) \quad \partial_p \frac{A(u, v)}{B(u, v)} = \frac{B(u, v) \partial_p A(u, v) - A(u, v) \partial_p B(u, v)}{B(u, v)B(pu, v)};$$

$$(24.6) \quad = \frac{B(pu, v) \partial_p A(u, v) - A(pu, v) \partial_p B(u, v)}{B(u, v)B(pu, v)};$$

and also the analogous relations for the q -partial derivative. If the series $A(u; p)$ and $B(u; p)$ are p -basic series of the variable u , we speak of p -derivatives. The above properties are similar and will not be restated.

Definition. The p -sine and p -cosine are defined by the identity

$$e_p(iu) =: \cos_p(u) + i \sin_p(u);$$

the p, q -sine and p, q -cosine by the identity

$$e_{p,q}(iu, iv) := e_p(iu)e_q(iv) =: \cos_{p,q}(u) + i \sin_{p,q}(u).$$

Proposition 24.1. *The following identities hold:*

$$(24.7) \quad e_{p,q}(-iu, -iv) = \cos_{p,q}(u, v) - i \sin_{p,q}(u, v);$$

$$(24.8) \quad \cos_{p,q}(u, v) = \frac{e_{p,q}(iu, iv) + e_{p,q}(-iu, -iv)}{2};$$

$$(24.9) \quad \sin_{p,q}(u, v) = \frac{e_{p,q}(iu, iv) - e_{p,q}(-iu, -iv)}{2i};$$

$$(24.10) \quad \partial_p e_{p,q}(\alpha u, v) = \alpha e_{p,q}(\alpha u, v);$$

$$(24.11) \quad \partial_p \sin_{p,q}(\alpha u, v) = \alpha \cos_{p,q}(\alpha u, v);$$

$$(24.12) \quad \partial_p \cos_{p,q}(\alpha u, v) = -\alpha \sin_{p,q}(\alpha u, v);$$

and analogous relations for the q -partial derivative. The identities for $e_p(-iu)$, $\sin_p(u)$, $\cos_p(u)$ are obtained by letting $v = 0$.

Proof. As $e_p(iu) = \cos_p(u) + i \sin_p(u)$, relation (25.7) is proved by direct calculation. Next, (24.8) and (24.9) are banal. For deriving (24.10) remember that $e_p(\alpha u) = \prod_{n \geq 0} (1 - \alpha up^n)^{-1}$, so that $e_p(p\alpha u) =$

$(1 - \alpha u)e_p(\alpha u)$. Then $\partial_p e_p(\alpha u) = \alpha e_p(\alpha u)$. The last two relations hold by linearity. \square

Definition. The p -tangent, p -secant, p, q -tangent and p, q -secant functions are defined by

$$(24.13) \quad \tan_p(u) := \frac{\sin_p(u)}{\cos_p(u)}; \quad \tan_{p,q}(u, v) := \frac{\sin_{p,q}(u, v)}{\cos_{p,q}(u, v)};$$

$$(24.14) \quad \sec_p(u) := \frac{1}{\cos_p(u)}; \quad \sec_{p,q}(u, v) := \frac{1}{\cos_{p,q}(u, v)}.$$

Also define

$$(24.15) \quad \text{Eul}(u; p) := \frac{1 + \sin_p(u)}{\cos_p(u)} = \sec_p(u) + \tan_p(u);$$

$$(24.16) \quad \text{Eul}(u, v; p, q) := \frac{1 + \sin_{p,q}(u, v)}{\cos_{p,q}(u, v)} = \sec_{p,q}(u, v) + \tan_{p,q}(u, v).$$

Theorem 24.2. *The following derivative relations hold:*

$$(24.17) \quad \partial_p \text{Eul}(u; p) = 1 + \tan_p(u) \text{Eul}(pu; p);$$

$$(24.18) \quad \partial_p \text{Eul}(u, v; p, q) = 1 + \tan_{p,q}(u, v) \text{Eul}(pu, v; p, q);$$

$$(24.19) \quad \partial_q \text{Eul}(u, v; p, q) = 1 + \tan_{p,q}(u, v) \text{Eul}(u, qv; p, q).$$

Proof. Relation (25.17) follows from (25.18) or can be proved in a straightforward manner. As $\text{Eul}(u, v; p, q) = \text{Eul}(v, u; q, p)$, it suffices to prove the first of those two relations. By (24.6)

$$\begin{aligned} & \partial_p \frac{1 + \sin_{p,q}(u, v)}{\cos_{p,q}(u, v)} \\ &= \frac{\cos_{p,q}(pu, v) \partial_p(1 + \sin_{p,q}(u, v)) - (1 + \sin_{p,q}(u, v)) \partial_p \cos_{p,q}(u, v)}{\cos_{p,q}(u, v) \cos_{p,q}(pu, q)} \\ &= \frac{\cos_{p,q}(pu, v) \cos_{p,q}(u, v) + \sin_{p,q}(u, v) + \sin_{p,q}(u, v) \sin_{p,q}(pu, v)}{\cos_{p,q}(u, v) \cos_{p,q}(pu, q)} \end{aligned}$$

Thus,

$$\begin{aligned} \partial_p \frac{1 + \sin_{p,q}(u, v)}{\cos_{p,q}(u, v)} &= 1 + \frac{\sin_{p,q}(u, v)}{\cos_{p,q}(u, v)} \frac{1 + \sin_{p,q}(pu, v)}{\cos_{p,q}(pu, v)} \\ &= 1 + \tan_{p,q}(pu, v) \text{Eul}(pu, v; p, q). \quad \square \end{aligned}$$

Denote the expansions of $\text{Eul}(u; p, q)$ and $\text{Eul}(u, v; p, q)$ by

$$(24.20) \quad \text{Eul}(u; p) := \sum_{n \geq 0} \text{Eul}_n(p) \frac{u^n}{(p; p)_n};$$

$$(24.21) \quad \text{Eul}(u, v; p, q) := \sum_{n \geq 0, m \geq 0} \text{Eul}_{n,m}(p, q) \frac{u^n}{(p; p)_n} \frac{v^m}{(q; q)_m};$$

so that

$$(24.22) \quad \tan_p(u) = \text{Eul}^{\text{odd}}(u; p) := \sum_{n \geq 0, n \text{ odd}} \text{Eul}_n(p) \frac{u^n}{(p; p)_n};$$

$$(24.23) \quad \tan_{p,q}(u) = \text{Eul}^{\text{odd}}(u, v; p, q) := \sum_{\substack{n \geq 0, m \geq 0 \\ n+m \text{ odd}}} \text{Eul}_{n,m}(p, q) \frac{u^n}{(p; p)_n} \frac{v^m}{(q; q)_m}.$$

Proposition 24.3. *The coefficients $\text{Eul}_n(p)$ ($n \geq 0$) and accordingly the function $\text{Eul}(u; p)$ are inductively defined by the following two relations:*

- (i) $\text{Eul}_0(p) = \text{Eul}_1(p) = 1$;
- (ii) *the recurrence formula valid for $n \geq 1$*

$$(24.24) \quad \text{Eul}_{n+1}(p) = \sum_{\substack{0 \leq a \leq n \\ a \text{ odd}}} \begin{bmatrix} n \\ a \end{bmatrix}_p p^{n-a} \text{Eul}_a(p) \text{Eul}_{n-a}(p).$$

The coefficients $\text{Eul}_{n,m}(p, q)$ ($n \geq 0, m \geq 0$) and accordingly the function $\text{Eul}(u, v; p, q)$ are inductively defined by the following three relations:

- (iii) $\text{Eul}_{0,0}(p) = \text{Eul}_{0,1}(p, q) = \text{Eul}_{1,0}(p, q) = 1$;
- (iv) *the recurrence formula valid for $n \geq 1$*

$$(24.25) \quad \text{Eul}_{n+1,0}(p, q) = \sum_{\substack{0 \leq a \leq n \\ a \text{ odd}}} \begin{bmatrix} n \\ a \end{bmatrix}_p p^{n-a} \text{Eul}_{a,0}(p, q) \text{Eul}_{n-a,0}(p, q).$$

- (v) *the recurrence formula valid for $n + m \geq 1$*

$$(24.26) \quad \text{Eul}_{n,m+1}(p, q) = \sum_{\substack{0 \leq a \leq n, 0 \leq b \leq m \\ a+b \text{ odd}}} \begin{bmatrix} n \\ a \end{bmatrix}_p \begin{bmatrix} m \\ b \end{bmatrix}_q q^{m-b} \text{Eul}_{a,b}(p, q) \text{Eul}_{n-a,m-b}(p, q).$$

Proof. Go back to identities (24.17) and (24.19), expand both sides and write that coefficients of the same monomial are equal on both sides. \square

When (24.18) is taken into account, conditions (iv) and (v) can be replaced by

(iv') the recurrence formula valid for $m \geq 1$

$$(24.25') \quad \text{Eul}_{0,m+1}(p, q) = \sum_{\substack{0 \leq b \leq m \\ b \text{ odd}}} \begin{bmatrix} m \\ b \end{bmatrix}_q q^{m-b} \text{Eul}_{0,b}(p, q) \text{Eul}_{0,m-b}(p, q).$$

(v') the recurrence formula valid for $n + m \geq 1$

$$(24.26') \quad \text{Eul}_{m+1,n}(p, q) = \sum_{\substack{0 \leq a \leq n, 0 \leq b \leq m \\ a+b \text{ odd}}} \begin{bmatrix} n \\ a \end{bmatrix}_p \begin{bmatrix} m \\ b \end{bmatrix}_q p^{n-a} \text{Eul}_{a,b}(p, q) \text{Eul}_{n-a,m-b}(p, q).$$

Finally, notice that both (24.25) and (24.25') are the same recurrence formulas as (24.24).

As $\text{Eul}_0(p) = \text{Eul}_1(p) = 1$, all the coefficients $\text{Eul}_n(p)$ can be determined by means of formula (24.24). The first values are the following:

$$\begin{aligned} \text{Eul}_0(p) &= \text{Eul}_1(p) = \text{Eul}_2(p) = 1, & \text{Eul}_3(p) &= p(1+p), \\ \text{Eul}_4(p) &= p(1+p)^2 + p^4, & \text{Eul}_5(p) &= p^2(1+p)^2(1+p^2)^2, \\ \text{Eul}_6(p) &= p^2(1+p)^2(1+p^2+p^4)(1+p+p^2+2p^3) + p^{12}, \\ \text{Eul}_7(p) &= p^3(1+p)^2(1+p^2)(1+p^3)(1+p+3p^2+2p^3+3p^4+2p^5+3p^6+p^7+p^8). \end{aligned}$$

The first polynomials $\text{Eul}_{n,m}(p, q)$ can be derived by means of Proposition 24.3. Because of the symmetry we only list the polynomials $\text{Eul}_{n,m}(p, q)$ such that $n \leq m$.

$$\begin{aligned} \text{Eul}_{0,0}(p, q) &= \text{Eul}_{0,1}(p, q) = \text{Eul}_{0,2}(p, q) = \text{Eul}_{1,1}(p, q) = 1; \\ \text{Eul}_{0,3}(p, q) &= q(1+q); & \text{Eul}_{1,2}(p, q) &= 1+q; \\ \text{Eul}_{0,4}(p, q) &= q(1+q)^2 + q^4; & \text{Eul}_{1,3}(p, q) &= 1+2q+2q^2; \\ \text{Eul}_{2,2}(p, q) &= 2+p+q+pq; \\ \text{Eul}_{0,5}(p, q) &= q^2(1+q)^2(1+q^2)^2; & \text{Eul}_{1,4}(p, q) &= 2q(1+q)^2(1+q^2); \\ \text{Eul}_{2,3}(p, q) &= (1+p)(1+q)^3; \\ \text{Eul}_{0,6}(p, q) &= q^2(1+q)^2(1+q^2+q^4)(1+q+q^2+2q^3) + q^{12}; \\ \text{Eul}_{1,5}(p, q) &= (3q^7+6q^6+10q^5+13q^4+12q^3+9q^2+6q+2)q; \\ \text{Eul}_{2,4}(p, q) &= 3q^5+3pq^5+7q^4+6pq^4+7pq^3+8q^3+9q^2+7pq^2+4pq+5q+p+1; \\ \text{Eul}_{3,3}(p, q) &= 2p^3q+2p^3q^2+p^3q^3+p^3+4p^2+8p^2q^2+8p^2q+2p^2q^3+4p+8pq^2+8pq+2pq^3+4q^2+4q+2+q^3; \\ \text{Eul}_{0,7}(p, q) &= q^3(1+q)^2(1+q^2)(1+q^3)(1+q+3q^2+2q^3+3q^4+2q^5+3q^6+q^7+q^8); \\ \text{Eul}_{1,6}(p, q) &= q^2(1+q)^2(1+q^2)(1+q^3)(3q^4+3q^3+5q^2+3q+3); \\ \text{Eul}_{2,5}(p, q) &= q(1+p)(1+q)^2(1+q^2)(3q^4+4q^3+3q^2+4q+3); \end{aligned}$$

$$\begin{aligned} \text{Eul}_{3,4}(p, q) &= (1 + p)(1 + q)^2(1 + q^2) \\ &\quad \times (q^2p^2 + pq^2 + q^2 + 5qp + 3qp^2 + 3q + p^2 + p + 1). \end{aligned}$$

24.2. *Alternating permutations.* We now use formula (24.24) to show that the polynomials $\text{Eul}_n(p)$ are generating polynomials for sets of permutations, called *alternating*, by the statistic “inv” (and also by “imaj”).

Definition. A *rising alternating* (resp. *falling alternating*) permutation is defined to be a permutation $\sigma = \sigma(1) \dots \sigma(n)$ having the following properties:

$$\begin{aligned} &\sigma(1) < \sigma(2), \sigma(2) > \sigma(3), \sigma(3) < \sigma(4), \text{ etc. in an alternating way} \\ &(\text{resp. } \sigma(1) > \sigma(2), \sigma(2) < \sigma(3), \sigma(3) > \sigma(4), \text{ etc. in an alternating way}). \end{aligned}$$

By \mathcal{DR}_n (resp. \mathcal{D}_n) is denoted the set of rising alternating (resp. falling alternating) permutations of order n .

One of the consequences of the following theorem is the fact that the number of rising alternating (resp. falling alternating) permutations of order n is equal to $\text{Eul}_n(1)$. Referring to the table of the polynomials $\text{Eul}_n(q)$ given above, we see that: $\text{Eul}_0(1) = \text{Eul}_1(1) = \text{Eul}_2(1) = 1$, $\text{Eul}_3(1) = 2$, $\text{Eul}_4(1) = 5$, $\text{Eul}_5(1) = 16$, $\text{Eul}_6(1) = 61$. The numbers $\text{Eul}_{2n}(1)$ (resp. $\text{Eul}_{2n+1}(1)$) are called *tangent numbers* (resp. *secant numbers*) and have been combinatorially studied for decades. The polynomials $\text{Eul}_n(q)$ appear as q -analogs of those numbers.

The rising alternating permutations for $n = 1, 2, 3, 4$ are the following: 1 ; 1, 2 ; 1, 3, 2 ; 2, 3, 1 ; 1, 3, 2, 4 ; 1, 4, 2, 3 ; 2, 3, 1, 4 ; 2, 4, 1, 3 ; 3, 4, 1, 2.

Theorem 24.4. *For every $n \geq 0$ the polynomial $\text{Eul}_n(p)$ is the generating function for the rising alternating permutations of length n by the number of inversions. In other words,*

$$\text{Eul}_n(p) = \sum_{\sigma \in \mathcal{DR}_n} p^{\text{inv } \sigma}.$$

When n is odd, we also have

$$\text{Eul}_n(p) = \sum_{\sigma \in \mathcal{D}_n} p^{\text{inv } \sigma}.$$

Proof. The result is banal for $n = 0, 1$ and 2. For $n \geq 2$ consider the set $S_{n+1, a+1}$ of rising alternating permutations σ of order $n + 1$ such that $n + 1$ is in the $(a + 1)$ -st position (i.e., $\sigma(a + 1) = n + 1$) with a odd. Such a permutation is characterized by the two rising alternating subpermutations $\sigma' = \sigma(1) \dots \sigma(a)$ and $\sigma'' = \sigma(a + 2) \dots \sigma(n + 1)$ that do not contain $n + 1$. The inversions of σ fall into four groups:

(a) the inversions that correspond to the pairs of letters whose first letter is in σ' and the second one in σ'' ; it follows from Proposition 4.3 that their generating polynomial is equal to $\begin{bmatrix} n \\ a \end{bmatrix}_p$;

(b) the $n - a$ inversions between $n + 1$ and each letter of σ'' ;

(c) the inversions inside σ' ; by induction their generating polynomial is equal to $\text{Eul}_a(p)$;

(d) the inversions inside σ'' ; their generating polynomial is equal to $\text{Eul}_{n-a}(p)$.

The generating polynomial for $S_{n+1,a+1}$ is then equal to

$$\sum_{\sigma \in S_{n+1,a+1}} q^{\text{inv } \sigma} = \begin{bmatrix} n \\ a \end{bmatrix} q^{n-a} \text{Eul}_a(p) \text{Eul}_{n-a}(p).$$

Hence,

$$\text{Eul}_{n+1}(p) = \sum_{0 \leq a \leq n, a \text{ odd}} \begin{bmatrix} n \\ a \end{bmatrix}_p p^{n-a} \text{Eul}_a(p) \text{Eul}_{n-a}(p),$$

that is exactly the induction formula (24.24).

When n is odd, the transformation \mathbf{rc} , already introduced (see the end of section 12), that sends the permutation σ onto the permutation $\mathbf{rc} \sigma$ defined by

$$\mathbf{rc} \sigma(i) := n + 1 - \sigma(n + 1 - i) \quad (1 \leq i \leq n),$$

maps \mathcal{DR}_n onto \mathcal{D}_n in a one-to-one manner. Moreover, under the transformation the number of inversions “inv” remains alike. This proves the second part of the theorem. \square

Saying that a permutation σ is rising alternating (resp. falling alternating) is equivalent to saying that $\text{Ligne } \sigma = \{2, 4, 6, \dots\}$ (resp. $\text{Ligne } \sigma = \{1, 3, 5, \dots\}$). Hence, Corollary 11.5 implies the following proposition.

Proposition 24.5. *For every $n \geq 1$ we have:*

$$\sum_{\sigma \in \mathcal{DR}_n} p^{\text{inv } \sigma} = \sum_{\sigma \in \mathcal{DR}_n} p^{\text{imaj } \sigma} \quad \text{and} \quad \sum_{\sigma \in \mathcal{D}_n} p^{\text{inv } \sigma} = \sum_{\sigma \in \mathcal{D}_n} p^{\text{imaj } \sigma}.$$

24.3. *Combinatorics of the bibasic secant and tangent.* In section 23 we have studied the class of the bi-indexed Eulerian polynomials defined by

$$A_{n,m}(t, p, q) := \sum_{\sigma \in \mathfrak{S}_{n+m}} t^{1+\text{des } \sigma} p^{\text{inv}_1 \sigma} q^{\text{inv}_2 \sigma}.$$

As the statistic “inv” (also the statistic “imaj”), when defined on the subset of alternating permutations, leads to a combinatorial interpretation of the q -tangent and q -secant numbers, the natural question arises: what can be said about the polynomials

$$(24.27) \quad D_{n,m}(p, q) := \sum_{\sigma \in \mathcal{DR}_{n+m}} p^{\text{inv}_1 \sigma} q^{\text{inv}_2 \sigma}, \quad E_{n,m}(p, q) := \sum_{\sigma \in \mathcal{D}_{n+m}} p^{\text{inv}_1 \sigma} q^{\text{inv}_2 \sigma}$$

when the pair $(\text{inv}_1, \text{inv}_2)$ is restricted to the subset \mathcal{DR}_{n+m} of the *rising* (resp. the subset \mathcal{D}_{n+m} of the *descending*) alternating permutations. No use keeping the variable t as all the permutations in \mathcal{DR}_{n+m} (resp. \mathcal{D}_{n+m}) have the same ligne of route, i.e., $\{2, 4, 6, \dots\}$ (resp. $\{1, 3, 5, \dots\}$).

Again, let (n, m) be an ordered pair of nonnegative integers. By convention, $D_{0,0}(p, q) := 1$. Let $n + m \geq 1$ and let σ be a permutation of order $n + m$. The statistics “ inv_1 ” and “ inv_2 ” (see Theorem 23.1) can be redefined as follows:

$$\begin{aligned} \text{inv}_1 \sigma &:= \#\{(i, j) : 1 \leq i < j \leq n + m, n \geq \sigma(i) > \sigma(j) \geq 1\}, \\ \text{inv}_2 \sigma &:= \#\{(i, j) : 1 \leq i < j \leq n + m, n + m \geq \sigma(i) > \sigma(j) \geq n + 1\}. \end{aligned}$$

To emphasize the fact that inv_1 and inv_2 are defined by means of the ordered pair (n, m) , we also use the notations:

$$\text{inv}_1^{(n,m)} := \text{inv}_1; \quad \text{inv}_2^{(n,m)} := \text{inv}_2.$$

As has been defined in § 24.1,

$$\begin{aligned} \text{Eul}(u, v; p, q) &= \sec_{p,q}(u, v) + \tan_{p,q}(u, v) \\ &= \sum_{n \geq 0, m \geq 0} \text{Eul}_{n,m}(p, q) \frac{u^n}{(p; p)_n} \frac{v^m}{(q; q)_m}. \end{aligned}$$

The purpose of this subsection is to prove the following theorem.

Theorem 24.6. *For each pair of nonnegative integers (n, m) we have:*

$$(24.28) \quad D_{n,m}(p, q) = \text{Eul}_{n,m}(p, q);$$

when $n + m$ is odd, we further have:

$$(24.29) \quad E_{n,m}(p, q) = \text{Eul}_{n,m}(p, q);$$

Before proving the theorem and making some comment about the symmetry of the polynomials $\text{Eul}_{n,m}(p, q)$ let us establish the following lemma.

Lemma 24.7. *Let σ be a permutation of order $n+m \geq 1$ and $\sigma' := \mathbf{rc}\sigma$. Then*

$$(\text{inv}_1^{(n,m)}, \text{inv}_2^{(n,m)})\sigma = (\text{inv}_2^{(m,n)}, \text{inv}_1^{(m,n)})\sigma'.$$

Proof. By definition of \mathbf{r} and \mathbf{c} we have

$$\sigma'(n+m+1-i) = n+m+1-\sigma(i) \quad (1 \leq i \leq m+n).$$

Hence the following statements are equivalent:

- (i) $1 \leq i < j \leq n+m$ & $n \geq \sigma(i) > \sigma(j) \geq 1$;
- (ii) $1 \leq n+m+1-j < n+m+1-i \leq n+m$
& $n+m \geq n+m+1-\sigma(j) > n+m+1-\sigma(i) \geq m+1$;
- (iii) $1 \leq j' < i' \leq n+m$ & $n+m \geq \sigma'(j') > \sigma'(i') \geq m+1$ with
 $j' := n+m+1-j, i' := n+m+1-i$;

Thus, $\text{inv}_1^{(n,m)}\sigma = \text{inv}_2^{(m,n)}\sigma'$. Also, $\text{inv}_2^{(n,m)}\sigma = \text{inv}_1^{(m,n)}\sigma'$ by permuting the roles of n and m . \square

If σ is a rising alternating permutation of order $n+m$, then $\sigma' = \mathbf{rc}\sigma$ is rising alternating or descending alternating, depending on whether $n+m$ is even or odd. By the previous lemma we then have

$$(24.30) \quad D_{n,m}(p, q) = \begin{cases} D_{m,n}(q, p), & \text{if } n+m \text{ is even;} \\ E_{m,n}(q, p), & \text{if } n+m \text{ is odd.} \end{cases}$$

Thus, when $n+m$ is even, the symmetry of the polynomials $D_{n,m}(p, q)$ is obvious combinatorially. When $n+m$ is *odd*, it is only a consequence of Theorem 24.6.

In order to prove identity (24.28) of Theorem 24.6 we show that the sequence $(D_{n,m}(p, q))$ satisfies the relations (iii), (iv), (v) of Proposition 24.3 that uniquely *define* the sequence $(\text{Eul}_{n,m}(p, q))$. This is the content of the next proposition.

Proposition 24.8. *Let $D_{n,m}(p, q)$ be the generating polynomial for the set \mathcal{DR}_{n+m} of the rising alternating permutations of order $n+m$ by the two-variable statistic $(\text{inv}_1, \text{inv}_2)$. Then the sequence $(D_{n,m}(p, q))$ ($n \geq 0, m \geq 0$) is inductively defined by the following three relations:*

- (iii) *the initial conditions $D_{0,0}(p, q) = D_{1,0}(p, q) = D_{0,1}(p, q) = 1$;*
- (iv) *the recurrence formula valid for $n \geq 1$:*

$$D_{n+1,0}(p, q) = \sum_{\substack{0 \leq a \leq n, \\ a \text{ odd}}} \begin{bmatrix} n \\ a \end{bmatrix}_p p^{n-a} D_{a,0}(p, q) D_{n-a,0}(p, q);$$

- (v) *and the following formula valid for $n+m \geq 1$:*

$$(24.31) \quad D_{n,m+1}(p, q) = \sum_{\substack{0 \leq a \leq n, 0 \leq b \leq m \\ a+b \text{ odd}}} \begin{bmatrix} n \\ a \end{bmatrix}_p \begin{bmatrix} m \\ b \end{bmatrix}_q q^{m-b} D_{a,b}(p, q) D_{n-a, m-b}(p, q).$$

Proof. Relation (iii) is banal, while relation (iv) is the recurrence formula for the p -tangent or p -secant number derived in the previous subsection.

Let $\sigma = \sigma(1)\sigma(2)\dots\sigma(n+m+1)$ be a rising alternating permutation of order $(n+m+1)$. As the permutation starts with a rise $\sigma(1) < \sigma(2)$, the letter $(n+m+1)$ in the word σ occurs in an even position, say, $2k+2$ ($k \geq 0$). Let

$$\begin{aligned} A &:= \{\sigma(l) : 1 \leq l \leq 2k+1, 1 \leq \sigma(l) \leq n\}; \\ B &:= \{\sigma(l) : 1 \leq l \leq 2k+1, n+1 \leq \sigma(l) \leq n+m\}; \\ C &:= \{\sigma(l) : 2k+3 \leq l \leq n+m+1, 1 \leq \sigma(l) \leq n\} = [n] \setminus A; \\ D &:= \{\sigma(l) : 2k+3 \leq l \leq n+m+1, n+1 \leq \sigma(l) \leq n+m\} \\ &= [n+1, n+m] \setminus B. \end{aligned}$$

Also, let $\#A := a$, $\#B := b$, $\#C := c = n - a$, $\#D := d = m - b$, so that $a + b = 2k + 1$ and $c + d = n + m - (2k + 2)$. The two words $\sigma' := \sigma(1)\dots\sigma(2k+1)$ and $\sigma'' := \sigma(2k+3)\dots\sigma(n+m+1)$ are alternating permutations of the sets $A + B$ and $C + D$, respectively. The *reduction* of σ' is defined to be the permutation $\tau' = \tau'(1)\dots\tau'(2k+1)$, where the letter $\tau'(l)$ is equal to m if and only if $\sigma(l)$ is the smallest m -th letter of $A + B$. The reduction $\tau'' = \tau''(1)\dots\tau''(n+m+1-2k-2)$ of σ'' is defined in the same way. Of course, both τ' and τ'' are rising alternating permutations.

Now σ is completely characterized by the four-sequence (A, B, τ', τ'') . The inversions of σ of the form $n \geq \sigma(i) > \sigma(j) \geq 1$ ($1 \leq i < j \leq n+m$) are of three kinds:

- (i) $1 \leq i < j \leq 2k+1$; they are counted by $\text{inv}_1^{(a,b)} \tau'$;
- (ii) $2k+3 \leq i < j \leq n+m$; they are counted by $\text{inv}_1^{(c,d)} \tau''$;
- (iii) $1 \leq i < 2k+2 < j \leq n+m+1$; using the notations of § 4.4 they are counted by $\text{inv}(\gamma(A)\gamma(C))$.

Thus,

$$\text{inv}_1^{(n,m)} = \text{inv}_1^{(a,b)} \tau' + \text{inv}_1^{(c,d)} \tau'' + \text{inv}(\gamma(A)\gamma(C));$$

and in the same way

$$\text{inv}_2^{(n,m)} = \text{inv}_2^{(a,b)} \tau' + \text{inv}_2^{(c,d)} \tau'' + \text{inv}(\gamma(B)\gamma(D)) + d.$$

The “ d ” that has been added takes the inversions $\sigma(2k+2) = n+m > \sigma(l) \geq n+1$ ($2k+3 \leq l \leq n+m+1$) into account.

When the pair (A, B) is fixed, so are the ordered partitions (A, C) and (B, D) of $[n]$ and $[n + 1, n + m]$, respectively. Hence

$$\sum_{\tau', \tau''} p^{\text{inv}_1^{(a,b)} \tau' + \text{inv}_1^{(c,d)} \tau''} q^{\text{inv}_2^{(a,b)} \tau' + \text{inv}_2^{(c,d)} \tau''} = D_{a,b}(p, q) D_{c,d}(p, q).$$

Now, when A (resp. B) runs over all subsets of $[n]$ (resp. of $[n + 1, n + m]$) of cardinality a (resp. b), the pairs (A, C) and (B, D) range over all ordered partitions of $[n]$ and of $[n + 1, n + m]$, respectively, into two blocks such that $\#A = a$, $\#C = c$, and $\#B = b$, $\#D = d$. By Proposition 4.3

$$\sum_{(A,B)} p^{\text{inv}(\gamma(A)\gamma(C))} q^{\text{inv}(\gamma(B)\gamma(D))+d} = \begin{bmatrix} n \\ a \end{bmatrix}_p \begin{bmatrix} m \\ b \end{bmatrix}_q q^d.$$

The final summation with respect to a, b yields (24.31). \square

For proving (24.29) consider (24.31) when $n + m$ is odd. Using (24.30) we can rewrite (24.31) as

$$E_{n+1,m}(p, q) = \sum_{\substack{0 \leq b \leq m, 0 \leq a \leq n \\ b+a \text{ odd}}} \begin{bmatrix} m \\ b \end{bmatrix}_q \begin{bmatrix} n \\ a \end{bmatrix}_p q^{m-b} E_{b,a}(q, p) E_{m-b, m-a}(q, p).$$

This is exactly the recurrence formula (24.16') written for $\text{Eul}_{n+1,m}(q, p)$. As the analogous relations (iii) and (iv') hold for $E_{m,n}(0, p)$, identity (24.29) is proved. \square

25. MacMahon's Master Theorem revisited

This theorem has been regarded as a keystone in Combinatorial Analysis. It was proved by MacMahon himself [Ma15, vol. 1, p. 93–98], who also gave numerous applications. Several other proofs are due to Good [Go62], Cartier [Ca72]. As the “Master Theorem” is a special case of the multivariable Lagrange Inversion Formula [Ge87], as was shown by Good [Go60] (see also Hofbauer [Ho82]), each proof of the latter formula yields a proof of this theorem. Its first noncommutative version can be found in [Fo65], which was later implemented into an appropriate algebraic set-up [CF69]. More recently Garoufalidis, Lê and Zeilberger derived another noncommutative version [GLZ06] (called “quantum Master Theorem”) by using difference operator techniques developed by Zeilberger [Z80].

Instead of reproducing a short version of the “Master Theorem,” in the commutative case only, we have preferred to put up an algebraic structure that can shelter the three versions of the theorem, the commutative one à

la MacMahon, the partially commutative one à la Cartier-Foata and the quantum one à la Garoufalidis-Lê-Zeilberger. After displaying the identity to be proved, and discussing properties of so-called circuits, the “Master Theorem” is established in the commutative case through several steps. The proof for the two non-commutative algebras will consist of verifying that each step is still valid in these two further environments.

25.1. *The “Master Theorem” identity.* Let r be a nonnegative integer; the set $\mathbb{A} = \{0, 1, \dots, r\}$ is referred to as being the underlying *alphabet*. Form the free monoid \mathbb{A}^* generated by \mathbb{A} , that is, the set of all finite words $w = x_1x_2 \cdots x_n$, whose letters x_1, x_2, \dots, x_n belong to \mathbb{A} . A *biword* on \mathbb{A} is defined to be a $2 \times n$ matrix $\alpha = \binom{w'}{w}$ ($n \geq 0$), where w' (the *top word*) and w (the *bottom word*) are two words with the same length n . The number n is also the length of α ; we write $\ell(\alpha) = n$. Let \mathcal{B} be the set of all biwords on \mathbb{A} . Each biword can also be viewed as a word of *biletters* $\binom{x}{a}$ written vertically. The *product* of two biwords is just the concatenation of them viewed as two words of biletters.

Let \mathbb{Z} be the ring of all integers. The set $\mathcal{A} = \mathbb{Z}\langle\langle\mathcal{B}\rangle\rangle$ of the formal sums $\sum_{\alpha} c(\alpha)\alpha$, where $\alpha \in \mathcal{B}$ and $c(\alpha) \in \mathbb{Z}$ for all $\alpha \in \mathcal{B}$, together with the above multiplication, the free addition and the free scalar product is an algebra over \mathbb{Z} , called the *free biword large \mathbb{Z} -algebra*.

For each word w let \bar{w} be its *non-decreasing* rearrangement, that is, the word derived from w by rearranging its letters in non-decreasing order. Using the terminology made popular in [GLZ05] define the *bosonic sum* as the infinite sum

$$(25.1) \quad \text{Bos} := \sum_w \binom{\bar{w}}{w},$$

where the sum is over all finite words w with letters from \mathbb{A} , *i.e.*, over the free monoid \mathbb{A}^* generated by \mathbb{A} .

The *fermionic sum* is defined to be

$$(25.2) \quad \text{Ferm} := \sum_{J \subset \mathbb{A}} (-1)^{|J|} \sum_{\sigma \in \mathcal{S}_J} (-1)^{\text{inv}\sigma} \binom{i_1 \quad i_2 \quad \cdots \quad i_l}{\sigma(i_1) \sigma(i_2) \cdots \sigma(i_l)},$$

where $J = \{i_1 < i_2 < \cdots < i_l\}$ and \mathcal{S}_J is the permutation group acting on the set J .

The purpose of this chapter is to show that the identity

$$(25.3) \quad \text{Ferm} \times \text{Bos} = 1$$

holds in three associative algebras, quotients of \mathcal{A} by ideals generated by commutation rules defined as follows:

(1) the *commutative* algebra, denoted by \mathcal{A}^{MM} , for which all billetters commute:

$$(25.4) \quad \binom{x}{a} \binom{y}{b} = \binom{y}{b} \binom{x}{a} \quad \text{for all } a, b, x, y \text{ from } \mathcal{A}.$$

This is the classical version that goes back MacMahon’s “Master Theorem,” although it was not stated as an identity of any sort of algebra, but as the fact that two multivariable series had identical coefficients; see Exercice 34.

(2) the *partially commutative* algebra à la Cartier-Foata, denoted by \mathcal{A}^{CF} , defined by the commutation rules:

$$(25.5) \quad \binom{xy}{ab} = \binom{yx}{ba} \quad \text{for all letters } a, b, x, y \text{ from } \mathbb{A} \text{ and } x \neq y.$$

See [CF69] and [Lo02, chap. 10] for more details.

(3) the *right quantum* algebra \mathcal{A}^{GLZ} à la Garoufalidis-Lê-Zeilberger defined by the commutation rules

$$(25.6) \quad \begin{aligned} \text{(R1)} \quad & \binom{xy}{aa} = \binom{yx}{aa} \quad \text{for all letters } a, x, y \text{ from } \mathbb{A} \text{ and } x \neq y \\ \text{(R2)} \quad & \binom{xy}{ab} + \binom{xy}{ba} = \binom{yx}{ab} + \binom{yx}{ba} \end{aligned}$$

for all letters a, b, x, y from \mathbb{A} and $x \neq y, a \neq b$.

See [FH07] for more details about the construction of the algebra and how can be shown that the algebra is associative.

In the commutative case the right-hand side of (25.2) can also be expressed as the determinant $\det(I - C_r)$ of the $(r + 1) \times (r + 1)$ matrix $(I - C_r)$, where I is the identity matrix and

$$(25.7) \quad C_r := \begin{pmatrix} \binom{0}{0} & \binom{0}{1} & \binom{0}{2} & \cdots & \binom{0}{r-1} & \binom{0}{r} \\ \binom{1}{0} & \binom{1}{1} & \binom{1}{2} & \cdots & \binom{1}{r-1} & \binom{1}{r} \\ \binom{2}{0} & \binom{2}{1} & \binom{2}{2} & \cdots & \binom{2}{r-1} & \binom{2}{r} \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ \binom{r-1}{0} & \binom{r-1}{1} & \binom{r-1}{2} & \cdots & \binom{r-1}{r-1} & \binom{r-1}{r} \\ \binom{r}{0} & \binom{r}{1} & \binom{r}{2} & \cdots & \binom{r}{r-1} & \binom{r}{r} \end{pmatrix}.$$

The formula $\det(I - C_r)$ is also valid when dealing with the above two non-commutative algebras, as will be further explained, so that the “Master Theorem” identity rewritten as

$$(25.8) \quad \det(I - C_r) \times \text{Bos} = 1$$

is to be established in each algebra \mathcal{A}^{MM} , \mathcal{A}^{CF} , \mathcal{A}^{GLZ} .

25.2. *Circuit Calculus.* A *circuit* is a biword whose top word is a rearrangement of its bottom word. Each formal sum $E = \sum_{\alpha} c(\alpha)\alpha \in \mathcal{A}$ is said to be *imaginary* (resp. *real*), if $c(\alpha) = 0$ for all circuits α (resp. non-circuits α). The following properties, although easy to verify, are essential for the proof of the theorem.

(P1) Each formal sum E can be decomposed in a unique way as a sum of a real expression $\Re(E)$ and an imaginary expression $\Im(E)$:

$$E = \Re(E) + \Im(E).$$

(P2) The real part operator \Re and the imaginary part operator \Im are linear. This means that for any two formal sums E , E' and scalars c , c' we have

$$\begin{aligned}\Re(cE + c'E') &= c\Re(E) + c'\Re(E'), \\ \Im(cE + c'E') &= c\Im(E) + c'\Im(E').\end{aligned}$$

(P3) The real part operator \Re and the imaginary part operator \Im are idempotent and orthogonal to each other. This means that for every formal sum E we have

$$\Re(\Re(E)) = \Re(E), \quad \Im(\Im(E)) = \Im(E) \quad \text{and} \quad \Re(\Im(E)) = \Im(\Re(E)) = 0.$$

(P4) If E , E' are two nonzero expressions from \mathcal{R} and if E is real, then $E \times E'$ is real (resp. imaginary) if and only if E' is real (resp. imaginary).

(P5) If E , E' are two nonzero formal sums and if E is real, then

$$\Re(E \times E') = E \times \Re(E').$$

Now, define the *universe* “Univ” to be the sum of all biwords whose top words are nondecreasing, that is,

$$(25.9) \quad \text{Univ} := \sum_{u,w} \binom{u}{w},$$

where u (resp. w) runs over the set of all *nondecreasing* words (resp. all words) and where u and w are of the same length: $\ell(u) = \ell(w)$. Both Boson and Fermion are sums of *circuits*. Moreover, $\Re(\text{Univ}) = \text{Bos}$. Hence, by (P5) identity (25.3) is equivalent to the following identity

$$(25.10) \quad \Re(\text{Ferm} \times \text{Univ}) = 1.$$

25.3. “Master Theorem” within the algebra \mathcal{A}^{MM} . For $i = 0, 1, \dots, r$ define:

$$(25.11) \quad S_i^{(r)} := \binom{i}{0} + \binom{i}{1} + \dots + \binom{i}{r};$$

$$(25.12) \quad K_i^{(r)} := \frac{1}{1 - S_i^{(r)}} = \sum_{n \geq 0} (S_i^{(r)})^n.$$

Step 1. As the product of the top words in the product $K_0^{(r)} K_1^{(r)} \dots K_r^{(r)}$ reads $00 \dots 0 11 \dots 1 \dots rr \dots r$, we have:

$$(25.13) \quad \text{Univ} = K_0^{(r)} K_1^{(r)} \dots K_r^{(r)}.$$

Step 2. Display $\det(I - C_r)$ with C_r defined in (25.7) and add its leftmost r columns to its rightmost column, to obtain

$$(25.14) \quad \det(I - C_r) = \det \begin{pmatrix} 1 - \binom{0}{0} & -\binom{0}{1} & \dots & 1 - S_0^{(r)} \\ -\binom{1}{0} & 1 - \binom{1}{1} & \dots & 1 - S_1^{(r)} \\ \vdots & \vdots & \ddots & \vdots \\ -\binom{r}{0} & -\binom{r}{1} & \dots & 1 - S_r^{(r)} \end{pmatrix}$$

Step 3. Multiply each entry of the rightmost column in the resulting determinant by $\text{Univ} = K_0^{(r)} K_1^{(r)} \dots K_r^{(r)}$ to get

$$(25.15) \quad \det(I - C_r) \times \text{Univ} \\ = \det \begin{pmatrix} 1 - \binom{0}{0} & -\binom{0}{1} & \dots & K_1^{(r)} K_2^{(r)} \dots K_r^{(r)} \\ -\binom{1}{0} & 1 - \binom{1}{1} & \dots & K_0^{(r)} K_2^{(r)} \dots K_r^{(r)} \\ \vdots & \vdots & \ddots & \vdots \\ -\binom{r}{0} & -\binom{r}{1} & \dots & K_0^{(r)} K_1^{(r)} \dots K_{r-1}^{(r)} \end{pmatrix}.$$

Step 4. Denote the minor of $I - C_r$ at position (i, j) by M_{ij} and derive the Laplace cofactor expansion of the determinant with respect to the rightmost column. This yields:

$$(25.16) \quad \det(I - C_r) \times \text{Univ} \\ = \sum_{0 \leq j \leq r} (-1)^{r+j} \det(M_{jr}) K_0^{(r)} \dots K_{j-1}^{(r)} K_{j+1}^{(r)} \dots K_r^{(r)}.$$

Let $0 \leq j \leq r - 1$. All the biwords occurring in the expansion of the minor M_{jr} have no letters equal to j in their top words, as no billetter $\binom{j}{i}$

($0 \leq i \leq r$) occurs in the minor. Furthermore, the sum $K_j^{(r)}$ being missing in the product $K_0^{(r)} \cdots K_{j-1}^{(r)} K_{j+1}^{(r)} \cdots K_r^{(r)}$, the same conclusion holds for its expansion. Consequently, for each biword occurring in $F_j := \det(M_{jr}) K_0^{(r)} \cdots K_{j-1}^{(r)} K_{j+1}^{(r)} \cdots K_r^{(r)}$ there is a letter j in the bottom, but none in the top. This means that F_j ($j \neq r$) does not contain any circuit. By applying the operator \mathfrak{R} to both members of (25.16) we get $\mathfrak{R}F_j = 0$ for every $j = 0, 1, \dots, r-1$ and then

$$(25.17) \quad \det(I - C_r) \times \mathfrak{R}(\text{Univ}) = \det(M_{rr}) \times \mathfrak{R}(K_0^{(r)} K_1^{(r)} \cdots K_{r-1}^{(r)}).$$

using Property (P5):

Step 5. When expanding the product $K_0^{(r)} K_1^{(r)} \cdots K_{r-1}^{(r)}$, the biwords containing biletters $\binom{i}{r}$ with $0 \leq i \leq r-1$ vanish under the application of \mathfrak{R} . After removing such biwords there only remain the biwords from the product $K_0^{(r-1)} K_1^{(r-1)} \cdots K_{r-1}^{(r-1)}$. As $M_{rr} = I - C_{r-1}$, we have by induction:

$$\begin{aligned} \det(I - C_r) \times \mathfrak{R}(\text{Univ}) &= \det(I - C_r) \times \mathfrak{R}(K_0^{(r)} K_1^{(r)} \cdots K_r^{(r)}) \\ &= \det(I - C_{r-1}) \times \mathfrak{R}(K_0^{(r-1)} K_1^{(r-1)} \cdots K_{r-1}^{(r-1)}) \\ &= \det(I - C_{r-2}) \times \mathfrak{R}(K_0^{(r-2)} K_1^{(r-2)} \cdots K_{r-2}^{(r-2)}) \\ &\quad \dots \\ &= \det(I - C_0) \times \mathfrak{R}(K_0^{(0)}) = (1 - \binom{0}{0}) \times \frac{1}{1 - \binom{0}{0}} = 1. \end{aligned}$$

In view of (25.15) this proves identity (25.8) in \mathcal{A}^{MM} .

25.4. Further matricidal properties of the algebras \mathcal{A}^{CF} and \mathcal{A}^{GLZ} . Let $A = (a_{i,j})_{1 \leq i,j \leq r}$ be a square matrix whose entries are elements from \mathcal{A}^{CF} (resp. \mathcal{A}^{GLZ}). Although these algebras are noncommutative, the *determinant* of A can be defined to be

$$(25.18) \quad \det(A) = \sum_{\sigma} (-1)^{\text{inv}\sigma} a_{\sigma_0,0} a_{\sigma_1,1} \cdots a_{\sigma_r,r},$$

where σ runs over the permutation group $\mathcal{S}_{\mathbb{A}}$ of the set \mathbb{A} , as long as the ordering of the factors $a_{\sigma_0,0}, a_{\sigma_1,1}, \dots, a_{\sigma_r,r}$ is taken into account. Several classical properties of the determinant still hold.

(P6) *Linearity.* When writing matrices as sequences of their $(r + 1)$ columns (c_0, c_1, \dots, c_r) , we have

$$\begin{aligned} \det(c_0, \dots, c_i, \dots, c_r) + \det(c_0, \dots, b c'_i, \dots, c_r) \\ = \det(c_0, \dots, c_i + b c'_i, \dots, c_r). \end{aligned}$$

(P7) *Cofactor expansion.* Let $A = (a_{i,j})_{0 \leq i, j \leq r}$ be a matrix with entries from \mathcal{A}^{CF} (resp. \mathcal{A}^{GLZ}) and A_{ij} be the matrix obtained from A by deleting the i -th row and j -th column. Then,

$$\det(A) = \sum_{i=0}^r (-1)^{r+1+i} \det(A_{ir}) a_{ir}.$$

This is simply the usual cofactor expansion of $\det(A)$ with respect of its *rightmost* column.

Let a_i, b_i, c_i ($i = 0, 1, \dots, r$) and x, y be scalars and form the $(r + 1) \times (r + 1)$ -matrix

$$A = \begin{pmatrix} \cdots & a_0 + b_0 \binom{0}{x} & c_0 + b_0 \binom{0}{y} & \cdots \\ \cdots & a_1 + b_1 \binom{1}{x} & c_1 + b_1 \binom{1}{y} & \cdots \\ \ddots & \vdots & \vdots & \ddots \\ \cdots & a_r + b_r \binom{r}{x} & c_r + b_r \binom{r}{y} & \cdots \end{pmatrix},$$

where besides the two *consecutive* columns explicitly displayed the other columns are arbitrary. Let B denote the matrix derived from A by transposing those two consecutive columns.

(P8) *Interchanging two columns.* Let A and B be the two matrices just defined. Then $\det A = -\det B$.

Property (P8) can be proved as follows. In the expansion of $\det A$ the sum of the two terms

$$S_{ij} := \pm \cdots (a_i + b_i \binom{i}{x})(c_j + b_j \binom{j}{y}) \cdots \mp \cdots (a_j + b_j \binom{j}{x})(c_i + b_i \binom{i}{y}) \cdots$$

may be put in a one-to-one correspondence with the sum

$$T_{ij} := \pm \cdots (c_i + b_i \binom{i}{y})(a_j + b_j \binom{j}{x}) \cdots \mp \cdots (c_j + b_j \binom{j}{y})(a_i + b_i \binom{i}{x}) \cdots$$

in the expansion of $\det B$. But

$$\begin{aligned}
 S_{ij} &= \pm \cdots \left(a_i c_j + a_i b_j \binom{j}{y} + c_j b_i \binom{i}{x} + b_i b_j \binom{ij}{xy} \right. \\
 &\quad \left. - a_j c_i - a_j b_i \binom{i}{y} - c_i b_j \binom{j}{x} - b_i b_j \binom{ji}{yx} \right) \cdots \\
 T_{ij} &= \pm \cdots \left(c_i a_j + c_i b_j \binom{j}{x} + b_i a_j \binom{i}{y} + b_i b_j \binom{ij}{yx} \right. \\
 &\quad \left. - c_j a_i - c_j b_i \binom{i}{x} - b_j a_i \binom{j}{y} - b_i b_j \binom{ji}{yx} \right) \cdots
 \end{aligned}$$

Within the algebra \mathcal{A}^{CF} we have $\binom{ij}{xy} = \binom{ji}{yx}$, so that $S_{ij} = -T_{ij}$. Within the algebra \mathcal{A}^{GLZ} the identity $\binom{ij}{xy} - \binom{ji}{yx} = -\left(\binom{ij}{yx} - \binom{ji}{xy}\right)$ holds. As the commutation relations can be made at *any* position within each biword (using the associativity property of the right quantum algebra) we also conclude that $S_{ij} = -T_{ij}$. \square

It is worth noticing that Property (P8) is only stated for very special matrices. In general, the column interchanging property does *not* hold for arbitrary matrices with entries from those two algebras.

As a consequence of Property (P8), we can state the following property.

(P9) *Two identical columns.* If the matrix A has the further property that two of its columns are identical, that is, if $a_i = a_j$ and $b_i = b_j$ for $0 \leq i < j \leq r$, then $\det A = 0$.

For the proof, it suffices to write: permute columns i and $i + 1$, then columns $i + 1$ and $i + 2$, \dots , finally columns $r - 2$ and $r - 1$. We obtain a matrix A' whose rightmost two columns are identical. Property (P8) implies that $\det(A') = 0$ and also $\det(A') = \pm \det(A) = 0$. \square

25.5. *The "Master Theorem" for the algebras \mathcal{A}^{CF} and \mathcal{A}^{GLZ} .* Starting with the definition of the determinant given in (26.18) and using Property 6 (linearity) the fermion matrix defined in (26.2) can also be expressed as

$$\text{Ferm} = \det(I - C_r) = \det \begin{pmatrix} 1 - \binom{0}{0} & -\binom{0}{1} & \cdots & -\binom{0}{r} \\ -\binom{1}{0} & 1 - \binom{1}{1} & \cdots & -\binom{1}{r} \\ \vdots & \vdots & \ddots & \vdots \\ -\binom{r}{1} & -\binom{r}{2} & \cdots & 1 - \binom{r}{r} \end{pmatrix}.$$

Now examine how each **Step 1**, \dots , **Step 5** can be validated for \mathcal{A}^{CF} and \mathcal{A}^{GJZ} . First, **Step 1** holds in both algebras. For **Step 2** designate the

$(r + 1)$ columns of the matrix $(I - C_r)$ by c_0, c_1, \dots, c_r . By Property 4 (Linearity) and (P9) (Two identical columns) we have:

$$\begin{aligned} \det(c_0, \dots, c_{r-1}, c_r + c_{r-1}) &= \det(c_0, \dots, c_{r-1}, c_{r-1}) + \det(c_0, \dots, c_{r-1}, c_r) \\ &= \det(c_0, \dots, c_{r-1}, c_r), \end{aligned}$$

By iteration,

$$\begin{aligned} \det(c_0, \dots, c_{r-1}, c_r + c_{r-1} + \dots + c_0) &= \det(c_0, \dots, c_{r-1}, c_r) \\ &= \det(I - C_r), \end{aligned}$$

so that Step 2 is correct.

In **Step 3** the multiplication (on the right) of the entry in the j -th row ($0 \leq j \leq r$) and r -th column by $\text{Univ} = K_0^{(r)} \dots K_j^{(r)} \dots K_r^{(r)}$ yields $(1/K_j^{(r)})K_0^{(r)} \dots K_j^{(r)} \dots K_r^{(r)}$, which is equal to $K_0^{(r)} \dots K_{j-1}^{(r)} K_{j+1}^{(r)} \dots K_r^{(r)}$ when the $K_i^{(r)}$'s commute with each other. It is obviously the case in \mathcal{L}^{CF} , because of the commutation rule (25.5) : $\binom{xy}{ab} = \binom{yx}{ba}$ if $x \neq y$. However, it requires a proof in the algebra \mathcal{A}^{GLZ} .

Lemma. In \mathcal{A}^{GLZ} we have: $S_i S_j = S_j S_i$ and $K_i K_j = K_j K_i$.

Proof. It suffices to prove that the S_i 's commute with each other. Grouping the biwords by pairs if necessary we have:

$$\begin{aligned} S_i S_j &= \sum_{a < b} \left(\binom{ij}{ab} + \binom{ij}{ba} \right) + \sum_a \binom{ij}{aa} \\ &= \sum_{a < b} \left(\binom{ji}{ab} + \binom{ji}{ba} \right) + \sum_a \binom{ji}{aa} \\ &= S_j S_i, \end{aligned}$$

by using the commutation rules (25.6). \square

Finally, **Step 4** and **Step 5** only require the validity of the cofactor expansion and manipulations with circuits, which are allowed in the the two algebras \mathcal{L}^{CF} and \mathcal{A}^{GLZ} . This completes the proof of the ‘‘Master Theorem.’’

26. The decrease value theorem

The *decrease value theorem*, further stated, makes the calculation of a certain multivariable statistical distribution on words possible. The multivariable statistic in question involves the basic notions of *decrease*,

increase and *record*, whose definitions are now introduced, together with the classical *descent* and *rise*.

Let $[0, r]^*$ be the set of all words, whose letters belong to the finite alphabet $[0, r] = \{0, 1, \dots, r\}$ and let $v = y_1 y_2 \cdots y_n$ be such a word. An integer $i \in [1, n]$ is said to be a *descent* (or *descent place*) of v if $1 \leq i \leq n - 1$ and $y_i > y_{i+1}$. The letter y_i (resp. y_{i+1}) is called a *descent value*. (See Fig. 26.1, where part of the graph of the mapping $i \mapsto y_i$ has been partially drawn.)




Fig. 26.1 : Graph of a descent




Fig. 26.2 : Graph of a decrease

The integer i is called a *decrease* of v if $y_i = y_{i+1} = \cdots = y_j > y_{j+1}$ for some j such that $i \leq j \leq n - 1$. The letter y_i (resp. y_{j+1}) is said to be a *decrease value* of v . The set of all decreases (resp. descents) is denoted by $\text{DEC}(v)$ (resp. $\text{DES}(v)$). Each descent is a decrease, so that $\text{DES}(v) \subset \text{DEC}(v)$.

In parallel with the notion of decrease, an integer $i \in [1, n]$ is said to be an *increase* (resp. a *rise*) of v if $y_i = y_{i+1} = \cdots = y_j < y_{j+1}$ for some j such that $i \leq j \leq n$ (resp. if $y_i < y_{i+1}$). By convention, $y_{n+1} = +\infty$. The letter y_i is said to be an *increase value* (resp. a *rise value*) of v . Thus, the rightmost letter y_n is always a rise and also an increase value. The set of all increases (resp. rises) of v is denoted by $\text{INC}(v)$ (resp. $\text{RISE}(v)$). Each rise is an increase, so that $\text{RISE}(v) \subset \text{INC}(v)$. (See Fig. 26.3 and 26.4.)




Fig. 26.3 : Graph of a rise




Fig. 26.4 : Graph of an increase

As, by convention, the rightmost letter of each word is always an increase value, we have $\text{DEC}(v) = \emptyset$ and $\text{INC}(v) = \{1\}$ for each single letter word v . Let $v = y_1 y_2 \cdots y_n$ be a word of length $n \geq 2$ and consider its right factor $y_{i+1} y_{i+2} \cdots y_n$ with $1 \leq i \leq n - 1$. If the right factor contains no rise value besides y_n and no descent value, then $y_i = y_{i+1} = \cdots = y_n$ and y_i is an increase value. Otherwise, consider the smallest integer k such that $i + 1 \leq k \leq n - 1$ and y_k is a rise (resp. descent) value, then $y_i = y_{i+1} = \cdots = y_k$ and $y_k < y_{k+1}$ (resp. $y_k > y_{k+1}$). Therefore, y_i is a rise (resp. descent) value. Thus, $\text{DEC}(v)$ and $\text{INC}(v)$ are disjoint and for each word v we have:

$$(26.5) \quad \text{DEC}(v) + \text{INC}(v) = [1, n].$$

Finally, a position i ($1 \leq i \leq n$) is said to be a *record* if $y_j \leq y_i$ for all j such that $1 \leq j \leq i - 1$. The letter y_i is said to be a *record value*. The set of all records of v is denoted by $\text{REC}(v)$.

Taking the two inclusions $\text{DES}(v) \subset \text{DEC}(v)$, $\text{RISE}(v) \subset \text{INC}(v)$ into account, the Venn diagram of the properties “DEC,” “INC,” “RISE,” “INC” attached to each word involves eight subsets, as shown in Fig. 26.6.




Fig. 26.6 : The Venn diagram of “DEC,” “INC,” “RISE,” “INC”

To each of the six components $\mathbf{X} \cup \mathbf{X}'$, \mathbf{Y} , \mathbf{Y}' , $\mathbf{Z} \cup \mathbf{Z}'$, \mathbf{T} , \mathbf{T}' , mutually disjoint, will be attached a family of commuting variables, as shown next:

$$\begin{aligned} \mathbf{X} \cup \mathbf{X}' &= \text{DES} \longleftarrow (X_i), \\ \mathbf{Y} &= \text{RISE} \setminus \text{REC} \longleftarrow (Y_i), \quad \mathbf{Y}' = \text{RISE} \cap \text{REC} \longleftarrow (Y'_i), \\ \mathbf{Z} \cup \mathbf{Z}' &= (\text{DEC} \setminus \text{DES}) \longleftarrow (Z_i), \\ \mathbf{T} &= (\text{INC} \setminus \text{RISE}) \setminus \text{REC} \longleftarrow (T_i), \quad \mathbf{T}' = (\text{INC} \setminus \text{RISE}) \cap \text{REC} \longleftarrow (T'_i), \end{aligned}$$

where $i = 1, 2, \dots$

The *weight* $\psi(v)$ of each word $v = y_1 y_2 \cdots y_n$ will be defined by;

$$(26.6) \quad \psi(v) := \prod_{i \in \text{DES}} X_{y_i} \prod_{i \in \text{RISE} \setminus \text{REC}} Y_{y_i} \prod_{i \in \text{DEC} \setminus \text{DES}} Z_{y_i} \\ \times \prod_{i \in (\text{INC} \setminus \text{RISE}) \setminus \text{REC}} T_{y_i} \prod_{i \in \text{RISE} \cap \text{REC}} Y'_{y_i} \prod_{i \in (\text{INC} \setminus \text{RISE}) \cap \text{REC}} T'_{y_i},$$

where the argument “(v)” has not been written for typographic reasons. For example, $i \in \text{RISE} \setminus \text{REC}$ stands for $i \in \text{RISE}(v) \setminus \text{REC}(v)$.

For instance, for the word $v = 3255886630014038$ the sets DES, DEC, INC, RISE, REC of v are indicated by bullets.

v	=	3	2	5	5	8	8	6	6	3	0	0	1	4	0	3	8	
DES	=	•					•	•	•	•					•			
DEC	=	•				•	•	•	•	•					•			
RISE	=		•		•							•	•			•	•	•
INC	=		•	•	•							•	•	•		•	•	•
REC	=	•		•	•	•	•											•
$\psi(v)$	=	X_3	Y_2	T'_5	Y'_5	Z_8	X'_8	Z_6	X_6	X_3	T_0	Y_0	Y_1	X_4	Y_0	Y_3	Y'_8	

The goal of this chapter is to prove the following theorem.

Theorem 26.1 (The Decrease Value Theorem). *With C being the $(r + 1) \times (r + 1)$ matrix*

$$C = \begin{pmatrix} 0 & \frac{X_1}{1 - Z_1} & \frac{X_2}{1 - Z_2} & \cdots & \frac{X_{r-1}}{1 - Z_{r-1}} & \frac{X_r}{1 - Z_r} \\ \frac{Y_0}{1 - T_0} & 0 & \frac{X_2}{1 - Z_2} & \cdots & \frac{X_{r-1}}{1 - Z_{r-1}} & \frac{X_r}{1 - Z_r} \\ \frac{Y_0}{1 - T_0} & \frac{Y_1}{1 - T_1} & 0 & \cdots & \frac{X_{r-1}}{1 - Z_{r-1}} & \frac{X_r}{1 - Z_r} \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ \frac{Y_0}{1 - T_0} & \frac{Y_1}{1 - T_1} & \frac{Y_2}{1 - T_2} & \cdots & 0 & \frac{X_r}{1 - Z_r} \\ \frac{Y_0}{1 - T_0} & \frac{Y_1}{1 - T_1} & \frac{Y_2}{1 - T_2} & \cdots & \frac{Y_{r-1}}{1 - T_{r-1}} & 0 \end{pmatrix}$$

the generating function for the set $[0, r]^*$ by the weight ψ is given by

$$(26.7) \quad \sum_{v \in [0, r]^*} \psi(v) = \frac{\prod_{0 \leq j \leq r} \left(1 + \frac{Y'_j}{1 - T'_j}\right)}{\det(I - C)}.$$

Of course, the expression $1/\det(I - C)$ is too redolent of the MacMahon Master Theorem [Mac13, p. 97-98] for not having it play a crucial role in the proof of Theorem 26.1. It does indeed. However, further tools are needed to complete the proof, in particular, the properties of the so-called *first fundamental transformation for arbitrary words*, as developed in Cartier-Foata [CaFo69] and also in Lothaire [Lo02, chap. 10].

26.1. *The first fundamental transformation for arbitrary words revisited.* This transformation denoted by \mathbf{F}_1 may be described as follows. Start with the word $w = x_1x_2 \cdots x_n$ from $[0, r]^*$ and let $\mathbf{x}_{i_1} \leq \mathbf{x}_{i_2} \leq \cdots \leq \mathbf{x}_{i_m}$ ($1 = i_1 < i_2 < \cdots < i_m \leq n$) be the non-decreasing sequence of its record values, from left to right. Next, cut the word w before each of these record values, to get the following factorization of the word w : $\mathbf{x}_{i_1}w_1 \mid \mathbf{x}_{i_2}w_2 \mid \cdots \mid \mathbf{x}_{i_m}w_m$. Then, form the *cyclic shifts* $w_1\mathbf{x}_{i_1}, w_2\mathbf{x}_{i_2}, \dots, w_m\mathbf{x}_{i_m}$, the *cycles* $\begin{pmatrix} w_1\mathbf{x}_{i_1} \\ \mathbf{x}_{i_1}w_1 \end{pmatrix}, \begin{pmatrix} w_2\mathbf{x}_{i_2} \\ \mathbf{x}_{i_2}w_2 \end{pmatrix}, \dots, \begin{pmatrix} w_m\mathbf{x}_{i_m} \\ \mathbf{x}_{i_m}w_m \end{pmatrix}$ and set up the factorization product $\mathbf{F}'_1(w) := \begin{pmatrix} w_1\mathbf{x}_{i_1} & w_2\mathbf{x}_{i_2} & \cdots & w_m\mathbf{x}_{i_m} \\ \mathbf{x}_{i_1}w_1 & \mathbf{x}_{i_2}w_2 & \cdots & \mathbf{x}_{i_m}w_m \end{pmatrix}$. Finally, rearrange the vertical biletters of that two-row matrix in such a way that the entries on the top row are in *non-decreasing order*, assuming that two biletters $\begin{pmatrix} a \\ a' \end{pmatrix}, \begin{pmatrix} b \\ b' \end{pmatrix}$ can commute only when $a \neq b$. We then

obtain the two-row matrix $\mathbf{F}_1(w) := \begin{pmatrix} \bar{v} \\ v \end{pmatrix} = \begin{pmatrix} z_1 & z_2 & \cdots & z_n \\ y_1 & y_2 & \cdots & y_n \end{pmatrix}$. The reverse transformation \mathbf{F}_1^{-1} is fully described in [Lo02, chap. 10], where it is also shown that to each integer i such that $1 \leq i \leq n - 1$ and $x_i > x_{i+1}$ there corresponds a unique j such that $z_j = x_{i+1} < x_i = y_j$ in a bijective manner. As a consequence, this latter property implies that descent and so-called exceedance numbers are equidistributed on every rearrangement class of words. A stronger property will be proved later when dealing with so-called horizontal and vertical rearrangements.

Example. Start with the word $w = 536533246612431$ and cut it before each record value to get $w = 53 \mid 653324 \mid 6 \mid 612431$. Form the cyclic shifts $35, 533246, 6, 124316$ and then the two-row matrix $\mathbf{F}'_1(w) = \begin{pmatrix} 355332466124316 \\ 536533246612431 \end{pmatrix}$. Finally, rearrange the vertical biletters as mentioned above to get $\mathbf{F}_1(w) := \begin{pmatrix} \bar{u} \\ u \end{pmatrix} = \begin{pmatrix} 11223334455666 \\ 63315542236461 \end{pmatrix}$.

26.2. *Horizontal derangements.* Call *h-derangement* (horizontal derangement) each word from $[0, r]^*$ that has no equal letters in succession.

If w is such a word, let $\begin{pmatrix} w_j & \mathbf{x}_{i_j} \\ \mathbf{x}_{i_j} & w_j \end{pmatrix}$ be one of its cycles attached to it (using the notation of the previous section). To avoid cumbersome sub-subscripts write it as $\begin{pmatrix} a_1 & a_2 & \cdots & a_{\ell-2} & a_{\ell-1} & \mathbf{x}_{i_j} \\ \mathbf{x}_{i_j} & a_1 & a_2 & \cdots & a_{\ell-2} & a_{\ell-1} \end{pmatrix}$. If this cycle is of length $\ell \geq 2$, then all its vertical biletters $\begin{pmatrix} a_1 \\ \mathbf{x}_{i_1} \end{pmatrix}, \begin{pmatrix} a_2 \\ a_1 \end{pmatrix}, \dots, \begin{pmatrix} \mathbf{x}_{i_1} \\ a_{\ell-1} \end{pmatrix}$ have different top and bottom letters: $a_1 \neq \mathbf{x}_{i_j}, a_2 \neq a_1, \dots, \mathbf{x}_{i_j} \neq a_{\ell-1}$.

On the contrary, if the cycle is of length $\ell = 1$, so that it is reduced to a single biletter $\begin{pmatrix} \mathbf{x}_{i_j} \\ \mathbf{x}_{i_j} \end{pmatrix}$, whose top and bottom letters are equal, then \mathbf{x}_{i_j} is both a rise and record value [in short, a *riserecord*] of the word w .

The equality $x_i = x_{i+1}$ occurs in the word $w = x_1 x_2 \cdots x_n$ if and only if x_i is either a decrease, but not a descent value, or an increase, but not a rise. Accordingly, if w is an *h-derangement*, then $(\text{DEC} \setminus \text{DES})(w) = \emptyset$, $(\text{INC} \setminus \text{RISE} \setminus \text{REC})(w) = \emptyset$, $(\text{INC} \setminus \text{RISE} \cap \text{REC})(w) = \emptyset$. If, furthermore, the *h-derangement* w has no riserecord, then $(\text{RISE} \cap \text{REC})(w) = \emptyset$, so that

$$(26.8) \quad \psi(w) = \prod_{a \in \text{DES}(w)} X_a \times \prod_{b \in (\text{RISE} \setminus \text{REC})(w)} Y_b,$$

For each pair of distinct integers (a, b) from $[0, r]$ define

$$(26.9) \quad \Phi \begin{pmatrix} a \\ b \end{pmatrix} := \begin{cases} X_b, & \text{if } b > a; \\ Y_b & \text{if } b < a; \end{cases}$$

where X_a and Y_b refer to the commuting variables introduced in (26.6), indicating that the letter a (resp. b) is a descent (resp. a rise, but not a record) value. Furthermore, extend Φ to an homomorphism defined on the free monoid $[0, r]^*$.

Proposition 26.2. *Let $w = x_1x_2 \cdots x_n$ be an h -derangement having no riserecords. Then, (26.8) holds and also*

$$(26.10) \quad \Phi(\mathbf{F}_1(w)) = \psi(w).$$

Proof. Consider the factorisation product

$$\mathbf{F}'_1(w) = \begin{pmatrix} w_1\mathbf{x}_{i_1} & w_2\mathbf{x}_{i_2} & \cdots & w_m\mathbf{x}_{i_m} \\ \mathbf{x}_{i_1}w_1 & \mathbf{x}_{i_2}w_2 & \cdots & \mathbf{x}_{i_m}w_m \end{pmatrix},$$

introduced in the construction of $\mathbf{F}_1(w)$, also written as $\mathbf{F}'_1(w) = \begin{pmatrix} c_1 & c_2 & \cdots & c_n \\ x_1 & x_2 & \cdots & x_n \end{pmatrix}$ and one of its cycles

$$\begin{pmatrix} w_j & \mathbf{x}_{i_j} \\ \mathbf{x}_{i_j} & w_j \end{pmatrix} = \begin{pmatrix} a_1 & a_2 & \cdots & a_{\ell-2} & a_{\ell-1} & \mathbf{x}_{i_j} \\ \mathbf{x}_{i_j} & a_1 & a_2 & \cdots & a_{\ell-2} & a_{\ell-1} \end{pmatrix}.$$

Next, set up the mapping $\begin{pmatrix} c_i \\ x_i \end{pmatrix} \mapsto x_i x_{i+1}$ for $i = 1, 2, \dots, n$, where $x_{n+1} = +\infty$ by convention.

Its action on the biletters of the cycle $\begin{pmatrix} w_j & \mathbf{x}_{i_j} \\ \mathbf{x}_{i_j} & w_j \end{pmatrix}$ reads: $\begin{pmatrix} a_1 \\ \mathbf{x}_{i_j} \end{pmatrix} \mapsto \mathbf{x}_{i_j} a_1$,

$\begin{pmatrix} a_2 \\ a_1 \end{pmatrix} \mapsto a_1 a_2, \dots, \begin{pmatrix} a_{\ell-1} \\ a_{\ell-2} \end{pmatrix} \mapsto a_{\ell-1} a_{\ell-2}, \begin{pmatrix} \mathbf{x}_{i_j} \\ a_{\ell-1} \end{pmatrix} \mapsto a_{\ell-1} \mathbf{x}_{i_{j+1}}$, where $\mathbf{x}_{i_{j+1}}$ is the record value following $a_{\ell-1}$ in the word w , or $+\infty$ when dealing with the rightmost cycle. In any case, both inequalities $a_{\ell-1} < \mathbf{x}_{i_j}$ and $a_{\ell-1} < \mathbf{x}_{i_{j+1}}$ hold, as $a_{\ell-1} < \mathbf{x}_{i_j} \leq \mathbf{x}_{i_{j+1}}$, since both \mathbf{x}_{i_j} and $\mathbf{x}_{i_{j+1}}$ are two successive records.

Consequently, $c_i \neq x_i$ for all $i = 1, 2, \dots, n$, as w has no equal letters in succession. Furthermore, for $1 \leq j \leq n - 1$ we have: $c_j < x_j$ (resp. $c_j > x_j$) if and only if $x_j > x_{j+1}$ (resp. $x_j < x_{j+1}$), that is, if and only if x_j is a descent (resp. a rise) value of the word $w = x_1x_2 \cdots x_n$.

As $\mathbf{F}_1(w) = \begin{pmatrix} z_1 & z_2 & \cdots & z_n \\ y_1 & y_2 & \cdots & y_n \end{pmatrix}$ is obtained from $\mathbf{F}'_1(w)$ by a well-defined permutation of its biletters, there corresponds to each descent value $x_i > x_{i+1}$ and variable X_i (resp. rise value $x_i < x_{i+1}$ and variable Y_i) of w a unique billetter $\begin{pmatrix} z_j \\ y_j \end{pmatrix}$ such that $z_j > y_j$ (resp. $z_j < y_j$). This establishes (26.10). \square

Example. The word $w = 5365324612431$ is a horizontal derangement. By cutting it before each record value one gets $w = 53 \mid 65324 \mid$

612431 =: $w_1 \mid w_2 \mid w_3$, so that w has no riserecord. Next,

$$\mathbf{F}'_1(w) = \begin{pmatrix} 3 & 5 & 5 & 3 & 2 & 4 & 6 & 1 & 2 & 4 & 3 & 1 & 6 \\ \wedge & \vee & \wedge & \wedge & \wedge & \vee & \vee & \wedge & \vee & \vee & \wedge & \wedge & \vee \\ 5 > 3 < 6 > 5 > 3 > 2 < 4 < 6 > 1 < 2 < 4 > 3 > 1 \end{pmatrix}.$$

Note that each right angle with three entries is of the form $\begin{pmatrix} a \\ \wedge \\ b > a \end{pmatrix}$ or $\begin{pmatrix} a \\ \vee \\ b < a \end{pmatrix}$; except the second one $\begin{pmatrix} 5 \\ \vee \\ 3 < 6 \end{pmatrix}$ and $\begin{pmatrix} 6 \\ \vee \\ 1 < +\infty \end{pmatrix}$ (by convention), but in each case the entry in the corner of the right angle is either greater, or smaller than both its ends.

By rearranging the vertical biletters of $\mathbf{F}'_1(w)$ in such a way that the entries on the top row are in nondecreasing order, assuming that two biletters $\begin{pmatrix} a \\ a' \end{pmatrix}$, $\begin{pmatrix} b \\ b' \end{pmatrix}$ can commute only when $a \neq b$, we obtain the two-row matrix $\mathbf{F}_1(w) = \begin{pmatrix} 1 & 1 & 2 & 2 & 3 & 3 & 3 & 4 & 4 & 5 & 5 & 6 & 6 \\ 6 & 3 & 3 & 1 & 5 & 5 & 4 & 2 & 2 & 3 & 6 & 4 & 1 \end{pmatrix}$. As the variables X_i and Y_j commute with each other, we have:

$$\begin{aligned} \psi(w) &= X_5 Y_3 X_6 X_5 X_3 Y_2 Y_4 X_6 Y_1 Y_2 X_4 X_3 Y_1; \\ &= X_6 X_3 X_3 Y_1 X_5 X_5 X_4 Y_2 Y_2 Y_3 X_6 Y_4 Y_1; \\ &= \Phi \begin{pmatrix} 1 \\ 6 \end{pmatrix} \Phi \begin{pmatrix} 1 \\ 3 \end{pmatrix} \Phi \begin{pmatrix} 2 \\ 3 \end{pmatrix} \Phi \begin{pmatrix} 2 \\ 1 \end{pmatrix} \cdots; \\ &= \Phi(\mathbf{F}_1(w)). \end{aligned}$$

26.3. Vertical derangements. Call *v-derangement* (vertical derangement) the image under the first fundamental transformation \mathbf{F}_1 of each *h-derangement* having no riserecords, so that the following definition can be stated.

Definition. A word $w = x_1 x_2 \cdots x_n$ from $[0, r]^*$ is said to be a *v-derangement* (vertical derangement), if the biword $\begin{pmatrix} \bar{w} \\ w \end{pmatrix} = \begin{pmatrix} \bar{x}_1 & \bar{x}_2 & \cdots & \bar{x}_n \\ x_1 & x_2 & \cdots & x_n \end{pmatrix}$, where $\bar{w} = \bar{x}_1 \bar{x}_2 \cdots \bar{x}_n$ is obtained from w by rearranging its letters in non-decreasing order, has no (vertical) biletter $\begin{pmatrix} \bar{x}_i \\ x_i \end{pmatrix}$ such that $\bar{x}_i = x_i$.

The set of all *h-derangements* (resp. all *v-derangements*) is denoted by $[0, r]_h^*$ (resp. by $[0, r]_v^*$). As proved in the previous Proposition, the transformation \mathbf{F}_1 provides a bijection of the set of all *h-derangements* having no riserecords onto $[0, r]_v^*$. The next step is to construct a bijection of all of $[0, r]_h^*$ onto the Cartesian product $\text{StrictInc}(r) \times [0, r]_v^*$, where $\text{StrictInc}(r)$ is the set of all the 2^{r+1} strictly increasing words $x_1 x_2 \cdots x_\ell$ such that $0 \leq \ell \leq r$ and $0 \leq x_1 < x_2 < \cdots < x_\ell \leq r$

Consider an h -derangement $w = x_1x_2 \cdots x_n$. Each riserecord of w has two characteristics: (i) there are no riserecord values to its left in w equal to it; (ii) the letter to its immediate right is necessarily a record value (rise or descent). Consequently, the subword $w_0 := x_{h_1}x_{h_2} \cdots x_{h_s}$ ($1 \leq h_1 < h_2 < \cdots < h_s \leq n$) of w , made of all the letters of w , which are riserecords, is *strictly increasing*.

Let v_0 be the word derived from w after removal of w_0 . Properties (i) and (ii) imply that to reconstruct the word w from the pair (w_0, v_0) it suffices to insert x_{h_s} just before the leftmost record of v_0 greater than x_{h_s} , assuming that $x_{n+1} = +\infty$ is a record by convention. Let $v_{0,1}$ be the word just obtained. Next, insert $x_{h_{s-1}}$ into $v_{0,1}$ just before the leftmost record of $v_{0,1}$ greater than $x_{h_{s-1}}$, and so on. Continue in the same way until the smallest riserecord x_{h_1} is inserted into the word $v_{0,s-1}$ derived by the procedure. Accordingly, the following propositions holds.

Proposition 26.3. *Let $w = x_1x_2 \cdots x_n$ be an h -derangement and $w_0 = x_{h_1}x_{h_2} \cdots x_{h_s}$ be the subword of w made of all the riserecord values of w . Let v_0 be the word derived from w after deleting the letters of w_0 and let*

$$\mathbf{F}_1(v_0) := \begin{pmatrix} \bar{u} \\ u \end{pmatrix}. \text{ Then,}$$

- (i) *the subword w_0 is strictly increasing: $x_{h_1} < x_{h_2} < \cdots < x_{h_s}$;*
- (ii) *the mapping $w \mapsto (w_0, u)$ provides a bijection of $[0, r]_h^*$ onto the Cartesian product $\text{StrictInc}(r) \times [0, r]_v^*$ having the property:*

$$(26.11) \quad \psi(w) = \psi(w_0)\psi(v_0),$$

also equal to

$$(26.12) \quad = \psi(w_0)\Phi \begin{pmatrix} \bar{u} \\ u \end{pmatrix}$$

by (26.10).

Example. Consider the following h -derangement w , where the riserecords have been reproduced in bold-face:

$$w = \mathbf{2} \mid 53 \mid \mathbf{5} \mid 65324 \mid 612431 \mid \mathbf{6}.$$

The pair (v_0, w_0) reads:

$$v_0 = \mathbf{256}, \quad w_0 = 5365324612431.$$

For the reinsertion of w_0 into v_0 insert 6 to the right of v_0 , next 5 just before the leftmost letter equal to 6; finally, 2 to the extreme left. The

biword $\mathbf{F}_1(v_0) = \begin{pmatrix} \bar{u} \\ u \end{pmatrix}$ has been derived in the previous two examples.

It follows from Proposition 26.3 (i) that all the letters of w_0 are riserecords, so that $\psi(w_0)$ only involves variables Y'_j . Thus,

$$(26.13) \quad \sum_{w_0 \in \text{StrictInc}(r)} \psi(w_0) = \prod_{0 \leq j \leq r} (1 + Y'_j).$$

Furthermore, Proposition 26.3 (ii) implies

$$(26.14) \quad \begin{aligned} \sum_{w \in [0, r]_h^*} \psi(w) &= \sum_{w_0 \in \text{StrictInc}(r)} \psi(w_0) \sum_{u \in [0, r]_v^*} \Phi \begin{pmatrix} \bar{u} \\ u \end{pmatrix} \\ &= \prod_{0 \leq j \leq r} (1 + Y'_j) \Phi \left(\sum_{u \in [0, r]_v^*} \begin{pmatrix} \bar{u} \\ u \end{pmatrix} \right). \end{aligned}$$

The calculation of the generating function $\sum_{u \in [0, r]_v^*} \begin{pmatrix} \bar{u} \\ u \end{pmatrix}$ for the vertical v -derangements was derived in Exercice 35 in the form

$$(26.15) \quad \sum_{u \in [0, r]_v^*} \begin{pmatrix} \bar{u} \\ u \end{pmatrix} = \frac{1}{\det(I - C'_r)},$$

where C'_r is the $((r + 1) \times (r + 1))$ matrix

$$C'_r = \begin{pmatrix} 0 & \binom{0}{1} & \binom{0}{2} & \cdots & \binom{0}{r-1} & \binom{0}{r} \\ \binom{1}{0} & 0 & \binom{1}{2} & \cdots & \binom{1}{r-1} & \binom{1}{r} \\ \binom{2}{0} & \binom{2}{1} & 0 & \cdots & \binom{2}{r-1} & \binom{2}{r} \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ \binom{r-1}{0} & \binom{r-1}{1} & \binom{r-1}{2} & \cdots & 0 & \binom{r-1}{r} \\ \binom{r}{0} & \binom{r}{1} & \binom{r}{2} & \cdots & \binom{r}{r-1} & 0 \end{pmatrix}.$$

26.4. *End of the proof of Theorem 26.1.* By definition of Φ (given in (26.9)) and (26.15) we conclude::

$$\begin{aligned} \sum_{w \in [0, r]_h^*} \psi(w) &= \prod_{0 \leq j \leq r} (1 + Y'_j) \Phi \left(\sum_{u \in [0, r]_v^*} \begin{pmatrix} \bar{u} \\ u \end{pmatrix} \right) \\ &= \prod_{0 \leq j \leq r} (1 + Y'_j) \Phi(1/\det(I - C'_r)) \\ &= \prod_{0 \leq j \leq r} (1 + Y'_j) (1/\det(I - C'')), \end{aligned}$$

where

$$C'' = \begin{pmatrix} 0 & X_1 & X_2 & \cdots & X_{r-1} & X_r \\ Y_0 & 0 & X_2 & \cdots & X_{r-1} & X_r \\ Y_0 & Y_1 & 0 & \cdots & X_{r-1} & X_r \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ Y_0 & Y_1 & Y_2 & \cdots & 0 & X_r \\ Y_0 & Y_1 & Y_2 & \cdots & Y_{r-1} & 0 \end{pmatrix}$$

Let $w = x_1x_2\cdots x_n$ be a word from $[0, r]^*$. The *key* of w is defined to be the h -derangement v derived from w by erasing all letters x_i such that $x_i = x_{i+1}$. For instance, the key of $w = 324455531114135$ is the h -derangement $v = 3245314135$.

Let β be the substitution of variables defined by

$$(26.16) \quad \beta := \{X_i \leftarrow X_i/(1 - Z_i), Y_i \leftarrow Y_i/(1 - T_i), Y'_i \leftarrow Y'_i/(1 - T'_i)\}.$$

Then, the generating function for the set of all w whose key is v by the weight ψ is given by

$$\sum_{w, \text{key}(w)=v} \psi(w) = \beta \psi(v).$$

Hence,

$$\begin{aligned} (26.17) \quad \sum_{w \in [0, r]^*} \psi(w) &= \sum_{v \in [0, r]_h^*} \sum_{\text{key}(w)=v} \psi(w) = \sum_{v \in [0, r]_h^*} \beta \psi(v) \\ &= \beta \left(\sum_{v \in [0, r]_h^*} \psi(v) \right) = \beta \frac{\prod_{0 \leq j \leq r} (1 + Y'_j)}{\det(I - C'')} \\ &= \frac{\prod_{0 \leq j \leq r} (1 + \beta Y'_j)}{\det(I - \beta C'')} = \frac{\prod_{0 \leq j \leq r} \left(1 + \frac{Y'_j}{1 - T'_j}\right)}{\det(I - C)}. \quad \square \end{aligned}$$

27. The Decrease Value Theorem; from words to permutations

The goal of this chapter is to make use of the various applications of the Decrease Value Theorem for calculating the distributions of several multivariable statistics, no longer on arbitrary words, but on classical groups, essentially, the symmetric group and possibly the hyperoctahedral group. The basic tool to transfer the calculation from words to groups is to

explore the properties of the Gessel-Reutenauer standardization [GeRe93], which is based on the theory of Lyndon words. An excellent review of this latter theory can be found in Lothaire’s book [Lo83, pp. 64–68].

27.1. *Lyndon words.* Let $l = x_1x_2 \cdots x_n$ be a nonempty word from $[0, r]^*$; it is said to be a *Lyndon word*, if either $n = 1$, or if $n \geq 2$ and, with respect to the lexicographic order, the inequality $x_1x_2 \cdots x_n > x_ix_{i+1} \cdots x_nx_1 \cdots x_{i-1}$ holds for every i such that $2 \leq i \leq n$. Classically, each Lyndon word is defined to be the minimum within its class of cyclic rearrangements. The modification in here is made for convenience.

Let w, w' be two nonempty primitive words (none of them can be expressed as v^b , where v is a word and b an integer greater than or equal to 2). We write $w \preceq w'$ if and only if $w^b \leq w'^b$, with respect to the lexicographic order, when b is large enough. The main result on Lyndon words is the following (see [Lo83, Theorem 5.1.5]):

Theorem 27.1. *Each nonempty word w , whose letters are nonnegative integers, can be written uniquely as a product $l_1l_2 \cdots l_k$, where each l_i is a Lyndon word and $l_1 \preceq l_2 \preceq \cdots \preceq l_k$.*

For instance, the factorization of the following word as a nondecreasing product of Lyndon words with respect to “ \preceq ” [in short, *Lyndon word factorization*] is indicated by vertical bars:

$$w = | 2 | 3 2 1 1 | 3 2 | 3 2 | 3 | 5 | 6 4 1 | 6 6 3 1 6 6 2 | 6 |$$

For instance, $6 6 3 1 6 6 2 \preceq 6$ holds, since $6 6 3 1 6 6 2 < 6 6 6 6 \cdots$

27.2. *The Gessel-Reutenauer standardization.* Recall that the number of descents of each permutation $\sigma \in \mathfrak{S}_n$ is denoted by “ $\text{des } \sigma$.” By $\mathfrak{S}_{n, \leq r}$ is meant the set of all $\sigma \in \mathfrak{S}_n$ such that $\text{des } \sigma \leq r$. Moreover, the set of all words from $[0, r]^*$ of length n is designated by $[0, r]^n$ and the subset of all its *non increasing* words $w = x_1x_2 \cdots x_n$ ($r \geq x_1 \geq x_2 \geq \cdots \geq x_n \geq 0$) by $\text{NIW}_n(r)$. The Gessel-Reutenauer standardization is a bijection

$$(27.1) \quad \begin{array}{ccc} w & \mapsto & (\sigma, c) \\ [0, r]^n & \longrightarrow & \{(\sigma, c) : \sigma \in \mathfrak{S}_{n, \leq r}, c \in \text{NIW}_n(r - \text{des } \sigma)\}. \end{array}$$

Example. For $r = 1$ and $n = 3$ the number of words from $[0, r]^n$ is equal to $2^3 = 8$ and $\mathfrak{S}_{n, \leq r}$ consists of the permutations 123, 213, 312, 132, 231. There are 8 pairs (σ, c) corresponding to words from $[0, 1]^3$ in (27.1), namely, (123, 000), (123, 100), (123, 110), (123, 111), (213, 000), (312, 000), (132, 000), (231, 000).

It is convenient to denote each non increasing word from $NIW_n(r)$, either as $\mathbf{m} = \bar{c}_1 \bar{c}_2 \cdots \bar{c}_n$, or as $m_1^{n_1} m_2^{n_2} \cdots m_s^{n_s}$ under the conditions:

$$(27.2) \quad r \geq m_1 > m_2 > \cdots > m_s \geq 0 \quad (n_1 + n_2 + \cdots + n_s = n),$$

$$n_1 \geq 1, n_2 \geq 1, \dots, n_s \geq 1.$$

Let $R(\mathbf{m})$ be the set of all the $n!/(n_1!n_2! \dots n_s!)$ rearrangements of \mathbf{m} . If w belongs to $R(\mathbf{m})$, we shall write: $\bar{c}(w) = \mathbf{m}$. Note that $[0, r]^n$ is the disjoint union of all the $R(\mathbf{m})$ with \mathbf{m} satisfying conditions (27.2).

The main idea behind was to imagine that the Lyndon word factorization of a word w from each $R(\mathbf{m})$ could serve to build up a permutation σ , whose cycles would have the same lengths as the lengths of the Lyndon factors of w , and provide a total ordering of all the letters of w . To accomplish this goal it suffices to start with the Lyndon word factorization (l_1, l_2, \dots, l_k) of w . If x is a letter of the factor $l_i = y_1 \cdots y_{j-1} x y_{j+1} \cdots y_h$, form the cyclic rearrangement $\text{cyc}(x) := x y_{j+1} \cdots y_h y_1 \cdots y_{j-1}$. When the Lyndon factor is the one-letter $l = x$, it is convenient to let the cyclic rearrangement be a word $x x \cdots x$ with an arbitrary number of x 's (see, for instance, the example at the end of the previous subsection). If x, y are two letters of w , say that x precedes y , if $\text{cyc } x \text{ succ cyc } y$, or if $\text{cyc } x = \text{cyc } y$ and the letter x is to the right of the letter y in the word w .

This defines a total ordering, as two letters from the same Lyndon factor l_i precede one another, because l_i is primitive, as well as two letters from different factors, because the Lyndon factors are themselves totally ordered. Accordingly, to each integer i ($1 \leq i \leq n$) there corresponds a unique integer $\check{\sigma}(i)$, which is the number of letters in the word $w = x_1 x_2 \cdots x_n$ preceding x_i plus one.

When replacing each letter x_i in the Lyndon word factorization of w by $\check{\sigma}(i)$, we obtain a word $\check{\sigma} := \check{\sigma}(1)\check{\sigma}(2) \cdots \check{\sigma}(n)$ which is a permutation of $1 2 \dots n$. Accordingly, for $i \neq j$, the letter $x_{\check{\sigma}^{-1}(i)}$ precedes $x_{\check{\sigma}^{-1}(j)}$ if and only if, either $\text{cyc}(x_{\check{\sigma}^{-1}(i)}) \text{ succ cyc}(x_{\check{\sigma}^{-1}(j)})$, or $\text{cyc}(x_{\check{\sigma}^{-1}(i)}) = \text{cyc}(x_{\check{\sigma}^{-1}(j)})$ and $i > j$. Moreover, the mapping $w \mapsto \check{\sigma}$ is an injection of $R(\mathbf{m})$ into \mathfrak{S}_n , and the word $x_{\check{\sigma}^{-1}(1)} x_{\check{\sigma}^{-1}(2)} \cdots x_{\check{\sigma}^{-1}(n)}$ is nothing but the monotonic *nonincreasing* rearrangement $\bar{c}(w) = \mathbf{m} = m_1^{n_1} m_2^{n_2} \cdots m_s^{n_s}$ of w .

Now, the word $\check{\sigma}$ can be split as a product $v_1 \mid v_2 \mid \cdots \mid v_k$, where each v_i has the same length as the Lyndon factor l_i ($1 \leq i \leq k$). As the Lyndon word factorization of w is a *non decreasing* product of factors, the splitting of the word $\check{\sigma}$ defines a *decreasing* product $v_1 > v_2 > \cdots > v_k$

with respect to the lexicographic order and each factor v_i starts with its *minimum* letter.

$$\begin{pmatrix} \text{Id} = & 1 & 2 & \cdots & n \\ w = & x_1 & x_2 & \cdots & x_n = l_1 & | & l_2 & | \cdots & | & l_k \\ \check{\sigma} = & \check{\sigma}(1) & \check{\sigma}(2) & \cdots & \check{\sigma}(n) = v_1 & | & v_2 & | \cdots & | & v_k \\ & & & & \sigma^{cyc} = & (v_1) & (v_2) & \cdots & & (v_k) \end{pmatrix}$$

Fig. 27.1 : The construction of the permutation σ .

The vertical bars can be further reinterpreted as defining the *cycle decomposition* of a new permutation, expressed as a *product of disjoint cycles*, as shown in Fig. 27.1, where the notation σ^{cyc} has been adopted, while the usual two-row matrix is denoted by $\sigma = \begin{pmatrix} 1 & 2 & \cdots & n \\ \sigma(1) & \sigma(2) & \cdots & \sigma(n) \end{pmatrix}$. If (v_i) reads $(i_1 i_2 \cdots i_m)$ as a cycle, then $\sigma(i_1) = i_2$, $\sigma(i_2) = i_3$, \dots , $\sigma(i_{m-1}) = i_m$, $\sigma(i_m) = \sigma(i_1)$. When going through the cycles of σ from left to right, the cycles start *with their minima* and they are in *decreasing* order. See Example 27.2 further displayed. Also, the mapping $\sigma = \sigma(1)\sigma(2)\cdots\sigma(n) \mapsto \check{\sigma} = \check{\sigma}(1)\check{\sigma}(2)\cdots\check{\sigma}(n)$ is nothing but the classical bijection, known as the *First Fundamental Transformation*, fully described in Lothaire's book [Lo83, § 16.2]. Hence, the composition product $w \mapsto \check{\sigma} \mapsto \sigma^{cyc} \mapsto \sigma$ is an *injection* of $R(\mathbf{m})$ into \mathfrak{S}_n we shall denote by $\phi : w \mapsto \sigma = \phi(w)$.

Property 27.2. *Let $\bar{c}(w) = \mathbf{m} = \bar{c}_1\bar{c}_2\cdots\bar{c}_n$ as defined in (27.2) and $\sigma = \phi(w)$. Then, $\sigma(i) > \sigma(i+1) \Rightarrow \bar{c}_i > \bar{c}_{i+1}$.*

Proof. Suppose that $\bar{c}_i = \bar{c}_{i+1}$, that is, $x_{\check{\sigma}^{-1}(i)} = x_{\check{\sigma}^{-1}(i+1)}$, so that we have

$$\left(\begin{array}{cccc} w = \cdots & x_{\check{\sigma}^{-1}(i)} & \cdots & x_{\check{\sigma}^{-1}(i+1)} & \cdots \\ \check{\sigma} = \cdots & i & \cdots & (i+1) & \cdots \end{array} \right) \text{ or } \left(\begin{array}{cccc} w = \cdots & x_{\check{\sigma}^{-1}(i+1)} & \cdots & x_{\check{\sigma}^{-1}(i)} & \cdots \\ \check{\sigma} = \cdots & (i+1) & \cdots & i & \cdots \end{array} \right)$$

The cycles $\text{cyc}(x_{\check{\sigma}^{-1}(i)})$ and $\text{cyc}(x_{\check{\sigma}^{-1}(i+1)})$ that serve to define the preceding order have both their first letters equal to \bar{c}_i . Also, by definition of σ , the cycles $\text{cyc}(x_{\check{\sigma}^{-1}(i)})$ and $\text{cyc}(x_{\check{\sigma}^{-1}(i+1)})$ may be written: $x_{\check{\sigma}^{-1}(i)} x_{\check{\sigma}^{-1}(\sigma(i))} u$ and $x_{\check{\sigma}^{-1}(i+1)} x_{\check{\sigma}^{-1}(\sigma(i+1))} v$, respectively, where u and v are two words, possibly empty. In case where $x_{\check{\sigma}^{-1}(j)}$ ($j = i$ or $(i+1)$) is a one-letter Lyndon factor, we simply have: $\text{cyc}(x_{\check{\sigma}^{-1}(j)}) = x_{\check{\sigma}^{-1}(j)} x_{\check{\sigma}^{-1}(j)}$, as $\sigma(j) = j$.

By deleting these first letters and adding $x_{\check{\sigma}^{-1}(i)}$ and $x_{\check{\sigma}^{-1}(i+1)}$ to the right of the cycles, respectively, we get :

$$\begin{aligned} \text{cyc}(x_{\check{\sigma}^{-1}(\sigma(i))}) &= x_{\check{\sigma}^{-1}(\sigma(i))} u x_{\check{\sigma}^{-1}(i)}; \\ \text{cyc}(x_{\check{\sigma}^{-1}(\sigma(i+1))}) &= x_{\check{\sigma}^{-1}(\sigma(i+1))} v x_{\check{\sigma}^{-1}(i+1)}; \end{aligned}$$

As $x_{\check{\sigma}^{-1}(i)}$ precedes $x_{\check{\sigma}^{-1}(i+1)}$ by definition of $\check{\sigma}$, the letter $x_{\check{\sigma}^{-1}(\sigma(i))}$ precedes $x_{\check{\sigma}^{-1}(\sigma(i+1))}$. If $\text{cyc}(x_{\check{\sigma}^{-1}(\sigma(i))}) \text{succ cyc}(x_{\check{\sigma}^{-1}(\sigma(i+1))})$, then $\sigma(i) < \sigma(i+1)$. Otherwise, $\text{cyc}(x_{\check{\sigma}^{-1}(\sigma(i))}) = \text{cyc}(x_{\check{\sigma}^{-1}(\sigma(i+1))})$ and $x_{\check{\sigma}^{-1}(\sigma(i))}$ is to the right of $x_{\check{\sigma}^{-1}(\sigma(i+1))}$ in w . As each Lyndon factor is primitive, the latter two letters cannot belong to the same Lyndon factor and the former is to the right of the latter one. Accordingly, $\sigma(i) < \sigma(i+1)$:

$$\left(\begin{array}{l} w = \cdots | \cdots x_{\check{\sigma}^{-1}(\sigma(i+1))} \cdots | \cdots x_{\check{\sigma}^{-1}(\sigma(i))} \cdots | \cdots \\ \check{\sigma} = \cdots | \cdots \check{\sigma}(i+1) \cdots | \cdots \check{\sigma}(i) \cdots | \cdots \\ \sigma^{\text{cyc}} = \cdots (\cdots \sigma(i+1) \cdots) (\cdots \sigma(i) \cdots) \cdots \quad \square \end{array} \right)$$

Now, let $\mathfrak{S}(n; n_1, n_2, \dots, n_s)$ be the subset of \mathfrak{S}_n of all permutations τ , whose descent sets “DES τ ” are *contained in* the set $\{n_1, n_1 + n_2, \dots, n_1 + n_2 + \dots + n_{s-1}\}$. This is equivalent to saying that the word $\tau(1)\tau(2) \cdots \tau(n)$ is the juxtaposition product of s *increasing* factors $\tau(1)\tau(2) \cdots \tau(n_1)$, $\tau(n_1 + 1)\tau(n_1 + 2) \cdots \tau(n_1 + n_2)$, \dots , $\tau(n - n_s + 1)\tau(n - n_s + 2) \cdots \tau(n)$; in particular, $\text{des } \tau \leq s - 1 \leq r$. But the number of such permutations is also equal to $n!/(n_1!n_2! \dots n_s!)$. On the other hand, Property 27.2 says that if $\bar{c}(w) = \mathbf{m}$ and $\phi(w) = \sigma$, then each inequality $\sigma(i) < \sigma(i+1)$ implies $\bar{c}_i = \bar{c}_{i+1}$, so that σ belongs to $\mathfrak{S}(n; n_1, n_2, \dots, n_s)$. Thus, $R(\mathbf{m})$ is mapped under the injection ϕ onto a set of equal cardinality. It is then a bijection of $R(\mathbf{m})$ onto $\mathfrak{S}(n; n_1, n_2, \dots, n_s)$. Moreover, if σ belongs to $\mathfrak{S}(n; n_1, n_2, \dots, n_s)$, then $\text{des } \sigma \leq s - 1 \leq r$. Thus, the union of all sets $\mathfrak{S}(n; n_1, n_2, \dots, n_s)$ where n_1, n_2, \dots, n_s are positive of sum n and $s \leq r + 1$ is simply the set $\mathfrak{S}_{n, \leq r}$. Note that the sets $\mathfrak{S}(n; n_1, n_2, \dots, n_s)$ are not disjoint from each other, as the identity permutation $12 \cdots n$ belongs to all of them.

Definition. Let \mathbf{m} be a non increasing word satisfying conditions (27.2) and $\sigma \in \mathfrak{S}_n$. Say that σ is *compatible with* \mathbf{m} [in short, “ σ comp. \mathbf{m} ”], if σ belongs to $\mathfrak{S}(n; n_1, n_2, \dots, n_s)$, or, equivalently, if $\text{DES } \sigma \subset \{n_1, n_1 + n_2, \dots, n_1 + n_2 + \dots + n_{s-1}\}$.

Altogether, the mapping

$$(27.3) \quad \begin{array}{ccc} w & \mapsto & (\phi(w), \bar{c}(w)) \\ [0, r]^n & \longrightarrow & \{(\sigma, \mathbf{m}) : \sigma \in \mathfrak{S}_{n, \leq r}, \mathbf{m} \in \text{NIW}_n(r), \sigma \text{ comp. } \mathbf{m}\} \end{array}$$

Is bijective. The latter condition on the pair (σ, \mathbf{m}) will be reinterpreted by using the descent coding $z(\sigma)$ introduced in Chapter 7, formula (5.2). For each permutation $\sigma \in \mathfrak{S}_n$ let $z(\sigma) = z_1 z_2 \cdots z_n$ be the word defined for $i = 1, 2, \dots, n$ by

$$(27.4) \quad z_i := \#\{j : i \leq j \leq n - 1, \sigma(j) > \sigma(j + 1)\}.$$

Each component z_i is the number of descents of σ , *i.e.*, the number of j such that $i \leq j \leq n - 1$ and $\sigma(j) > \sigma(j + 1)$ within the right factor $\sigma(i)\sigma(i + 1) \cdots \sigma(n)$. In particular, $z_1 = \text{des } \sigma$.

Example. The bijection defined in (27.3) is next illustrated with the case $r = 1, n = 3$. As further explained, $c := \mathbf{m} - z$ componentwise.

w	$\check{\sigma}$	σ	\mathbf{m}	z	c
0 0 0	3 2 1	1 2 3	0 0 0	0 0 0	0 0 0
1 0 0	1 3 2	3 1 2	1 0 0	1 0 0	0 0 0
0 1 0	3 1 2	2 1 3	1 0 0	1 0 0	0 0 0
0 0 1	3 2 1	1 2 3	1 0 0	0 0 0	1 0 0
1 1 0	1 2 3	2 3 1	1 1 0	1 1 0	0 0 0
1 0 1	2 3 1	1 3 2	1 1 0	1 1 0	0 0 0
0 1 1	3 2 1	1 2 3	1 1 0	0 0 0	1 1 0
1 1 1	3 2 1	1 2 3	1 1 1	0 0 0	1 1 1

Fig. 27.2 : The bijections $w \mapsto (\sigma, \mathbf{m}) \mapsto (\sigma, c)$ for $r = 1, n = 3$.

Property 27.3. Let $\mathbf{m} = \bar{c}_1 \bar{c}_2 \cdots \bar{c}_n = m_1^{n_1} m_2^{n_2} \cdots m_s^{n_s}$ satisfy conditions (27.2) and σ be compatible with \mathbf{m} . In particular, $\text{des } \sigma \leq r$. Define

$$(27.5) \quad c_i := \bar{c}_i - z_i \quad (i = 1, 2, \dots, n) \quad \text{and} \quad c := c_1 c_2 \cdots c_n.$$

Then, the word c belongs to $\text{NIW}(r - \text{des } \sigma)$.

Conversely, if $z(\sigma) = z_1 z_2 \cdots z_n$ is the word defined by (27.4), corresponding to a permutation σ such that $\text{des } \sigma \leq r$ and if c belongs to $\text{NIW}(r - \text{des } \sigma)$, then $\bar{c} = \bar{c}_1 \bar{c}_2 \cdots \bar{c}_n$ defined by $\bar{c} = z + c$ componentwise, belongs to $\text{NIW}_n(r)$. Moreover, if $\bar{c} = m_1^{n_1} m_2^{n_2} \cdots m_s^{n_s}$, when expressed with notations (27.2), then σ belongs to $\mathfrak{S}(n; n_1, n_2, \dots, n_s)$.

Proof. In the interval $\{i, i + 1, \dots, n\}$ there are exactly z_i integers j for which $\sigma(j) > \sigma(j + 1)$, and then z_j integers such that $\bar{c}_j \geq \bar{c}_{j+1} + 1$. Hence, $\bar{c}_i \geq \bar{c}_n + z_i \geq z_i$. If i is a descent, then $\bar{c}_i \geq \bar{c}_{i+1} + 1$ and $z_i = z_{i+1} + 1$, so that $c_i = \bar{c}_i + z_i \geq \bar{c}_{i+1} - z_{i+1} = c_{i+1}$; otherwise, $\bar{c}_i \geq \bar{c}_{i+1}$ and $z_i = z_{i+1}$, and the same consequence prevails. In particular, $c_1 = \bar{c}_1 - z_1 \leq r - \text{des } \sigma$. Finally, $c_n = \bar{c}_n - z_n = \bar{c}_n \geq 0$.

For proving the converse note that $\bar{c} = z + c$ is obviously non increasing, as sum of two such words. Next, if $\sigma(i) > \sigma(i + 1)$ and so $z_i = z_{i+1} + 1$. This implies that $\check{c}_i = z_i + c_i = z_{i+1} + 1 + c_{i+1} > z_{i+1} + c_{i+1} = \check{c}_{i+1}$, which implies that σ belongs to $\mathfrak{S}(n; n_1, n_2, \dots, n_s)$. \square

Property 27.4. The mapping

$$(27.6) \quad \begin{array}{l} (\sigma, \mathbf{m}) \\ \{(\sigma, \mathbf{m}) : \sigma \in \mathfrak{S}_{n, \leq r}, \\ \mathbf{m} \in \text{NIW}_n(r), \sigma \text{ comp. } \mathbf{m}\} \end{array} \quad \begin{array}{l} \mapsto \\ \longrightarrow \end{array} \quad \begin{array}{l} (\sigma, c) \\ \{(\sigma, c) : \sigma \in \mathfrak{S}_{n, \leq r}, \\ c \in \text{NIW}_n(r - \text{des } \sigma)\} \end{array}$$

is bijective.

Proof. Let $(\sigma, c), (\sigma, c')$ be the images of the pairs $(\sigma, \mathbf{m}), (\sigma, \mathbf{m}')$ and let $\mathbf{m} = \bar{c}_1\bar{c}_2\cdots\bar{c}_n \neq \mathbf{m}' = \bar{c}'_1\bar{c}'_2\cdots\bar{c}'_n$, so that we may assume that $\bar{c}_1\bar{c}_2\cdots\bar{c}_j = \bar{c}'_1\bar{c}'_2\cdots\bar{c}'_j$, but $\bar{c}_{j+1} > \bar{c}'_{j+1}$ for some j ($1 \leq j \leq n - 1$),. By (27.5) this implies $c_{j+1} > c'_{j+1}$ and then $c \neq c'$. The mapping is truly injective. Furthermore, the converse of Proposition 27.4 says that the mapping is also surjective. \square

With the composition product $w \mapsto (\sigma, \mathbf{m}) \mapsto (\sigma, c)$ of the two bijections displayed in (27.3) and (27.6) the Gessel-Reutenauer standardization has been fully described, as announced in (27.1).

27.3. *The construction of the bijection $w \mapsto (c, \sigma)$.* It is illustrated in the next example: start with a word w from $[0, r]^*$ and make up all the steps previously defined, that is,

$$(27.7) \quad w \mapsto (l_1, l_2, \dots, l_k) \mapsto (\check{\sigma}, \mathbf{m}) \mapsto (\sigma^{cyc}, \mathbf{m}) \mapsto (\sigma, \mathbf{m}) \mapsto (\sigma, c),$$

$$\left(\begin{array}{l} w = 2 \mid 3 \ 2 \ 1 \ 1 \mid 3 \ 2 \mid 3 \ 2 \mid 3 \mid 5 \mid 6 \ 4 \ 1 \mid 6 \ 6 \ 3 \ 1 \ 6 \ 6 \ 2 \mid 6 \\ \check{\sigma} = 17 \mid 12 \ 18 \ 22 \ 21 \mid 11 \ 16 \mid 10 \ 15 \mid 9 \mid 7 \mid 4 \ 8 \ 20 \mid 2 \ 5 \ 13 \ 19 \ 3 \ 6 \ 14 \mid 1 \\ \sigma^{cyc} = (17) (12 \ 18 \ 22 \ 21) (11 \ 16) (10 \ 15) (9) (7) (4 \ 8 \ 20) (2 \ 5 \ 13 \ 19 \ 3 \ 6 \ 14) (1) \\ \sigma = \left(\begin{array}{cccccccccccccccccccc} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 & 10 & 11 & 12 & 13 & 14 & 15 & 16 & 17 & 18 & 19 & 20 & 21 & 22 \\ 1 & 5 & 6 & 8 & 13 & 14 & 7 & 20 & 9 & 15 & 16 & 18 & 19 & 2 & 10 & 11 & 17 & 22 & 3 & 4 & 12 & 21 \end{array} \right) \\ \mathbf{m} = 6 \ 6 \ 6 \ 6 \ 6 \ 6 \ 5 \ 4 \ 3 \ 3 \ 3 \ 3 \ 3 \ 2 \ 2 \ 2 \ 2 \ 2 \ 1 \ 1 \ 1 \ 1 \ 1 \\ z = 4 \ 4 \ 4 \ 4 \ 4 \ 4 \ 3 \ 3 \ 2 \ 2 \ 2 \ 2 \ 2 \ 1 \ 1 \ 1 \ 1 \ 1 \ 0 \ 0 \ 0 \ 0 \\ c = 2 \ 2 \ 2 \ 2 \ 2 \ 2 \ 2 \ 2 \ 1 \ 1 \ 1 \ 1 \ 1 \ 1 \ 1 \ 1 \ 1 \ 1 \ 1 \ 1 \ 1 \ 1 \end{array} \right)$$

Example 27.2 : The bijection $w \mapsto (c, \sigma)$

In the example $n = 22$. The first row contains the starting word w , whose Lyndon factorization has been indicated by vertical bars. In the second row each letter x of w is replaced by its rank with respect to the “ \prec ” order. For instance, the rightmost letter 6, whose cyclic rearrangement 666... is the greatest one, gets rank 1. The word w has two Lyndon factors equal to (32); the letters in the left factor get ranks greater than the letters in the right one. The third row has the same block splitting as the second row, but the blocks, now between parentheses, serve to define a permutation σ^{cyc} as a product of cycles. The next row shows the same permutation, expressed as a two-row matrix. Next, comes the non increasing rearrangement $\bar{c}(w) = \mathbf{m}$ of w , then the word z defined in (27.4) and finally the word $c := \bar{c} - z$.

The reverse bijection $(\sigma, c) \mapsto w$ consists of starting with a permutation σ from $\mathfrak{S}_{n, \leq r}$ and a word $c = c_1c_2\cdots c_n$ from $\text{NIW}_n(r - \text{des } \sigma)$. Next, display the word $z = z_1z_2\cdots z_n$ defined by (27.4) and add it to c componentwise. This yields a non increasing word $\mathbf{m} = \bar{c}_1\bar{c}_2\cdots\bar{c}_n$ from $\text{NIW}_n(r)$. Moreover, since $z_i > z_{i+1}$ if and only if i is a descent of σ , we have: $\bar{c}_i = \bar{c}_{i+1} \Rightarrow \sigma(i) < \sigma(i + 1)$, that is, σ comp. \mathbf{m} . Next, the

bijection $\sigma \mapsto \sigma^{cyc}$ maps the two-matrix σ onto the product σ^{cyc} of its disjoint cycles, with the convention that *each cycle starts with its minimum element, and these minima are in decreasing order when reading the whole product from left to right*. This is the essential step in the construction. Removing the parentheses of the cycles leads to the word $\check{\sigma} = \check{\sigma}(1)\check{\sigma}(2) \cdots \check{\sigma}(n)$. As $\text{des } \sigma \leq r$,. There corresponds a unique word w from $[0, r]^*$ to σ under the the inverse of the bijection displayed in (27.3), given by $w = x_1x_2 \cdots x_n = \bar{c}_{\check{\sigma}(1)}\bar{c}_{\check{\sigma}(2)} \cdots \bar{c}_{\check{\sigma}(n)}$.

27.4. *A further property of the bijection $w \mapsto (c, \sigma)$.* As defined in Chapter 26, the integer i ($1 \leq i \leq n - 1$) is said to be a *decrease* (resp. *increase*) of the word $w = x_1x_2 \cdots x_n$, if $x_i = x_{i+1} = \cdots = x_j > x_{j+1}$ for some j such that $i \leq j \leq n - 1$ (resp. if $x_i = x_{i+1} = \cdots = x_j < x_{j+1}$ for some j such that $i \leq j \leq n$). By convention, $x_{n+1} = +\infty$. The letter x_i is then said to be a *decrease value* (resp. *increase value*). Note that the rightmost letter x_n is always an increase value. The *number* of decrease (resp. of increase) values in w is denoted by $\text{dec } w$ (resp. by $\text{inc } w$), while the *set* of all its decrease (resp. increase) values by $\text{REC } w$ (resp. $\text{INC } w$). Also, x_i ($1 \leq i \leq n$) is said to be a *record value* of w if $x_j \leq x_i$ for all j such that $1 \leq j \leq i - 1$. Finally, the number of letters of w , which are both increase *and* record values is denoted by $\text{inrec } w$.

Beside the number of descents “ $\text{des } \sigma$ ” of each permutation σ from \mathfrak{S}_n already introduced, mention the number or *rises* “ $\text{rise } \sigma$,” simply equal to $n - 1 - \text{des } \sigma$ ($n \geq 2$), and recall that the *number of excedances*, “ $\text{exc } \sigma$,” is defined to be the number of i ’s such that $1 \leq i \leq n - 1$ and $i < \sigma(i)$. Finally, the number of *fixed points* of σ (number of i ’s such that $\sigma(i) = i$) is denoted by $\text{fix } \sigma$.

Property 27.5. *Along the chain of bijections $w \mapsto (\check{\sigma}, \mathbf{m}) \mapsto (\sigma, c)$ hold the following properties:*

- (i) $\text{DEC } w = \{i : \check{\sigma}(i) < \sigma(\check{\sigma}(i))\}$; (ii) $\text{dec } w = \text{rise } \check{\sigma} = \text{exc } \sigma$;
- (iii) $\text{INC } w = \{i : \check{\sigma}(i) \geq \sigma(\check{\sigma}(i))\}$; (iv) $\text{inrec } w = \text{fix } \sigma$.

Proof. The letter x_i is a decrease value of $w = x_1x_2 \cdots x_n$, if and only if $x_i = x_{i+1} = \cdots = x_j > x_{j+1}$ for some j such that $i \leq j \leq n - 1$. This is equivalent to saying that $x_i \text{succ } x_{i+1} \text{succ } \cdots \text{succ } x_j > x_{j+1}$ and then $\check{\sigma}(i) < \check{\sigma}(i + 1) < \cdots < \check{\sigma}(x_j) < \check{\sigma}(j + 1)$ by definition of $\check{\sigma}$. Next, $\check{\sigma}(i) < \check{\sigma}(i + 1)$ if and only if $\check{\sigma}(i) < \sigma(\check{\sigma}(i))$, as necessarily both letters $\check{\sigma}(i)$ and $\check{\sigma}(i + 1)$ belong to the same factor, say, v_i , which determines the cycle (v_i) of σ (see Fig. 27.1). Consequently, both (i) and (ii) hold. Note that the second identity in (ii) reproduces a classical property of the First Fundamental Transformation (*op. cit.*). Formula (iii) is a simple consequence of (i).

Next, x_i is an increase and record value of w if and only if $x_k = x_{k+1} = \dots = x_{i-1} = x_i < x_{i+1}$ for $1 \leq k \leq i$ and x_k greater than all x_j 's such that $1 \leq j \leq k-1$. This means that the Lyndon factorization of w reads: $w = \dots | x_k | x_{k+1} | \dots | x_{i-1} | x_i | x_{i+1} \dots$ and $\check{\sigma}$ is the product of factors $\check{\sigma} = \dots | \check{\sigma}(k) | \check{\sigma}(k+1) | \dots | \check{\sigma}(i-1) | \check{\sigma}(i) | \check{\sigma}(i+1) | \dots$. In particular, $\check{\sigma}(i)$ is a fixed point of σ : $\sigma(\check{\sigma}(i)) = \check{\sigma}(i)$. \square

27.5. Generating functions for permutations. As an application of the Decrease Value Theorem, developed in Exercices 40 and 41, the generating function for the set of *words* $[0, r]^*$ by a certain weight ψ_R has been calculated and led to a closed expression. The next step is then to see whether there is also a parallel identity in the *permutation* environment.

For each word $w = x_1 x_2 \dots x_n$ from $[0, r]^*$ the weight ψ_R has been introduced as:

$$(27.8) \quad \psi_R(w) = R^{\text{inrec } w} s^{\text{dec } w} \prod_{i \in \text{DEC}} Y_{x_{i-1}} \prod_{i \in \text{INC}} Y_{x_i}$$

and the following identity derived

$$(27.9) \quad \sum_{w \in [0, r]^*} \psi_R(w) = \frac{(1-s)H_{r+1}(RY)}{H_r(sY) - sH_r(Y)},$$

where

$$H_r(Y) := \prod_{0 \leq i \leq r-1} (1 - Y_i)^{-1} \quad (r \geq 0).$$

Let (σ, c) be the pair associated with the word w under the Gessel-Reutenauer standardization. As $w = x_1 x_2 \dots x_n = \bar{c}_{\check{\sigma}(1)} \bar{c}_{\check{\sigma}(2)} \dots \bar{c}_{\check{\sigma}(n)}$, we may write using Property 27.8:

$$\begin{aligned} \psi_R(w) &= R^{\text{fix } \sigma} s^{\text{exc } \sigma} \prod_{\check{\sigma}(i) < \sigma(\check{\sigma}(i))} Y_{\bar{c}_{\check{\sigma}(i)}-1} \prod_{\check{\sigma}(i) \geq \sigma(\check{\sigma}(i))} Y_{\bar{c}_{\check{\sigma}(i)}} \\ &= R^{\text{fix } \sigma} s^{\text{exc } \sigma} \prod_{j < \sigma(j)} Y_{\bar{c}_j-1} \prod_{j \geq \sigma(j)} Y_{\bar{c}_j}. \end{aligned}$$

Therefore, in view of (27.5)

$$(27.10) \quad \psi_R(w) = R^{\text{fix } \sigma} s^{\text{exc } \sigma} \prod_{j < \sigma(j)} Y_{c_j+z_j-1} \prod_{j \geq \sigma(j)} Y_{c_j+z_j}.$$

The last two monomials depend only on the pair (σ, c) . Let NIW_n denote the collection of all non increasing words of length n , whose letters are integers. With each pair $(\sigma, c) \in \mathfrak{S}_n \times \text{NIW}_n$ associate the monomial

$$Y_{(\sigma, c)} := \prod_{j < \sigma(j)} Y_{c_j+z_j-1} \times \prod_{j \geq \sigma(j)} Y_{c_j+z_j}.$$

By summing (27.7) over all words from $[0, r]^*$ we get

$$(27.11) \quad \sum_{w \in [0, r]^*} \psi_R(w) = \sum_{n \geq 0} \sum_{\substack{\sigma \in \mathfrak{S}_n \\ \text{des } \sigma \leq r}} R^{\text{fix } \sigma} s^{\text{exc } \sigma} \sum_{c \in \text{NIW}_n(r - \text{des } \sigma)} Y_{(\sigma, c)}.$$

The following theorem is a banal consequence of (27.9) and (27.11).

Theorem 27.6. *We have:*

$$(27.12) \quad \sum_{n \geq 0} \sum_{\substack{\sigma \in \mathfrak{S}_n \\ \text{des } \sigma \leq r}} R^{\text{fix } \sigma} s^{\text{exc } \sigma} \sum_{c \in \text{NIW}_n(r - \text{des } \sigma)} Y_{(\sigma, c)} = \frac{(1-s)H_{r+1}(RY)}{H_r(sY) - sH_r(Y)}.$$

There is a graded form of the latter identity, as stated next, which also takes the number of descents “des” into account. Let

$$(27.13) \quad Y(\sigma; t) := \sum_{k \geq 0} t^k \sum_{c \in \text{NIW}_n(k)} Y_{(\sigma, c)}.$$

Theorem 27.7. *The graded form of (27.12) reads:*

$$(27.14) \quad \sum_{n \geq 0} \sum_{\sigma \in \mathfrak{S}_n} R^{\text{fix } \sigma} s^{\text{exc } \sigma} t^{\text{des } \sigma} Y(\sigma; t) = \sum_{r \geq 0} t^r \frac{(1-s)H_{r+1}(RY)}{H_r(sY) - sH_r(Y)}.$$

Proof. Multiply both sides of (27.12) by t^r and sum over $r \geq 0$. We obtain:

$$\begin{aligned} & \sum_{r \geq 0} t^r \frac{(1-s)H_{r+1}(RY)}{H_r(sY) - sH_r(Y)} \\ &= \sum_{r \geq 0} t^r \sum_{n \geq 0} \sum_{\substack{\sigma \in \mathfrak{S}_n \\ \text{des } \sigma \leq r}} R^{\text{fix } \sigma} s^{\text{exc } \sigma} \sum_{c \in \text{NIW}_n(r - \text{des } \sigma)} Y_{(\sigma, c)} \\ &= \sum_{n \geq 0} \sum_{r \geq 0} t^r \sum_{0 \leq j \leq r} \sum_{\substack{\sigma \in \mathfrak{S}_n \\ \text{des } \sigma = r - j}} R^{\text{fix } \sigma} s^{\text{exc } \sigma} \sum_{c \in \text{NIW}_n(j)} Y_{(\sigma, c)} \\ &= \sum_{n \geq 0} \sum_{j \geq 0} t^j \sum_{r \geq j} t^{r-j} \sum_{\substack{\sigma \in \mathfrak{S}_n \\ \text{des } \sigma = r - j}} R^{\text{fix } \sigma} s^{\text{exc } \sigma} \sum_{c \in \text{NIW}_n(j)} Y_{(\sigma, c)} \\ &= \sum_{n \geq 0} \sum_{j \geq 0} t^j \sum_{k \geq 0} t^k \sum_{\substack{\sigma \in \mathfrak{S}_n \\ \text{des } \sigma = k}} R^{\text{fix } \sigma} s^{\text{exc } \sigma} \sum_{c \in \text{NIW}_n(j)} Y_{(\sigma, c)} \\ &= \sum_{n \geq 0} \sum_{\sigma \in \mathfrak{S}_n} R^{\text{fix } \sigma} s^{\text{exc } \sigma} t^{\text{des } \sigma} \sum_{j \geq 0} t^j \sum_{c \in \text{NIW}_n(j)} Y_{(\sigma, c)} \\ &= \sum_{n \geq 0} \sum_{\sigma \in \mathfrak{S}_n} R^{\text{fix } \sigma} s^{\text{exc } \sigma} t^{\text{des } \sigma} Y(\sigma; t). \quad \square \end{aligned}$$

27. THE DECREASE VALUE THEOREM; FROM WORDS TO PERMUTATIONS

27.6. *A quadruple q -series for permutations.* Recall the traditional notation of the q -ascending factorial

$$(a; q)_n = \begin{cases} 1, & \text{if } n = 0; \\ (1 - a)(1 - aq) \cdots (1 - aq^{n-1}), & \text{if } n \geq 1. \end{cases}$$

The next theorem will appear as is a simple consequence of Theorem 27.7 by applying the homomorphism ϕ generated by $\phi(Y_j) := uq^j$ ($j \geq 0$) and $\phi(s) := sq$ to both sides of (27.11). Recall that the statistic “maj”, the *major index*, is defined as the sum of all descent values of a permutation $\sigma \in \mathfrak{S}_n$, that is, the sum of all integers i such that $1 \leq i \leq n - 1$ and $\sigma(i) > \sigma(i + 1)$. It is also the sum “tot z ” of all the z_i ’s introduced in (27.3). Note that in the next proof “tot c ” is a shorthand for $c_1 + c_2 + \cdots + c_n$.

Theorem 27.8. *The factorial generating function for the distribution of the vector (fix, exc, des, maj) over the symmetric groups \mathfrak{S}_n is given by*

$$(27.15) \quad \sum_{n \geq 0} \frac{u^n}{(t; q)_{n+1}} \sum_{\sigma \in \mathfrak{S}_n} R^{\text{fix } \sigma} s^{\text{exc } \sigma} t^{\text{des } \sigma} q^{\text{maj } \sigma} \\ = \sum_{r \geq 0} t^r \frac{1}{(uR; q)_{r+1}} \frac{(1 - sq)(u; q)_r (usq; q)_r}{((u; q)_r - sq(usq; q)_r)}.$$

Proof. First, $\phi H_r(Y) = \prod_{0 \leq j \leq r-1} (1 - uq^j)^{-1} = 1/(u; q)_r$ and

$$\phi \frac{(1 - s)H_{r+1}(RY)}{H_r(sY) - sH_r(Y)} = \frac{(1 - sq)/(uR; q)_{r+1}}{1/(usq; q)_r - sq/(u; q)_r} \\ = \frac{(1 - sq)(u; q)_r (usq; q)_r}{(uR; q)_{r+1} ((u; q)_r - sq(usq; q)_r)}.$$

Then, for $(\sigma, c) \in \mathfrak{S}_n \times \text{NIW}_n$

$$\phi Y_{(\sigma, c)} = u^n q^{\text{tot } c + \text{tot } z - \text{exc } \sigma} = q^{\text{maj } \sigma - \text{exc } \sigma} u^n q^{\text{tot } c};$$

$$\phi Y(\sigma; t) = q^{\text{maj } \sigma - \text{exc } \sigma} u^n \sum_{j \geq 0} t^j \sum_{c \in \text{NIW}_n(j)} q^{\text{tot } c} = q^{\text{maj } \sigma - \text{exc } \sigma} \frac{u^n}{(t; q)_{n+1}};$$

so that

$$\phi(R^{\text{fix } \sigma} s^{\text{exc } \sigma} t^{\text{des } \sigma} Y(\sigma; t)) = R^{\text{fix } \sigma} (sq)^{\text{exc } \sigma} t^{\text{des } \sigma} q^{\text{maj } \sigma - \text{exc } \sigma} \frac{u^n}{(t; q)_{n+1}}.$$

Hence, the image of identity (27.14) under ϕ gives back (27.15). \square

Multiplying Identity (27.15) by $(1 - t)$ and letting $t = 1$ yields to

$$\sum_{n \geq 0} \frac{u^n}{(q; q)_n} \sum_{\sigma \in \mathfrak{S}_n} R^{\text{fix } \sigma} s^{\text{exc } \sigma} q^{\text{maj } \sigma} = \frac{1}{(uR; q)_\infty} \frac{(1 - sq)(u; q)_\infty (usq; q)_\infty}{((u; q)_\infty - sq(usq; q)_\infty)},$$

which is an identity derived by Shareshian and Wachs [SW07] by means of a symmetric function argument. For $s = 1$ Identity (27.15) reads

$$\sum_{n \geq 0} \frac{u^n}{(t; q)_{n+1}} \sum_{\sigma \in \mathfrak{S}_n} R^{\text{fix } \sigma} t^{\text{des } \sigma} q^{\text{maj } \sigma} = \sum_{r \geq 0} t^r \left(1 - u \sum_{i=0}^r q^i\right)^{-1} \frac{(u; q)_{r+1}}{(uR; q)_{r+1}},$$

an identity derived by Gessel and Reutenauer [GeRe93].

Exercises and examples

1. *Rogers-Szegö Polynomials.* The product $\exp(xu) \exp(u)$ is the exponential generating function for polynomials $H_n(x)$, whose expression is easy to derive.

In the same manner, the product of the two q -exponentials $e_q(xu) e_q(u)$ is the factorial generating function $\sum H_n(x, q) u^n / (q; q)_n$ for polynomials $H_n(x, q)$, called the Rogers-Szegö polynomials. They can be expressed by means of the q -binomial coefficients.

We have $H_{2n+1}(-1, q) = 0$, $H_{2n}(-1; q) = (q; q^2)_n$, then $H_n(q^{1/2}, q) = (-q^{1/2}; q^{1/2})_n$ and finally the induction formula

$$H_{n+1}(x, q) = (1+x)H_n(x, q) - (1-q^n)x H_{n-1}(x, q).$$

2. *The Ramanujan sum.* The sum in question is:

$$\sum_{n=-\infty}^{+\infty} \frac{(a; q)_n}{(b; q)_n} u^n = \frac{(au; q)_\infty (qa^{-1}u^{-1}; q)_\infty (q; q)_\infty (ba^{-1}; q)_\infty}{(u; q)_\infty (ba^{-1}u^{-1}; q)_\infty (b; q)_\infty (qa^{-1}; q)_\infty}.$$

On the left-hand side each term is a series in the variable q and is defined for each integer, positive and negative. In the proof of the identity there will be no ambiguity in the definition of the sum.

Start with the sum

$$h(b) = \sum_{n=-\infty}^{+\infty} \frac{(bq^n; q)_\infty}{(aq^n; q)_\infty} u^n = \frac{(b; q)_\infty}{(a; q)_\infty} \sum_{n=-\infty}^{+\infty} \frac{(a; q)_n}{(b; q)_n} u^n$$

and use the trivial relation $(bq^n; q)_\infty = (1 - bq^n)(bq^{n+1}; q)_\infty$. We obtain the q -recurrence

$$h(b) = (1 - ba^{-1}) h(bq) + ba^{-1}u^{-1} h(b),$$

and by itération

$$h(b) = h(bq^n) \frac{(ba^{-1}; q)_n}{(ba^{-1}u^{-1}; q)_n}$$

for every $n \geq 0$. As soon as $n \geq j + 1$ the latter identity implies that the coefficients of $u^i q^j$

$$\text{in } h(bq^n) \frac{(ba^{-1}; q)_n}{(ba^{-1}u^{-1}; q)_n} \quad \text{and in } h(0) \frac{(ba^{-1}; q)_\infty}{(ba^{-1}u^{-1}; q)_\infty}$$

are identical. Hence, $h(b) = h(0)(ba^{-1}; q)_\infty / (ba^{-1}u^{-1}; q)_\infty$ and also $h(q) = h(0)(qa^{-1}; q)_\infty / (qa^{-1}u^{-1}; q)_\infty$. Going back to the original definition of $h(b)$ and using the q -binomial theorem we obtain

$$h(q) = \frac{(q; q)_\infty (au; q)_\infty}{(a; q)_\infty (u; q)_\infty}.$$

By combining the last three formulas in an evident manner we obtain the Ramanujan sum.

3. *A maj-inv bijection for permutations.* Let $x = (x_1, x_2, \dots, x_n)$ be the inv-coding of a permutation $\sigma \in \mathfrak{S}_n$. With $x_{n+1} := \sigma(n+1) = 0$ define a sequence $y = (y_1, y_2, \dots, y_n)$ by $y_i := x_i - x_{i+1} + i \chi(\sigma(i) > \sigma(i+1))$ for each $i = 1, 2, \dots, n$. Then the sequence y is subexcedent, the mapping $x \mapsto y$ is bijective and $\text{tot } y = y_1 + y_2 + \dots + y_n$ is equal to $\text{maj } \sigma$. Let σ' be the permutation whose inv-coding is y . Then $\sigma \mapsto \sigma'$ is a bijection of \mathfrak{S}_n onto itself having the property: $\text{maj } \sigma = \text{inv } \sigma'$.

4. *A numerical example.* Calculate the inv-coding and the maj-coding of $\sigma = 259478361$ and the number of inversions and the Major Index of σ .

Find the permutations having for inv-coding and maj-coding the subexcedent sequence $x = 001304645$, respectively.

5. *Another maj-coding.* Let $\sigma = \sigma(1)\sigma(2)\dots\sigma(n)$ be a permutation. With $\sigma(n+1) := +\infty$ define a sequence $z = z_1 z_2 \dots z_n$ by

$$z_i := \#\{1 \leq j < i \mid \sigma(j) \in \llbracket \sigma(i), \sigma(i+1) \rrbracket\},$$

using the notation $\llbracket \dots \rrbracket$ for a cyclic interval (see section 2).

- (a) The sequence z is subexcedent.
- (b) Determine the sequence z that corresponds to $\sigma = 259478361$.
- (c) The transformation $\sigma \mapsto z$ is invertible. Find the inverse of $z = 001304645$.
- (d) Let $x = x_1 x_2 \dots x_n$ be the inv-coding of σ and $x_{n+1} = 0$. The following identity holds:

$$z_i = x_i - x_{i+1} + i \chi(\sigma_i > \sigma_{i+1}).$$

- (e) We have: $z_1 + z_2 + \dots + z_n = \text{maj } \sigma$. Thus, the transformation $\sigma \mapsto y$ defines another maj-coding.

6. *A counter-example.* For each permutation $\sigma = \sigma(1)\sigma(2)\cdots\sigma(n)$ define a sequence $y = y_1y_2\cdots y_n$ by $y_i := (i-1)\chi(\sigma(i-1) > \sigma(i))$. The transformation $\sigma \mapsto y$ is not a maj-coding although $\text{tot } y = \text{maj } \sigma$.

7. *Number of vector subspaces.* Let q be the power of a prime number and let \mathbb{F}_q denote the field having q elements. The number of vector subspaces of dimension n of the vector space \mathbb{F}_q^{N+n} is equal to $\begin{bmatrix} N+n \\ n \end{bmatrix}$.

First, the number of sets $\{v_1, v_2, \dots, v_n\}$ of n vectors v_1, v_2, \dots, v_n of \mathbb{F}_q^{N+n} , that are linearly independent, is equal to

$$q^{(N+n)(N+n-1)/2}(q^{N+n} - 1)(q^{N+n-1} - 1)\cdots(q^{N+1} - 1);$$

then, the number of such sets that generate a given vector subspace of dimension n in \mathbb{F}_q^{N+n} is equal to $q^{n(n-1)/2}(q^n - 1)(q^{n-1} - 1)\cdots(q - 1)$.

8. *The use of the q -Pascal Triangle.* Derive the expansions of $1/(u; q)_N$ and of $(-u; q)_n$ (formulas (3.9) and (3.10)) by means of the q -Pascal Triangle formulas (3.5) and (3.6).

9. *Nondecreasing sequences of integers.* Calculate the generating function for nondecreasing sequences of N integers at most equal to n by “tot” (formula (4.5)) by using the q -Pascal Triangle formulas (3.5) and (3.6)

10. *Binary words.* Calculate the generating function for binary words of length $(N+n)$ having N letters equal to 1 and n letters equal to 0 by their number of inversions (formula (4.17)) by using the q -Pascal Triangle formulas (3.5) and (3.6)

11. *Partitions of integers and q -binomial coefficients.* The interpretation of the q -binomial coefficients in terms of partitions of integers (see § 4.1) makes it easy to prove the following identities.

(a) By classifying the partitions in at most n parts, all being at most equal to N , according the size of the smallest n -th part (possibly 0) we get:

$$\sum_{j=0}^N \begin{bmatrix} N-j+n-1 \\ N-j \end{bmatrix} q^{jn} = \begin{bmatrix} N+n \\ N \end{bmatrix}.$$

(b) Also

$$\begin{aligned} \sum_{j=0}^N \begin{bmatrix} n+j \\ j \end{bmatrix} \begin{bmatrix} m-1+N-j \\ N-j \end{bmatrix} q^{jm} &= \begin{bmatrix} m+n+N \\ N \end{bmatrix} \\ &= (-1)^N q^{N(n+m)+N(N+1)/2} \begin{bmatrix} -m-n-1 \\ N \end{bmatrix}. \end{aligned}$$

EXERCISES AND EXAMPLES

12. *The Eulerian numbers.* They are denoted by $A_{n,k}$ and are defined by the recurrence relation (see (10.8))

$$(E12.1) \quad A_{n,k} = (k+1)A_{n-1,k} + (n-k)A_{n-1,k-1} \quad (1 \leq k \leq n-1);$$

$$A_{n,0} = 1 \quad (n \geq 0); \quad A_{n,k} = 0 \quad (k \geq n).$$

Their first values are shown in Table E12.1.

k=	0	1	2	3	4	5	6
n=1	1						
2	1	1					
3	1	4	1				
4	1	11	11	1			
5	1	26	66	26	1		
6	1	57	302	302	57	1	
7	1	120	1191	2416	1191	120	1

Table E12.1

(a) For each $n \geq 0$ we have $\sum_{k \geq 0} A_{n,k} = n!$ and $A_{n,k} = A_{n,n-1-k}$.

(b) The number of descents, $\text{des } \sigma$, of a permutation $\sigma = \sigma(1) \dots \sigma(n)$ is defined to be the number of integers j such that $1 \leq j \leq n-1$ and $\sigma(j) > \sigma(j+1)$. For every $k \geq 0$ and every $n \geq 0$ the number $A_{n,k}$ is equal to the number of permutations $\sigma \in \mathfrak{S}_n$ having k descents.

(c) The *Eulerian polynomial* $A_n(t)$ is defined by $A_n(t) := \sum_{0 \leq k \leq n-1} A_{n,k} t^k$.

As said in Definition 10.2, relations (10.7) and (10.8) are equivalent.

13. *The Eulerian Polynomials.* Consider the sequence $(A_n(t))$ ($n \geq 0$) of formal series in the variable t (they will appear to be *polynomials* and more precisely the Eulerian Polynomials defined in Exercice 12) defined by

$$(E13.1) \quad \frac{A_n(t)}{(1-t)^{n+1}} := \sum_{j \geq 0} t^j (j+1)^n.$$

(a) If D designates the derivative-operator for the formal series, we have: $A_n(t) = (1 + (n-1)t)A_{n-1}(t) + t(1-t).DA_{n-1}(t)$ ($n \geq 1$).

(b) For $n \geq 0$ let $A_n(t) := \sum_{k \geq 0} A_{n,k} t^k$. Then the relations listed in (E12.1) hold.

(c) It follows from the definition $A_n(t) = \sum_{k \geq 0} A_{n,k} t^k$ and (E13.1) that

$$A_{n,k} = \sum_{0 \leq i \leq k} (-1)^i (k - i + 1)^n \binom{n+1}{i}.$$

(d) The “inverse” of the previous formula is the *Worpitzky* formula:

$$x^n = \sum_{0 \leq k \leq n-1} \binom{x+k}{n} A_{n,k}.$$

[Start with the expansion of $A_n(t)(1-t)^{-(n+1)}$, then use Exercise 12 (a) and (d) to recover (E12.1).]

(e) The exponential generating function

$$\sum_{n \geq 0} A_n(t) \frac{u^n}{n!} = \frac{1-t}{-t + \exp(u(t-1))}$$

is a consequence of (E13.1). Thus, the five definitions (10.5)—(10.9) are shown to be equivalent.

14. *A less unwieldy definition for the Denert statistic.* The Denert statistic “den” has been defined in section 2 by means of the cyclic intervals. There is an alternative definition that is the following. Let $\sigma = \sigma(1)\sigma(2)\dots\sigma(n)$ be a permutation of order n . If $1 \leq i \leq n-1$ and $\sigma(i) > i$, say that i is an *excedence-place* for σ , and $\sigma(i)$ is an *excedence-letter* for σ . Let $i_1 < i_2 < \dots < i_k$ be the increasing sequence of the excedence places and $j_1 < j_2 < \dots < j_{n-k}$ the increasing sequence of *non-excedence places*. The subwords $\text{Exc } \sigma = \sigma(i_1)\dots\sigma(i_k)$ and $\text{Nexc } \sigma = \sigma(j_1)\dots\sigma(j_{n-k})$ are referred to as the *excedence-letter* and *non-excedence-letter* subwords.

Let $\sigma = \sigma(1)\sigma(2)\dots\sigma(n)$ be a permutation. Let i_1, \dots, i_k be its excedence-place sequence. Then the Denert statistic of σ is also given by

$$\text{den } \sigma = i_1 + i_2 + \dots + i_k + \text{inv Exc } \sigma + \text{inv Nexc } \sigma.$$

For instance, for the permutation $\sigma = \begin{pmatrix} 123456789 \\ 715492638 \end{pmatrix}$ shown in § 2.3 we have $i_1 = 1, i_2 = 3, i_3 = 5, \text{Exc } \sigma = 7, 5, 9, \text{Nexc } \sigma = 1, 4, 2, 6, 3, 8, \text{inv Exc } \sigma = 1, \text{inv Nexc } \sigma = 3$, and then $\text{den } \sigma = 1 + 3 + 5 + 1 + 3 = 13$.

15. *Euler-Mahonian Statistics.* In Definition 10.1 the q -maj-Eulerian $\text{maj}A(t, q) = \sum_{k \geq 0} t^k \text{maj}A_{n,k}(q)$ have been introduced in four different ways.

EXERCISES AND EXAMPLES

To simplify the notations the superscript “maj” will be deleted in this Exercise. For an easy reference rewrite the recurrence for the coefficients $A_{n,k}(q) = \text{maj}A_{n,k}(q)$:

$$(E15.1) \quad A_{n,k}(q) = [k+1]_q A_{n-1,k}(q) + q^k [n-k]_q A_{n-1,k-1}(q),$$

for $1 \leq k \leq n-1$ with the initial conditions $A_{n,0}(q) = 1$ for $n \geq 0$ and $A_{n,k}(q) = 0$ for $k \geq n$.

Let $E = (E_n)$ ($n \geq 0$) be a family of finite sets such that $\text{card } E_n = n!$ for all $n \geq 0$. A family $(f, g) = (f_n, g_n)$ ($n \geq 0$) is said to be *Euler-Mahonian* on E , if $f_0 = g_0 = 0$, $f_1 = g_1 = 0$ and if for every $n \geq 2$ both f_n and g_n are integral-valued mappings defined on E_n and if there exists a bijection $\psi_n : (w', j) \mapsto w$ of $E_{n-1} \times [0, n-1]$ onto E_n having the following properties:

$$(E15.2) \quad \begin{aligned} g_n(w) &= g_{n-1}(w') + j; \\ f_n(w) &= \begin{cases} f_{n-1}(w'), & \text{if } 0 \leq j \leq f_{n-1}(w'); \\ f_{n-1}(w') + 1, & \text{if } f_{n-1}(w') + 1 \leq j \leq n-1. \end{cases} \end{aligned}$$

Every such a pair (f_n, g_n) is called a *Euler-Mahonian* statistic on E_n .

(a) Let (f, g) be a Euler-Mahonian family on E and for each triple (n, k, l) let $A_{n,k,l}$ be the number of elements $w \in E_n$ such that $f_n(w) = k$ and $g_n(w) = l$ and let $A_{n,k}(q) := \sum_l A_{n,k,l} q^l$. Then $(A_{n,k}(q))$ satisfies recurrence (E15.1).

(b) On the set SE_n of the subexcedent sequences of length n (see Proposition 2.1) we know that “tot” is Mahonian. Further, define the *Eulerian value* “eul x ” of a sequence $x = (x_1, \dots, x_n) \in \text{SE}_n$ by $\text{eul } x = 0$ if x is of length 1 and for $n \geq 2$

$$\text{eul } x := \begin{cases} \text{eul}(x_1, \dots, x_{n-1}), & \text{if } x_n \leq \text{eul}(x_1, \dots, x_{n-1}); \\ \text{eul}(x_1, \dots, x_{n-1}) + 1, & \text{if } x_n \geq \text{eul}(x_1, \dots, x_{n-1}) + 1. \end{cases}$$

Then the pair (eul, tot) is a Euler-Mahonian statistic on the family (SE_n) ($n \geq 0$).

(c) The pair (des, maj) is a Euler-Mahonian statistic on the family (\mathfrak{S}_n) ($n \geq 0$).

(d) Let $n \geq 2$ and $\sigma' = x_1 \dots x_{n-1}$ be a permutation having k excedences, that is, there are k integers i such that $1 \leq i \leq n-1$ and $i < x_i$. In short, $\text{exc } \sigma' = k$. Let $(x_{i_1} > \dots > x_{i_k})$ be the *decreasing* sequence of the excedence values $x_i > i$ and let $(x_{i_{k+1}} < \dots < x_{i_{n-1}})$ be the *increasing* sequence of the non-excedence values $x_i \leq i$. By convention, $x_{i_0} := n$.

Define $\psi_n(\sigma', 0) := x_1 x_2 \dots x_{n-1} n$. If $1 \leq j \leq n-1$, let $y_j := \min\{x_{i_m} : x_{i_m} \geq x_{i_j}\}$. Replace each letter x_{i_m} ($1 \leq m \leq j$) in σ' such that $x_{i_m} \geq x_{i_j}$ by $x_{i_{m-1}}$, leave the other letters alike and insert y_j into the x_{i_j} -th position in σ' . Let $\sigma = \psi_n(\sigma', j)$ denote the permutation derived by that procedure.

For example, $\sigma' = 32541$ has $k = 2$ excedences $x_3 = 5 > 3$, $x_1 = 3 > 1$ (in decreasing order) and three non-excedences $x_5 = 1 \leq 5$, $x_2 = 2 \leq 2$, $x_4 = 4 \leq 4$ (in increasing order), so that $(i_1, i_2, i_3, i_4, i_5) = (3, 1, 5, 2, 4)$. With $j = 1$ we have $i_j = 3$ and $x_3 = 5$. For getting $\psi_6(\sigma', 1)$ replace $x_{i_1} = 5$ by $x_{i_0} = 6$, leave the other letters alike and insert $x_{i_1} = 5$ into the x_{i_1} -th = 5-th position. Thus, $\psi_6(\sigma', 1) = 326451$.

For $j = 3$ we have $i_j = 5$ and $x_5 = 1$. As $j = 3 > k = 2$, replace $x_{i_1} = x_3$ by $x_{i_0} = 6$, then $x_{i_2} = x_1 = 3$ by $x_{i_1} = 5$, leave the other letters alike and insert $x_{i_k} = x_{i_2} = x_1 = 3$ into the x_{i_3} -th = 1-st position to obtain $\psi_6(\sigma', 3) = 352641$.

With $(f, g) = (\text{exc}, \text{den})$ (see section 2) properties (E15.2) hold for ψ_n , so that (exc, den) is a Euler-Mahonian statistic on the family (\mathfrak{S}_n) ($n \geq 0$).

Let us illustrate the latter property with the running example. The statistic “den” is calculated by using the definition of Exercice 14. We have:

$\sigma' = \begin{pmatrix} \mathbf{1} \mathbf{2} \mathbf{3} \mathbf{4} \mathbf{5} \\ \mathbf{3} \mathbf{2} \mathbf{5} \mathbf{4} \mathbf{1} \end{pmatrix}$, so that $\text{exc } \sigma' = 2$ and $\text{den } \sigma' = (\mathbf{1} + \mathbf{3}) + 0 + 2 = 6$. Next $\psi_6(\sigma', 1) = \begin{pmatrix} \mathbf{1} \mathbf{2} \mathbf{3} \mathbf{4} \mathbf{5} \mathbf{6} \\ \mathbf{3} \mathbf{2} \mathbf{6} \mathbf{4} \mathbf{5} \mathbf{1} \end{pmatrix}$, so that $\text{exc } \psi_6(\sigma', 1) = 2$ and $\text{den } \psi_6(\sigma', 1) = (\mathbf{1} + \mathbf{3}) + 0 + 3 = 7 = \text{den } \sigma' + 1$. Finally, $\psi_6(\sigma', 3) = \begin{pmatrix} \mathbf{1} \mathbf{2} \mathbf{3} \mathbf{4} \mathbf{5} \mathbf{6} \\ \mathbf{3} \mathbf{5} \mathbf{2} \mathbf{6} \mathbf{4} \mathbf{1} \end{pmatrix}$, so that $\text{exc } \psi_6(\sigma', 3) = 3$ and $\text{den } \psi_6(\sigma', 3) = (\mathbf{1} + \mathbf{2} + \mathbf{4}) + 0 + 2 = 9 = \text{den } \sigma' + 3$.

(e) Let (f, g) be a Euler-Mahonian family on $E = (E_n)$. For each $w \in E_n$ ($n \geq 2$) let $\psi_n^{-1}(w) := (w', j_n)$, $\psi_{n-1}^{-1}(w') := (w'', j_{n-1})$, \dots , $\psi_2^{-1}(w^{(n-2)}) := (w^{(n-1)}, j_2)$ and $j_1 := 0$; the sequence $\Psi(w) := (j_1, j_2, \dots, j_{n-1}, j_n)$ is subexcedent and Ψ is a bijection if E_n onto SE_n such that $f(w) = \text{tot } \Psi(w)$ and $g(w) = \text{eul } \Psi(w)$. The bijection Ψ is said to be an (f, g) -coding E_n .

Let $\Psi_{(\text{des}, \text{maj})}$ (resp. $\Psi_{(\text{exc}, \text{den})}$) be the (des, maj) -coding (see question (c)) (resp. the (exc, den) -coding (see question (d)) of \mathfrak{S}_n . Then $\Theta := \Psi_{(\text{des}, \text{maj})}^{-1} \circ \Psi_{(\text{exc}, \text{den})}$ is a bijection of \mathfrak{S}_n onto itself having the property: $(\text{exc}, \text{den}) w = (\text{des}, \text{maj}) \Theta(w)$.

16. *Binary words.* Let $\text{BW}(N, n)$ be the set of binary words of length $(N + n)$ containing N times 1 and n times 0. If $x = x_1 x_2 \dots x_{N+n}$ is such a word, define

$$\text{rise } x := \sum_{1 \leq i \leq N+n-1} \chi(x_i < x_{i+1}) \quad \text{and} \quad \text{rmaj } x := \sum_{1 \leq i \leq N+n-1} i \chi(x_i < x_{i+1}).$$

EXERCISES AND EXAMPLES

Next, let $\text{DES } x := \text{des } x + \text{rise } x$ and $\text{MAJ } x := \text{maj } x + \text{rmaj } x$.

(a) For each of the words in $\text{BW}(2, 3)$ write the values of the six statistics “des”, “maj”, “rise”, “rmaj”, “DES”, “MAJ”.

$$(b) \sum_{x \in \text{BW}(N, n)} q^{\text{rmaj } x} = \begin{bmatrix} N+n \\ n \end{bmatrix}.$$

$$(c) \sum_{x \in \text{BW}(N, n)} t^{\text{rise } x} q^{\text{rmaj } x} = \sum_{x \in \text{BW}(N, n)} t^{\text{des } x} q^{\text{maj } x}.$$

(d) For $x \in \text{BW}(N, n)$ let $x' := x_1 x_2 \cdots x_{N+n-1}$. Then

$$\begin{cases} \text{maj } x = \text{rise } x + N = \text{rmaj } x' + N, & \text{if } x_{N+n} = 1; \\ \text{rmaj } x = \text{maj } x + n = \text{rmaj } x' + n, & \text{if } x_{N+n} = 0. \end{cases}$$

$$(e) \sum_{x \in \text{BW}(N, n)} q^{\text{MAJ } x} = \begin{bmatrix} N+n \\ n \end{bmatrix}_{q^2} \frac{q^N + q^n}{1 + q^{N+n}}.$$

17. *The Z-statistic is a Mahonian statistic.* Two transformations are described, the *global cycling* “gycy” for manipulating the Major Index and the *local cycling* “lcyc” for dealing with the Z-statistic itself.

(a) Let $\mathbf{m} = (m_1, m_2, \dots, m_r)$ be a *multiplicity*, that is, a sequence of positive integers and $R(\mathbf{m})$ be the class of all the rearrangements of the word $1^{m_1} 2^{m_2} \dots r^{m_r}$. Let \mathbf{n} be a rearrangement of \mathbf{m} . Construct a bijection $\theta_{\mathbf{m}, \mathbf{n}}$, defined on $R(\mathbf{m})$, with values in $R(\mathbf{n})$, preserving “maj.” [It suffices to give the construction when \mathbf{m} and \mathbf{n} differ by two *consecutive* letters x and $y = x + 1$; in other words, construct a bijection θ of $R(m_1, \dots, m_x, m_y, \dots, m_r)$ onto $R(m_1, \dots, m_y, m_x, \dots, m_r)$.]

(b) The Z-statistic is defined, for each word $w = x_1 x_2 \dots x_m$, by

$$Z(w) := \sum_{i < j} \text{maj } w_{ij},$$

where w_{ij} stands for the subword of w made of all the letters i and j . For example, for the word $w = 2412131242$, the subwords $w_{12} = 2121122$, $w_{13} = 1131, \dots$, are to be considered, the Major Indices are to be calculated and their sum to be added up. Calculate $Z(2412131242)$.

(c) For each word $w = x_1 x_2 \cdots x_m$ in $R(\mathbf{m})$ and each letter $x = 1, 2, \dots, r$ the *global cycling* $\text{gycy}_x(w) = y_1 y_2 \cdots y_m$ and the *local cycling* $\text{lcyc}_x(w) = z_1 z_2 \cdots z_m$ are defined by

$$y_i := \begin{cases} x_i - x, & \text{si } x_i > x; \\ x_i - x + r, & \text{sinon.} \end{cases} \quad z_i := \begin{cases} x_i, & \text{si } x_i < x; \\ x_i - 1, & \text{si } x_i > x; \\ r, & \text{si } x_i = x. \end{cases}$$

Let \mathbf{m}^x denote the multiplicity of $\text{gcyc}_x(w)$ and \mathbf{m}_x the multiplicity of $\text{lcyc}_x(w)$. Characterize those two multiplicities, that is to say, express \mathbf{m}^x and \mathbf{m}_x as rearrangements of the sequence $\mathbf{m} = (m_1, m_2, \dots, m_r)$.

(d) For the word $w = 2412131242$ calculate the differences $\text{maj } w - \text{maj gcyc}_2(w)$ et $Z(w) - Z(\text{lcyc}_2(w))$.

(e) If $x = x_m$ is the last letter of w , show that

$$\text{maj } w - \text{maj}(\text{gcyc}_x(w)) = m_{x+1} + m_{x+2} + \dots + m_r.$$

(f) If $x = x_m$ is the last letter of w , show that

$$Z(w) - Z(\text{lcyc}_x(w)) = m_{x+1} + m_{x+2} + \dots + m_r.$$

(g) Construct a bijection of $R(\mathbf{m})$ onto itself having the property

$$\text{maj } w = Z(\Phi(w)),$$

thus proving that the Z -statistic is *Mahonian* on each class of rearrangements.

18. *The $t = 1$ Lemma.* Let (b_r) ($r \geq 0$) be a sequence of formal series belonging to an algebra \mathfrak{A} of formal series in one or several variables. Let t be a new variable; we can form the series $b(t) := \sum_{r \geq 0} b_r t^r$, that belongs to the algebra $\mathfrak{A}[[t]]$. ‘Let $t = 1$ ’ in $b(t)$ does not always make sense. However, if the series $\sum_r b_r$ converges for the topology of the formal series in \mathfrak{A} , that is, if there exists $a \in \mathfrak{A}$ such that the order $o((b_0 + b_1 + \dots + b_r) - a)$ tends to $+\infty$ with r , we can *define* ‘let $t = 1$ in $b(t)$ ’ and then $b(1)$ by: $b(1) := a = \sum_{r \geq 0} b_r$.

(a) Let (a_r) ($r \geq 0$) be a sequence of formal series in \mathfrak{A} such that $\lim_r a_r = a$, that is, such that $o(a - a_r)$ tends to infinity with r . We define: $b(t) := (1 - t) \cdot \sum_{r \geq 0} a_r t^r$. Then $b(1) = a$.

(b) Deduce (6.13) from (7.8).

(c) Deduce (12.3) from (13.7).

19. *The dihedral group.* On the group \mathfrak{S}_n of the permutations on order n three transformations \mathbf{i} , \mathbf{r} and \mathbf{c} can be defined in the following way. First, \mathbf{i} is the bijection that maps each permutation σ onto its *inverse* σ^{-1} . Write σ as a linear word $\sigma = \sigma(1) \dots \sigma(n)$. Then, define

$$\begin{aligned} \mathbf{c} \sigma &:= (n + 1 - \sigma(1))(n + 1 - \sigma(2)) \dots (n + 1 - \sigma(n)); \\ \mathbf{r} \sigma &:= \sigma(n) \dots \sigma(2)\sigma(1). \end{aligned}$$

We say that \mathbf{c} is the *complement to $(n + 1)$* and \mathbf{r} the *reverse image*.

EXERCISES AND EXAMPLES

- (a) $\mathbf{r} = \mathbf{ic i}$.
- (b) The group acting on \mathfrak{S}_n generated by $\{\mathbf{i}, \mathbf{c}\}$ is isomorphic to the dihedral group D_4 of order 8 of the rotations of the square.
- (c) $\mathbf{rc} = \mathbf{cr}$, $\mathbf{ir} = \mathbf{ci}$ et $\mathbf{irc} = \mathbf{rci}$.
- (d) For each $\sigma \in \mathfrak{S}_n$ the following relations hold: $\text{Ligne } \mathbf{c}\sigma = [n - 1] \setminus \text{Ligne } \sigma$ and $\text{Ligne } \mathbf{r}\mathbf{c}\sigma = n - \text{Ligne } \sigma = \{n - i : i \in \text{Ligne } \sigma\}$.

20. *Action of the dihedral group.* By means of an example the action of the dihedral group on several set-valued statistics, such as “Ligne,” “Iligne,” the various extremum letter or place subsets (see § 11.4) is examined.

In Fig. E20 the dihedral group of order 8 is acting on the permutation $\sigma = 2, 6, 8, 1, 7, 4, 3, 5$, whose graph appears on the first square entitled “Id”. The ligne of route, $\text{Ligne } \sigma$, of σ (the set of the “places” where a descent occurs) is denoted by A . Here $A = \{3, 5, 6\}$. The inverse ligne of route, $\text{Iligne } \sigma$, of σ (the set of the “letters” i occurring to the right of the letters $(i + 1)$) is denoted by B ; here $B = \{1, 3, 5, 7\}$.

The **R**ight to left **M**aximum letter set, $\text{Rmals } \sigma$, of σ (the set of the letters greater than all the letters to their right) is denoted by C . The elements of $\text{Rmals } \sigma$ are the ordinates of the bullets “•”. Here $C = \{5, 7, 8\}$.

The **R**ight to left **m**inimum letter set, $\text{Rmil } \sigma$, of σ (the set of the letters less than all the letters to their right) is denoted by D . The elements of $\text{Rmals } \sigma$ are the ordinates of the crosses “×”. Here $D = \{1, 3, 5\}$.

Each graph corresponds to the action of an element $\text{Id}, \mathbf{r}, \mathbf{c}, \mathbf{rc}, \mathbf{i}, \mathbf{ir}, \mathbf{ic}, \mathbf{irc}$ of the dihedral group on the permutation σ . Notice that $n - A$ is to be understood as the set $\{n - x : x \in A\}$. Under each graph the ligne of route and the inverse ligne of route have been determined. Under graph \mathbf{rc} , for instance, $\text{Ligne} = n - A$ and $\text{Iligne} = n - B$ are to be read $\text{Ligne } \mathbf{rc}\sigma = n - A$ and $\text{Iligne } \mathbf{rc}\sigma = n - B$.







Fig. E20




Ligne = $[n - 1] \setminus A$
 Iligne = $[n - 1] \setminus (n - B)$
 Rmil = $n + 1 - C$
 Rmals = $n + 1 - D$




Ligne = $n - A$
 Iligne = $n - B$
 Lmil = $n + 1 - C$
 Lmals = $n + 1 - D$




Ligne = B
 Iligne = A
 Rmap = C
 Lmap = D



Ligne = $[n - 1] \setminus (n - B)$
 Iligne = $[n - 1] \setminus A$
 Lmap = $n + 1 - C$
 Rmap = $n + 1 - D$



Ligne = $[n - 1] \setminus B$
 Iligne = $[n - 1] \setminus (n - A)$
 Rmip = C
 Lmip = D



Ligne = $n - B$
 Iligne = $n - A$
 Lmip = $n + 1 - C$
 Rmip = $n + 1 - D$

EXERCISES AND EXAMPLES

Notice that whenever the transformation “**i**” is applied the “**l**” for “letter” is replaced by the “**p**” for “place”. Under graph **i**, for instance, $\text{Rmap} = C$ means that the **R**ight-to-left **M**aximum **p**lace of **i** σ is equal to C , i.e., $\text{Rmap } \mathbf{i}\sigma = C$.

21. *Variations on the cycle number polynomial.* Let $\text{cyc } \sigma$ denote the number of cycles of a permutation σ and let $C_n(x) := \sum_{\sigma} x^{\text{cyc } \sigma}$ ($\sigma \in \mathfrak{S}_n$).

(a) Then, $C_n(x) = x(x+1)\cdots(x+n-1)$ ($n \geq 1$).

(b) As in Exercice 20 let $\text{Rmals } \sigma$ (resp. $\text{Rmil } \sigma$) be the *set* of the *letters* greater than (resp. smaller than) all the letters located on their right in $\sigma = \sigma(1)\cdots\sigma(n)$. Let $\text{Rmap } \sigma$ (resp. $\text{Rmip } \sigma$) be the *set* of the *places* of those letters, respectively. Recall that $\# \text{Rmals } \sigma$ (resp. $\# \text{Rmil } \sigma$) designates the *number* of those letters. For instance, with $\sigma = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 \\ 3 & 1 & 5 & 6 & 2 & 4 \end{pmatrix}$ we have $\text{Rmals } \sigma = \{6, 4\}$, $\text{Rmap } \sigma = \{4, 6\}$. $\text{Rmil } \sigma = \{1, 2, 4\}$, $\text{Rmip } \sigma = \{2, 5, 6\}$, so that $\# \text{Rmals } \sigma = \# \text{Rmap } \sigma = 2$, $\# \text{Rmil } \sigma = \# \text{Rmip } \sigma = 3$. For each $n \geq 1$ we have: $C_n(x) = \sum_{\sigma} x^{\# \text{Rmals } \sigma} = \sum_{\sigma} x^{\# \text{Rmap } \sigma} = \sum_{\sigma} x^{\# \text{Rmil } \sigma} = \sum_{\sigma} x^{\# \text{Rmip } \sigma}$ ($\sigma \in \mathfrak{S}_n$).

(c) Let $C_n(x, y, q) := \sum_{\sigma} x^{\# \text{Rmap } \sigma} y^{\# \text{Rmip } \sigma} q^{\text{inv } \sigma}$ ($\sigma \in \mathfrak{S}_n$). Then, $C_n(x, y, q) = xy(y+qx)(y+q+q^2x)\cdots(y+q+q^2+\cdots+q^{n-2}+q^{n-1}x)$.

(d) Let $w = x_1x_2\dots x_n$ be the maj-coding of $\sigma = \sigma(1)\sigma(2)\dots\sigma(n)$. Then, $\text{Rmil } \sigma = \{i : 1 \leq i \leq n, x_i = 0\}$.

(e) Let $D_n(y, q) := \sum_{\sigma} y^{\# \text{Rmil } \sigma} q^{\text{maj } \sigma}$ ($\sigma \in \mathfrak{S}_n$). Then, $D_n(y, q) = y(y+q)(y+q+q^2)\cdots(y+q+q^2+\cdots+q^{n-1})$.

(f) Starting with the expression obtained in (c) for $C_n(x, y, q)$ derive

$$\sum_{n \geq 0} C_n(q, x, y) \frac{u^n}{(q; q)_n} = 1 - \frac{xy}{x+y-1} + \frac{xy}{x+y-1} \frac{\left(\frac{u}{1-q} - ux; q\right)_{\infty}}{\left(uy + \frac{uq}{1-q}; q\right)_{\infty}}.$$

22. *A lower-record extension of the q-maj-Eulerian polynomial.* For each $n \geq 0$ let $A_n(t, q, y) := \sum_{w \in \mathfrak{S}_n} t^{\text{des } w} q^{\text{maj } w} y^{\# \text{Rmil } \sigma} = \sum_k t^k A_{n,k}$.

(a) The following induction formula holds

$$A_{n,k} = (y+q+q^2+\cdots+q^k)A_{n-1,k} + (q^k+\cdots+q^{n-1})A_{n-1,k-1},$$

that implies

$$\begin{aligned} (1-q)A_n(t, q, 1, y) &= (y(1-q) + q - tq^n)A_{n-1}(t, q, 1, y) - q(1-t)A_{n-1}(tq, q, 1, y), \end{aligned}$$

for $n \geq 1$ and $A_0(t, q, 1, y) = 0$.

(b) Next, derive

$$\sum_{n \geq 0} A_n(t, q, y) \frac{u^n}{(t; q)_{n+1}} = \sum_{s \geq 0} t^s \frac{\left(\frac{uq}{1 - q + uq^{s+1}}; q\right)_{s+1}}{\left(u \frac{y(1 - q) + q}{1 - q + uq^{s+1}}; q\right)_{s+1}}$$

that specializes into the generating function (10.2a) for the q -maj-Eulerian polynomials ${}^{\text{maj}}A_n(t, q)$ when $y = 1$.

23. *The tableau emptying-filling involution.* Consider a standard Young tableau P of order n whose entries are the integers $1, 2, \dots, n$, for example the following one with $n = 5$ and consider the tableaux successively derived from P :

$$\begin{aligned} P &= \begin{array}{|c|c|} \hline 3 & 4 \\ \hline 1 & 2 & 5 \\ \hline \end{array} \\ \mapsto \begin{array}{|c|c|} \hline 3 & 4 \\ \hline \bullet & 2 & 5 \\ \hline \end{array} & \mapsto \begin{array}{|c|c|} \hline 3 & 4 \\ \hline 2 & \bullet & 5 \\ \hline \end{array} & \mapsto \begin{array}{|c|c|} \hline 3 & \bullet \\ \hline 2 & 4 & 5 \\ \hline \end{array} & \mapsto \begin{array}{|c|c|} \hline 3 & 5 \\ \hline 2 & 4 & 5 \\ \hline \end{array} & \text{(new entry: 5)} \\ & \mapsto \begin{array}{|c|c|} \hline 3 & 5 \\ \hline \bullet & 4 & 5 \\ \hline \end{array} & \mapsto \begin{array}{|c|c|} \hline \bullet & 5 \\ \hline 3 & 4 & 5 \\ \hline \end{array} & \mapsto \begin{array}{|c|c|} \hline 4 & 5 \\ \hline 3 & 4 & 5 \\ \hline \end{array} & \text{(new entry: 4)} \\ \mapsto \begin{array}{|c|c|} \hline 4 & 5 \\ \hline \bullet & 4 & 5 \\ \hline \end{array} & \mapsto \begin{array}{|c|c|} \hline 4 & 5 \\ \hline 4 & \bullet & 5 \\ \hline \end{array} & \mapsto \begin{array}{|c|c|} \hline 4 & 5 \\ \hline 4 & 5 & \bullet \\ \hline \end{array} & \mapsto \begin{array}{|c|c|} \hline 4 & 5 \\ \hline 4 & 5 & 3 \\ \hline \end{array} & \text{(new entry: 3)} \\ & \mapsto \begin{array}{|c|c|} \hline 4 & 5 \\ \hline \bullet & 5 & 3 \\ \hline \end{array} & \mapsto \begin{array}{|c|c|} \hline 4 & 5 \\ \hline 5 & \bullet & 3 \\ \hline \end{array} & \mapsto \begin{array}{|c|c|} \hline 4 & 5 \\ \hline 5 & 2 & 3 \\ \hline \end{array} & \text{(new entry: 2)} \\ & \mapsto \begin{array}{|c|c|} \hline 4 & 5 \\ \hline \bullet & 2 & 3 \\ \hline \end{array} & \mapsto \begin{array}{|c|c|} \hline 4 & 5 \\ \hline 1 & 2 & 3 \\ \hline \end{array} =: P^J & \text{(new entry: 1)} \end{aligned}$$

The entries are taken from the following set $\{1, 2, 3, 4, 5, \mathbf{1}, \mathbf{2}, \mathbf{3}, \mathbf{4}, \mathbf{5}, \bullet\}$. When a tableau contains a bullet “ \bullet ,” adopt the convention that the *neighbors* of the bullet are the *roman typed* entries just above or on the right of it. There are zero, one or two such neighbors. The passage “ \mapsto ” to the next tableau is defined as follows:

(i) if the tableau has no bullet, either the tableau has no *roman typed* entries and the final tableau P^J is reached, or there is such an entry, in particular on the leftmost bottom corner; call it x , then replace x by the bullet “ \bullet ”;

(ii) if the tableau contains a bullet, consider the neighbors of the bullet; if there is no neighbor and the number of boldface typed entries is i , replace the bullet by the boldface typed $(\mathbf{n} - \mathbf{i})$; if there is one neighbor, say, x ,

permute the two entries “•” and “ x ”; if there are two neighbors, let x be the smallest entry of the two and again permute the two entries “•” and “ x .”

It is obvious that P^J is a *standard Young tableau* having the *same shape* as P . Let $P \mapsto P^T$ be the *transposition* operation acting on tableaux. Further properties are the following:

- (i) *the map $P \mapsto P^J$ is an involution (called the tableau emptying-filling involution) having the property: $\text{ligne } P^J = n - \text{ligne } P$;*
- (ii) *J and T commute;*
- (iii) *if $\sigma \mapsto PQ$ under the Robinson-Schensted correspondence, then $\mathbf{r} \sigma \mapsto P^T Q^{JT}$.*

(The operations \mathbf{r} , as well as \mathbf{c} and \mathbf{i} , are defined in Exercice 19.)

Properties (i) and (iii) are not straightforward, the J -algorithm being not easy to handle (see [Sch63, 77] for a proof or use the *jeu de taquin* approach as developed, for instance, in [Lo02, chap. 5]).

(a) Using the above notations, if $\sigma \mapsto PQ$ (and $\mathbf{r} \sigma \mapsto P^T Q^{JT}$ by (iii)), then $\mathbf{c} \sigma \mapsto P^{JT} Q^T$.

(b) Define the involutions \mathbf{t} and \mathbf{j} of \mathfrak{S}_n by $\sigma \mapsto PQ \mapsto P^T Q^T \mapsto \mathbf{t} \sigma$, and $\sigma \mapsto PQ \mapsto P Q^J \mapsto \mathbf{j} \sigma$, where, in each case, the last arrow “ \mapsto ” refers to the inverse of the Robinson-Schensted correspondence. Then $\mathbf{i}^2 = \mathbf{j}^2 = \mathbf{t}^2 = (\mathbf{i}\mathbf{j})^4 = 1$, $\mathbf{i}\mathbf{t} = \mathbf{t}\mathbf{i}$, $\mathbf{j}\mathbf{t} = \mathbf{t}\mathbf{j}$. Also $\mathbf{r} = \mathbf{t}\mathbf{j} = \mathbf{j}\mathbf{t}$, $\mathbf{c} = \mathbf{i}\mathbf{j}\mathbf{t}\mathbf{i}$, so that the group generated by $\{\mathbf{i}, \mathbf{t}, \mathbf{j}, \mathbf{r}, \mathbf{c}\}$ is of order 16, in fact, the direct product of the dihedral group D_4 generated by $\{\mathbf{i}, \mathbf{j}\}$ by the group of two elements $\{1, \mathbf{t}\}$.

(c) Let $PQ = \begin{array}{|c|c|} \hline 3 & 4 \\ \hline 1 & 2 & 5 \\ \hline \end{array} \begin{array}{|c|c|} \hline 2 & 4 \\ \hline 1 & 3 & 5 \\ \hline \end{array}$ be the pair of standard Young tableaux associated with the permutation $\sigma = 3\ 1\ 4\ 2\ 5$ under the Robinson-Schensted correspondence. Determine P^J , Q^J , P^T , Q^T , P^{JT} , Q^{JT} and the sixteen pairs of standard Young tableaux associated with the sixteen permutations in the orbit of σ with respect to the group defined in (b).

(d) As already described in section 19 the following two properties hold: $\text{Ligne } \mathbf{j} \sigma = n - \text{Ligne } \sigma$ and $\text{ligne } \mathbf{j} \sigma = \text{ligne } \sigma$.

24. *A classical tool for symmetry proving.* Let σ be a permutation of $1, \dots, r$ and $\mathbf{m} = (m_1, \dots, m_r)$ be a sequence of nonnegative integers. Let $R(\mathbf{m})$ (resp. $R(\sigma\mathbf{m})$) denote the class of the rearrangements of $1^{m_1} \dots r^{m_r}$ (resp. the rearrangements of $1^{m_{\sigma(1)}} \dots r^{m_{\sigma(r)}}$). As noticed in Remark 8.2, $A_{\mathbf{m}}(t, q) = A_{\sigma\mathbf{m}}(t, q)$. A combinatorial proof of that property can be made as follows.

It suffices to prove the property when σ is a transposition $(i, i + 1)$ of two adjacent integers ($1 \leq i \leq r - 1$). Consider a word w in $R(\mathbf{m})$ and write all its factors of the form $(i + 1)i$ in bold-face; then replace

all the maximal factors of the form $i^p(i+1)^q$ that do not involve any boldface letters by $i^q(i+1)^q$. Finally, rewrite all the boldface letters in roman type. Clearly, the transformation is a bijection that maps each word w in $R(\mathbf{m})$ onto a word w' in $R((i, i+1)\mathbf{m})$ with the property that $(\text{des}, \text{maj}) w = (\text{des}, \text{maj}) w'$.

If we take the statement of Theorem 17.1 as the definition of the Schur function $s_\lambda(x)$, the symmetric nature of $s_\lambda(x)$ does not appear. To still prove that the Schur function is symmetric, we can use a similar argument.

25. *Line of route-indexed Eulerian polynomials.* Let $L = \{\ell_1 < \dots < \ell_k\}$ be a subset of $\{1, 2, \dots, n-1\}$ with the conventions $\ell_0 := 0$ and $\ell_{k+1} := n$. The L -indexed Eulerian polynomial $A_L(t, q)$ is defined to be

$$A_L(t, q) := \sum_{\sigma} t^{\text{idex } \sigma} q^{\text{imaj } \sigma} \quad (\sigma \in \mathfrak{S}_n, \text{Ligne } \sigma = L).$$

Form the $(k+1) \times (k+1)$ matrix $N_r(L, n)$:

$$N_r(L, n) = \begin{pmatrix} \begin{bmatrix} \ell_1 - \ell_0 + r \\ r \\ 1 \end{bmatrix} & \begin{bmatrix} \ell_2 - \ell_0 + r \\ r \\ \ell_2 - \ell_1 + r \\ r \end{bmatrix} & \cdots & \begin{bmatrix} \ell_k - \ell_0 + r \\ r \\ \ell_k - \ell_1 + r \\ r \end{bmatrix} & \begin{bmatrix} \ell_{k+1} - \ell_0 + r \\ r \\ \ell_{k+1} - \ell_1 + r \\ r \end{bmatrix} \\ 0 & 1 & \cdots & \begin{bmatrix} \ell_k - \ell_2 + r \\ r \end{bmatrix} & \begin{bmatrix} \ell_{k+1} - \ell_2 + r \\ r \end{bmatrix} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \cdots & 1 & \begin{bmatrix} \ell_{k+1} - \ell_k + r \\ r \end{bmatrix} \end{pmatrix}.$$

In particular, $N_r(\emptyset, n) = \begin{bmatrix} n+r \\ r \end{bmatrix}$. The purpose of this exercise is to prove the formula:

$$(E22.1) \quad \frac{A_L(t, q)}{(t; q)_{n+1}} = \sum_{r \geq 0} t^r \det N_r(L, n).$$

(a) Let $W_r(L, n)$ denote the set of the words $w = w_1 w_2 \dots w_n$, of length n , whose letters are nonnegative integers satisfying the identities

$$\begin{aligned} (*)_{(L, n)} \quad & r \geq w_1 \geq \dots \geq w_{\ell_1} \geq 0; \quad r \geq w_{\ell_1+1} \geq \dots \geq w_{\ell_2} \geq 0; \quad \dots \\ & r \geq w_{\ell_k+1} \geq \dots \geq w_n \geq 0; \\ & w_{\ell_1} < w_{\ell_1+1}, \quad w_{\ell_2} < w_{\ell_2+1}, \quad \dots, \quad w_{\ell_k} < w_{\ell_k+1}. \end{aligned}$$

Then $\sum_{w \in W_r(L, n)} q^{\text{tot } w} = \det N_r(L, n)$. [By induction on k .]

EXERCISES AND EXAMPLES

(b) Let $S_r(L, n)$ be the set of all pairs (σ, s) , where σ is a permutation of $1, 2, \dots, n$, of ligne of route L , and where $s = s_1 s_2 \dots s_n$ is a nonincreasing word having the properties:

$$(E22.2) \quad r \geq s_1 \geq s_2 \geq \dots \geq s_n \geq 0;$$

$$\text{Ligne } \sigma = L; \quad i \in \text{Iligne } \sigma \Rightarrow s_i > s_{i+1}.$$

Each word $w = w_1 w_2 \dots w_n \in W_r(L, n)$ is mapped onto a pair $(\sigma, s) \in S_r(L, n)$ in the following way: suppose that the nondecreasing rearrangement of w is of the form $i_1^{a_1} \dots i_m^{a_m}$, with $i_1 < \dots < i_m$ and $a_1 \geq 1, \dots, a_m \geq 1$. Read w from left to right and give the labels $1, 2, \dots, a_m$ to the a_m letters equal to i_m ; continue, again from left to right, giving the labels $a_m+1, \dots, a_m+a_{m-1}$ to the a_{m-1} letters equal to i_{m-1} and so on. Reading those labels from left to right yields a permutation $\sigma = \sigma(1)\sigma(2) \dots \sigma(n)$. The word s is the nonincreasing rearrangement of w .

Example. Let $n = 9, r = 8, L = \{2, 5, 7\}$. The word $w \in N_8(L, 9)$ and the associated pair $(\sigma, s) \in S_8(L, 9)$ are shown in the following table:

$$\begin{pmatrix} L = \\ w = \\ \sigma = \\ \text{Iligne} = \\ s = \end{pmatrix} \begin{pmatrix} 2 & 5 & 7 \\ 5 & 5 & 7 & 4 & 1 & 3 & 0 & 4 & 4 \\ 2 & 3 & 1 & 4 & 8 & 7 & 9 & 5 & 6 \\ \mathbf{1} & & & & & & \mathbf{6} & \mathbf{7} & \\ 7 & 5 & 5 & 4 & 4 & 4 & 3 & 1 & 0 \end{pmatrix}.$$

The map $w \mapsto (\sigma, s)$ is a bijection of $W_r(L, n)$ onto $S_r(L, n)$.

(c) The following identity holds

$$\frac{\sum_{\sigma, \text{Ligne } \sigma=L} t^{\text{idex } \sigma} q^{\text{imaj } \sigma}}{(t; q)_{n+1}} = \sum_{r \geq 0} t^r \sum_{w \in W_r(L, n)} q^{\text{tot } w},$$

so identity (E22.1) also holds by (a) and (c).

(d) Let $A_L(q) := \sum_{\sigma} q^{\text{imaj } \sigma}$ (Ligne $\sigma = L$). Then $A_L(q)/(q; q)_n = \det N$, where $N = (N_{i,j})$ ($1 \leq i, j \leq k+1$) is the $(k+1) \times (k+1)$ -matrix

$$N_{i,j} := \begin{cases} 1/(q; q)_{\ell_j - \ell_{i-1}}, & \text{if } i \leq j; \\ 1, & \text{if } i = j + 1; \\ 0, & \text{else.} \end{cases}$$

(e) We also have $A_L(q) := \sum_{\sigma} q^{\text{inv } \sigma}$ (Ligne $\sigma = L$).

26. *Partitions, permutations and descents.* The statistic “ligne of route” (“Ligne”) is defined in § 11.4. The notations used in (b) refer to § 14.1.

(a) Let (a_1, a_2, \dots, a_k) be a *composition* of the integer n , that is, an ordered sequence of integers such that $a_1 \geq 1, a_2 \geq 1, \dots, a_k \geq 1$ and $a_1 + a_2 + \dots + a_k = n$. The number of permutations $\sigma = \sigma(1)\sigma(2) \cdots \sigma(n)$, of order n , such that $\text{Ligne } \sigma \subset \{a_1, a_1 + a_2, \dots, a_1 + a_2 + \dots + a_{k-1}\}$ is equal to $\binom{n}{a_1, a_2, \dots, a_k} = n! / (a_1! a_2! \cdots a_k!)$.

(b) For each permutation $\sigma = \sigma(1)\sigma(2) \cdots \sigma(n)$ and each integer $i = 1, 2, \dots, n$ let $d_i(\sigma)$ denote the number of *descents* ($\sigma(j) > \sigma(j+1)$) occurring in the *right factor* $\sigma(i)\sigma(i+1) \cdots \sigma(n)$ and let

$$\mathbf{q}^{\mathbf{d}(\sigma)} := q_1^{d_1(\sigma)} q_2^{d_2(\sigma)} \cdots q_n^{d_n(\sigma)},$$

where q_1, q_2, \dots, q_n is a finite sequence of variables. Notice that $d_1(\sigma)$ is the usual number of descents of σ . Furthermore, if $\lambda = (\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n \geq 0)$ is a partition, whose number $l(\lambda)$ of *positive* parts is at most equal to n , let $m_i(\lambda)$ designate its number of parts equal to i , for $i = 0, 1, \dots, n$. Then, define

$$\binom{n}{\mathbf{m}(\lambda)} := \binom{n}{m_0(\lambda), m_1(\lambda), \dots, m_n(\lambda)}, \quad \mathbf{q}^\lambda := q_1^{\lambda_1} q_2^{\lambda_2} \cdots q_n^{\lambda_n}.$$

The following identity holds:

$$\sum_{l(\lambda) \leq n} \binom{n}{\mathbf{m}(\lambda)} \mathbf{q}^\lambda = \frac{\sum_{\sigma \in \mathfrak{S}_n} \mathbf{q}^{\mathbf{d}(\sigma)}}{(1 - q_1)(1 - q_1 q_2) \cdots (1 - q_1 q_2 \cdots q_n)}.$$

[Expand the right-side of the previous identity and use (a).]

27. *A further extension of the MacMahon Verfahren.* Let $U := (S_<, S_\leq, L_<, L_\leq)$ be a partition of the alphabet $X = \{1, \dots, r\}$ such that $S_< \cup S_\leq = \{1, \dots, h\}$ (the *small* letters) and $L_< \cup L_\leq = \{h + 1, \dots, r\}$ (the *large* letters) for a certain h ($0 \leq h \leq r$). Let $w = x_1 x_2 \cdots x_m$ be a word in the alphabet and let $x_{m+1} := h + \frac{1}{2}$. An integer i such that $1 \leq i \leq m$ is said to be a *U-descent* in w , if either $x_i > x_{i+1}$, or $x_i = x_{i+1}$ and $x_i \in S_\leq \cup L_\leq$. Because of the convention $x_{m+1} := h + \frac{1}{2}$ there is a *U-descent* in position m iff $x_m \in L_< \cup L_\leq$, so that the four blocks of the partition determine different kinds of descents.

Let $\text{des}_U w$ (resp. $\text{maj}_U w$) denote the *number* (resp. the *sum*) of the *U*-descents in w . For each sequence $\mathbf{m} = (m_1, \dots, m_r)$ consider the generating polynomial for the class $R(\mathbf{m})$ by the pair $(\text{des}_U, \text{maj}_U)$, i.e.,

EXERCISES AND EXAMPLES

$A_{\mathbf{m}}^U(t, q) = \sum_w t^{\text{des}_U w} q^{\text{maj}_U w}$ ($w \in R(\mathbf{m})$) (see section 6). Then the extensions of identities (7.7), (7.8) and (6.13) read:

$$(E27.1) \quad \frac{1}{(t; q)_{1+\|\mathbf{m}\|}} A_{\mathbf{m}}^U(t, q) = \sum_{s \geq 0} t^s \prod_{i \in S_{<}} \begin{bmatrix} m_i + s \\ m_i \end{bmatrix} \prod_{i \in S_{\leq}} q^{\binom{m_i}{2}} \begin{bmatrix} s + 1 \\ m_i \end{bmatrix} \\ \times \prod_{i \in L_{<}} q^{m_i} \begin{bmatrix} m_i + s - 1 \\ m_i \end{bmatrix} \prod_{i \in L_{\leq}} q^{\binom{m_i + 1}{2}} \begin{bmatrix} s \\ m_i \end{bmatrix};$$

$$(E27.2) \quad \sum_{\mathbf{m}} A_{\mathbf{m}}^U(t, q) \frac{\mathbf{u}^{\mathbf{m}}}{(t; q)_{1+\|\mathbf{m}\|}} = \sum_{s \geq 0} t^s \frac{\prod_{i \in S_{\leq}} (-u_i; q)_{s+1} \prod_{i \in L_{\leq}} (-q u_i; q)_s}{\prod_{i \in S_{<}} (u_i; q)_{s+1} \prod_{i \in L_{<}} (q u_i; q)_s};$$

$$(E27.3) \quad \sum_{\mathbf{m}} A_{\mathbf{m}}^U(1, q) \frac{\mathbf{u}^{\mathbf{m}}}{(q; q)_{\|\mathbf{m}\|}} = \frac{\prod_{i \in S_{\leq}} (-u_i; q)_{\infty} \prod_{i \in L_{\leq}} (-q u_i; q)_{\infty}}{\prod_{i \in S_{<}} (u_i; q)_{\infty} \prod_{i \in L_{<}} (q u_i; q)_{\infty}};$$

where $\|\mathbf{m}\| = m_1 + \dots + m_r$.

28. A *maj-inv transformation for signed words*. The purpose is to extend the transformation, introduced in section 11, to signed words. A first extension has already been given in § 20.5 for signed permutations. Keep the notations of Exercise 27, in particular the definitions of des_U and of maj_U , and for each word $w = x_1 x_2 \dots x_m$ with letters in $S_{<} \cup S_{\leq} \cup L_{<} \cup L_{\leq}$ define

$$\text{inv}_U w := \sum_{i < j} \left(\chi(x_i > x_j) + \chi(x_i = x_j \in S_{\leq} \cup L_{\leq}) \right) + \#\{i : x_i \in L_{<} \cup L_{\leq}\}.$$

There are then four kinds of U -inversions. Let $\text{inv} A_{\mathbf{m}}^U(q) := \sum_{w \in R(\mathbf{m})} q^{\text{inv}_U w}$.

(a) The following identity holds

$$\text{inv} A_{\mathbf{m}}^U(q) = \begin{bmatrix} \|\mathbf{m}\| \\ m_1, \dots, m_r \end{bmatrix} \prod_{i \in S_{\leq}} q^{\binom{m_i}{2}} \prod_{i \in L_{<}} q^{m_i} \prod_{i \in L_{\leq}} q^{m_i + \binom{m_i}{2}}.$$

It then follows that the generating function $\sum_{\mathbf{m}} \text{inv} A_{\mathbf{m}}^U(q) \mathbf{u}^{\mathbf{m}} / (q; q)_{\|\mathbf{m}\|}$ is equal to the right-hand side of (E27.3). Thus, $\text{inv} A_{\mathbf{m}}^U(q) = A_{\mathbf{m}}^U(1, q)$, the generating polynomial for $R(\mathbf{m})$ by maj_U .

(b) By using the transformation Φ of Theorem 11.3 build a bijection Ψ of $R(\mathbf{m})$ onto itself having the property: $\text{maj}_U w = \text{inv}_U \Psi(w)$.

(c) Let $S_{<} = \{1, 3\}$, $S_{\leq} = \{2, 4\}$, $L_{<} = \{5, 7\}$, $L_{\leq} = \{6, 8\}$ and consider the word $w = 6, 2, 2, 3, 1, 7, 7, 4, 8, 6$ whose U -descents occur at positions 1, 2, 4, 7, 9 and 10, so that $\text{maj}_U w = 1 + 2 + 4 + 7 + 9 + 10 = 33$. Notice that the factor 2, 2 involves a descent because $2 \in S_{\leq}$, but not the factor 7, 7 because $7 \in L_{<}$. There is a descent at the right end because $6 \in L_{\leq}$. Determine $\Psi(w)$ and verify that $\text{inv}_U \Psi(w) = 33$.

29. *The Brenti homomorphism.* The homomorphism materializes the similarity between the classical identities for the symmetric functions and the identities found for the Eulerian polynomials and their extensions. Recall that the generating functions for the elementary symmetric functions, on the one hand, and for homogeneous symmetric functions, on the other hand, read (see (14.1) and (14.2))

$$(E24.1) \quad E(u) = \sum_{r \geq 0} e_r u^r = \prod_{i \geq 1} (1 + x_i u);$$

$$(E24.2) \quad H(u) = \sum_{r \geq 0} h_r u^r = \prod_{i \geq 1} (1 - x_i u)^{-1};$$

so that

$$(E24.3) \quad E(-u)H(u) = 1.$$

On the other hand, the exponential generating for the Eulerian polynomials $A_n(t)$ (see Exercise 13) can be put into the form:

$$(E24.4) \quad \left(1 + \sum_{k \geq 1} \frac{1 - t^{k-1}}{k!} (-u)^k\right) \left(\sum_{k \geq 0} \frac{A_k(t)}{k!} u^k\right) = 1.$$

The comparison of (E24.3) and (E24.4) shows that the following two statements are equivalent:

$$\begin{aligned} \xi(e_k) &= \frac{1 - t^{k-1}}{k!} \quad (k \geq 1), & \xi(e_0) &= 1; \\ \xi(h_k) &= \frac{A_k(t)}{k!} \quad (k \geq 0). \end{aligned}$$

The mapping ξ is called the *Brenti homomorphism*.

(a) For each sequence $\lambda = (\lambda_1, \dots, \lambda_l)$ of positive integers such that $\lambda_1 + \dots + \lambda_l = n$ (in particular, for each partition of n) and each permutation $\sigma = \sigma(1) \dots \sigma(n)$ define $\text{des}_\lambda \sigma$ to be the number of descents $\sigma(i) > \sigma(i+1)$ with i different from $\lambda_1, \lambda_1 + \lambda_2, \dots, \lambda_1 + \dots + \lambda_l$. Then form $A_\lambda(t) := \sum_{\sigma} t^{\text{des}_\lambda \sigma}$ ($\sigma \in \mathfrak{S}_n$). Then $\xi(h_\lambda) = A_\lambda(t)/n!$

EXERCISES AND EXAMPLES

(b) The factorial generating function for the polynomials ${}^{\text{inv}}A_n(t, q)$ (see (10.10)) may be rewritten as:

$$\left(1 + \sum_{k \geq 1} \frac{(1 - t^{k-1})q^{\binom{k}{2}}}{(q; q)_k} (-u)^k\right) \left(\sum_{k \geq 0} \frac{{}^{\text{inv}}A_k(t, q)}{(q; q)_k} u^k\right) = 1.$$

so that the following two statements are equivalent:

$$\begin{aligned} \xi_q(e_k) &= \frac{(1 - t^{k-1})q^{\binom{k}{2}}}{(q; q)_k} \quad (k \geq 1), \quad \xi_q(e_0) = 1; \\ \xi_q(h_k) &= \frac{A_k(t, q)}{(q; q)_k} \quad (k \geq 0). \end{aligned}$$

Determine $\xi_q(h_\lambda)$.

(c) Let $B_n(t, Q, q) = \sum_{\Sigma, \sigma} t^{\text{ddes}(\Sigma, \sigma)} Q^{\text{inv}\Sigma} q^{\text{coinv}\sigma}$ $((\Sigma, \sigma) \in \mathfrak{S}_n \times \mathfrak{S}_n)$ be the generating polynomial for the pairs of permutations by the three-variable statistic (ddes, inv, coinv). By specializing Theorem 21.1 we have

$$\frac{1 - t}{-t + \mathbf{J}((1 - t)u; Q, q)} = \sum_{n \geq 0} \frac{B_n(t, Q, q)}{(Q; q)_n (q; q)_n} u^n.$$

The following two statements are also equivalent:

$$\begin{aligned} \xi_{Q, q}(e_k) &= \frac{(1 - t^{k-1})Q^{\binom{k}{2}}}{(Q; q)_k (q; q)_k} \quad (k \geq 1), \quad \xi_{Q, q}(e_0) = 1; \\ \xi_{Q, q}(h_k) &= \frac{B_k(t, Q, q)}{(Q; q)_k (q; q)_k} \quad (k \geq 0). \end{aligned}$$

(d) Let $P(u) = \sum_{r \geq 1} p_r u^{r-1}$ be the generating series for the power symmetric functions. Using identity $E'(u)/E(u) = P(-u)$ derived in Theorem 14.2 (vi), determine $\xi(p_{k+1})$ $(k \geq 0)$.

(e) Using the relation $P(u) = E'(u)H(u)$ determine $\xi_q(p_n)$ and $\xi_{Q, q}(p_n)$ $(n \geq 1)$.

(f) By Theorem 20.1 the exponential generating function for the generating polynomials $B_n(X, Y, t)$ for *signed permutations* by number of descents is

$$\frac{(1 - t) \exp((t - 1)X)}{-t + \exp((t - 1)(X + Y))} = \sum_{n \geq 0} \frac{1}{n!} B_n(X, Y, t),$$

a relation that may be written as

$$\left(1 + \sum_{n \geq 1} \frac{(1 - t)^{n-1}}{n!} (-tX^n + (-Y)^n)\right) \cdot \sum_{n \geq 0} \frac{1}{n!} B_n(X, Y, t) = 1.$$

Make the changes $X \leftarrow XY$, $Y \leftarrow Y$. Then $-tX^n + (-Y)^n$ is changed into $(-Y)^n((-1)^{n-1}tX^n + 1)$, and $B_n(X, Y, t)$ into $Y^n B_n(X, 1, t)$, so that the following two statements are equivalent:

$$\begin{aligned} \xi_B(e_n) &= \frac{(1-t)^{n-1}}{n!}((-1)^{n-1}tX^n + 1) \quad (n \geq 1), \quad \xi_B(e_0) = 1; \\ \xi_B(h_n) &= \frac{B_n(X, 1, t)}{n!} \quad (n \geq 0). \end{aligned}$$

With the changes $X \leftarrow -X$, $Y \leftarrow XY$ the two statements are equivalent:

$$\begin{aligned} \xi_b(e_n) &= \frac{(1-t)^{n-1}}{n!}(Y^n - t) \quad (n \geq 1), \quad \xi_b(e_0) = 1; \\ \xi_b(h_n) &= \frac{B_n(-1, Y, t)}{n!} \quad (n \geq 0). \end{aligned}$$

30. *The des-length distribution for signed permutations.* The length l_{Cox} of a signed permutation (σ, ε) is defined by

$$\begin{aligned} l_{\text{Cox}}(\sigma, \varepsilon) &:= \#\{i < j : \varepsilon(i) = \varepsilon(j), \sigma(i) > \sigma(j)\} \\ &\quad + \#\{i < j : \varepsilon(i) = x, \varepsilon(j) = y\} + \sum_{\varepsilon(i)=x} \cdot \sigma(i) \end{aligned}$$

Let B''_n be the generating function for the group of the signed permutations of order n by the pair $(\text{des}, l_{\text{Cox}})$ expressed in the form

$$B''_n = \sum_{(\sigma, \varepsilon)} X^{\ell(\varepsilon|x)} t^{\text{des}(\sigma, \varepsilon)} q^{l_{\text{Cox}}(\sigma, \varepsilon)}.$$

Then

$$(*) \quad \frac{1-t}{1-t e_q(u(1-t))} \sum_{n \geq 0} \frac{(u(1-t))^n}{(-Xq; q)_n (q; q)_n} = \sum_{n \geq 0} \frac{u^n}{(-Xq; q)_n (q; q)_n} B''_n.$$

The first values of B''_n are the following:

$$\begin{aligned} B''_1 &= 1 + tX; \quad B''_2 = (t+q) + tq(1+q)^2X + tq^3(t+q)X^2; \\ B''_3 &= (t^2q^3 + 2tq^2 + 2tq + 1) + t(1+q+q^2)(1+q+2q^2+tq+ tq^3)X \\ &\quad + t^2(1+q+q^2)(1+q^2+2tq+ tq^2+ tq^3)X^2 \\ &\quad + tq^6(t^2q^3 + 2tq^2 + 2tq + 1)X^3. \end{aligned}$$

Introduce the polynomials

$$A_n^k(t, q) := \sum_{\sigma \in \mathfrak{S}_n} t^{\text{des}_k \sigma} q^{\text{inv} \sigma} \quad (0 \leq k \leq n),$$

where

$$\text{des}_k \sigma := \begin{cases} \text{des} \sigma, & \text{if } \sigma(n) \leq n-k; \\ 1 + \text{des} \sigma, & \text{if } \sigma(n) \geq n-k+1. \end{cases}$$

With the notations of section 23

$$A_n^n(t, q) = \text{inv}A_n(t, q).$$

An easy derivation yields

$$B_n'' = \sum_{0 \leq k \leq n} X^k q^{\binom{k+1}{2}} \begin{bmatrix} n \\ k \end{bmatrix}_q A_n^k(t, q).$$

By Theorem 10.1 the left-hand side of (*) can be rewritten as

$$\sum_{l \geq 0} \frac{u^l}{(q; q)_l} A_l^l(t, q) \sum_{m \geq 0} \frac{(u(1-t))^m}{(-Xq; q)_m (q; q)_m},$$

that is denoted by

$$\sum_{n \geq 0} \frac{u^n}{(-Xq; q)_n (q; q)_n} B_n^{(3)}.$$

Proving (*) amounts to proving the identity $B_n'' = B_n^{(3)}$.

31. *The Wachs Involution.* Let ${}^0A_{2n}(t) := \sum_{\sigma} t^{1+\text{des } \sigma}$ and $\text{sgn}A_{2n}(t) := \sum_{\sigma} \text{sgn } \sigma t^{1+\text{des } \sigma}$ ($\sigma \in \mathfrak{S}_{2n}$) be the two polynomials introduced in Theorem 23.7, where the two identities $\text{sgn}A_{2n}(t) = (1-t)^n {}^0A_n(t)$ and $\text{sgn}A_{2n+1}(t) = (1-t)^n {}^0A_{n+1}(t)$ were proved analytically. A combinatorial proof can be derived by imagining an involution on \mathfrak{S}_{2n} preserving the ligne of route of each permutation and changing the signature of some of them in such a way that the generating polynomial for the remaining permutations is precisely $(1-t)^n {}^0A_n(t)$. This is accomplished by the *Wachs involution* ϕ defined as follows: consider a permutation $\sigma = \sigma(1)\sigma(2) \cdots \sigma(2n)$ and let j be the *smallest* integer such that $1 \leq j \leq n$ and the letters $2j-1$ and $2j$ are *not* adjacent in the word σ . Then define

$$\phi(\sigma) := \begin{cases} (2j-1, 2j) \sigma, & \text{if such an integer } j \text{ exists;} \\ \sigma, & \text{otherwise.} \end{cases}$$

The involution preserves the ligne of route and changes the signature, so that if \mathfrak{F}_{2n} designates the set of its fixed points, we have: $\text{sgn}A_{2n}(t) = \sum_{\sigma} \text{sgn } \sigma t^{1+\text{des } \sigma}$ ($\sigma \in \mathfrak{F}_{2n}$). Now, if σ is in \mathfrak{F}_{2n} , there is a unique permutation $\tau = \tau(1)\tau(2) \cdots \tau(n)$ of order n such that for each $i = 1, 2, \dots, n$ the *unordered* pairs $\{\sigma(2i-1), \sigma(2i)\}$ and $\{2\tau(i), 2\tau(i)-1\}$ are identical. Let $E(\sigma)$ be the set of all integers i such that the two-letter factor $2\tau(i), (2\tau(i)-1)$ occurs in σ (and *not* the factor $(2\tau(i)-1), 2\tau(i)$). Then $\sigma \mapsto (\tau, E(\sigma))$ is a bijection of \mathfrak{F}_{2n} onto the Cartesian product $\mathfrak{S}_n \times \mathfrak{P}(\{1, 2, \dots, n\})$, where $\mathfrak{P}(\{1, 2, \dots, n\})$ denotes the power set of $\{1, 2, \dots, n\}$, such that

$$\begin{aligned} \text{inv } \sigma &= 4 \text{inv } \tau + \#E(\sigma), \\ \text{Ligne } \sigma &= \{2i : i \in \text{Ligne } \tau\} \cup \{2i-1 : i \in E(\sigma)\}. \end{aligned}$$

Accordingly,

$$\begin{aligned} \operatorname{sgn}A_{2n}(t) &= \sum_{\sigma \in \mathfrak{S}_{2n}} \operatorname{sgn} \sigma t^{1+\operatorname{des} \sigma} = \sum_{\tau \in \mathfrak{S}_n, EC\{1, \dots, n\}} (-1)^{\#E} t^{\operatorname{des} \tau + \#E} \\ &= \sum_{EC\{1, \dots, n\}} (-t)^{\#E} \sum_{\tau \in \mathfrak{S}_n} t^{1+\operatorname{des} \tau} = (1-t)^n {}^0A_n(t). \end{aligned}$$

Let $\operatorname{maj}A_n(t, q)$ be the q -maj-Eulerian polynomial (see section 10) and $\operatorname{sgn}A_n(t, q) := \sum_{\sigma} \operatorname{sgn} \sigma t^{1+\operatorname{des} \sigma} q^{\operatorname{maj} \sigma}$ ($\sigma \in \mathfrak{S}_n$) be the *signed q -Eulerian polynomial*. Using the above procedure it is shown that

$$\operatorname{sgn}A_{2n}(t, q) = (tq; q^2)_n t^{\operatorname{maj}A_n(t, q^2)}.$$

32. *The divisibility of the q -tangent coefficients.* As introduced in section 24 the coefficients of the q -tangent functions are denoted by $D_{2n+1}(q)$ ($n \geq 0$). Each integer $n \geq 1$ can be written as $n = m2^l$, where m is odd and $l \geq 0$. Define $Ev_n(q) := \prod_{0 \leq j \leq l} (1 + q^{m2^j})$ and also

$$F_n(q) := \begin{cases} \prod_{1 \leq i \leq n} Ev_i(q), & \text{if } n \text{ is odd;} \\ (1 + q^2) \prod_{1 \leq i \leq n} Ev_i(q), & \text{if } n \text{ is even.} \end{cases}$$

In an equivalent manner, let $Ev_0(q) = F_0(q) := 1$, $F_1(q) := 1 + q$, $F_2(q) := (1 + q)^2(1 + q^2)^2$ and $F_n(q) := F_{n-2}(q) Ev_{n-1}(q) Ev_n(q)$ for $n \geq 3$. Define the exponents $a(n, i)$ by $F_n(q) := \prod_{1 \leq i \leq n} (1 + q^i)^{a(n, i)}$.

- (a) Give the table of the coefficients $a(n, i)$ ($1 \leq i \leq n$) for $1 \leq n \leq 8$.
- (b) For $i \geq 1$ let $\Phi_i(q)$ be the i -th cyclotomic polynomial with $\phi_1(q) = 1 - q$, so that $1 - q^i = \prod_{d|i} \phi_d(q)$. For $n = m2^l$ (m odd, $l \geq 0$) and $j = 0, 1, \dots, l$ let

$$A_j := \{d : d \mid m2^{j+1}, d \nmid m2^j\}, \quad B := \{d : d \mid m2^{l+1}, d \text{ even}\}.$$

Then $1 + q^{m2^j} = \prod_{d \in A_j} \phi_d(q)$ ($0 \leq j \leq l$).

- (c) Also $Ev_n(q) = \prod_d \phi_d(q)$ ($d \mid 2n, d$ even).
- (d) For each $n \geq 1$ let $Od_n(q) := \prod_d \phi_d(q)$ ($d \mid 2n, d$ odd), so that $1 - q^{2n} = Od_n(q) Ev_n(q)$. Using that factorization we see that the product $\left[\begin{smallmatrix} 2n \\ 2k+1 \end{smallmatrix} \right] \frac{Ev_0(q) Ev_1(q) \dots Ev_k(q)}{Ev_{n-k}(q) Ev_{n-k+1}(q) \dots Ev_n(q)}$ is a polynomial in q .
- (e) Finally, $F_n(q)$ divides $D_{2n+1}(q)$.

EXERCISES AND EXAMPLES

33. *Congruences for the q -secant coefficients.* The purpose is to prove the congruence

$$D_{2n}(q) \equiv q^{2n(n-1)} \pmod{(q+1)^2}$$

by using a combinatorial argument.

(a) Let $1 \leq i \leq 2n - 1$. A rising alternating permutation $\sigma = \sigma(1)\sigma(2) \dots \sigma(n)$ is said to be *i -balanced* if the word σ contains the factor $i(i+1)$, or if the letter $(i+1)$ lies on the left of i in σ and not adjacent to i . For example, the permutation $\sigma = 263415$ is i -balanced only for $i = 1, 3, 5$. If σ is i -balanced for every $i = 1, 2, \dots, (2n - 1)$, it is said to be *balanced*.

The permutation $(2n - 1)2n(2n - 3)(2n - 2) \dots 3412$ is the unique balanced rising alternating permutation of order $2n$.

(b) Let σ be a nonbalanced rising alternating permutation. The greatest integer i such that σ is not i -balanced is denoted by i' . Then σ is necessarily of the form $\sigma = w i' w' (i' + 1) w''$, where w' is a nonempty factor. The permutation $\Phi(\sigma) := w (i' + 1) w' i' w''$ is still rising alternating and $\text{inv}\Phi(\sigma) = \text{inv}\sigma + 1$.

(c) The polynomial $D_{2n}(q)$ is monic of degree $2n(n - 1)$.

(d) For $n \geq 2$ let A_{2n} (resp. B_{2n}) be the set of all rising alternating permutations of length $2n$ beginning with the factor $(2n - 1)2n$ (resp. ending with 12). By induction on n ,

$$\sum_{\sigma \in A_{2n} \cup B_{2n}} q^{\text{inv}\sigma} \equiv q^{2n(n-1)} \pmod{(q+1)^2}.$$

(e) Let E_{2n} be the complement of $A_{2n} \cup B_{2n}$. Then

$$\sum_{\sigma \in E_{2n}} q^{\text{inv}\sigma} \equiv 0 \pmod{(q+1)^2}.$$

34. *The “Master Theorem” traditional statement.* *Mutatis mutandis*, as stated in [Ma15, p.97–98], this theorem says:

Let $X_i = \sum_{0 \leq j \leq r} a_{i,j} x_j$ ($0 \leq i \leq r$) and let n_0, n_1, \dots, n_r be nonnegative integers. Then, the coefficient of $x_0^{n_0} x_1^{n_1} \dots x_r^{n_r}$ in the expansion of the product $X_0^{n_0} X_1^{n_1} \dots X_r^{n_r}$ is equal to the coefficient of $x_0^{n_0} x_1^{n_1} \dots x_r^{n_r}$ in the expansion of the inverse of the determinant $\det(\delta_{i,j} - a_{i,j} x_j)_{0 \leq i,j \leq r}$.

Accordingly, if $C(n_0, n_1, \dots, n_r)$ designates the coefficient in the expansion of the product previously defined, the following identity holds

$$\sum_{n_0 \geq 0, n_1 \geq 0, \dots, n_r \geq 0} C(n_0, n_1, \dots, n_r) x_0^{n_0} x_1^{n_1} \dots x_r^{n_r} = 1 / \det(\delta_{i,j} - a_{i,j} x_j)_{0 \leq i,j \leq r}.$$

Rewrite formula (25.8) as $\text{Bos} = 1/\det(I - C_r)$ and apply the homomorphism $\Phi\left(\binom{i}{j}\right) := a_{i,j}x_j$.

35. *The generating function for vertical derangements.* By *vertical derangement* it is meant a word $w = x_1x_2 \cdots x_n$ from $[0, r]^*$ having the property that the biword $\binom{\bar{w}}{w} = \binom{\bar{x}_1 \bar{x}_2 \cdots \bar{x}_n}{x_1 x_2 \cdots x_n}$, where $\bar{w} = \bar{x}_1 \bar{x}_2 \cdots \bar{x}_n$ is the non-decreasing rearrangement of w , has no (vertical) biletter $\binom{\bar{x}_i}{x_i}$ such that $\bar{x}_i = x_i$. The set of all vertical derangements is denoted by $[0, r]_v^*$.

Let C'_r be the matrix $\left((1 - \delta_{i,j})\binom{i}{j}\right)_{(0 \leq i, j \leq r)}$, which only differs from the matrix C_r by the diagonal entries which are all null. Then,

$$\sum_{w \in [0, r]_v^*} \binom{\bar{w}}{w} = \frac{1}{\det(I - C'_r)},$$

an identity which will be used in Chapter 26.

36. *The generating function for words by exceedance number.* For each word $w = x_1x_2 \cdots x_n$ from $[0, r]^*$ let $\bar{w} = \bar{x}_1 \bar{x}_2 \cdots \bar{x}_n$ be its non-decreasing rearrangement and let $\text{exc } w := \sum_{1 \leq i \leq n-1} \chi(x_i > \bar{x}_i)$ denote its *number of exceedances*. If $\bar{w} = 0^{n_0} 1^{n_1} \cdots r^{n_r}$ and if u_0, u_1, \dots, u_r are $(r + 1)$ commuting variables, let $\mathbf{u}(w) := u_0^{n_0} u_1^{n_1} \cdots u_r^{n_r}$. Also, let

$$D := \det \begin{pmatrix} 1 - u_0 & -u_1 & \cdots & -u_{r-1} & -u_r \\ -tu_0 & 1 - u_1 & \cdots & -u_{r-1} & -u_r \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ -tu_0 & -tu_1 & \cdots & 1 - u_{r-1} & -u_r \\ -tu_0 & -tu_1 & \cdots & -tu_{r-1} & 1 - u_r \end{pmatrix}.$$

(a) The generating function for all words from $[0, r]^*$ by “exc” is given by

$$\sum_{w \in [0, r]^*} t^{\text{exc } w} \mathbf{u}(w) = 1/D,$$

(b) We also have: $D = 1 - e_1(\mathbf{u}) + (1 - t)e_2(\mathbf{u}) - \cdots + (-1)^r (1 - t)^{r-1} e_r(\mathbf{u})$, where the $e_i(\mathbf{u})$'s are the elementary symmetric function in the u_i 's, and the further formulas:

$$\sum_{w \in [0, r]^*} t^{\text{exc } w} \mathbf{u}(w) = \frac{1 - t}{(1 - u_0(1 - t))(1 - u_1(1 - t)) \cdots (1 - u_r(1 - t)) - t},$$

and by the change of variables $u_i \leftarrow u_i(1 - t)$ ($i = 0, 1, \dots, r$)

$$\sum_{w \in [0, r]^*} t^{\text{exc } w} \frac{\mathbf{u}(w)}{(1-t)^{1+|w|}} = \sum_{s \geq 0} \frac{t^s}{(1-u_0)^{s+1} (1-u_1)^{s+1} \dots (1-u_r)^{s+1}},$$

(c) Identity (7.8), the last formula in Chapter 7, provides the generating function for $[0, r]^*$ by the pair (des, maj) (descent number and major index):

$$\sum_{w \in [0, r]^*} t^{\text{des } w} q^{\text{maj } w} \frac{\mathbf{u}(w)}{(t; q)_{1+|w|}} = \sum_{s \geq 0} \frac{t^s}{(u_0; q)_{s+1} (u_1; q)_{s+1} \dots (u_r; q)_{s+1}}.$$

By letting $q = 1$ in this latter identity we get

$$\sum_{w \in [0, r]^*} t^{\text{exc } w} \frac{\mathbf{u}(w)}{(1-t)^{1+|w|}} = \sum_{w \in [0, r]^*} t^{\text{des } w} \frac{\mathbf{u}(w)}{(1-t)^{1+|w|}}.$$

For each non-decreasing word $\mathbf{n} = 0^{n_0} 1^{n_1} \dots r^{n_r}$ let $R(\mathbf{n})$ be the set of all rearrangements of \mathbf{n} . It follows from the last identity that “exc” and “des” are equidistributed on $R(\mathbf{n})$. This is essentially the spirit of the analytical proof originally obtained by MacMahon. The construction of a bijection of each $R(\mathbf{n})$ onto itself, which shows the equidistribution property can be found in [Lo02, chap. 10].

37. *Minimality of the quantum algebra commutation rules.* Rules (R1) and (R2), valid for the algebra \mathcal{A}^{GLZ} , also hold in the two algebras \mathcal{A}^{MM} and \mathcal{A}^{CF} . It would have been sufficient to establish the “Master Theorem” identity only for the latter algebra. This arises the problem to see whether an associative algebra $\overline{\mathcal{A}}$, which is a quotient of the free biword large algebra \mathcal{A} by a set of commutation rules and in which the “Master Theorem” identity holds, also admits the commutation rules (R1) and (R2). And it does.

To prove it take the original alphabet $\{1, 2\}$ and expand the product $\text{Ferm} \times \text{Bos}^{(3)}$ in \mathcal{A} , where $\text{Bos}^{(3)}$ is the formal sum of all the biwords in Bos of length 0, 1, 2, 3. There are 15 terms in $\text{Bos}^{(3)}$ and 17 in the product $\text{Ferm} \times \text{Bos}^{(3)}$ after cancellation. See that Rules (R1) and (R2) necessarily hold if the latter product equals 1, as wanted.

38. *Generating function for the horizontal derangements.* As mentioned in Chap. 26.2, each word w is a horizontal derangement, if it has no equal letters in succession; accordingly, if $(\text{DEC} \setminus \text{DES})(w) = \emptyset$, $(\text{INC} \setminus \text{RISE} \setminus \text{REC})(w) = \emptyset$, $(\text{INC} \setminus \text{RISE} \cap \text{REC})(w) = \emptyset$. Its weight $\psi(w)$ is then equal to

$$\psi(w) = \prod_{a \in \text{DES}(w)} X_a \times \prod_{b \in (\text{RISE} \setminus \text{REC})(w)} Y_b \times \prod_{c \in (\text{RISE} \cap \text{REC})(w)} Y'_c,$$

and the generating function for the set $[0, r]_h^*$ of all horizontal derangements is derived from (26.7) by letting all the Z_i 's, T_i 's and T_i' 's be equal to 0. Thus,

$$\sum_{w \in [0, r]_h^*} \psi(w) = \frac{\prod_{0 \leq j \leq r} (1 + Y_j')}{D},$$

where

$$D = \begin{vmatrix} 1 & -X_1 & -X_2 & \cdots & -X_{r-1} & -X_r \\ -Y_0 & 1 & -X_2 & \cdots & -X_{r-1} & -X_r \\ -Y_0 & -Y_1 & 1 & \cdots & -X_{r-1} & -X_r \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ -Y_0 & -Y_1 & -Y_2 & \cdots & 1 & -X_r \\ -Y_0 & -Y_1 & -Y_2 & \cdots & -Y_{r-1} & 1 \end{vmatrix}.$$

This generating function can also be expressed as:

$$\sum_{w \in [0, r]_h^*} \psi(w) = \frac{\prod_{0 \leq j \leq r} \frac{1 + Y_j'}{1 + X_j}}{1 - \sum_{0 \leq l \leq r} \prod_{0 \leq j \leq l-1} \frac{1 + Y_j}{1 + X_j} \frac{X_l}{1 + X_l}}.$$

39. *Non-determinantal expressions for the Decrease Value Theorem Identity.* We also have:

$$\sum_{w \in [0, r]^*} \psi(w) = \frac{\prod_{0 \leq j \leq r} \frac{1 - Z_j}{1 - Z_j + X_j} / \prod_{0 \leq j \leq r} \frac{1 - T_j'}{1 - T_j' + Y_j'}}{1 - \sum_{0 \leq l \leq r} \frac{\prod_{0 \leq j \leq l-1} \frac{1 - Z_j}{1 - Z_j + X_j}}{\prod_{0 \leq j \leq l-1} \frac{1 - T_j}{1 - T_j + Y_j}} \frac{X_l}{1 - Z_l + X_l}}.$$

Another non-determinantal expression can be obtained by replacing the summation $\sum_{0 \leq l \leq r}$ by $\sum_{1 \leq l \leq r}$.

40. *A specialization of the Decrease Value Theorem.* Let γ be the homomorphism defined by the following substitutions of variables:

$$\gamma := \{X_j \leftarrow s Y_{j-1}, \quad Z_j \leftarrow s Y_{j-1}, \quad T_j \leftarrow Y_j, \quad T_j' \leftarrow Y_j'\}.$$

For each word $w = x_1 x_2 \cdots x_n \in [0, r]^*$ we have:

$$(*) \quad \gamma \psi(w) = \prod_{x_i \in \text{INC} \cap \text{REC}} Y_{x_i}' \times \prod_{x_i \in \text{DEC}} s Y_{x_i-1} \times \prod_{x_i \in \text{INC} \setminus \text{REC}} Y_{x_i}.$$

Using the identity of Exercice 39, second form, we get

$$(**) \sum_{w \in [0, r]^*} \gamma \psi(w) = \frac{\prod_{1 \leq j \leq r} (1 - sY_{j-1}) / \prod_{0 \leq j \leq r} (1 - Y'_j)}{1 - \sum_{1 \leq l \leq r} sY_{l-1} \prod_{1 \leq j \leq l-1} (1 - sY_{j-1}) / \prod_{0 \leq j \leq l-1} (1 - Y'_j)},$$

and finally

$$\sum_{w \in [0, r]^*} \gamma \psi(w) = \frac{(1-s) \prod_{0 \leq j \leq r-1} (1 - Y_j) \prod_{0 \leq j \leq r-1} (1 - sY_j)}{\prod_{0 \leq j \leq r} (1 - Y'_j) \left(\prod_{0 \leq j \leq r-1} (1 - Y_j) - s \prod_{0 \leq j \leq r-1} (1 - sY_j) \right)}.$$

41. *A further specialization of the Decrease Value Theorem.* For each word $w = x_1 x_2 \cdots x_n$ let $\text{inrec } w$ denote the *number* of letters of w , which are increase *and* record values. Make the substitution $Y'_j \leftarrow RY_j$, where R is a new variable and let $\psi_R := \gamma \psi |_{Y'_j \leftarrow RY_j}$, so that formula (*) in Exercise 40 becomes:

$$\psi_R(w) = R^{\text{inrec } w} s^{\text{dec } w} \prod_{x_i \in \text{DEC}} Y_{x_i-1} \times \prod_{x_i \in \text{INC}} Y_{x_i}.$$

Also, let

$$H(Y) := \prod_{i \geq 0} (1 - Y_i)^{-1}; \quad H_r(Y) := \prod_{0 \leq i \leq r-1} (1 - Y_i)^{-1} \quad (r \geq 0).$$

Using Exercise 40 we obtain:

$$(***) \sum_{w \in [0, r]^*} \psi_R(w) = \frac{(1-s)H_{r+1}(RY)}{H_r(sY) - sH_r(Y)}.$$

42. *Two ultimate specialisations of the Decrease Value Theorem.* In the last formula of Exercise 40 replace Y_j, Y'_j ($j \geq 0$) by u . Then,

$$\sum_{w \in [0, r]^*} s^{\text{dec } w} u^{|w|} = \frac{1-s}{(1-u)^{r+1}(1-us)^{-r} - s(1-u)},$$

where $|w|$ denotes the length of w .

Next, replace X_j ($j \geq 0$) by us and the other variables by u in the identity displayed in Exercise 39. We recover the *classical* generating function for words by number of descents:

$$\sum_{w \in [0, r]^*} s^{\text{des } w} u^{\lambda w} = \frac{1-s}{(1-u+us)^{r+1} - s}.$$

43. *A Krattenthaler telescoping technique.* For each word w let $\text{inrec } w$ denote the *number* of its letters, which are increase *and* record values, let $\text{dec } w$ be the number its decreases, and $|w|$ (resp. $\text{tot } w$) *the number* (resp. *the sum*) of its letters. In the identity displayed in Exercise 39 make the substitutions $X_j \leftarrow usq^j, Y_j \leftarrow uq^j, Z_j \leftarrow usq^j, T_j \leftarrow uq^j, Y'_j \leftarrow uRq^j,$

$T'_j \leftarrow uRq^j$. The weight $\psi(w)$ becomes $s^{\text{dec } w} R^{\text{inrec } w} u^{\lambda w} q^{\text{tot } w}$. This yields the identity

$$\sum_{w \in [0, r]^*} s^{\text{dec } w} R^{\text{inrec } w} u^{\lambda w} q^{\text{tot } w} = \frac{(us; q)_{r+1}}{(uR; q)_{r+1}} / \left(1 - \sum_{0 \leq l \leq r} \frac{(us; q)_l}{(u; q)_l} usq^l \right).$$

By using the following q -telescoping argument

$$\frac{(us; q)_l}{(u; q)_l} usq^l = \frac{sq}{1 - sq} \left(\frac{(us; q)_{l+1}}{(u; q)_l} - \frac{(us; q)_l}{(u; q)_{l-1}} \right) \quad (1 \leq l \leq r),$$

one gets:

$$\sum_{w \in [0, r]^*} s^{\text{dec } w} R^{\text{inrec } w} u^{|w|} q^{\text{tot } w} = \frac{1}{(uR; q)_{r+1}} \frac{(1 - sq)(u; q)_r (usq; q)_r}{((u; q)_r - sq(usq; q)_r)}.$$

Answers to the exercises

1. Clearly, $\exp(xu) \exp(u) = \exp((1+x)u)$ is the exponential generating function for the polynomials $H_n(x) = (1+x)^n = \sum_{0 \leq j \leq n} \binom{n}{j} x^j$. In the case of q -series the calculation is to be made explicitly:

$$e_q(xu) e_q(u) = \sum_{n \geq 0} \sum_{0 \leq j \leq n} \frac{(xu)^j}{(q; q)_j} \frac{u^{n-j}}{(q; q)_{n-j}} = \sum_{n \geq 0} \frac{u^n}{(q; q)_n} \sum_{0 \leq j \leq n} \binom{n}{j} x^j$$

$$= \sum_{n \geq 0} \frac{u^n}{(q; q)_n} H_n(x, q). \text{ With } x = -1 \text{ we get } \sum_{n \geq 0} \frac{u^n}{(q; q)_n} H_n(-1, q) =$$

$$\frac{1}{(-u; q)_\infty (u; q)_\infty} = \frac{1}{(-u^2; q^2)_\infty} = \sum_{n \geq 0} (-1)^n \frac{u^{2n}}{(q^2; q^2)_n}, \text{ hence}$$

$$H_{2n+1}(-1, q) = 0 \text{ and } H_{2n}(-1, q) = (q; q)_{2n} / (q^2; q^2)_n = (q; q^2)_n. \text{ With}$$

$$x = q^{1/2} \text{ it turns into } \sum_{n \geq 0} \frac{u^n}{(q; q)_n} H_n(q^{1/2}, q) = \frac{1}{(uq^{1/2}; q)_\infty (u; q)_\infty} =$$

$$\frac{1}{(u; q^{1/2})_\infty} = \sum_{n \geq 0} \frac{u^n}{(q^{1/2}; q^{1/2})_n}, \text{ hence}$$

$$H_n(q^{1/2}; q) = \frac{(q; q)_n}{(q^{1/2}; q^{1/2})_n} = \frac{(q; q)_n (-q^{1/2}; q^{1/2})_n}{(q^{1/2}; q^{1/2})_n (-q^{1/2}; q^{1/2})_n} =$$

$$(q^{-1/2}; q^{1/2})_n. \text{ Start with } f(x, u) = \sum_{n \geq 0} \frac{u^n}{(q; q)_n} H_n(x, q). \text{ Then}$$

$$f(x, u) - f(x, qu) = x \sum_{n \geq 0} \frac{u^n}{(q; q)_n} H_{n+1}(x, q). \text{ On the other hand,}$$

$$f(x, u) - f(x, qu) = \frac{1}{(u; q)_\infty} \frac{1}{(xu; q)_\infty} (1 - (1-u)(1-xu)) =$$

$$f(x, u)u(1+x-xu). \text{ Therefore } \sum_{n \geq 0} \frac{u^n}{(q; q)_n} H_{n+1}(x, q) = (1+x)$$

$$\times \sum_{n \geq 0} \frac{u^n}{(q; q)_n} H_n(x, q) - \sum_{n \geq 0} \frac{u^{n+1}}{(q; q)_n} x H_n(x, q). \text{ Hence the recurrence}$$

for $n \geq 0$ with $H_{-1}(x, q) = 0$.
2. No further comment.
3. By definition of y we have $y_i = i - \sum_{1 \leq j \leq i} \chi(\sigma(i) > \sigma(j) > \sigma(i+1))$ or $\sum_{1 \leq j \leq i-1} \chi(\sigma(i+1) > \sigma(j) > \sigma(i))$, depending on whether $\sigma(i) > \sigma(i+1)$ or $\sigma(i) < \sigma(i+1)$. The sequence y is then subexcedent. Now, given a subexcedent sequence y we can reconstruct the permutation σ in a unique way, starting with $\sigma(n)$ from $y_n = n - \sum_{1 \leq j \leq n} \chi(\sigma(n) > \sigma(j))$.

The mapping $y \mapsto \sigma \mapsto x$ is then bijective. Finally,

$$\text{tot } y = \sum_{1 \leq i \leq n} (x_i - x_{i+1}) + i\chi(\sigma(i) > \sigma(i+1)) = \sum_{1 \leq i \leq n} i\chi(\sigma(i) > \sigma(i+1)) = \text{maj } \sigma, \text{ since } x_0 = x_{n+1} = 0.$$

4. We have $\text{inv-coding}(\sigma) = (0, 0, 0, 2, 1, 1, 5, 3, 8)$ and $\text{inv}\sigma = 20$; furthermore, $\text{maj-coding}(\sigma) = (0, 1, 1, 3, 4, 1, 2, 2, 3)$, $\text{maj}\sigma = 17$. Finally, $\text{inv-coding}(687293154) = x$ and $\text{maj-coding}(938164275) = x$.

5. (a) $z_i \leq \#\{1 \leq j < i\} = i$; (b) $z = 001102238$; (c) $\sigma = 756312984$.
 (d) Distinguish two cases. If $\sigma(i) > \sigma(i+1)$, the components of this new maj-coding satisfy the relations:

$$\begin{aligned} x_i - x_{i+1} + i\chi(\sigma(i) > \sigma(i+1)) &= \{j < i \mid \sigma(j) > \sigma(i)\} - \{j < i+1 \mid \sigma(j) > \sigma(i+1)\} + i \\ &= \{j < i \mid \sigma(j) > \sigma(i)\} + \{j < i+1 \mid \sigma(j) \leq \sigma(i+1)\} \\ &= \{j < i \mid \sigma(j) > \sigma(i)\} + \{j < i \mid \sigma(j) \leq \sigma(i+1)\} \\ &= \#\{1 \leq j < i \mid \sigma(j) \in \llbracket \sigma(i), \sigma(i+1) \rrbracket\} = z_i. \end{aligned}$$

If $\sigma(i) < \sigma(i+1)$, then

$$\begin{aligned} x_i - x_{i+1} + i\chi(\sigma(i) > \sigma(i+1)) &= \{j < i \mid \sigma(j) > \sigma(i)\} - \{j < i+1 \mid \sigma(j) > \sigma(i+1)\} \\ &= \{j < i \mid \sigma(j) > \sigma(i)\} - \{j < i \mid \sigma(j) > \sigma(i+1)\} \\ &= \#\{1 \leq j < i \mid \sigma(j) \in \llbracket \sigma(i), \sigma(i+1) \rrbracket\} = z_i. \end{aligned}$$

6. The subsequence y is subexcedent and $y_1 + y_2 + \dots + y_n = \text{maj}\sigma$. However the transformation $\sigma \mapsto y$ is not bijective; for instance, the two permutations 213 and 312 have the same image 010.

7. The vectors of a set $\{v_1, v_2, \dots, v_n\}$ are linearly independent, first if v_1 is not zero (there are $(q^{N+n} - 1)$ possible choices), then if v_2 is not proportional to v_1 (there are $(q^{N+n} - q)$ possible choices), more generally, if for $i = 2, \dots, n$ the vector v_i does not belong to the subspace of dimension $(i - 1)$ generated by v_1, v_2, \dots, v_{i-1} (there are $(q^{N+n} - q^{i-1})$ possible choices). The number of such sets $\{v_1, v_2, \dots, v_n\}$ is then equal to $(q^{N+n} - 1)(q^{N+n} - q) \dots (q^{N+n} - q^{n-1})$. The second enumeration is derived by the same reasoning. Finally, the number of sets $\{v_1, v_2, \dots, v_n\}$ having only linearly independent vectors is equal to the number of subspaces of dimension n , multiplied by the number of such sets that span a given space of dimension n .

8. The two formulas are banal for $N = 1$. For proving (3.9) use (3.6) with the substitutions $n \leftarrow N + n$, $i \leftarrow n$, so that $(1 - uq^N) \sum_{n \geq 0} \begin{bmatrix} N+n \\ n \end{bmatrix} u^n = \sum_{n \geq 0} (\begin{bmatrix} N+n \\ n \end{bmatrix} - q^N \begin{bmatrix} N+n-1 \\ n-1 \end{bmatrix}) u^n = \sum_{n \geq 0} \begin{bmatrix} N+n-1 \\ n \end{bmatrix} u^n = (u; q)_N^{-1}$ (by induction on N). Hence $\sum_{n \geq 0} \begin{bmatrix} N+n \\ n \end{bmatrix} u^n = (u; q)_N^{-1} (1 - uq^N)^{-1} = (u; q)_{N+1}^{-1}$. For (3.10) make use of (3.5) with the substitutions $n \leftarrow N + 1$, $i \leftarrow n$, so that $(-u; q)_{N+1} = (\sum_{0 \leq n \leq N} q^{n(n-1)/2} \begin{bmatrix} N \\ n \end{bmatrix} u^n) \cdot (1 + uq^N) = 1 + \sum_{1 \leq n \leq N} (\begin{bmatrix} N \\ n \end{bmatrix} q^{n(n-1)/2} + \begin{bmatrix} N \\ n-1 \end{bmatrix} q^{(n-1)(n-2)/2+N}) u^n + q^{(N+1)N/2} u^{N+1} = 1 + \sum_{1 \leq n \leq N} (\begin{bmatrix} N \\ n \end{bmatrix} + q^{(N+1)-n} \begin{bmatrix} N \\ n-1 \end{bmatrix}) q^{n(n-1)/2} u^n + q^{(N+1)N/2} u^{N+1} = \sum_{0 \leq n \leq N+1} \begin{bmatrix} N+1 \\ n \end{bmatrix} q^{n(n-1)/2} u^n$.

9. By induction on $N + n$. Let $\text{ND}(N, n; q) := \sum_{b \in \text{ND}(N, n)} q^{\text{tot } b}$. Then $\text{ND}(N, n; q) = \sum_{b_1=0} q^{\text{tot } b} + \sum_{b_1 \geq 1} q^{\text{tot } b}$. Let $a_i := b_i - 1$ ($i = 1, \dots, N$) in the second sum. With the convention that $\text{ND}(N, n; q) = 1$ when $N = 0$ we get:

$$\begin{aligned} \text{ND}(N, n; q) &= \sum_{0 \leq b_2 \leq \dots \leq b_N \leq n} q^{\|b\|} + \sum_{0 \leq a_1 \leq \dots \leq a_N \leq n-1} q^{N+\|a\|} \\ &= \text{ND}(N-1, n; q) + q^N \text{ND}(N, n-1; q) \\ &= \begin{bmatrix} N+n-1 \\ n \end{bmatrix} + q^N \begin{bmatrix} N+n-1 \\ n-1 \end{bmatrix} = \begin{bmatrix} N+n \\ n \end{bmatrix}. \end{aligned}$$

10. Proceed by induction on $N + n$. Let $\text{BW}(N, n; q) := \sum_{x \in \text{BW}(N, n)} q^{\text{inv } x}$. Then $\text{BW}(N, n; q) = \sum_{x_1=0} q^{\text{inv } x} + \sum_{x_1=1} q^{\text{inv } x}$. Let $y_i := x_{i+1}$ ($i = 1, \dots, N + n - 1$) in each of the sums and let y be the word $y_1 y_2 \dots y_{N+n-1}$. In the first (resp. the second) sum we have $\text{inv } x = \text{inv } y$ and $y \in \text{BW}(N, n-1)$ (resp. $\text{inv } x = n + \text{inv } y$ and $y \in \text{BW}(N-1, n)$). With the convention that $\text{BW}(N, n; q) = 1$ when $N = 0$,

$$\begin{aligned} \text{BW}(N, n; q) &= \sum_{y \in \text{BW}(N, n-1)} q^{\text{inv } y} + \sum_{y \in \text{BW}(N-1, n)} q^{n+\text{inv } y} \\ &= \text{BW}(N, n-1; q) + q^n \text{BW}(N-1, n; q) \\ &= \begin{bmatrix} N+n-1 \\ n-1 \end{bmatrix} + q^n \begin{bmatrix} N+n-1 \\ n \end{bmatrix} = \begin{bmatrix} N+n \\ n \end{bmatrix}. \end{aligned}$$

11. (a) As shown in Fig. A1, for each partition π in at most n parts, all at most equal to N denote the size of the n -th part of π (the smallest one being possibly 0) by j . Withdraw j from all the parts of π : there remains a partition in at most $(n - 1)$ parts, all at most equal to $(N - j)$. The latter partitions have a generating function equal to $\begin{bmatrix} N-j+n-1 \\ N-j \end{bmatrix}$.




Fig. A1




Fig. A2

- (b) Report to Fig. A2. For each partition π in at most $(n + m)$ parts, all at most equal to N , denote the size of the smallest $(n + 1)$ -st part by j . To such a partition π associate the partition π' in at most n parts, all at most equal to j and a partition π'' in at most $(m - 1)$ parts, all at most equal to $(N - j)$.

The two q -binomial coefficients occurring on the left-hand side of the identity are precisely the generating functions for the partitions π' and π'' , respectively. The identity is then a consequence of the fact that the weight of π is equal to the sum of the weights of π' and of π'' , plus the size of the rectangle of dimension $j \times m$.

12. (a) By induction.
 (b) If n is inserted onto the left of a permutation $\sigma' = \sigma'(1) \dots \sigma'(n - 1)$ or between two letters $\sigma(i), \sigma(i + 1)$ such that $\sigma(i) < \sigma(i + 1)$, the number of descents increases by one. In the other case the number of descents remains alike. To obtain the set $S_{n,k}$ of all permutations of order n having k descents it suffices
 either to take each permutation belonging to $S_{n-1,k-1}$ and insert n into the $(n - 1) - (k - 1)$ positions that create a new descent; we then obtain $(n - k)A_{n-1,k-1}$ permutations of order n ;

or to take each permutation belonging to $S_{n-1,k}$ and insert n into the $(k+1)$ positions that create no descent; $(k+1)A_{n-1,k}$ permutations of order n having k descents are derived in that way.

13. (a) The right-hand side of the identity to be proved is equal to:

$$(1-t)^n \sum_{n \geq 0} t^j (j+1)^{n-1} (1+(n-1)t-nt+(1-t)j) = (1-t)^n \sum_{n \geq 0} t^j (j+1)^{n-1} (1-t)(j+1) = (1-t)^{n+1} \sum_{n \geq 0} t^j (j+1)^n = A_n(t).$$
- (b) First $A_0(t) = (1-t)/(1-t) = 1$ and the constant coefficient of each series $A_n(t)$ is $A_{n,0} = 1$. Consider the coefficient of t^k ($k \geq 1$) on each of sides of (E13.1). Then $A_{n,k} = A_{n-1,k} + (n-1)A_{n-1,k-1} + kA_{n-1,k} - (k-1)A_{n-1,k-1}$, which is the desired recurrence relation. Finally, the recurrence implies that $A_{n,k} = 0$ for $k \geq n \geq 1$.
- (c) It suffices to calculate the coefficient of t^k in $(1-t)^{n+1} \sum_{n \geq 0} t^n (j+1)^n$.
- (d) $A_n(t)(1-t)^{-(n+1)} = \sum_{k \geq 0} \binom{n+k}{n} t^k \sum_{0 \leq l \leq n-1} A_{n,l} t^l = \sum_{j \geq 0} t^j \sum_{0 \leq l \leq \min(j, n-1)} A_{n,l} \binom{n+j-l}{n} = \sum_{j \geq 0} t^j \sum_{0 \leq l \leq n-1} A_{n, n-1-l} \binom{(j+1)+(n-l-1)}{n} = \sum_{j \geq 0} t^j (j+1)^n$, using $\binom{n+j-l}{n} = 0$ when $j \leq n-2$ and $l = j+1, \dots, n-1$ and the Worpitzky for the last step.
- (e) From (E13.1) $\sum_{n \geq 0} A_n(t)(1-t)^{-(n+1)} u^n / n! = \sum_{n \geq 0} u^n / n! \sum_{j \geq 0} t^j (j+1)^n = \sum_{j \geq 0} t^j \sum_{n \geq 0} (u(j+1))^n / n! = e^u \sum_{j \geq 0} (te^u)^j = e^u / (1-te^u)$. With $u := v(1-t)$ we get $\sum_{n \geq 0} A_n(t)v^n / n! = (1-t) \exp(v(1-t)) / (1-t \exp(v(1-t)))$ and the desired generating function.

14. Let $\sigma = \sigma(1) \dots \sigma(n)$ be a permutation. If $\sigma(i) < i$, say that i is an *accedence-letter* of σ . Let $A(\sigma)$ (resp. $E(\sigma)$) denote the set of all *accedence-letters* (resp. *excedence-letters*) of σ . First, verify that for each $j = 1, \dots, n$

$$\#\{i \in A(\sigma) : j \in]\sigma(i), i]\} = \#\{i \in E(\sigma) : j \in]i, \sigma(i)]\}.$$

Next, verify that

$$\begin{aligned} \#\{(l, k) : l, k \in E(\sigma), k \in [l, \sigma(l)]\} \\ = \#\{(l, k) : l, k \in E(\sigma), \sigma(k) \in]l, \sigma(l)]\}; \end{aligned}$$

so that

$$\begin{aligned} \#\{(l, k) : l \in A(\sigma), k \in E(\sigma), \sigma(k) \in]\sigma(l), l]\} \\ = \#\{(l, k) : l, k \in E(\sigma), k \in [l, \sigma(l)]\}; \end{aligned}$$

This is sufficient to show that

$$\begin{aligned} & \#\{(l, k) : l < k, \sigma(k) < \sigma(l) \leq k\} + \#\{(l, k) : l < k, \sigma(l) \leq k < \sigma(k)\} \\ & = \#\{(l, k) : l < k, \sigma(k) < \sigma(l) \leq l\} + \#\{(l, k) : l \leq k, k < \sigma(k)\}. \end{aligned}$$

Merge the third term into the first and the fourth into the second:

$$\begin{aligned} & \#\{(l, k) : l < \sigma(l), \sigma(k) < k, \sigma(k) < \sigma(l) \leq k\} \\ & = \#\{(l, k) : l < \sigma(k), k < \sigma(l), l \leq k < \sigma(l)\}. \end{aligned}$$

15. (a) and (b) are easy; (c) is a rewriting of the maj-coding discussed in § 2.2. The difficult part is (d) and requires a deeper study. A numerical example is discussed in details inviting the reader to reconstruct the proof himself. For a complete formal proof see [Ha90b]. Finally, (e) is merely making the appropriate composition products.

16. (a) The table of the six statistics for $\text{BW}(2, 3)$:

	des	maj	rise	rmaj	DES	MAJ
11000	1	2	0	0	1	2
10100	2	4	1	2	3	6
10010	2	5	1	3	3	8
01100	1	3	1	1	2	4
01010	2	6	2	4	4	10
00110	1	4	1	2	2	6
10001	1	1	1	4	2	5
01001	1	2	2	5	3	7
00101	1	3	2	6	3	9
00011	0	0	1	3	1	3

- (b) The bijection $x \mapsto y$ of $\text{BW}(N, n)$ onto $\text{BW}(n, N)$ is defined as follows. For $1 \leq i \leq N + n$ let $y_i = 1 - x_i$. Then $\text{rmaj } x = \text{maj } y$ and

$$\sum_{x \in \text{BW}(N, n)} q^{\text{rmaj } x} = \sum_{y \in \text{BW}(n, N)} q^{\text{maj } y} = \begin{bmatrix} N + n \\ n \end{bmatrix}.$$

- (c) A bijection $x \mapsto z$ of $\text{BW}(N, n)$ onto itself is constructed as follows. First, for each $x \in \text{BW}(N, n)$ apply the bijection of (b) to get $y \in \text{BW}(n, N)$. Then factorize y in a unique manner under the form $y = v_1 10 v_2 10 \cdots v_{k-1} 10 v_k$, where v_i is a nondecreasing binary word of the form $0^a 1^b$; moreover, the length $l(v_i)$ of v_i satisfies $l(v_i) \geq 1$ for $2 \leq i \leq k - 1$ and $l(v_1) \geq 0, l(v_k) \geq 0$. To each nondecreasing

word $v_i = 0^a 1^b$ associate the nondecreasing word $u_i = 0^b 1^a$. Then, let $z := u_1 10 u_2 10 \cdots u_{k-1} 10 u_k$.

For example with $x = 10110001011 \in \text{BW}(6, 5)$ we get $y = 01001110100 \in \text{BW}(5, 6)$. The factorization of $y = 0(10)011(10)(10)0$ yields $z = 1(10)001(10)(10)1 = 11000110101 \in \text{BW}(6, 5)$. We verify that $\text{rmaj } x = \text{maj } y = \text{maj } z$ and $\text{rise } x = \text{des } y = \text{des } z$.

(d) Look for the difference between rmaj and maj .

$$\begin{aligned}
 \text{(e)} \quad \sum_{x \in \text{BW}(n, N)} q^{\text{MAJ } x} &= \sum_{x \in \text{BW}(n, N), x_{n+N}=0} q^{\text{MAJ } x} + \sum_{x \in \text{BW}(n, N), x_{n+N}=1} q^{\text{MAJ } x} \\
 &= \sum_{x \in \text{BW}(n-1, N)} q^{2 \text{maj}' x + N} + \sum_{x \in \text{BW}(n, N-1)} q^{2 \text{maj } x + n} \\
 &= q^N \begin{bmatrix} n + N - 1 \\ n - 1, N \end{bmatrix}_{q^2} + q^n \begin{bmatrix} n + N - 1 \\ n, N - 1 \end{bmatrix}_{q^2} \\
 &= q^N \begin{bmatrix} n + N \\ n, N \end{bmatrix}_{q^2} \frac{1 - q^{2n}}{1 - q^{2N+2n}} + q^n \begin{bmatrix} n + N \\ n, N \end{bmatrix}_{q^2} \frac{1 - q^{2N}}{1 - q^{2N+2n}} \\
 &= \begin{bmatrix} n + N \\ n, N \end{bmatrix}_{q^2} \frac{q^N(1 - q^{2n}) + q^n(1 - q^{2N})}{1 - q^{2N+2n}} \\
 &= \begin{bmatrix} n + N \\ n, N \end{bmatrix}_{q^2} \frac{q^n + q^N}{1 + q^{n+N}}.
 \end{aligned}$$

17. (a) Let w be a word of $R(\mathbf{m})$ and let $1 \leq x < y = x + 1 \leq r$. Replace all the factors yx of w by a special letter “ \sim ”. In the new word the *maximal* factors containing the two letters x and y are of the form $x^a y^b$ ($a \geq 0, b \geq 0$). Change all those factors into $x^b y^a$ and replace every “ \sim ” by yx to obtain a word w' of the second set. For instance, with $w = 122322233243213 \in R(2, 7, 5, 1)$, $x = 2$ and $y = 3$ we successively get:

$$\begin{aligned}
 w &= 122322233243213 \mapsto 122 \sim 223 \sim 4 \sim 13 \\
 &\mapsto 133 \sim 233 \sim 4 \sim 12 \mapsto 133322333243212 = w' \in R(2, 5, 7, 1).
 \end{aligned}$$

That transformation is bijective and $\text{maj } w = \text{maj } w'$ holds.

- (b) $Z(w) = \text{maj}(2121122) + \text{maj}(1131) + \text{maj}(41114) + \text{maj}(22322) + \text{maj}(242242) + \text{maj}(434) = 4 + 3 + 1 + 3 + 7 + 1 = 19$.
- (c) The multiplicities of $\text{gcyc}_x(w)$ and $\text{lcyc}_x(w)$ are respectively:
 $\mathbf{m}^x = (m_{x+1}, m_{x+2}, \dots, m_r, m_1, m_2, \dots, m_{x-1}, m_x)$ and
 $\mathbf{m}_x = (m_1, m_2, \dots, m_{x-1}, m_{x+1}, m_{x+2}, \dots, m_r, m_x)$.
- (d) The global and local cycling are equal to $\text{gcyc}_2(w) = 4234313424$ and $\text{lcyc}_2(w) = 4214121434$. But $\text{maj } \text{gcyc}_2(w) = 18$ and $Z(\text{lcyc}_2(w)) = 16$. As $\text{maj } w = 21$ and $Z(w) = 19$, the two differences are equal to 3.

- (e) Consider the unique factorization $w = p_0 q_1 p_1 q_2 p_2 \dots q_s p_s$ of w , where the p_i 's are words all letters of which are at most equal to x and where the q_i 's are words whose letters are all greater than x with $|p_i| \geq 1$, $|q_i| \geq 1$ for all $i \geq 1$ and $|p_0| \geq 0$. The last letter of each factor p_i (resp. q_i) is less (resp. greater) than the first letter of the following factor q_{i+1} (resp. p_i). Say that there is a *rise* at the end of the factor p_i and a *descent* at the end of the factor q_i . In the word $\text{gcyc}_x(w)$ those rises become descents and the descents become rises, the other rises or descents remaining invariant. Therefore $\text{maj } w - \text{maj}(\text{gcyc}_x(w)) = |q_1| + |q_2| + \dots + |q_s| = m_{x+1} + m_{x+2} + \dots + m_r$.
- (f) Let $a = 1, 2, \dots, r$ be a letter occurring in w and denote the image of a under the local cycling $u := \text{lcyc}_x(w)$ by a' . If $i < j$ and $i \neq x$, the subwords w_{ij} and $u_{i'j'}$ are identical up to the reduction. On the other hand, for $x < j$, item (e) implies $\text{maj } w_{xj} - \text{maj}(\text{gcyc}_x(w))_{x'j'} = m_j$. As the two cyclings are identical up to the reduction for the words with two letters, it follows that $Z(w) - Z(u) = \sum_{x < j} (\text{maj}(w_{xj}) - \text{maj}(u_{x'j'})) = m_{x+1} + m_{x+2} + \dots + m_r$.
- (g) The desired bijection Φ is defined, for each word $w \in R(\mathbf{m})$ and each letter x , by the composition product:

$$\Phi(wx) := (\text{lcyc}_x^{-1} \circ \Phi \circ \theta_{\mathbf{m}^x, \mathbf{m}_x} \circ \text{gcyc}_x(w))x .$$

It is readily verified that $\Phi(wx)$ is a rearrangement of wx ; furthermore, $\text{maj } w = Z(\Phi(w))$ holds.

18. (a) Let $b := \sum_{r \geq 0} b_r t^r$. Then $b_0 = a_0$ and $b_r = a_r - a_{r-1}$ for $r \geq 1$. But $a - (b_0 + b_1 + \dots + b_r) = a - (a_0 + a_1 - a_0 + \dots + a_r - a_{r-1}) = a - a_r$ and by assumption the order of that difference tends to infinity with r .
- (b) The right-hand side of (7.8) is of the form $\sum_s b_s t^s$, where $b_s = 1/(\mathbf{u}; q)_{s+1}$. But $\lim_s b_s = 1/(\mathbf{u}; q)_\infty$. Then let $b(t) := (1-t) \cdot \sum_s b_s t^s$. It follows that $b(1) = 1/(\mathbf{u}; q)_\infty$. Multiply the left-hand side of (7.9) by $(1-t)$ and put $t = 1$ afterwards. It is found that $\sum_{\mathbf{m}} A_{\mathbf{m}}(t = 1, q) \mathbf{u}^{\mathbf{m}} / (q; q)_m$. This yields (6.13).
- (c) This time (a) is to be applied twice in succession. The coefficient of t_1^r , that is, $\sum_s t_2^s / (u; q_1, q_2)_{r+1, s+1}$, tends to $\sum_s t_2^s / (u; q_1, q_2)_{\infty, s+1}$ (obvious definition) when r tends to infinity. Multiply the left-hand side of (13.7) by $(1-t_1)$ and put $t_1 = 1$; we get the expression $\sum_s t_2^s / (u; q_1, q_2)_{\infty, s+1}$. As $1/(u; q_1, q_2)_{\infty, s+1}$ tends to $1/(u; q_1, q_2)_{\infty, \infty}$, when s tends to infinity when the previous sum is multiplied by $(1-t_2)$ and when t_2 is replaced by 1, we get: $1/(u; q_1, q_2)_{\infty, \infty}$. By

multiplying the right-hand side by $(1 - t_1)$ and $(1 - t_2)$ successively and by making the substitutions $t_1 = 1$, then $t_2 = 1$, we obtain: $\sum_n A_n(q_1, q_2) u^n / ((q_1; q_1)_n (q_2; q_2)_n)$.

19. (a) If $\sigma(i) = j$, then $\mathbf{i}\sigma(j) = i$ and $\mathbf{c}\mathbf{i}\sigma(j) = (n + 1 - i)$. Hence $\mathbf{i}\mathbf{c}\mathbf{i}\sigma(n + 1 - i) = j$ and $\mathbf{r}\sigma(n + 1 - i) = \sigma(i) = j$.
- (b) Write the n points $(1, \sigma(1)), (2, \sigma(2)), \dots, (n, \sigma(n))$ onto the square $\{1, \dots, n\} \times \{1, \dots, n\}$. The reflection of the square about the horizontal (resp. vertical, resp. major diagonal) axis transforms the graph of σ onto the graph of the permutation $\mathbf{c}\sigma$ (resp. $\mathbf{r}\sigma$, resp. $\mathbf{i}\sigma$). But those reflections generate the dihedral group D_4 and there is a one-to-one correspondence between those reflections and the operations \mathbf{i} , \mathbf{c} and \mathbf{r} .
- (c) The relations can be verified either by computation, or geometrically.
- (d) Lets $\mathbf{c}\sigma := \sigma'$ and $\mathbf{r}\mathbf{c}\sigma := \sigma''$. Then $j \in \text{Ligne}\sigma$ if and only if $j \in [n - 1]$ and $\sigma(j) > \sigma(j + 1)$; hence $\sigma'(j) < \sigma'(j + 1)$ and also $\sigma''(j'') < \sigma''(j'' + 1)$ with $j'' = n + 1 - j - 1$. Consequently, $j \in \text{Ligne}\sigma$ if and only if $j \in [n - 1]$ and, in an equivalent manner, $j \notin \text{Ligne}\mathbf{r}\sigma$ or $n - j \in \text{Ligne}\mathbf{r}\mathbf{c}\sigma$.
20. The verification of all those properties is easy.
21. (a) First $C_1(x) = x$; by induction $x C_{n-1}(x)$ is the generating polynomial for the permutations having n as a unit cycle; moreover, $(n - 1)C_{n-1}(x)$ is the generating polynomial for the other permutations, as the element n can be inserted in $(n - 1)$ manners into each permutation of order $(n - 1)$ to create a permutation of order n , where n is not a unit cycle.
- (b) Several proofs can be made. If we use $\# \text{Rmals}$, then $x C_{n-1}(x)$ is the generating polynomial for the permutations ending by 1. Moreover, $(n - 1)C_{n-1}(x)$ is the generating polynomial for the other permutations, because the insertion of 1 into each one of the $(n - 1)$ possible slots (excluding the rightmost one) of a permutation of $2, 3, \dots, n$, does not modify the right-to-left maximum letter set.
- (c) Consider the following modification of the Lehmer-coding: if $\sigma = \sigma(1) \dots \sigma(n)$, is a permutation, let $w := x_1 x_2 \dots x_n$ be the (modified) Lehmer coding defined by $x_i := \#\{j : i + 1 \leq j \leq n, \sigma(i) > \sigma(j)\}$. Then, $\text{inv}\sigma = \text{tot}w$, $\text{Rmip}\sigma = \{i : 1 \leq i \leq n, x_i = 0\}$ and $\text{Rmap}\sigma = \{i : 1 \leq i \leq n, x_i = n - i\}$. First, $C_1(x, y, q) = xy$;

then, by induction,

$$\begin{aligned}
 C_n(x, y, q) &= \sum_{w \in \text{SE}_n} x^{\sum_{1 \leq i \leq n} \chi(x_i = n-i)} y^{\sum_{1 \leq i \leq n} \chi(x_i = 0)} q^{\sum_{1 \leq i \leq n} x_i} \\
 &= \sum_{x_1=0}^{n-1} x^{\chi(x_1 = n-1)} y^{\chi(x_1 = 0)} q^{x_1} \\
 &\quad \times \sum_{w' \in \text{SE}_{n-1}} x^{\sum_{2 \leq i \leq n} \chi(x_i = n-i)} y^{\sum_{2 \leq i \leq n} \chi(x_i = 0)} q^{\sum_{2 \leq i \leq n} x_i} \\
 &= (y + q + q^2 + \cdots + q^{n-2} + q^{n-1}x)C_{n-1}(x, y, q).
 \end{aligned}$$

- (d) In the maj-coding $x_1 x_2 \dots x_n$ of a permutation σ we have $x_i = 0$ if and only if i is on the right of the subpermutation of σ reduced to the letters $1, 2, \dots, (i-1)$, i.e., if and only if $i \in \text{Rmals } \sigma$.
- (e) First, $D_1(y, q) = y$ and for $n \geq 2$

$$\begin{aligned}
 D_n(y, q) &= \sum_{w \in \text{SE}_n} y^{\sum_{1 \leq i \leq n} \chi(x_i = 0)} q^{\text{tot } w} \\
 &= \sum_{w' \in \text{SE}_{n-1}} y^{\sum_{1 \leq i \leq n-1} \chi(x_i = 0)} q^{\text{tot } w'} \sum_{x_n=0}^{n-1} y^{\chi(x_n = 0)} q^{x_n} \\
 &= D_{n-1}(y, q)(y + q + q^2 + \cdots + q^{n-2} + q^{n-1}).
 \end{aligned}$$

- (f) First, notice that, for $k \geq 1$,

$$y + q + q^2 + \cdots + q^k x = \frac{y(1-q) + q}{1-q} \left(1 - q^k \frac{1-x(1-q)}{y(1-q) + q} \right)$$

and $\frac{y(1-q) + q}{1-q} \left(1 - \frac{1-x(1-q)}{y(1-q) + q} \right) = x + y - 1$. Therefore,

$C_n(q, x, y) = \left(\frac{1-x(1-q)}{y(1-q) + q}; q \right)_n \left(\frac{y(1-q) + q}{1-q} \right)^n \frac{xy}{x+y-1}$, so that

$$\begin{aligned}
 &\sum_{n \geq 0} C_n(q, x, y) \frac{u^n}{(q; q)_n} \\
 &= 1 + \frac{xy}{x+y-1} \sum_{n \geq 1} \left(\frac{1-x(1-q)}{y(1-q) + q}; q \right)_n \left(u \frac{y(1-q) + q}{1-q} \right)^n \\
 &= 1 + \frac{xy}{x+y-1} \left(\left(\frac{u}{1-q} - ux; q \right)_\infty - 1 \right),
 \end{aligned}$$

by using the q -binomial theorem (Theorem 1.1).

22. (a) The polynomial $A_{n,k}$ is the generating function for the permutations of order n having k descents by (maj, # Rmil). When the letter n is inserted into a permutation of order $(n-1)$, the number of descents remains alike, if n is inserted into a descent and is increased by one otherwise; moreover, the statistic “# Rmil” is increased by one, if n is placed at the end of the permutation.
- (b) Form the two factorial generating functions:

$$A(t, q, y; u) := \sum_{n \geq 0} A_n(t, q, y) \frac{u^n}{(t; q)_n};$$

$$B(t, q, y; u) := \sum_{n \geq 0} A_n(t, q, y) \frac{u^n}{(t; q)_{n+1}}.$$

From (a) we deduce:

$$A(t, q, y; u) = 1 + yuB(t, q, y; u) + \frac{uq}{1-q}A(t, q, y; u) - \frac{uq}{1-q}A(tq, q, y; u).$$

As $A(t, q, y; u) = B(t, q, y; u) - tB(t, q, y; qu)$, we get

$$\begin{aligned} \left(1 - \frac{uq}{1-q} - uy\right)B(t, q, y; u) + \left(-t + \frac{utq}{1-q}\right)B(t, q, y; qu) \\ + \frac{uq}{1-q}B(tq, q, y; u) - \frac{utq^2}{1-q}B(tq, q, y; qu) = 1. \end{aligned}$$

Let $B(t, q, y; u) = \sum_{s \geq 0} t^s G_s(q, y; u)$ and work out a recurrence for the coefficients $G_s(q, y; u)$:

$$\begin{aligned} (1 - q - uq - yu + yqu) \sum_{s \geq 0} t^s G_s(q, y; u) \\ + (-1 + q + qu) \sum_{s \geq 1} t^s G_{s-1}(q, y; qu) \\ + uq \sum_{s \geq 0} t^s q^s G_s(q, y; u) - u \sum_{s \geq 1} t^s q^{s+1} G_{s-1}(q, y; qu) = 1 - q. \end{aligned}$$

By taking the coefficients of t^s on both sides we obtain:

$$G_0(q, y; u) = \frac{1}{1-yu};$$

$$G_s(q, y; u) = \frac{1 - u(q + q^2 + \cdots + q^s)}{1 - u(y + q + q^2 + \cdots + q^s)} G_{s-1}(q, y; qu) \quad (s \geq 1),$$

so that $G_s(q, y; u) = \prod_{0 \leq k \leq s} \frac{1 - u(q^{k+1} + \dots + q^s)}{1 - u(yq^k + q^{k+1} + \dots + q^s)}$. As

$$1 - u(yq^k + q^{k+1} + \dots + q^s) = \frac{1 - q + uq^{s+1}}{1 - q} \left(1 - q^k u \frac{y(1 - q) + q}{1 - q + uq^{s+1}} \right),$$

for $k = 0, 1, \dots, s$, we get

$$G_s(q, y; u) = \prod_{0 \leq k \leq s} \frac{1 - q^k u \frac{q}{1 - q + uq^{s+1}}}{1 - q^k u \frac{y(1 - q) + q}{1 - q + uq^{s+1}}} = \frac{\left(\frac{uq}{1 - q + uq^{s+1}}; q \right)_{s+1}}{\left(u \frac{y(1 - q) + q}{1 - q + uq^{s+1}}; q \right)_{s+1}}.$$

23. (a) As $\mathbf{c} = \mathbf{i} \mathbf{r} \mathbf{i}$, properties (i)—(iii) and relation (19.4) imply that if $\sigma \mapsto PQ$, then $\mathbf{i} \sigma \mapsto QP$, $\mathbf{r} \mathbf{i} \sigma \mapsto Q^T P^{JT}$ and $\mathbf{c} \sigma = \mathbf{i} \mathbf{r} \mathbf{i} \sigma \mapsto P^{JT} Q^T$.

(b) All the relations are proved by direct calculation.

(c) $P^J = \begin{array}{|c|c|c|} \hline 4 & 5 & \\ \hline 1 & 2 & 3 \\ \hline \end{array}$, $Q^J = \begin{array}{|c|c|c|} \hline 3 & 5 & \\ \hline 1 & 2 & 4 \\ \hline \end{array}$, $P^T = \begin{array}{|c|c|} \hline 5 & \\ \hline 2 & 4 \\ \hline 1 & 3 \\ \hline \end{array}$, $Q^T = \begin{array}{|c|c|} \hline 5 & \\ \hline 3 & 4 \\ \hline 1 & 2 \\ \hline \end{array}$,

$P^{JT} = \begin{array}{|c|c|} \hline 3 & \\ \hline 2 & 5 \\ \hline 1 & 4 \\ \hline \end{array}$, $Q^{JT} = \begin{array}{|c|c|} \hline 4 & \\ \hline 2 & 5 \\ \hline 1 & 3 \\ \hline \end{array}$. Notice that \mathbf{i} , \mathbf{c} , \mathbf{r} correspond to *geometric* transformations on permutations, but neither \mathbf{t} , nor \mathbf{j} .

Tableaux	Permutations	Tableaux	Permutations
PQ	$\sigma = 31425$	$P^T Q^T$	$\mathbf{t} \sigma = 25143$
PQ^J	$\mathbf{j} \sigma = 34152$	$P^T Q^{JT}$	$\mathbf{r} \sigma = 52413$
$P^J Q$	41523	$P^{JT} Q^T$	$\mathbf{c} \sigma = 35241$
$P^J Q^J$	14253	$P^{JT} Q^{JT}$	32514
QP	$\mathbf{i} \sigma = 24135$	$Q^T P^T$	31542
$Q^J P$	35124	$Q^{JT} P^T$	42531
QP^J	24513	$Q^T P^{JT}$	53142
$Q^J P^J$	13524	$Q^{JT} P^{JT}$	42153

24. For each Ferrers diagram λ with m boxes and each vector $\mathbf{m} = (m_1, m_2, \dots, m_r)$ of positive integers such that $m_1 + m_2 + \dots + m_r = m$ let $\mathcal{K}(\lambda, \mathbf{m})$ denote the set of Young tableaux containing m_1 1's, m_2 2's, \dots , m_r r 's. Let σ be a permutation of $1, 2, \dots, r$.

We construct a one-to-one correspondence between $\mathcal{K}(\lambda, \mathbf{m})$ and $\mathcal{K}(\lambda, \sigma \mathbf{m})$ as follows. Let $\mathbf{m} = (m_1, \dots, m_i, m_{i+1}, \dots, m_r)$ and $\mathbf{m}' = (m_1, \dots, m_{i+1}, m_i, \dots, m_r)$ differ only by a transposition of two adjacent terms and consider a tableau T in $\mathcal{K}(\lambda, \mathbf{m})$ in its planar representation. Write all the pairs $i, i+1$ in boldface whenever those two integers occur in the same column with $(i+1)$ just above i . The remaining i 's and $(i+1)$'s in T occur as horizontal blocks $i^a j^b$ ($a \geq 0, b \geq 0$). We define a bijection $T \mapsto T'$ of $\mathcal{K}(\lambda, \mathbf{m})$ onto $\mathcal{K}(\lambda, \mathbf{m}')$ by replacing each block $i^a j^b$ in T by $i^b j^a$ and rewriting the vertical pairs $i, i+1$ in roman type.

25. (a) The identity is true for $k = 1$. For $k \geq 2$ expand the determinant of $N_r(L, n)$ by the cofactors of the last row. By induction

$$\det N_r(L, n) = \det N_r(\{\ell_1, \dots, \ell_{k-1}\}, \ell_k) \begin{bmatrix} \ell_{k+1} - \ell_k + r \\ r \end{bmatrix} - \det N_r(\{\ell_1, \dots, \ell_{k-1}\}, n).$$

The first (resp. second) term on the right-hand side is the generating polynomial for the words $w = w_1 w_2 \dots w_n$ satisfying the conditions $(*)_{(L, n)}$, with the possible (resp. sole) exception of the subcondition $w_{\ell_k} < w_{\ell_k+1}$.

- (b) The construction of the bijections may be regarded as another version of the *MacMahon Verfahren*.
 (c) Similar to the proof of Theorem 18.2. Start with a pair (σ, s) having the properties (E22.2) and let $d = d_1 d_2 \dots d_n$ be the word whose letters d_i are defined by

$$d_i := \begin{cases} s_i - s_{i+1} - \chi(i \in \text{ligne } \sigma), & \text{if } 1 \leq i \leq n-1; \\ s_n, & \text{if } i = n. \end{cases}$$

Then $d_1 + \dots + d_n + \text{ides } \sigma = s_1$, so that it makes sense to put $d_0 := r - s_1 \geq 0$ and $d_0 + d_1 + \dots + d_n + \text{ides } \sigma = r$. On the other hand, $1 \cdot d_1 + 2 \cdot d_2 + \dots + n \cdot d_n + \text{imaj } \sigma = \text{tot } s$. The mapping $(\sigma, s) \mapsto (\sigma, (d_0, d_1, \dots, d_n))$ is a bijection of the pairs (σ, s) satisfying (E22.2) onto the pairs $(\sigma, (d_0, d_1, \dots, d_n))$ such that $\text{Ligne } \sigma = L$ and where the integers d_0, d_1, \dots, d_n are nonnegative and of sum $d_0 + d_1 + \dots + d_n = r - \text{ides } \sigma$. Hence,

$$\begin{aligned} \sum_{r \geq 0} t^r \det N_r(L, n) &= \sum_{r \geq 0} t^r \sum_{w \in W_r(L, n)} q^{\text{tot } w} = \sum_{r \geq 0} t^r \sum_{(\sigma, s)} q^{\text{tot } s} \end{aligned}$$

$$\begin{aligned}
 &= \sum_{r \geq 0} \sum_{\sigma} \sum_{d_0, \dots, d_n} t^{d_0 + \dots + d_n + \text{idess } \sigma} q^{1.d_1 + \dots + n.d_n + \text{imaj } \sigma} \\
 &= \sum_{\sigma} t^{\text{idess } \sigma} q^{\text{imaj } \sigma} \sum_{r \geq \text{idess } \sigma} \sum_{\substack{d_0 + \dots + d_n \\ = r - \text{idess } \sigma}} t^{d_0} (tq)^{d_1} \dots (tq^n)^{d_n} \\
 &= \sum_{\sigma} t^{\text{idess } \sigma} q^{\text{imaj } \sigma} \frac{1}{(t; q)_{n+1}}.
 \end{aligned}$$

- (c) This is the specialization for $t = 1$.
- (d) A direct proof can be found (see [St86, p. 70]); it is also a consequence of Corollary 11.5.

26. (a) Distribute the n integers $1, 2, \dots, n$ into k classes of sizes a_1, a_2, \dots, a_k . Write the elements in each class in increasing order and juxtapose those class increasing sequences in the following order: first class 1, then class 2, \dots , class k . This yields a permutation σ having the following property: $\text{Ligne } \sigma \subset \{a_1, a_1 + a_2, \dots, a_1 + a_2 + \dots + a_{k-1}\}$. Furthermore, there are $n! / (a_1! a_2! \dots a_k!)$ ways of making up such permutations.

(b) The expansion of the right-hand side reads $\sum_{\sigma, \mathbf{b}} q_1^{b_1(\sigma)} q_2^{b_2(\sigma)} \dots q_n^{b_n(\sigma)}$,

where the summation is over all permutations σ and the sequences of integers $\mathbf{b} = (b_1(\sigma) \geq b_2(\sigma) \geq \dots \geq b_n(\sigma) \geq 0)$ such that the following property holds: $k \in \text{Ligne } \sigma \Rightarrow b_k(\sigma) > b_{k+1}(\sigma)$. But the sequence \mathbf{b} may be written as $i_1^{a_1} i_2^{a_2} \dots i_k^{a_k}$, where $i_1 > i_2 > \dots > i_k \geq 0$ and $a_1 \geq 1, a_2 \geq 1, \dots, a_k \geq 1$, so that the monomial $q_1^{b_1(\sigma)} q_2^{b_2(\sigma)} \dots q_n^{b_n(\sigma)}$ is still equal to (*) $q_1^{i_1} \dots q_{a_1}^{i_1} q_{a_1+1}^{i_2} \dots q_{a_1+a_2}^{i_2} \dots q_{a_1+\dots+a_{k-1}+1}^{i_k} \dots q_{a_1+\dots+a_k}^{i_k}$. But the sequence $i_1^{a_1} i_2^{a_2} \dots i_k^{a_k}$ is nothing but a partition $\lambda = (\lambda_1 \geq \lambda_2 \geq \dots)$ having at most n positive parts, and a_1 parts equal to i_1, a_2 parts equal to i_2, \dots, a_k parts equal to i_k , so that the monomial (*) is also equal to \mathbf{q}^λ . Hence,

$$\begin{aligned}
 &\sum_{\sigma, \mathbf{b}} q_1^{b_1(\sigma)} q_2^{b_2(\sigma)} \dots q_n^{b_n(\sigma)} \\
 &= \sum_{\lambda} \mathbf{q}^\lambda \#\{\sigma : \text{Ligne } \sigma \subset \{a_1, a_1 + a_2, \dots, a_1 + \dots + a_{k-1}\}\} \\
 &= \sum_{\lambda, l(\lambda) \leq n} \mathbf{q}^\lambda \frac{n!}{a_1! a_2! \dots a_k!} = \sum_{\lambda, l(\lambda) \leq n} \mathbf{q}^\lambda \binom{n}{\mathbf{m}(\lambda)}.
 \end{aligned}$$

27. The left-hand side of (E27.1) is equal to the sum of the series $\sum t^{s' + \text{des}_U w} q^{\|\mathbf{a}\| + \text{maj}_U w}$ over all triples (s', \mathbf{a}, w) . By using Proposition 3.1 the right-hand side is equal to the sum of the series

$\sum t^s q^{\|\mathbf{a}^{(1)}\| + \dots + \|\mathbf{a}^{(r)}\|}$, where each $\mathbf{a}^{(i)} = (a_{i,1}, \dots, a_{i,c_i})$ is a sequence of integers satisfying $s \geq a_{i,1} \geq \dots \geq a_{i,c_i} \geq 0$, if $i \in S_{<}$; $s \geq a_{i,1} > \dots > a_{i,c_i} \geq 0$, if $i \in S_{\leq}$; $s \geq a_{i,1} \geq \dots \geq a_{i,c_i} \geq 1$, if $i \in L_{<}$; $s \geq a_{i,1} > \dots > a_{i,c_i} \geq 1$, if $i \in L_{\leq}$. The bijection $(s', \mathbf{a}, w) \mapsto (s, \mathbf{a}^{(1)}, \dots, \mathbf{a}^{(r)})$ such that $s = s' + \text{des}_U w$ and $\|\mathbf{a}\| + \text{maj}_U w = \|\mathbf{a}^{(1)}\| + \dots + \|\mathbf{a}^{(r)}\|$ can be constructed by rewriting the *MacMahon Verfahren* developed in sections 6 and 7 almost verbatim. To obtain (E27.2) use the manipulation on q -series as at the end of section 7 for going from identity (7.7) to (7.8). To derive (E27.3) multiply (E27.2) by $(1 - t)$ and make $t = 1$.

28. (a) The identity follows from Theorem 5.1. The letters belonging to $S_{<}$ bring no further U -inversions; those belonging to $S_{\leq} \cup L_{\leq}$ bring $\prod_{i \in S_{\leq} \cup L_{\leq}} q^{\binom{m_i}{2}}$ extra U -inversions when they are compared between themselves; finally, the U -inversion number must be increased by the number of letters belonging to $L_{<} \cup L_{\leq}$, that is, $\sum_{i \in L_{<} \cup L_{\leq}} m_i$. To show that $\text{inv} A_{\mathbf{m}}^U(q) = A_{\mathbf{m}}^U(1, q)$ is equal to the right-hand side of (E27.3) it suffices to expand the q -multinomial coefficient and to make use of (3.7) and (3.8).
- (b) Let $w = x_1 x_2 \dots x_m$ be a word in $R(\mathbf{m})$. For each $i \in S_{\leq} \cup L_{\leq}$ replace the m_i occurrences of the letter i in w by the letters $(i, m_i), (i, m_i - 1), \dots, (i, 1)$, when reading the word *from left to right*. For each $i \in S_{<} \cup L_{<}$ replace all the m_i occurrences of i by the same letter $(i, 1)$. Finally, let $*$:= $(h + \frac{1}{2}, 1)$. By convention, $(i, k) < (j, l)$ if either $i < j$, or $i = j$ and $k < l$, so that the new word, say, \bar{w} , has its letters in a totally ordered alphabet. Apply the transformation Φ to $\bar{w}*$ to obtain a word of the form $\Phi(\bar{w}*) = \bar{w}'*$. In \bar{w}' replace each letter (i, k) by the letter i , to obtain a word w' in the class $R(\mathbf{m})$. Define $\Psi(w) = w'$.
- (c) Apply the algorithm described in (b). In the first operation write i_k instead of (i, k) and obtain $\bar{w} = 6_1, 2_2, 2_1, 3_1, 1_1, 7_1, 7_1, 4_1, 8_1, 5_1$. Here $*$ = $4 + \frac{1}{2}$; apply Φ to $\bar{w}*$, then derive $\Psi(w)$:

$$\begin{aligned} \bar{w} * &= 6_1, 2_2, 2_1, 3_1, 1_1, 7_1, 7_1, 4_1, 8_1, 5_1, * \\ &\quad | 6_1 \\ &6_1 | 2_2 \\ &6_1, 2_2 | 2_1 \\ &6_1, 2_2 | 2_1 | 3_1 \\ &2_2, 6_1, 2_1, 3_1 | 1_1 \\ &2_2, 6_1, 2_1, 3_1, 1_1 | 7_1 \\ &2_2, 6_1, 2_1, 3_1, 1_1, 7_1 | 7_1 \end{aligned}$$

$$\begin{aligned}
 & 2_2, 6_1 \mid 2_1, 3_1, 1_1, 7_1 \mid 7_1 \mid 4_1 \\
 & 6_1, 2_2, 7_1, 2_1, 3_1, 1_1, 7_1, 4_1 \mid 8_1 \\
 & 6_1 \mid 2_2, 7_1 \mid 2_1, 3_1, 1_1, 7_1 \mid 4_1, 8_1 \mid 5_1 \\
 & 6_1 \mid 7_1 \mid 2_2, 7_1 \mid 2_1, 3_1, 1_1, 8_1 \mid 4_1, 5_1 \mid * \\
 \Phi(\bar{w} *) &= 6_1, 7_1, 7_1, 2_2, 8_1, 2_1, 3_1, 1_1, 5_1, 4_1, * \\
 \Psi(w) &= 6, 7, 7, 2, 8, 2, 3, 1, 5, 4
 \end{aligned}$$

The number of U -inversions is $0 + 0 + 0 + 3 + 0 + 5 + 4 + 7 + 4 + 5 = 28$, plus the number of elements in $L_{<} \cup L_{\leq}$, which is 5. Thus, $\text{inv}_U \Psi(w) = 33$.

29. (a) By induction on the number l of parts of λ .
 (b) Again by induction on l we get $\xi_q(h_\lambda) = \text{inv}A_\lambda(t, q)/(q; q)_n$, where $\text{inv}A_\lambda := \sum_{\sigma} t^{\text{des}_\lambda \sigma} q^{\text{inv} \sigma}$ ($\sigma \in \mathfrak{S}_n$).
 (c) Same derivation as in (a).
 (d) Let $\mathcal{E}(u) := (-t + \exp(u(t-1)))/(1-t) = 1 + \sum_{k \geq 1} (1-t)^{k-1} u^k/k!$
 Then $\mathcal{E}(u)(1 + \sum_{k \geq 1} t A_k(t)(-u)^k/k!) = \mathcal{E}'(u)$ (the derivative of $\mathcal{E}(u)$).
 (e) $\xi_q(p_n) = \frac{1}{(q; q)_n} \sum_{1 \leq k \leq n} q^{\binom{k}{2}} \begin{bmatrix} n \\ k \end{bmatrix} k(t-1)^{k-1} \text{inv}A_{n-k}(t, q);$
 $\xi_{Q,q}(p_n) = \frac{1}{(Q; Q)_n (q; q)_n} \sum_{1 \leq k \leq n} Q^{\binom{k}{2}} \begin{bmatrix} n \\ k \end{bmatrix}_Q \begin{bmatrix} n \\ k \end{bmatrix}_q k(t-1)^{k-1} B_{n-k}(t, Q, q).$
 (f) No more hints.

30. The generating function for the $B_n^{(3)}$ is equivalent to the recurrence formula

$$B_n^{(3)} = \sum_{0 \leq k \leq n} X^k q^{\binom{k+1}{2}} \sum_{0 \leq m \leq n-k} \begin{bmatrix} n \\ m \end{bmatrix} A_{n-m}^{n-m}(t, q) \begin{bmatrix} n-m \\ k \end{bmatrix} q^{mk} (1-t)^m.$$

Hence, to show that $B_n''' = B_n^{(3)}$, it suffices for $0 \leq k \leq n$ to prove the recurrence formula

$$A_n^k(t, q) = \sum_{0 \leq m \leq n-k} \begin{bmatrix} n-k \\ m \end{bmatrix}_q A_{n-m}^{n-m}(t, q) q^{mk} (1-t)^m.$$

Use the iterative method already illustrated in § 21.3. If a permutation of order n ends with a term at most equal to $(n-k)$, we can

consider its longest increasing rightmost factor (*l.i.r.f.*). If the permutation ends with a term greater than or equal to $(n - k + 1)$, make the convention that its *l.i.r.f.* is of length zero.

For $m = 0, 1, \dots, n - k$ designate the generating function for those permutations whose *l.i.r.f.* is of length m by F_m and let $G_m := F_m + \dots + F_{n-k}$. Then $u_m = \begin{bmatrix} n-k \\ m \end{bmatrix}_q A_{n-m}^{n-m}(t, q) q^{mk}$ is the generating function for the permutations whose *l.i.r.f.* is of length at least equal to m , adding a supplementary descent when the *l.i.r.f.* is of length greater than or equal to $(m + 1)$. We then have

$$u_m = F_m + tG_{m+1};$$

and

$$G_m = u_m + (1 - t)G_{m+1}.$$

As $A_{n,k}(t, q) = G_0$ and as $G_{n-k} = A_{k,k}(t, q) q^{(n-k)k}$, we obtain the recurrence formula by iteration starting with G_0 .

31. No comment for the first part. The relation involving $\text{sgn}A_{2n+1}(t)$ is treated in the same way. For the last part notice that $\text{maj } \sigma = 2 \text{ maj } \tau + \sum_{i \in E(\sigma)} (2i - 1)$. Hence

$$\begin{aligned} \text{sgn}A_n(t, q) &= \sum_{\sigma} \text{sgn } \sigma t^{1+\text{des } \sigma} q^{\text{maj } \sigma} \\ &= \sum_{(\tau, E(\sigma))} (-1)^{\#E(\sigma)} t^{1+\text{des } \tau + \#E(\sigma)} q^{2 \text{ maj } \tau + \sum_{i \in E(\sigma)} (2i - 1)} \\ &= \sum_{E \subset \{1, \dots, n\}} (-t)^{\#E} q^{\sum_{i \in E} (2i - 1)} \sum_{\tau \in \mathfrak{S}_n} t^{1+\text{des } \tau} q^{2 \text{ maj } \tau} \\ &= (tq; q^2)_n t^{\text{maj}} A_n(t, q^2). \end{aligned}$$

32. (a) The table of the first $a(n, i)$ is the following:

$i =$	1	2	3	4	5	6	7	8
$n = 1$	1							
2	2	2						
3	2	1	1					
4	3	3	1	1				
5	3	2	1	1	1			
6	3	3	2	1	1	1		
7	3	2	2	1	1	1	1	
8	4	4	2	2	1	1	1	1

- (b) By definition $1 - q^i = \prod\{\Phi_d(q) : d \mid i\}$ for each $i \geq 1$. As $(1 - q^{2i}) = (1 - q^i)(1 + q^i)$, we have $1 + q^i = \prod\{\Phi_d(q) : d \mid 2i, d \nmid i\}$. In particular, if $0 \leq j \leq l$, then $1 + q^{m2^j} = \prod\{\Phi_d(q) : d \mid m2^{j+1}, d \nmid m2^j\} = \prod\{\Phi_d(q) : d \in A_j\}$.
- (c) As the sets A_j are two by two disjoint, it suffices to show that B is the union of the A_j 's. But if $d \mid m2^{j+1}$, $d \nmid m2^j$ for some j with $0 \leq j \leq l$, then $d \mid 2n$ (equal to $m2^{l+1}$) and d is even. Thus d belongs to B . Conversely, suppose $d \mid 2n$ and d even. Then $d = m'2^{j+1}$ with m' odd, $m' \mid m$ and $0 \leq j \leq l$. Consequently, d is an element of A_j .
- (d) The product is zero if $k \geq n$. When $0 \leq k \leq n - 1$, the Gaussian polynomial is the product of the two factors

$$P = \frac{Od_n(q)(1 - q^{2n-1})Od_{n-1}(q) \cdots (1 - q^{2n-2k+1})Od_{n-k}(q)}{(1 - q^{2k+1})Od_k(q)(1 - q^{2k-1}) \cdots Od_1(q)(1 - q)}$$
 and

$$Q = \frac{Ev_n(q)Ev_{n-1}(q) \cdots Ev_{n-k}(q)}{Ev_k(q)Ev_{k-1}(q) \cdots Ev_1(q)}$$
. When numerators and denominators are expressed in terms of cyclotomic polynomials, then P and Q involve cyclotomic polynomials Φ_d with d odd for P and d even for Q . As $\begin{bmatrix} 2n \\ 2k+1 \end{bmatrix}$ is a polynomial and the cyclotomic polynomials are irreducible, each of those two factors is also a polynomial. But P is precisely equal to the expression under consideration.
- (e) Rewrite the recurrence relation (24.5)

$$D_{2n+1}(q) = \sum_{0 \leq k \leq n-1} \begin{bmatrix} 2n \\ 2k+1 \end{bmatrix} q^{2k+1} D_{2k+1}(q) D_{2n-2k-1}(q).$$

First, $D_1(q) = 1$. Proceed by induction on $n \geq 1$. For $0 \leq k \leq n - 1$ the product $\begin{bmatrix} 2n \\ 2k+1 \end{bmatrix} \frac{D_{2k+1}(q)D_{2n-2k-1}(q)}{Ev_1(q) \cdots Ev_n(q)}$ is a polynomial because it may be factorized as $\begin{bmatrix} 2n \\ 2k+1 \end{bmatrix} \frac{Ev_0(q) \cdots Ev_k(q)}{Ev_{n-k}(q) \cdots Ev_n(q)} \times \frac{D_{2k+1}(q)}{Ev_0(q) \cdots Ev_k(q)} \times \frac{D_{2n-2k-1}(q)}{Ev_1(q) \cdots Ev_{n-k-1}(q)}$. The first factor is a polynomial by (d), so are the other two by the induction hypothesis. Hence, when n is odd, each term in the sum in the quadratic recurrence above is divisible by $F_n(q) = Ev_1(q)Ev_2(q) \cdots Ev_n(q)$. Thus $F_n(q) \mid D_{2n+1}(q)$. When n is even, rewrite the recurrence formula by grouping the terms two by two to give

$$D_{2n+1}(q) = \sum_k \begin{bmatrix} 2n \\ 2k+1 \end{bmatrix} q^{2k+1} (1 + q^{2(n-2k-1)}) D_{2k+1}(q) D_{2n-2k-1}(q),$$

where this time k runs over the interval $[0, n/2 - 1]$. As n is even, the binomial $1 + q^{2(n-2k-1)}$ is divisible by $1 + q^2$; and the product $\begin{bmatrix} 2n \\ 2k+1 \end{bmatrix} D_{2k+1}(q) D_{2n-2k-2}(q)$ is divisible by $Ev_1(q) \cdots Ev_n(q)$. Hence, each term in the sum is divisible by $F_n(q) = (1+q^2)Ev_1(q) \cdots Ev_n(q)$.

33. (a) Let $\sigma = x_1 x_2 \dots x_{2n}$ be a balanced rising alternating permutation (*b.r.a.p.*) of order $2n$. As $2n$ is necessarily a peak of σ , it must be the leftmost one, i.e., $x_2 = 2n$. If $x_1 = i$ with $i \leq 2n - 2$, then $(i + 1)$ would occur to the right of i and σ would not be balanced. Thus σ is of the form $\sigma = (2n - 1)(2n)x_3 \cdots x_{2n-1}x_{2n}$ and is balanced if and only if the right factor $x_3 \cdots x_{2n-1}x_{2n}$ is i -balanced for every $i = 1, 2, \dots, 2n - 3$. By induction, $(2n - 3)(2n - 2) \cdots 3412$ is the only *b.r.a.p.* of order $2n - 2$, so that the unique *b.r.a.p.* of order $2n$ is $(2n - 1)(2n)(2n - 3)(2n - 2) \cdots 3412$.
- (b) If σ contains the factor $(i' + 1)i'$, then all the letters greater than $(i' + 1)$ are to the left of $(i' + 1)$. But as σ is alternating of even length, the letter i' is between two letters greater than i' , which is a contradiction.
- (c) The relation $\text{inv}\Phi(\sigma) = \text{inv}\sigma + 1$ implies that a rising alternating permutation with a maximal number of inversions is necessarily balanced. As there is only one such a permutation whose number of inversions is equal to $2n(n - 1)$, the property is proved.
- (d) As $\sum\{q^{\text{inv}\sigma} : \sigma \in A_{2n}\} = \sum\{q^{\text{inv}\sigma} : \sigma \in B_{2n}\} = q^{4(n-1)}E_{2n-2}(q)$ and $\sum\{q^{\text{inv}\sigma} : \sigma \in A_{2n} \cap B_{2n}\} = q^{4(n-2)+4(n-1)}E_{2n-4}(q)$, the induction on n implies $\sum\{q^{\text{inv}\sigma} : \sigma \in A_{2n} \cup B_{2n}\} \equiv 2q^{4(n-1)}q^{2(n-1)(n-2)} - q^{4(n-2)+4(n-1)}q^{2(n-2)(n-3)} \equiv q^{2n(n-1)} \pmod{(q+1)^2}$.
- (e) Let C_{2n} be the complement of $A_{2n} \cup B_{2n}$, that is, the set of rising alternating permutations of length $2n$ having $(2n)$ and $(2n - 1)$ among their peaks and 1 and 2 among their troughs. There remains to show that $\sum\{q^{\text{inv}\sigma} : \sigma \in C_{2n}\} \equiv 0 \pmod{(q+1)^2}$. Let τ, τ' be the transpositions $(2n - 1, 2n)$ and $(1, 2)$, respectively, and G be the group of order 4 generated by $\{\tau, \tau'\}$. The group G acts on C_{2n} and the generating polynomial for the four elements in each orbit by the number of inversions is divisible by $(q+1)^2$. Therefore, the generating polynomials for all elements of C_{2n} is also divisible by $(q+1)^2$.
34. No comment.
35. Rewrite identity (25.3) as $\sum_{w \in [0, r]^*} \binom{\bar{w}}{w} = 1/\det(I - C_r)$, where C_r is the matrix displayed in (25.4) and let \mathfrak{V} be the homomorphism of

the algebra \mathcal{A}^{MM} generated by $\mathfrak{A} \binom{a}{b} = \begin{cases} \binom{a}{b}, & \text{if } a \neq b; \\ 0, & \text{if } a = b. \end{cases}$ This yields the corresponding formula for the vertical derangements.

36. For getting (i) apply the homomorphism $\Phi \binom{i}{j} := \begin{cases} tu_j, & \text{if } i < j; \\ u_j, & \text{if } i \geq j; \end{cases}$ to the identity $\det(I - C_r) \times \text{Bos} = 1$.

Let $f_{r+1}(u_0, \dots, u_r)$ denote the right-hand side of formula (ii). Then $f_r((1-t)u_1, \dots, (1-t)u_r) = \prod_{1 \leq i \leq r} (1 - (1-t)u_i)$ by Theorem 14.2;

also, $f_{r+1}(u_0, \dots, u_r) = f_r(u_1, \dots, u_r) - u_0 f_r(((1-t)u_1, \dots, (1-t)u_r)$. Next, let $g_{r+1}(u_0, \dots, u_r)$ denote the value of the determinant D . By subtracting the rightmost column from each of the other columns we get for $r \geq 1$:

$$g_{r+1}(u_0, \dots, u_r) = \det \begin{pmatrix} 1 & 0 & \dots & 0 & -u_0 \\ (1-tu_1) & 1 & \dots & -u_1 & -u_1 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ (1-t)u_{r-1} & (1-t)u_{r-1} & \dots & 1 & -u_{r-1} \\ (1-t)u_r - 1 & (1-t)u_r - 1 & \dots & (1-t)u_r - 1 & 1 - u_r \end{pmatrix}.$$

Expanding the determinant along the first row yields

$$g_{r+1}(u_0, \dots, u_r) = g_r(u_1, \dots, u_r) + (-1)^{r+1} u_0 h_r(u_1, \dots, u_r) \text{ with } h_r(u_1, \dots, u_r) = \det \begin{pmatrix} (1-tu_1) & 1 & \dots & -u_1 \\ \vdots & \vdots & \ddots & \vdots \\ (1-t)u_{r-1} & (1-t)u_{r-1} & \dots & 1 \\ (1-t)u_r - 1 & (1-t)u_r - 1 & \dots & (1-t)u_r - 1 \end{pmatrix}.$$

But

$$\begin{aligned} h_r(u_1, \dots, u_r) &= ((1-t)u_r - 1) h_{r-1}(u_1, \dots, u_{r-1}) \\ &= \prod_{1 \leq i \leq r} ((1-t)u_i - 1) \\ &= (-1)^r f_r((1-t)u_1, \dots, (1-t)u_r), \end{aligned}$$

and then

$$g_{r+1}(u_0, \dots, u_r) = g_r(u_1, \dots, u_r) - u_0 f_r((1-t)u_1, \dots, (1-t)u_r).$$

As $g(u_0) = 1 - u_0 = f_1(u_0)$ and $g_2(u_0, u_1) = \det \begin{pmatrix} 1 & -u_0 \\ (1-t)u_1 - 1 & 1 - u_1 \end{pmatrix} = f_2(u_0, u_1)$, identity (ii) holds. For (iii) use Theorem 4.2 and (iv) is banal.

37. We have: $\text{Ferm} = \det \begin{pmatrix} 1 - \binom{1}{1} & -\binom{1}{2} \\ -\binom{2}{1} & 1 - \binom{2}{2} \end{pmatrix} = 1 - \binom{1}{1} - \binom{2}{2} + \binom{12}{12} - \binom{21}{12}$ and $\text{Bos}^{(3)} = 1 + \binom{1}{1} + \binom{2}{2} + \binom{11}{11} + \binom{12}{12} + \binom{12}{21} + \binom{12}{22} + \binom{111}{111} + \binom{112}{112} +$

$\binom{112}{121} + \binom{112}{211} + \binom{122}{122} + \binom{122}{212} + \binom{122}{221} + \binom{222}{222}$. Making all the cancellations the product $\text{Ferm} \times \text{Bos}^{(3)}$ becomes $1 + S_1 + S_2 + S_3 + S_4$, where $S_1 = \binom{12}{12} + \binom{12}{21} - \binom{21}{12} - \binom{21}{21}$, $S_2 = \binom{112}{211} + \binom{121}{121} - \binom{211}{211} - \binom{211}{121}$, $S_3 = \binom{122}{122} + \binom{122}{212} - \binom{212}{122} - \binom{212}{212}$, $S_4 = \binom{122}{221} - \binom{212}{221}$. As by assumption $\text{Ferm} \times \text{Bos}^{(3)} = 1$, we necessarily have: $S_1 = 0$, $S_2 = 0$ and $S_3 + S_4 = 0$. The identity $S_1 = 0$ implies that rule (R2) holds in $\overline{\mathcal{A}}$. But if (R2) holds, S_2 is also null, as the billetter $\binom{2}{2}$ can be factorized out. In its turn $S_4 = 0$, and so $\binom{12}{22} - \binom{21}{22} = 0$ and this means that (R1) also holds in $\overline{\mathcal{A}}$.

38. In the determinant D subtract the r -th row from the $(r + 1)$ -st one; then the $(r - 1)$ -st from the r -th row; \dots , the first row from the second, to obtain:

$$D = \begin{vmatrix} 1 & -X_1 & -X_2 & \cdots & -X_{r-1} & -X_r \\ -1 - Y_0 & 1 + X_1 & 0 & \cdots & 0 & 0 \\ 0 & -1 - Y_1 & 1 + X_2 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & 1 + X_{r-1} & 0 \\ 0 & 0 & 0 & \cdots & -1 - Y_{r-1} & 1 + X_r \end{vmatrix}.$$

Next, make the cofactor expansion of D with respect to the *first row*:

$$\begin{aligned} D &= \prod_{1 \leq j \leq r} (1 + X_j) - \sum_{1 \leq l \leq r} \left(\prod_{0 \leq j \leq l-1} (1 + Y_j) \prod_{l+1 \leq j \leq r} (1 + X_j) \right) X_l \\ &= \prod_{1 \leq j \leq r} (1 + X_j) + X_0 \prod_{1 \leq j \leq r} (1 + X_j) \\ &\quad - \sum_{0 \leq l \leq r} \left(\prod_{0 \leq j \leq l-1} (1 + Y_j) \prod_{l \leq j \leq r} (1 + X_j) \right) \frac{X_l}{1 + X_l}. \end{aligned}$$

Hence,

$$\sum_{w \in [0, r]_h^*} \psi(w) = \frac{\prod_{0 \leq j \leq r} \frac{(1 + Y'_j)}{(1 + X_j)}}{1 - \sum_{0 \leq l \leq r} \prod_{0 \leq j \leq l-1} \frac{(1 + Y_j)}{(1 + X_j)} \frac{X_l}{1 + X_l}}.$$

39. As $\sum_{w \in [0, r]^*} \psi(w) = \sum_{v \in [0, r]_h^*} \beta \psi(v)$ by (26.17) with β defined in (26.16), we get the desired expression. To get the second formula note that by definition each letter equal to 0 cannot be a decrease value. Consequently, the weight $\psi(w)$ of each word w must not contain

the variables X_0, Z_0 . There is then another expression for the right-hand side of the formula, which does not involve the variables X_0, Z_0 . To obtain factor out $(1 - Z_0)/(1 - Z_0 + X_0)$ from both numerator and denominator of the right-hand side.

40. First, check that

$$\begin{aligned} & \prod_{0 \leq j \leq r} (1 - sY_j) - \prod_{0 \leq j \leq r} (1 - Y_j) \\ = & \sum_{0 \leq l \leq r} \prod_{0 \leq j \leq l} (1 - sY_j) \prod_{l+1 \leq j \leq r} (1 - Y_j) - \sum_{0 \leq l \leq r} \prod_{0 \leq j \leq l-1} (1 - sY_j) \prod_{l \leq j \leq r} (1 - Y_j) \\ = & (1 - s) \sum_{0 \leq l \leq r} Y_l \prod_{0 \leq j \leq l-1} (1 - sY_j) \prod_{l+1 \leq j \leq r} (1 - Y_j). \end{aligned}$$

Using identity (*) we get:

$$\begin{aligned} & \frac{1}{1 - s \sum_{0 \leq l \leq r-1} Y_l \prod_{0 \leq j \leq l-1} (1 - sY_j) / \prod_{0 \leq j \leq l} (1 - Y_j)} \\ = & \frac{\prod_{0 \leq j \leq r-1} (1 - Y_j)}{\prod_{0 \leq j \leq r-1} (1 - Y_j) - s \sum_{0 \leq l \leq r-1} Y_l \prod_{0 \leq j \leq l-1} (1 - sY_j) \prod_{l+1 \leq j \leq r-1} (1 - Y_j)} \\ = & \frac{\prod_{0 \leq j \leq r-1} (1 - Y_j)}{\prod_{0 \leq j \leq r-1} (1 - Y_j) - s \left(\prod_{0 \leq j \leq r-1} (1 - sY_j) - \prod_{0 \leq j \leq r-1} (1 - Y_j) \right) / (1 - s)} \\ = & \frac{(1 - s) \prod_{0 \leq j \leq r-1} (1 - Y_j)}{\prod_{0 \leq j \leq r-1} (1 - Y_j) - s \prod_{0 \leq j \leq r-1} (1 - sY_j)}. \end{aligned}$$

41. No comment.

42. No comment.

43. No comment.

44. (a) The permutations compatible with 321^2 are the following: 1123 (no descent); 2134, 3124, 4123 (one descent at position 1); 1324, 2314, 1423, 3412, 2413 (one descent at position 2); 4213, 4312, 3214 (two descents at positions 1 and 2).

(b) We have

ANSWERS TO THE EXERCISES

$$\left(\begin{array}{cccccc} w = 1|1|2|3 & 1|1|3|2 & 1|2|1|3 & 1|2|3|1 & 1|3|1|2 & 1|3|2|1 \\ \check{\sigma} = 4|3|2|1 & 4|3|1|2 & 4|2|3|1 & 4|2|1|3 & 4|1|3|2 & 4|1|2|3 \\ \sigma = 1234 & 2134 & 1324 & 3214 & 3124 & 2314 \\ z = 0000 & 1000 & 1100 & 2100 & 1000 & 1100 \\ \mathbf{m} = 3211 & 3211 & 3211 & 3211 & 3211 & 3211 \\ c = 3211 & 2211 & 2111 & 1111 & 2211 & 2111 \end{array} \right)$$

$$\left(\begin{array}{cccccc} w = 211|3 & 21|3|1 & 2|3|1|1 & 311|2 & 312|1 & 321|1 \\ \check{\sigma} = 243|1 & 24|1|3 & 2|1|4|3 & 143|2 & 142|3 & 124|3 \\ \sigma = 1423 & 341|2 & 421|3 & 412|3 & 431|2 & 241|3 \\ z = 1100 & 1100 & 2100 & 1000 & 2100 & 1100 \\ \mathbf{m} = 3211 & 3211 & 3211 & 3211 & 3211 & 3211 \\ c = 2111 & 2111 & 1111 & 2211 & 1111 & 2111 \end{array} \right)$$

45. No comment.

Notes

The use of the algebra of q -series in Combinatorics goes back to MacMahon [Mac13, 15, 78], who realized that certain closed formulas in Enumeration could only be expressed in that context. He had a great talent for deriving many of his results in a very intuitive manner. In our to-day's more systematic approach a memoir on q -series in Combinatorics has to begin with a chapter on basic hypergeometric series, at least on the fundamental result of that theory, which is the q -binomial theorem. More material can be found in Gasper and Rahman [GaRa90], also in the old book by Slater [Sl66], or in the more recent one by Fine [Fi88]. The second chapter in Andrews' book [An76] also covers all that is needed on this subject. The combinatorial aspects of the hypergeometric series identities are developed in [JoSt87], but not discussed in those Notes. Published in the seventies Gessel's, Ph.D. thesis [Ge77] must be viewed as an excellent memoir on q -series and combinatorics.

Coding permutations by sequences of integers for computer purposes goes back to Lehmer [Le60]. The inv-coding, as such, belongs to him. The maj-coding is implicit in the early papers by Carlitz [Ca54, 59, 75] on Eulerian numbers and made explicit in Rawlings [Ra79, 80]. In studying the genus zeta function of local minimal hereditary orders, Denert [Den90] introduced a new permutation statistic, which was later christened "den" and was shown to be equidistributed with the major index or inversion number. The den-coding in § 2.3 is taken from [FoZe90].

The algebra of the q -binomial coefficients is classical; see, e.g. [An71, § 3.3]. It was convenient to devote a full chapter, essentially chap. 4, to presenting the main combinatorial structures counted by those coefficients. Chapters 5 and 6 may be regarded as an extension of section 3.4 of Andrews's monograph [An71]. However, the proof of the fact that the Major Index is a q -multinomial statistic, a result that is due to MacMahon [Mac13], is given in greater detail; it involves the so-called *MacMahon Verfahren*, that can be viewed as a transformation on two-row matrices. That transformation is based upon a *commutation rule* that preserves the statistics under study. Notice that the *MacMahon Verfahren* has been extended by Stanley [St72] in his theory of (P, ω) -partitions.

As is often the case, an explicit combinatorial tool like the *MacMahon Verfahren* opens the way to significant extensions, such as the study of bivariable polynomials $A_{\mathbf{m}}(t, q)$, indexed by sequences $\mathbf{m} = (m_1, \dots, m_r)$ of r nonnegative integers. Formula (7.7) already appears in MacMahon [Mac15, vol. 2, p. 211]. Chapters 7, 8 and 9 should be regarded as a systematic approach to combinatorial q -calculus. The material has been taken from [ClFo95a and b, Fo95, FoKr95].

Several sources have been used for chapter 10: [Ri58, p. 38–39 and 213–216, FoSc70] for the traditional Eulerian polynomials, [ChMo71, St76] for the q -inv Eulerian polynomials, [Fo76, Ga79] for the joint study of the two q -extensions. The *iteration method* developed for Lemma 10.2 appears in several different forms, for instance in [Ge82, Ze80a, FoZe91].

The joint combinatorial study of the statistics “maj” and “inv” on a class of rearrangements of an arbitrary sequence, as written in chapter 11, is borrowed from [Fo68, FoSc78]. However the construction of the fundamental transformation is made in a very different manner, as there is a canonical way of extending each bijection valid for binary words to a bijection over an arbitrary class of rearrangements. Notice that Björner and Wachs have proposed an interesting extension to Poset Theory [BjW88] and derived further properties of the fundamental transformation. Properties (d) and (e) in Theorem 11.3 are basically theirs.

The expansion of the *infinite* product $1/\prod_{i \geq 0, j \geq 0} (1 - uq_1^i q_2^j)$, made in chapter 12, can be found in [Ca56] and also in [Ro74], where a first combinatorial interpretation was given. See also [St76].

The expansion of the *finite* product $1/\prod_{0 \leq i \leq r, 0 \leq j \leq s} (1 - uq_1^i q_2^j)$ naturally leads to the study of a four-variable generating polynomial for the permutation group. Identity (13.7) first appeared in [GG78]. Other proofs can be found in [Ra80, DeFo85]. Specializations had been anticipated in [Ca76].

There is no originality in discussing some basic facts on symmetric functions, as done in chapters 14, 15, 16. The best source of study remains Macdonald’s book [Ma95]. To-day we rejoice at the coming out of Lascoux’s treatise [La03] with its creative approach to the subject, as started in [LaSch81, 84].

The combinatorial definition of the Schur function given in chapter 17 is taken from Proctor [Pr89], but has been rewritten in a more systematic way, under the superb guidance of our friend Jean-Pierre Jouanolou.

Corollary 18.3 can be found in [St76], also in [DeFo85]. The idea of keeping a several-variable statistic y instead of the one-variable “maj” is due to Adin et al. [ABR01]. This yields Theorem 18.2, but the resulting identities involve a *non-ring* linear homomorphism.

The Robinson-Schensted correspondence [Ro38, Sc61], that has been so popular in the seventies and eighties, is described in the more general set-up developed by Knuth [Kn70]. The geometric properties of that correspondence are due to Schützenberger [Sch73, 77]. An excellent exposition of its various properties is given in [Kn70, p. 48–72]. Theorem 19.4 may be regarded as an Adin-Brenti-Roichman extension of identity (13.7).

The generating function for the signed permutations by number of descents (Theorem 20.1) has been calculated by various authors: [St92, StE93, Br94, Re93a, Re93b, Re93c, Re95a, ClFo94]. The proof of Theorem 20.1 is taken from the last reference. Theorem 20.3 is due to Carlitz et al. [Ca76], Theorem 21.1 to [FoHa97]; however its specialization to the symmetric group appears already in [St76]. The idea of taking *finite analogs* of the Bessel functions for enumerating pairs of permutations is due to Fedou and Rawlings [Fe95, FeRa94, FeRa95]. The extension to the group of signed permutations and accordingly Theorem 22.5 can be found in [FoHa96]. The proof of the theorem relies upon a very convenient inversion formula, which has appeared in various contexts [GoJa83, p. 131, St86, p. 266, Vi86, HuWi75, Ze81]. The inversion formula stated here (Lemma 22.3) is borrowed from [FeRa94, FeRa95].

The proof of Theorem 23.1 is taken from [FoZe81]. The *Désarménien Verfahren* was developed in the two papers [De82 and 83]. Lemma 23.2, Proposition 23.3, as well congruences (23.19) and (23.27) are due to Désarménien. Congruences (23.20) and (23.28) are apparently new. Section 23.4 on signed Eulerian numbers are taken from [DeFo92]. Those numbers were introduced by Loday [Lod89] in a study of the cyclic homology of commutative algebras.

The theory of basic trigonometric functions is due to Jackson [Ja04] (see the detailed bibliography in Gasper and Rahman’s book [GaRa90].) The study of *basic* trigonometric functions proposed in this memoir, as well as its combinatorial counterpart, are apparently new.

The purpose of Chapter 25 has been to make a joint study of the celebrated MacMahon’s “Master Theorem,” in its various forms, the original commutative one, the Cartier-Foata partially commutative and Garoufalidis-Lê-Zeilberger ones.

The decrease value theorem, developed in the last two chapters, may be regarded as an extension of the “Master Theorem” in the sense that it takes a multivariate statistic into account, involving the basic notions of decrease, increase, record, rise and descent. It can be used to derive several explicit distributions of permutation statistics.

For Ex. 1 see [De82a] or [An76, chap. 3, Examples 1–6]. The proof of the Ramanujan sum (Ex. 2) reproduces Ismail’s derivation [Is77]. The

maj-inv bijection for permutations described in Ex. 3 belongs to common knowledge. For Ex. 12 and 13 refer to [Ri58, p. 38–39 and 213–216, FoSc70], as mentioned above. In Ex. 14 the solution made by Clarke [Cl95] has been borrowed. Ex. 14 reproduces the techniques developed in [Ha90a, 90b]. Notice that the study of the Denert statistics for arbitrary words, as developed in [Ha94, 95] has not been touched in those Notes. See, e.g. [Lo02, chap. 10].

The Z -statistic has been introduced in [ZeBr85] for the proof of the Andrews q -Dyson conjecture. The combinatorial approach in Ex. 17 is due to Han [Ha92]. The *tableau emptying-filling involution* was invented by Schützenberger [Sch73] and used in [FoSc78]. Ex. 20 is taken from [FoSc78]. Ex. 25 can be regarded as the t -extension of § 11.4 in the book by Lothaire [Lo02]. The identity of Ex. 26 is taken from [ABR01]. The calculation in Ex. 27 was made in [FoKr95], as well as the content of Ex. 28.

The Brenti homomorphism (see Ex. 29) has been used in [BecRe95], other applications of the Brenti homomorphism are proposed in the Exercise. The formula used for the length in the group of the signed permutations is due to Brenti [Br94], the generating function for the pair (length, number of descents) derived in Ex. 30 is due to Reiner [Re95a]. The involution in Ex. 31 belongs to Wachs [Wa92]. Finally, Ex. 32 and 33 are taken from [Fo81] and [AnFo82]. The remaining exercises make use of the various forms of the “Master Theorem” and the Decrease Value Theorem.

The algebra of symmetric functions, especially the Cauchy identity for Schur functions, has been a powerful tool for deriving various generating counting series. We have not mentioned other tools, such as the Hook Young diagrams developed by Berele, Regev and Remmel [BeRe85, 87] and the derived (k, l) -Schur functions [Rem83, 84, 87]. In most cases it seems that the Schur function model suffices for the derivations [DeFo91].

We have not discussed operator techniques, as developed in [An71] or [Ze80b]. The subject of permutation statistics is in full expansion, so that the present Notes can only be regarded as a partial aspect of to-day’s state of the art. There have been interesting studies by the Californian school (see [Bec95, 96]) and by the vigorous Israeli school ([AR01, ABR01, ReRo03a, ReRo03b]). The goal is to find the most appropriate maj-analogs for the further Weyl groups, or to work out with set-valued statistics [FoHa02]. The first results found in [AR01] are very promising. Finally, let us mention the paper by Babson and Steingrímsson [BaSt00] on the classification of the patterns that lead to a Mahonian distribution.

REFERENCES

- [AR01] Ron M. Adin and Yuval Roichman, The flag major index and group actions on polynomial rings, *Europ. J. Combin.*, vol. **22**, 2001, p. 431–446.
- [ABR01] Ron M. Adin, Francesco Brenti and Yuval Roichman, Descent Numbers and Major Indices for the Hyperoctahedral Group, *Adv. in Appl. Math.*, vol. **27**, 2001, p. 210–224.
- [An71] George E. Andrews, On the Foundations of Combin. Theory V, Eulerian Differential Operators, *Studies in Appl. Math.*, vol. **50**, 1971, p. 345–375.
- [An76] George E. Andrews, *The Theory of Partitions*. London, Addison-Wesley, 1976 (*Encyclopedia of Math. and Its Appl.*, **2**).
- [AnDo80] George E. Andrews and Dominique Foata, Congruences for the q -secant number, *Europ. J. Combin.*, vol. **1**, 1980, p. 283–287.
- [BaSt00] Eric Babson and Einar Steingrímsson, Generalized permutation patterns and a classification of the Mahonian statistics, *Sém. Lothar. Combin.* [<http://mat.univie.ac.at/~slc/>] **44**, 2000, B44b, 18 pp.
- [Bec96] Desirée Beck, The Combinatorics of Symmetric Functions and Permutation Enumeration of the Hyperoctahedral Group, *Discrete Math.*, vol. **163**, 1997, p. 13–45.
- [BecRe95] Desirée A. Beck and Jeff B. Remmel, Permutation Enumeration of the Symmetric Group and the Combinatorics of Symmetric Functions, *J. Combin. Theory, Ser. A*, vol. **72**, 1995, p. 1–49.
- [BeRe85] A. Berele and Jeff B. Remmel, Hook flag characters and their combinatorics, *J. Pure and Appl. Algebra*, vol. **35**, 1985, p. 222–245.
- [BeRe87] A. Berele and Amitai Regev, Hook Young diagrams with applications to combinatorics and to representations of Lie superalgebras, *Adv. in Math.*, vol. **64**, 1987, p. 118–175.
- [BjW88] Anders Björner and Michelle L. Wachs, Permutation Statistics and Linear Extensions of Posets, *J. Combin. Theory, Ser. A*, vol. **58**, 1991, p. 85–114.
- [Br94] Francesco Brenti, q -Eulerian Polynomials Arising from Coxeter Groups, *Europ. J. Combinatorics*, vol. **15**, 1994, p. 417–441.
- [Ca54] L. Carlitz, q -Bernoulli and Eulerian numbers, *Trans. Amer. Math. Soc.*, vol. **76**, 1954, p. 332–350.
- [Ca56] L. Carlitz, The Expansion of certain Products, *Proc. Amer. Math. Soc.*, vol. **7**, 1956, p. 558–564.
- [Ca59] L. Carlitz, Eulerian numbers and polynomials, *Math. Magazine*, vol. **33**, 1959, p. 247–260.
- [Ca66] L. Carlitz, D. P. Roselle and R. A. Scoville, Permutations and Sequences with Repetitions by Number of Increases, *J. Combin. Theory*, vol. **1**, 1966, p. 350–374.
- [Ca75] L. Carlitz, A combinatorial property of q -Eulerian numbers, *Amer. Math. Monthly*, vol. **82**, 1975, p. 51–54.
- [Ca76] L. Carlitz, R. Scoville and T. Vaughan, Enumeration of pairs of permutations, *Discrete Math.*, vol. **14**, 1976, p. 215–239.
- [Ca72] P. Cartier, *La série génératrice exponentielle*, *Applications probabilistes et algébriques*. Publ. I.R.M.A., Université Louis Pasteur, Strasbourg, 1972, 68 pages.
- [CF69] P. Cartier, D. Foata, *Problèmes combinatoires de commutations et réarrangements*. Berlin, Springer-Verlag, 1969 (*Lecture Notes in Math.*, **85**).
- [ChMo71] M. S. Cheema and T. S. Motzkin, Multipartitions and Multipermutations, *Combinatorics* [Los Angeles. 1968], p. 39–70. Providence, Amer. Math. Soc., 1971 (*Proc. Symposia in Pure Math.*, **19**).

- [Cl95] R. J. Clarke, A short proof of a result of Foata and Zeilberger, *Adv. Appl. Math.*, vol. **16**, 1995, p. 129–131.
- [ClFo94] R. J. Clarke and D. Foata, Eulerian Calculus, I: Univariable Statistics, *Europ. J. Combinatorics*, vol. **15**, 1994, p. 345–362.
- [ClFo95a] R. J. Clarke and D. Foata, Eulerian Calculus, II: An Extension of Han’s Fundamental Transformation, *Europ. J. Combinatorics*, vol. **16**, 1995, p. 221–252.
- [ClFo95b] R. J. Clarke and D. Foata, Eulerian Calculus, III: The Ubiquitous Cauchy Formula, *Europ. J. Combinatorics*, vol. **16**, 1995, p. 329–355.
- [Co70] Louis Comtet, *Analyse Combinatoire*, vol. 1 and 2. Paris, Presses Universitaires de France, 1970. (English edition *Advanced Combinatorics*, D. Reidel, Dordrecht, 1974).
- [Den90] M. Denert, The genus zeta function of hereditary orders in central simple algebras over global fields, *Math. Comp.*, vol. **54**, 1990, p. 449–465.
- [De82] Jacques Désarménien, Un analogue des congruences de Kummer pour les q -nombres d’Euler, *Europ. J. Combin.*, vol. **3**, 1982, p. 19–28.
- [De82a] Jacques Désarménien, Les q -analogues des polynômes d’Hermite, *Sém. Lothar. Combin.*, vol. **06b**, 12 p. (<http://www.mat.univie.ac.at/~slc>).
- [De83] Jacques Désarménien, Fonctions symétriques associées à des suites classiques de nombres, *Ann. scient. Éc. Norm. Sup.*, vol. **16**, 1983, p. 271–304.
- [DeFo85] Jacques Désarménien et Dominique Foata, Fonctions symétriques et séries hypergéométriques basiques multivariées, *Bull. Soc. Math. France*, vol. **113**, 1985, p. 3–22.
- [DeFo91] Jacques Désarménien et Dominique Foata, Statistiques d’ordre sur les permutations colorées, *Discrete Math.*, vol. **87**, 1991, p. 133–148.
- [DeFo92] Jacques Désarménien et Dominique Foata, The signed Eulerian Numbers, *Discrete Math.*, vol. **99**, 1992, p. 49–58.
- [DeWa93] Jacques Dsarmnien, Michelle L. Wachs, Descent classes of permutations with a given number of fixed points, *J. Combin. Theory, Ser. A*, vol. **64**, 1993, p. 311–328.
- [Fe95] J.-M. Fedou, Sur les fonctions de Bessel, *Discrete Math.*, vol. **139**, 1995, p. 473–480.
- [FeRa94] J.-M. Fedou and D. Rawlings, More Statistics on Permutations Pairs, *Electronic J. Combin.*, vol. **1**, 1994, R11, 17 pp. [<http://www.emis.de/journals/EJC>].
- [FeRa95] J.-M. Fedou and D. Rawlings, Adjacencies in words, *Advances in Appl. Math.*, vol. **16**, 1995, p. 206–218.
- [Fi88] Nathan J. Fine, *Basic Hypergeometric Series and Applications*. Providence, Amer. Math. Soc., 1988.
- [Fo65] Dominique Foata, Étude algébrique de certains problèmes d’analyse combinatoire et du calcul des probabilités, *Publ. Inst. Statist. Univ. Parts*, vol. **14**, 1965, p. 81–241.
- [Fo68] Dominique Foata, On the Netto inversion number of a sequence, *Proc. Amer. Math. Soc.*, vol. **19**, 1968, p. 236–240.
- [Fo76] Dominique Foata, Distributions eulériennes et mahoniennes sur le groupe des permutations, in *Higher Combinatorics* [M. Aigner, ed., Berlin. 1976], p. 27–49. D. Reidel, Amsterdam, 1977 (Proc. NATO Adv. Study Inst.).
- [Fo81] Dominique Foata, Further divisibility properties of the q -tangent numbers, *Proc. Amer. Math. Soc.*, vol. **81**, 1981, p. 143–148.
- [Fo95] Dominique Foata, Les distributions Euler-Mahoniennes sur les mots, *Discrete Math.*, vol. **139**, 1995, p. 167–188.
- [FoHa97] Dominique Foata et Guo-Niu Han, Calcul basique des permutations signées, I : longueur et nombre d’inversions, *Adv. in Appl. Math.*, vol. **18**, 1997, p. 489–509.
- [FoHa96] Dominique Foata et Guo-Niu Han, Calcul basique des permutations signées, II: analogues finis des fonctions de Bessel, Strasbourg, 28 p., 1996.
- [FoHa02] Dominique Foata et Guo-Niu Han, Une nouvelle transformation pour les statistiques Euler-Mahoniennes ensembliste, [Paris, École Normale Sup., Colloque en

REFERENCES

- l'honneur de Pierre Cartier, 17–18 June 2002], 24 p., to appear, (<http://www-irma.u-strasbg.fr/~foata/paper/>).
- [FH07] Dominique Foata, Guo-Niu Han, A new proof of the Garoufalidis-Le-Zeilberger Quantum Master Theorem, *Journal of Algebra*, vol. **307**, 2007, p. 424–31.
- [FoKr95] Dominique Foata and Christian Krattenthaler, Graphical Major Indices, II, *Sém. Lothar. Combin.*, **B34k**, 1995, 16 pp. (<http://www.mat.univie.ac.at/~slc>).
- [FoSc70] Dominique Foata and Marcel-Paul Schützenberger, *Théorie géométrique des polynômes eulériens*. Berlin, Springer-Verlag, 1970 (*Lecture Notes in Math.*, **138**).
- [FoSc78] Dominique Foata and Marcel-Paul Schützenberger, Major Index and Inversion number of Permutations, *Math. Nachr.*, vol. **83**, 1978, p. 143–159.
- [FoZe90] Dominique Foata and Doron Zeilberger, Denert’s Permutation Statistic Is Indeed Euler-Mahonian, *Studies in Appl. Math.*, vol. **83**, 1990, p. 31–59.
- [FoZe91] Dominique Foata and Doron Zeilberger, Multibasic Eulerian Polynomials, *Trans. Amer. Math. Soc.*, vol. **329**, 1991, p. 843–862.
- [Fkes76] Herbert O. Foulkes, Tangent and secant numbers and representations of symmetric groups, *Discrete Math.*, vol. **15**, 1976, p. 311–324.
- [Fkes80] Herbert O. Foulkes, Eulerian numbers, Newcomb’s problem and representations of symmetric groups, *Discrete Math.*, vol. **30**, 1980, p. 3–49.
- [Ga79] Adriano M. Garsia, On the “maj” and “inv” q -analogues of Eulerian Polynomials, *Linear and Multilinear Algebra*, vol. **8**, 1979, p. 21–34.
- [GLZ06] Stavros Garoufalidis, Thang TQ L, Doron Zeilberger, The quantum MacMahon master theorem, *Proceedings of the National Academy of Sciences*, vol. **103**, 2006, p. 13928–13931.
- [GG78] Adriano M. Garsia and Ira Gessel, Permutations Statistics and Partitions, *Adv. in Math.*, vol. **31**, 1979, p. 288–305.
- [GaRa90] George Gasper and Mizan Rahman, *Basic Hypergeometric Series*. London, Cambridge Univ. Press, 1990 (*Encyclopedia of Math. and Its Appl.*, **35**).
- [Ge77] Ira Gessel, Generating functions and enumeration of sequences, Ph. D. thesis, Dept. Math., M.I.T., Cambridge, Mass., 111 p., 1977.
- [Ge82] Ira M. Gessel, A q -analogue of the exponential formula, *Discrete Math.*, vol. **40**, 1982, p. 69–80.
- [Ge87] Ira M. Gessel, A combinatorial proof of the multivariable Lagrange inversion formula, *J. Combin. Theory Ser. A*, vol. **45**, 1987, p. 78–195.
- [GeRe93] I. M. Gessel, Christophe Reutenauer, Counting Permutations with Given Cycle Structure and Descent Set, *Journal of Combinatorial Theory, Ser. A*, vol. **64**, 1993, p. 189–215.
- [Go60I] I. J. Good, A short proof of MacMahon’s “Master Theorem”, *Proc. Cambridge Philos. Soc.*, vol. **58**, 1962, p. 360.
- [GoJa83] I. P. Goulden and D. M. Jackson, *Combinatorial Enumeration*. New York, John Wiley, 1983.
- [Ha90a] Guo-Niu Han, Distribution Euler-mahonienne: une correspondance, *C. R. Acad. Sci. Paris*, vol. **310**, 1990, p. 311–314.
- [Ha90b] Guo-Niu Han, Une nouvelle bijection pour la statistique de Denert, *C. R. Acad. Sci. Paris*, vol. **310**, 1990, p. 493–496.
- [Ha92] Guo-Niu Han, Une courte démonstration d’un résultat sur la Z-statistique, *C. R. Acad. Sci. Paris*, vol. **314**, 1992, p. 969–971.
- [Ha94] Guo-Niu Han, Une transformation fondamentale sur les réarrangements de mots, *Advances in Math.*, vol. **105**, 1994, p. 26–41.
- [Ha95] Guo-Niu Han, The k -Extension of a Mahonian Statistic, *Adv. Appl. Math.*, vol. **16**, 1995, p. 297–305.
- [Ho82] J. Hofbauer, Lagrange Inversion, *Séminaire Lotharingien de Combinatoire*, **B06a**, 1982, 28 pages, (<http://www.mat.univie.ac.at/~slc>).
- [HuWi75] J. P. Hutchinson and H.S. Wilf, On Eulerian circuits and words with prescribed adjacency patterns, *J. Combin. Theory A*, vol. **18**, 1975, p. 80–87.
- [Is77] Mourad E. H. Ismail, A simple proof of Ramanujan’s ${}_1\psi_1$ sum, *Proc. Amer. Math. Soc.*, vol. **63**, 1977, p. 185–186.

- [Is82] Mourad E. H. Ismail, The zeros of basic Bessel functions, *J. Math. Anal. and Appl.*, vol. **86**, 1982, p. 1–19.
- [JaKe81] Gordon James and Adalbert Kerber, *The Representation Theory of the Symmetric Group*. Reading, Mass., Addison-Wesley, 1981 (*Encyclopedia of Math. and Its Appl.*, **16**).
- [JoSt87] J. T. Joichi and Dennis Stanton, Bijective proofs of basic hypergeometric series identities, *Pacific J. Math.*, vol. **127**, 1987, p. 103–120.
- [Kn70] Donald E. Knuth, Permutations, matrices, and generalized Young tableaux, *Pacific J. Math.*, vol. **34**, 1970, p. 709–727.
- [Kn73] Donald E. Knuth, *The Art of Computer Programming*, vol. 3, *Sorting and Searching*. Addison-Wesley, Reading, 1973.
- [La03] Alain Lascoux, *Symmetric Functions and Combinatorial Operators on Polynomials*. Amer. Math. Soc., Providence (CBMS no. **99**, Conference on Algebraic Combinatorics), 268 p., 2003.
- [LaSch81] Alain Lascoux and Marcel-Paul Schützenberger, Le monoïde plaxique, *Non-commutative Structures in Algebra and geometric Combinatorics* [A. de Luca, ed., Napoli. 1978], p. 129–156. Roma, Consiglio Nazionale delle Ricerche, 1981 (*Quaderni de “La Ricerca Scientifica”*, **109**).
- [LaSch84] Alain Lascoux and Marcel-Paul Schützenberger, Formulaire raisonné des fonctions symétriques, L.I.T.P., U.E.R. Math., Univ. Paris VII, 138 p., 1984.
- [Le60] D. H. Lehmer, Teaching combinatorial tricks to a computer, *Proc. Sympos. Appl. Math.*, vol. **10**, 1960, p. 179–193. Amer. Math. Soc., Providence, R.I.
- [Lod89] Jean-Louis Loday, Opérations sur l’homologie cyclique des algèbres commutatives, *Inv. Math.*, vol. **96**, 1989, p. 205–230.
- [Lo02] M. Lothaire, *Algebraic Combinatorics on Words*. Cambridge Univ. Press, 2002 (*Encyclopedia of Math. and its Appl.*, **90**).
- [Ma95] Ian G. Macdonald, *Symmetric Functions and Hall Polynomials*, Second Edition. Clarendon Press, Oxford, 1995.
- [Mac13] Percy Alexander MacMahon, The indices of permutations and the derivation therefrom of functions of a single variable associated with the permutations of any assemblage of objects, *Amer. J. Math.*, vol. **35**, 1913, p. 314–321.
- [Mac15] Percy Alexander MacMahon, *Combinatory Analysis*, vol. 1 and 2. Cambridge, Cambridge Univ. Press, 1915, (Reprinted by Chelsea, New York, 1955).
- [Mac78] Percy Alexander MacMahon, “Collected Papers, vol. 1” [G.E. Andrews, ed.], Cambridge, Mass., The M.I.T. Press, 1978.
- [Pr89] Robert A. Proctor, Equivalence of the Combinatorial and the Classical Definitions of Schur Functions, *J. Combin. Theory Ser. A*, vol. **51**, 1989, p. 135–137.
- [Ra79] Don P. Rawlings, Permutation and Multipermutation Statistics, Ph. D. thesis, Univ. Calif. San Diego. *Publ. I.R.M.A. Strasbourg*, 49/P-23, 1979.
- [Ra80] Don P. Rawlings, Generalized Worpitzky identities with applications to permutation enumeration, *Europ. J. Comb.*, vol. **2**, 1981, p. 67–78.
- [Ra84] Don P. Rawlings, Enumeration of permutations by descents, idescents, imajor index and basic components, *J. Combin. Theory, Ser. A*, vol. **36**, 1984, p. 1–14.
- [Ra90] Don P. Rawlings, Multicolored Simon Newcomb Problems, *J. Combin. Theory, Ser. A*, vol. **53**, 1990, p. 53–67.
- [Re93a] V. Reiner, Signed permutation statistics, *Europ. J. Combinatorics*, vol. **14**, 1993, p. 553–567.
- [Re93b] V. Reiner, Signed permutation statistics and cycle type, *Europ. J. Combinatorics*, vol. **14**, 1993, p. 569–579.
- [Re93c] V. Reiner, Upper binomial posets and signed permutation statistics, *Europ. J. Combinatorics*, vol. **14**, 1993, p. 581–588.
- [Re95a] V. Reiner, Descents and one-dimensional characters for classical Weyl groups, *Discrete Math.*, vol. **140**, 1995, p. 129–140.
- [Re95b] V. Reiner, The distribution of descents and length in a Coxeter group, *Electronic J. Combin.*, vol. **2**, 1995, # R25.
[<http://www.emis.de/journals/EJC/>].

REFERENCES

- [Rem83] Jeff B. Remmel, A bijective proof of a factorization theorem for (k, l) -hook Schur functions, *Combinatorics and Algebra* [Boulder, Colo., 1983], p. 253–287. *Contemp. Math.*, **34**, Amer. Math. Soc., Providence, R.I., 1984.
- [Rem84] Jeff B. Remmel, The combinatorics of (k, l) -hook Schur functions, *Cont. Math.*, vol. **34**, 1984, p. 253–287.
- [Rem87] Jeff B. Remmel, Permutation Statistics and (k, l) -hook Schur functions, *Discrete Math.*, vol. **67**, 1987, p. 271–298.
- [ReRo03a] Amitai Regev and Yuval Roichman, Permutation Statistics on the Alternating Group, Weizmann Institute, preprint 45 p.
- [ReRo03a] Amitai Regev and Yuval Roichman, q -Statistics on S_n and Pattern Avoidance, Weizmann Institute, preprint 39 p.
- [Ri58] John Riordan, *An Introduction to Combinatorial Analysis*. New York, J. Wiley, 1958.
- [Ro38] Gilbert de Beauregard Robinson, On the Representations of the Symmetric Group, *Amer. J. Math.*, vol. **60**, 1938, p. 745–760.
- [Ro74] David P. Roselle, Coefficients associated with the Expansion of certain Products, *Proc. Amer. Math. Soc.*, vol. **45**, 1974, p. 144–150.
- [Sc61] C. Schensted, Longest Increasing and Decreasing Sequences, *Canad. J. Math.*, vol. **13**, 1961, p. 179–192.
- [Sch63] Marcel-Paul Schützenberger, Quelques remarques sur une construction de Schensted, *Math. Scand.*, vol. **12**, 1963, p. 117–128.
- [Sch77] Marcel-Paul Schützenberger, La correspondance de Robinson, *Combinatoire et représentation du groupe symétrique* [Actes Table Ronde C.N.R.S., Strasbourg, 1976], p. 59–113. Berlin, Springer-Verlag, 1977 (*Lecture Notes in Math.*, **579**).
- [SW07] John Shareshian, Michelle L. Wachs, q -Eulerian polynomials: exceedance number and major inde, *Electronic Research Announcements of the Amer. Math. Soc.*, vol. **13**, 2007, p. 33–45.
- [Sl66] Lucy Joan Slater, *Generalized Hypergeometric Functions*. Cambridge Univ. Press, 1966.
- [St72] Richard P. Stanley, *Ordered Structures and Partitions*. Providence, R.I., Amer. Math. Soc., 1972 (*Memoirs Amer. Math. Soc.*, **119**).
- [St76] Richard P. Stanley, Binomial posets, Möbius inversion, and permutation enumeration, *J. Combin. Theory Ser. A*, vol. **20**, 1976, p. 336–356.
- [St86] Richard P. Stanley, *Enumerative Combinatorics*, vol. 1. Wadsworth & Brooks, 1986.
- [StE93] Einar Steingrímsson, Permutation Statistics of Indexed Permutations, *Europ. J. Combinatorics*, vol. **15**, 1994, p. 187–205.
- [StJ92] John Stembridge, Eulerian numbers, tableaux, and the Betti numbers of a toric variety, *Discrete Math.*, vol. **99**, 1992, p. 307–320.
- [Vi86] Xavier Gérard Viennot, Heaps of pieces, I: basic definitions and combinatorial lemmas, in *Combinatoire Énumérative* [G. Labelle and P. Leroux, eds. Montréal, 1985]. *Lecture Notes in Math.*, **1234**, Berlin, Springer-Verlag, 1986.
- [Wa92] Michelle L. Wachs, An involution for signed Eulerian numbers, *Discrete Math.*, vol. **99**, 1992, p. 59–62.
- [Ze80a] Doron Zeilberger, A lattice walk approach to the counting of multiset permutations, *J. Math. Anal. Appl.*, vol. **74**, 1980, p. 192–199.
- [Ze80b] Doron Zeilberger, Partial difference equations in $m_1 \geq m_2 \geq \dots \geq m_n$ and their applications to combinatorics, *Discrete Math.*, vol. **31**, 1980, p. 65–77.
- [Ze81] Doron Zeilberger, Enumeration of words by their number of mistakes, *Discrete Math.*, vol. **34**, 1981, p. 89–92.
- [ZeBr85] Doron Zeilberger and David M. Bressoud, A proof of Andrews’ q -Dyson conjecture, *Discrete Math.*, vol. **54**, 1985, p. 201–224.