

PROOF OF THE CONTINUED FRACTION CONJECTURE FOR

$$(a, b) = (z^3 + z^2 + 1, z + 1)$$

GUO-NIU HAN* AND YINING HU*

ABSTRACT. The continued fraction conjecture claims that the continued fraction defined by the (a, b) -Thue-Morse sequence is algebraic over $\mathbb{F}_2(x)$ for all pairs of distinct elements (a, b) in $\mathbb{F}_2[x] \setminus \mathbb{F}_2$. In this paper we prove the conjecture for $(a, b) = (z^3 + z^2 + 1, z + 1)$.

1. INTRODUCTION

Let $\mathbb{F}_2$ be the finite field of cardinality 2. Given a sequence of polynomials $\mathbf{a} = (a_0(z), a_1(z), a_2(z), \dots)$ of elements from $\mathbb{F}_2[z] \setminus \mathbb{F}_2$, we define the infinite continued fraction

$$(1.1) \quad \text{CF}(\mathbf{a}(z)) = \frac{1}{a_0(z) + \frac{1}{a_1(z) + \frac{1}{a_2(z) + \frac{1}{\ddots}}}}$$

as the limit of the finite continued fractions

$$(1.2) \quad \text{CF}_n(\mathbf{a}(z)) = \frac{1}{a_0(z) + \frac{1}{a_1(z) + \frac{1}{\ddots + \frac{1}{a_n(z)}}}} \in \mathbb{F}_2((1/z)).$$

Conversely, each power series in $\mathbb{F}_2[[1/z]]$ without constant term

$$(1.3) \quad c_{-1}z^{-1} + c_{-2}z^{-2} + c_{-3}z^{-3} + \dots$$

admits a continued fraction expansion of form (1.1).

Let a and b be two distinct elements from $\mathbb{F}_2[z] \setminus \mathbb{F}_2$. We consider the continued fraction (1.1) associated with the (a, b) -Thue-Morse sequence

$$\mathbf{t} = (t_0(z), t_1(z), t_2(z), \dots) = (a, b, b, a, b, a, a, b, \dots)$$

that we shall denote

$$(1.4) \quad \text{CF}(a, b) := \text{CF}(\mathbf{t}).$$

Date: 9/04/2020 00:12:09 .

2010 Mathematics Subject Classification. 11B85, 11J70, 11B50, 11Y65, 05A15.

Key words and phrases. automatic sequence, continued fraction, Thue-Morse sequence.

* This paper was automatically generated from a computer program developed by the authors, with 10.1 minutes CPU time used.

Theorem 1.1. *When $(a, b) = (z^3 + z^2 + 1, z + 1)$, the two power series $\text{CF}(a, b)$ and $\text{CF}(b, a)$ are algebraic over $\mathbb{F}_2(z)$, with minimal polynomials of the form*

$$p_4(z)y^4 + p_3(z)y^3 + p_2(z)y^2 + p_1(z)y + p_0(z) = 0$$

where, for $\text{CF}(a, b)$,

$$\begin{aligned} p_0(z) &= z^{12} + z^9 + z^8 + z^7 + z^6 + z^4 + z + 1, \\ p_1(z) &= z^{15} + z^{14} + z^{13} + z^{12} + z^{11} + z^6 + z^3 + z, \\ p_2(z) &= z^{16} + z^{14} + z^{12} + z^8 + z^6 + z^4 + z^2 + 1, \\ p_3(z) &= z^{15} + z^{14} + z^{13} + z^{12} + z^{11} + z^6 + z^3 + z, \\ p_4(z) &= z^{14} + z^{12} + z^9 + z^7 + z^2 + z + 1, \end{aligned}$$

and for $\text{CF}(b, a)$,

$$\begin{aligned} p_0(z) &= z^9 + z^7 + z^6 + z + 1, \\ p_1(z) &= z^{15} + z^{14} + z^{13} + z^{12} + z^{11} + z^6 + z^3 + z, \\ p_2(z) &= z^{16} + z^{14} + z^{12} + z^8 + z^6 + z^4 + z^2 + 1, \\ p_3(z) &= z^{15} + z^{14} + z^{13} + z^{12} + z^{11} + z^6 + z^3 + z, \\ p_4(z) &= z^{14} + z^9 + z^8 + z^7 + z^4 + z^2 + z + 1. \end{aligned}$$

2. PROOF

Define the sequences of polynomials $(P_n(z))_{n \geq 0}$ and $(Q_n(z))_{n \geq 0}$ by

$$\begin{pmatrix} P_n(z) & Q_n(z) \\ P_{n-1}(z) & Q_{n-1}(z) \end{pmatrix} := \begin{pmatrix} t_n(z) & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} t_{n-1}(z) & 1 \\ 1 & 0 \end{pmatrix} \cdots \begin{pmatrix} t_0(z) & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}.$$

By the basic properties of continued fractions, we have

$$(2.1) \quad \text{CF}_n(a, b) = \frac{P_n(z)}{Q_n(z)} \in \mathbb{F}_2((1/z)).$$

As formula (2.1) is not efficient for calculating $\text{CF}_n(a, b)$, a fast method is to be derived and described as follows. Define

$$\begin{aligned} M_n(x) &= x^{\deg(t_{2^n-1})} \begin{pmatrix} t_{2^n-1}(1/x) & 1 \\ 1 & 0 \end{pmatrix} \times x^{\deg(t_{2^n-2})} \begin{pmatrix} t_{2^n-2}(1/x) & 1 \\ 1 & 0 \end{pmatrix} \times \\ &\quad \cdots \times x^{\deg(t_0)} \begin{pmatrix} t_0(1/x) & 1 \\ 1 & 0 \end{pmatrix}, \\ W_n(x) &= x^{\deg(\bar{t}_{2^n-1})} \begin{pmatrix} \bar{t}_{2^n-1}(1/x) & 1 \\ 1 & 0 \end{pmatrix} \times x^{\deg(\bar{t}_{2^n-2})} \begin{pmatrix} \bar{t}_{2^n-2}(1/x) & 1 \\ 1 & 0 \end{pmatrix} \times \\ &\quad \cdots \times x^{\deg(\bar{t}_0)} \begin{pmatrix} \bar{t}_0(1/x) & 1 \\ 1 & 0 \end{pmatrix}, \end{aligned}$$

where $\bar{t}$ is the (b, a) -Thue-Morse sequence. By the properties of the Thue-Morse sequence, we have for $n \geq 0$

$$\begin{aligned} M_{n+1}(x) &= W_n(x) \cdot M_n(x), \\ W_{n+1}(x) &= M_n(x) \cdot W_n(x). \end{aligned}$$

Let $x := 1/z$. For each non-zero polynomial $P(z)$, define $\tilde{P}(x)$ to be $P(1/x)$. Then,

$$\text{CF}_n(a, b) = \frac{P_n(z)}{Q_n(z)} = \frac{\tilde{P}_n(x)}{\tilde{Q}_n(x)} \in \mathbb{F}_2((x)) = \mathbb{F}_2((1/z)).$$

Comparing the definition of $M_n(x)$ with the definition of $P_n(x)$ and $Q_n(x)$, we see that

$$\begin{aligned} M_n(x)_{0,1} &= x^{d_n} \tilde{P}_{2^n-1}(x), \\ M_n(x)_{0,0} &= x^{d_n} \tilde{Q}_{2^n-1}(x), \end{aligned}$$

for some positive integer d_n . Hence,

$$(2.2) \quad \text{CF}_{2^{2^n}-1}(a, b) = \frac{\tilde{P}_{2^{2^n}-1}(x)}{\tilde{Q}_{2^{2^n}-1}(x)} = \frac{M_{2^n}(x)_{0,1}}{M_{2^n}(x)_{0,0}}.$$

Therefore, by the convergence theorem of continued fraction, the algebraicity of $\text{CF}(a, b)$ will be established if it is shown that both $M_{2^n}(x)_{0,1}$ and $M_{2^n}(x)_{0,0}$ converge to algebraic series in $\mathbb{F}_2[[x]]$.

Actually, we will prove that for all $0 \leq i, j \leq 1$ the four sequences $(M_{2^n}(x)_{i,j})_n$, $(M_{2^{n+1}}(x)_{i,j})_n$, $(W_{2^n}(x)_{i,j})_n$, and $(W_{2^{n+1}}(x)_{i,j})_n$ converge to algebraic series in $\mathbb{F}_2[[x]]$. For this purpose, we define four 2×2 matrices M^e, M^o, W^e, W^o as follows: For each $T \in \{M^e, M^o, W^e, W^o\}$ and $0 \leq i, j \leq 1$, $T_{i,j}$ is defined to be the unique solution in $\mathbb{F}_2[[x]]$ of the polynomial $\phi(T, i, j)$ under certain initial conditions; the polynomials $\phi(T, i, j)$ and initial conditions are given in Section 3. We will prove that these four matrices, whose components are algebraic by definition, are the limits of $(M_{2^n}(x))_n$, $(M_{2^{n+1}}(x))_n$, $(W_{2^n}(x))_n$, and $(W_{2^{n+1}}(x))_n$.

Let us explain how the polynomials $\phi(T, i, j)$ and initial conditions are found, and why the solutions exist and are unique. For $0 \leq i, j \leq 1$, the coefficients of the polynomial $\phi(M^e, i, j)$ (resp. $\phi(M^o, i, j)$, $\phi(W^e, i, j)$, and $\phi(W^o, i, j)$) are the Padé-Hermite approximants of type

$$(100, 100, 100, 100, 100)$$

of the vector

$$(1, T^3, T^6, T^9, T^{12}),$$

where $T = M_{12,i,j}$ (resp. $M_{11,i,j}$, $W_{12,i,j}$, and $W_{11,i,j}$), found by the Derksen algorithm. We take the first 8 terms of $M_{12,i,j}$ (resp. $M_{11,i,j}$, $W_{12,i,j}$, and $W_{11,i,j}$) as initial conditions for $\phi(M^e, i, j)$ (resp. $\phi(M^o, i, j)$, $\phi(W^e, i, j)$, and $\phi(W^o, i, j)$). The following fact will be used to ensure that the solution exists and is unique: let $P(x, y) \in \mathbb{F}_2[x, y]$ and for each series $f(x) = \sum_{n=0}^{\infty} a_n x^n \in \mathbb{F}_2[[x]]$ denote the partial sum $\sum_{j=0}^{n-1} a_j x^j$ by $f_n(x)$ for $n \geq 0$. If for some $n \geq 0$ and $a_0, a_1, \dots, a_{n-1} \in \mathbb{F}_2$ $P(x, \sum_{j=0}^{n-1} a_j x^j) = O(x^n)$ and $Q(x, y) := P(x, \sum_{j=0}^{n-1} a_j x^j + x^n y)$ can be written as $x^m \sum_{j=0}^{\infty} q_j(x) y^j$ where $q_j(x)$ are polynomials for $j \geq 0$, $q_1(0) = 1$, and $q_j(0) = 0$ for $j > 1$, then there exists a unique solution $f(x) \in \mathbb{F}_2[[x]]$ of $P(x, f(x)) = 0$ that satisfies the initial condition $f_n(x) = \sum_{j=0}^{n-1} a_j x^j$.

We state two lemmas concerning the four matrices M^e, M^o, W^e, W^o . The first one is about relations between them; the second, about the structure of the each matrix.

Lemma 2.1. *We have*

$$(2.3) \quad M^e = W^o \cdot M^o,$$

$$(2.4) \quad M^o = W^e \cdot M^e,$$

$$(2.5) \quad W^e = M^o \cdot W^o,$$

$$(2.6) \quad W^o = M^e \cdot W^e.$$

Proof. We give the proof of the identity

$$M_{0,0}^e = W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o,$$

the proofs of the others being similar.

First, we compute the minimal polynomials of $W_{0,0}^o M_{0,0}^o$ and $W_{0,1}^o M_{1,0}^o$. We know that

$$P(x, y) = \text{Res}_z (\phi(W^o, 0, 0)(x, z), z^{12} \cdot \phi(M^o, 0, 0)(x, y/z))$$

is an annihilating polynomial of $W_{0,0}^o M_{0,0}^o$. We use Padé-Hermite approximation to find a candidate for the minimal polynomial of $W_{0,0}^o M_{0,0}^o$, that will be called $\phi_0(x, y)$. To prove that $\phi_0(x, y)$ is indeed the minimal polynomial, it suffices to prove that it is an irreducible factor of $P(x, y)$ of multiplicity m and that $Q(x, y) := P(x, y)/\phi_0(x, y)^m$ is not an annihilating polynomial of $W_{0,0}^o M_{0,0}^o$. We verify the first point directly. For the second point, we truncate $W_{0,0}^o M_{0,0}^o$ to order 360 and substitute it for y in $Q(x, y)$. We get a series of valuation less than 360, which proves that $Q(x, y)$ is not an annihilating polynomial of $W_{0,0}^o M_{0,0}^o$. We find the minimal polynomial $\phi_1(x, y)$ of $W_{0,1}^o M_{1,0}^o$ in a similar way.

Now we prove that $\phi(M^e, 0, 0)$ is the minimal polynomial of $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$. We know that

$$S(x, y) = \text{Res}_z (\phi_0(x, z), \phi_1(x, y + z))$$

is an annihilating of $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$. We verify that $\phi(M^e, 0, 0)$ is an irreducible factor of $S(x, y)$ of multiplicity μ , and that $Q(x, y) := S(x, y)/\phi(M^e, 0, 0)^\mu$ is not an annihilating polynomial of $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$. To see the last point, we truncate $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$ to order 440 and substitute it for y in $Q(x, y)$. We get a series of valuation less than 440, and therefore $Q(x, y)$ is not an annihilating polynomial of $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$.

Finally, the first 8 terms of $M_{0,0}^e$ and $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$ coincide. As these first terms determine a unique solution of $\phi(M^e, 0, 0)$, we know that the two series are one and the same. $\square$

Define

$$R^e = \begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix} \quad \text{and} \quad R^o = \begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix}.$$

Lemma 2.2. *For all integers $k \geq 2$ and $u = 2^{2k-1}$, the following identities hold.*

$$\begin{aligned} M^e[:8u] &= M^e[:4u] + x^{3u} M^e[:u] + x^{3u} M^e[:4u] + x^{6u} M^e[:u] + x^{5u} M^e[:3u] \\ &\quad + x^{6u} M^e[:2u] + (x^{4u} + x^{7u}) R^e, \\ W^e[:8u] &= W^e[:4u] + x^{3u} W^e[:u] + x^{3u} W^e[:4u] + x^{6u} W^e[:u] + x^{5u} W^e[:3u] \\ &\quad + x^{6u} W^e[:2u] + (x^{4u} + x^{7u}) R^e. \end{aligned}$$

For all integers $k \geq 2$ and $u = 2^{2k}$, the following identities hold.

$$M^o[:8u] = M^o[:4u] + x^{3u} M^o[:u] + x^{3u} M^o[:4u] + x^{6u} M^o[:u] + x^{5u} M^o[:3u]$$

$$\begin{aligned}
& + x^{6u}M^o[:2u] + (x^{4u} + x^{7u})R^o, \\
W^o[:8u] &= W^o[:4u] + x^{3u}W^o[:u] + x^{3u}W^o[:4u] + x^{6u}W^o[:u] + x^{5u}W^o[:3u] \\
& + x^{6u}W^o[:2u] + (x^{4u} + x^{7u})R^o.
\end{aligned}$$

Proof. To prove Lemma 2.2 we first construct an automaton for each sequence concerned, and then transform the conditions on infinitely many k 's into finitely many conditions on the states of the automaton. In the following, we will prove that for $T = M_{0,0}^o$, for all integer $k \geq 2$ and $u = 2^{2k}$,

$$\begin{aligned}
T[:8u] &= T[:4u] + x^{3u}T[:u] + x^{3u}T[:4u] + x^{6u}T[:u] + x^{5u}T[:3u] + x^{6u}T[:2u] \\
& + (x^{4u} + x^{7u}).
\end{aligned}$$

The proof of the other 15 cases are similar. We break down the above identity into 4 parts:

$$\begin{aligned}
x^4u &= x^{3u}T[u:2u] + T[4u:5u], \\
0 &= x^{5u}T[:u] + x^{3u}T[2u:3u] + T[5u:6u], \\
0 &= x^{5u}T[u:2u] + x^{3u}T[3u:4u] + T[6u:7u], \\
x^7u &= x^{6u}T[u:2u] + x^{5u}T[2u:3u] + T[7u:8u],
\end{aligned}$$

which can be rewritten as

$$\begin{aligned}
(2.7) \quad & 0 = T[[1w]_2] + T[[100w]_2], \\
(2.8) \quad & 0 = T[[w]_2] + T[[10w]_2] + T[[101w]_2], \\
(2.9) \quad & 0 = T[[1w]_2] + T[[11w]_2] + T[[110w]_2], \\
(2.10) \quad & 0 = T[[1w]_2] + T[[10w]_2] + T[[111w]_2],
\end{aligned}$$

for all binary word w of length $2k$ and $w \neq 0^{2k}$, and

$$\begin{aligned}
(2.11) \quad & 1 = T[[1w]_2] + T[[100w]_2], \\
(2.12) \quad & 0 = T[[w]_2] + T[[10w]_2] + T[[101w]_2], \\
(2.13) \quad & 0 = T[[1w]_2] + T[[11w]_2] + T[[110w]_2], \\
(2.14) \quad & 1 = T[[1w]_2] + T[[10w]_2] + T[[111w]_2],
\end{aligned}$$

for $w = 0^{2k}$.

First we calculate an 2-automaton that generates T from its minimal polynomial and its first terms. This automaton has 493 states; its transition function and output function can be found in the annex. Let $A(s, w)$ denote the state reached after reading w from right to left starting from the state s , and τ the output function. Define

$$E_{2k} = \{A(i, w) : |w| = 2k, w \neq 0^{2k}\}.$$

Identities (2.7) through (2.14) can be written as

$$\begin{aligned}
(2.15) \quad & 0 = \tau(A(s, 1)) + \tau(A(s, 100)), \\
(2.16) \quad & 0 = \tau(A(s, \epsilon)) + \tau(A(s, 10)) + \tau(A(s, 101)), \\
(2.17) \quad & 0 = \tau(A(s, 1)) + \tau(A(s, 11)) + \tau(A(s, 110)), \\
(2.18) \quad & 0 = \tau(A(s, 1)) + \tau(A(s, 10)) + \tau(A(s, 111)),
\end{aligned}$$

for all $s \in E_{2k}$, and

$$(2.19) \quad 1 = \tau(A(s, 1)) + \tau(A(s, 100)),$$

$$(2.20) \quad 0 = \tau(A(s, \epsilon)) + \tau(A(s, 10)) + \tau(A(s, 101)),$$

$$(2.21) \quad 0 = \tau(A(s, 1)) + \tau(A(s, 11)) + \tau(A(s, 110)),$$

$$(2.22) \quad 1 = \tau(A(s, 1)) + \tau(A(s, 10)) + \tau(A(s, 111)),$$

for $s = A(i, 0^{2k})$. We find that $(A(i, 0^{24}), E_{24}) = (A(i, 0^{12}), E_{12})$, so that we only have to verify that identities (2.15) through (2.22) hold for $2 \leq k \leq 12$, which turns out to be true. $\square$

In the following lemma, we express M_{2k} , M_{2k+1} , W_{2k} , and W_{2k+1} in terms of M^e , M^o , W^e , and W^o .

Lemma 2.3. *For all integer $k \geq 2$, and $u = 2^{2k-1}$,*

$$M_{2k} = M^e[:4u] + x^{3u}M^e[:u] + x^{4u}R^e,$$

$$W_{2k} = W^e[:4u] + x^{3u}W^e[:u] + x^{4u}R^e.$$

For all integer $k \geq 2$, and $u = 2^{2k}$,

$$M_{2k+1} = M^o[:4u] + x^{3u}M^o[:u] + x^{4u}R^o,$$

$$W_{2k+1} = W^o[:4u] + x^{3u}W^o[:u] + x^{4u}R^o.$$

Proof. We call the four identities in Lemma 2.3 also by the name M_{2k} , W_{2k} , M_{2k+1} , and W_{2k+1} . For $n = 2$, the identities can be verified directly. For $n \geq 2$, we claim that

$$\begin{aligned} M_{2k} \wedge W_{2k} &\Rightarrow M_{2k+1} \wedge W_{2k+1}, \\ M_{2k+1} \wedge W_{2k+1} &\Rightarrow M_{2k+2} \wedge W_{2k+2}. \end{aligned}$$

We give the proof of

$$M_{2k} \wedge W_{2k} \Rightarrow M_{2k+1},$$

the other ones being similar. Set $u = 2^{2k}$ and $v = 2^{2k-1}$. By definition and induction hypothesis, the left side of identity M_{2k+1} is equal to

$$(2.23) \quad \begin{aligned} W_{2k}M_{2k} &= (W^e[:4v] + x^{3v}W^e[:v] + x^{4u}R^e) \times (M^e[:4v] + x^{3v}M^e[:v] \\ &\quad + x^{4u}R^e); \end{aligned}$$

Call this expression *lhs*. Note that both sides of identity M_{2k+1} have the same term of highest degree $x^{8v}R^o$. Therefore we only need to prove that their difference is $O(x^{8v})$. Using Lemma 2.1 it can be seen that the right side of identity M_{2k+1} is congruent, modulo x^{8v} , to

$$W^e[:8v]M^e[:8v] + x^{6v}W^e[:2v]M^e[:2v].$$

For all $n \leq 8$, replace the occurrences of $W^e[:n \cdot v]$ and $M^e[:n \cdot v]$ in the above expression by the reduction modulo $x^{n \cdot v}$ of the right side of the corresponding identity in Lemma 2.2 and get a new expression, which we call *rhs*. Define

$$\begin{aligned} X &:= x^v, \\ a_n &:= W^e[n \cdot v : (n+1) \cdot v] / X^n, \\ b_n &:= M^e[n \cdot v : (n+1) \cdot v] / X^n, \\ c &:= R^e. \end{aligned}$$

Using the notation introduced above, we can represent the expressions *lhs* (2.23) and *rhs* as polynomials in $\mathbb{F}_2[a_1, \dots, a_8, b_1, \dots, b_8, c][X]$. Note that it is not a problem that a_j commutes with b_k while W^e does not commute with M^e , because in the expressions concerned, the products of W^e -terms and M^e -terms are always in the same order. We let the computer do the simplification and check that the difference between these two polynomials is indeed $O(X^8)$, which completes the proof. $\square$

Proof of Theorem 1.1. We prove the theorem for $\text{CF}(a, b)$; for $\text{CF}(b, a)$, the proof is similar. By Lemma 2.3, we have For all $0 \leq j, k \leq 1$,

$$\begin{aligned} \lim_{n \rightarrow \infty} M_{2n,j,k} &= M_{j,k}^e, \\ \lim_{n \rightarrow \infty} M_{2n+1,j,k} &= M_{j,k}^o, \\ \lim_{n \rightarrow \infty} W_{2n,j,k} &= W_{j,k}^e, \\ \lim_{n \rightarrow \infty} W_{2n+1,j,k} &= W_{j,k}^o. \end{aligned}$$

Let $z = 1/x$. By the convergence theorem and identity (2.2),

$$\text{CF}(a, b) = \frac{M_{0,1}^e(x)}{M_{0,0}^e(x)}.$$

By definition, $\phi(M^e, 0, 1)$ and $\phi(M^e, 0, 0)$ are minimal polynomials of $M_{0,1}^e$ and $M_{0,0}^e$. Therefore

$$P(x, y) = \text{Res}_t (\phi(M^e, 0, 1)(x, t), y^{12} \phi(M^e, 0, 0)(x, t/y))$$

is an annihilating polynomial of $f(x) = M_{0,1}^e/M_{0,0}^e$.

Define

$$Q(x, y) = q_4(x)y^4 + q_3(x)y^3 + q_2(x)y^2 + q_1(x)y + q_0(x),$$

where

$$\begin{aligned} q_0(x) &= x^{16} + x^{15} + x^{12} + x^{10} + x^9 + x^8 + x^7 + x^4, \\ q_1(x) &= x^{15} + x^{13} + x^{10} + x^5 + x^4 + x^3 + x^2 + x, \\ q_2(x) &= x^{16} + x^{14} + x^{12} + x^{10} + x^8 + x^4 + x^2 + 1, \\ q_3(x) &= x^{15} + x^{13} + x^{10} + x^5 + x^4 + x^3 + x^2 + x, \\ q_4(x) &= x^{16} + x^{15} + x^{14} + x^9 + x^7 + x^4 + x^2. \end{aligned}$$

The polynomial $Q(x, y)$ is the candidate for the minimal polynomial of $f(x)$ found by Padé-Hermite approximation. To prove that it is indeed the minimal polynomial of $f(x)$, we only need to prove that it is an irreducible factor of $P(x, y)$ of multiplicity m and $R(x, y) := P(x, y)/Q(x, y)^m$ is not an annihilating polynomial of $f(x)$. We verify the first point directly. For the second point, we find that when we truncate $f(x)$ to order 126, and substitute it for y in $R(z, y)$, we get a series with valuation smaller than 126, which proves that $R(z, y)$ is not an annihilating polynomial of $f(x)$. Finally, $z^{12}Q(1/z, y)$ is the minimal polynomial of $\text{CF}(a, b) = f(1/z)$. $\square$

3. ANNEXE

All 16 polynomials are of the form

$$p_0(x) + p_3(x)y^3 + p_6(x)y^6 + p_9(x)y^9 + p_{12}(x)y^{12}.$$

The coefficients $p_j(x)$ and the 8 initial terms to determine the solutions uniquely are given below.

For $\phi(M^e, 0, 0)$:

$$\begin{aligned}
p_0(x) &= x^{78} + x^{76} + x^{74} + x^{72} + x^{70} + x^{62} + x^{60} + x^{56} + x^{54} + x^{50} + x^{46} \\
&\quad + x^{44} + x^{40} + x^{38} + x^{36} + x^{34} + x^{26} + x^{24} + x^{16} + x^{14} + x^{12}, \\
p_3(x) &= x^{76} + x^{74} + x^{72} + x^{70} + x^{68} + x^{64} + x^{62} + x^{60} + x^{58} + x^{56} + x^{52} \\
&\quad + x^{50} + x^{40} + x^{34} + x^{30} + x^{28} + x^{26} + x^{22} + x^{18} + x^{14} + x^{10} + x^8, \\
p_6(x) &= x^{76} + x^{72} + x^{70} + x^{66} + x^{64} + x^{60} + x^{58} + x^{52} + x^{50} + x^{48} + x^{44} \\
&\quad + x^{42} + x^{28} + x^{22} + x^{18} + x^{12} + x^8 + x^6 + x^4 + 1, \\
p_9(x) &= x^{76} + x^{74} + x^{70} + x^{68} + x^{66} + x^{64} + x^{58} + x^{56} + x^{52} + x^{48} + x^{46} \\
&\quad + x^{44} + x^{42} + x^{34} + x^{20} + x^{14} + x^8 + x^6 + x^4 + x^2, \\
p_{12}(x) &= x^{78} + x^{74} + x^{72} + x^{70} + x^{66} + x^{64} + x^{30} + x^{26} + x^{24} + x^{22} + x^{18} \\
&\quad + x^{16} + x^{14} + x^{10} + x^8 + x^6 + x^2 + 1,
\end{aligned}$$

and the initial terms are $[1, 0, 1, 0, 0, 0, 0, 0]$.

For $\phi(M^e, 0, 1)$:

$$\begin{aligned}
p_0(x) &= x^{84} + x^{83} + x^{82} + x^{81} + x^{77} + x^{76} + x^{75} + x^{74} + x^{73} + x^{70} + x^{69} \\
&\quad + x^{68} + x^{67} + x^{65} + x^{63} + x^{62} + x^{59} + x^{58} + x^{57} + x^{55} + x^{53} + x^{51} \\
&\quad + x^{48} + x^{47} + x^{43} + x^{42} + x^{39} + x^{36} + x^{35} + x^{29} + x^{28} + x^{27} + x^{25} \\
&\quad + x^{23} + x^{18}, \\
p_3(x) &= x^{75} + x^{74} + x^{72} + x^{70} + x^{68} + x^{67} + x^{65} + x^{64} + x^{63} + x^{59} + x^{58} \\
&\quad + x^{56} + x^{55} + x^{51} + x^{50} + x^{48} + x^{47} + x^{39} + x^{38} + x^{36} + x^{34} + x^{33} \\
&\quad + x^{27} + x^{26} + x^{24} + x^{22} + x^{20} + x^{18} + x^{17} + x^{15} + x^{14} + x^{12} + x^{10} \\
&\quad + x^9, \\
p_6(x) &= x^{78} + x^{77} + x^{76} + x^{74} + x^{68} + x^{67} + x^{65} + x^{64} + x^{63} + x^{59} + x^{57} \\
&\quad + x^{56} + x^{55} + x^{54} + x^{53} + x^{49} + x^{48} + x^{45} + x^{44} + x^{42} + x^{40} + x^{39} \\
&\quad + x^{38} + x^{36} + x^{28} + x^{27} + x^{26} + x^{23} + x^{22} + x^{19} + x^{18} + x^{16} + x^{12} \\
&\quad + x^{11} + x^9 + x^6, \\
p_9(x) &= x^{81} + x^{80} + x^{78} + x^{77} + x^{69} + x^{68} + x^{67} + x^{65} + x^{64} + x^{63} + x^{59} \\
&\quad + x^{58} + x^{56} + x^{55} + x^{53} + x^{52} + x^{51} + x^{49} + x^{48} + x^{46} + x^{44} + x^{42} \\
&\quad + x^{41} + x^{35} + x^{34} + x^{32} + x^{30} + x^{28} + x^{26} + x^{25} + x^{23} + x^{22} + x^{20} \\
&\quad + x^{18} + x^{16} + x^{15} + x^{13} + x^{12} + x^{10} + x^8 + x^7 + x^5 + x^4 + x^3, \\
p_{12}(x) &= x^{84} + x^{83} + x^{82} + x^{79} + x^{78} + x^{75} + x^{74} + x^{72} + x^{68} + x^{67} + x^{66} \\
&\quad + x^{64} + x^{36} + x^{35} + x^{34} + x^{31} + x^{30} + x^{27} + x^{26} + x^{24} + x^{16} + x^{15} \\
&\quad + x^{14} + x^{11} + x^{10} + x^8 + x^4 + x^3 + x^2 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 1, 1, 0, 1, 1]$.

For $\phi(M^e, 1, 0)$:

$$\begin{aligned} p_0(x) = & x^{84} + x^{83} + x^{82} + x^{81} + x^{77} + x^{76} + x^{75} + x^{74} + x^{73} + x^{70} + x^{69} \\ & + x^{68} + x^{67} + x^{65} + x^{63} + x^{62} + x^{59} + x^{58} + x^{57} + x^{55} + x^{53} + x^{51} \\ & + x^{48} + x^{47} + x^{43} + x^{42} + x^{39} + x^{36} + x^{35} + x^{29} + x^{28} + x^{27} + x^{25} \\ & + x^{23} + x^{18}, \end{aligned}$$

$$\begin{aligned} p_3(x) = & x^{75} + x^{74} + x^{72} + x^{70} + x^{68} + x^{67} + x^{65} + x^{64} + x^{63} + x^{59} + x^{58} \\ & + x^{56} + x^{55} + x^{51} + x^{50} + x^{48} + x^{47} + x^{39} + x^{38} + x^{36} + x^{34} + x^{33} \\ & + x^{27} + x^{26} + x^{24} + x^{22} + x^{20} + x^{18} + x^{17} + x^{15} + x^{14} + x^{12} + x^{10} \\ & + x^9, \end{aligned}$$

$$\begin{aligned} p_6(x) = & x^{78} + x^{77} + x^{76} + x^{74} + x^{68} + x^{67} + x^{65} + x^{64} + x^{63} + x^{59} + x^{57} \\ & + x^{56} + x^{55} + x^{54} + x^{53} + x^{49} + x^{48} + x^{45} + x^{44} + x^{42} + x^{40} + x^{39} \\ & + x^{38} + x^{36} + x^{28} + x^{27} + x^{26} + x^{23} + x^{22} + x^{19} + x^{18} + x^{16} + x^{12} \\ & + x^{11} + x^9 + x^6, \end{aligned}$$

$$\begin{aligned} p_9(x) = & x^{81} + x^{80} + x^{78} + x^{77} + x^{69} + x^{68} + x^{67} + x^{65} + x^{64} + x^{63} + x^{59} \\ & + x^{58} + x^{56} + x^{55} + x^{53} + x^{52} + x^{51} + x^{49} + x^{48} + x^{46} + x^{44} + x^{42} \\ & + x^{41} + x^{35} + x^{34} + x^{32} + x^{30} + x^{28} + x^{26} + x^{25} + x^{23} + x^{22} + x^{20} \\ & + x^{18} + x^{16} + x^{15} + x^{13} + x^{12} + x^{10} + x^8 + x^7 + x^5 + x^4 + x^3, \end{aligned}$$

$$\begin{aligned} p_{12}(x) = & x^{84} + x^{83} + x^{82} + x^{79} + x^{78} + x^{75} + x^{74} + x^{72} + x^{68} + x^{67} + x^{66} \\ & + x^{64} + x^{36} + x^{35} + x^{34} + x^{31} + x^{30} + x^{27} + x^{26} + x^{24} + x^{16} + x^{15} \\ & + x^{14} + x^{11} + x^{10} + x^8 + x^4 + x^3 + x^2 + 1, \end{aligned}$$

and the initial terms are $[0, 0, 0, 1, 1, 0, 1, 1]$.

For $\phi(M^e, 1, 1)$:

$$\begin{aligned} p_0(x) = & x^{90} + x^{82} + x^{80} + x^{76} + x^{74} + x^{72} + x^{68} + x^{66} + x^{64} + x^{54} + x^{52} \\ & + x^{40} + x^{38} + x^{36} + x^{34} + x^{32} + x^{30} + x^{26} + x^{24}, \end{aligned}$$

$$\begin{aligned} p_3(x) = & x^{88} + x^{80} + x^{68} + x^{64} + x^{60} + x^{54} + x^{52} + x^{50} + x^{48} + x^{46} + x^{44} \\ & + x^{40} + x^{38} + x^{30} + x^{26} + x^{20} + x^{18} + x^{16} + x^8 + x^6, \end{aligned}$$

$$\begin{aligned} p_6(x) = & x^{88} + x^{80} + x^{76} + x^{66} + x^{62} + x^{56} + x^{54} + x^{38} + x^{34} + x^{32} + x^{28} \\ & + x^{22} + x^{18} + x^{12} + x^{10} + 1, \end{aligned}$$

$$\begin{aligned} p_9(x) = & x^{88} + x^{80} + x^{74} + x^{68} + x^{64} + x^{62} + x^{60} + x^{52} + x^{46} + x^{44} + x^{36} \\ & + x^{34} + x^{30} + x^{26} + x^{24} + x^{22} + x^{20} + x^{16} + x^{12} + x^6 + x^4 + x^2, \end{aligned}$$

$$\begin{aligned} p_{12}(x) = & x^{90} + x^{88} + x^{74} + x^{72} + x^{66} + x^{64} + x^{42} + x^{40} + x^{18} + x^{16} + x^{10} + x^8 \\ & + x^2 + 1, \end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 0, 1, 0]$.

For $\phi(W^e, 0, 0)$:

$$p_0(x) = x^{90} + x^{86} + x^{84} + x^{82} + x^{74} + x^{70} + x^{66} + x^{64} + x^{58} + x^{56} + x^{52}$$

$$\begin{aligned}
& + x^{42} + x^{40} + x^{38} + x^{20} + x^{14} + x^{12}, \\
p_3(x) &= x^{88} + x^{84} + x^{82} + x^{80} + x^{74} + x^{72} + x^{70} + x^{66} + x^{64} + x^{62} + x^{60} \\
& + x^{58} + x^{56} + x^{48} + x^{34} + x^{32} + x^{30} + x^{28} + x^{22} + x^{20} + x^{14} + x^8, \\
p_6(x) &= x^{88} + x^{86} + x^{82} + x^{80} + x^{78} + x^{74} + x^{72} + x^{66} + x^{64} + x^{58} + x^{54} \\
& + x^{48} + x^{40} + x^{38} + x^{36} + x^{34} + x^{32} + x^{24} + x^{16} + x^{14} + x^{12} + x^8 \\
& + x^4 + 1, \\
p_9(x) &= x^{88} + x^{80} + x^{74} + x^{68} + x^{64} + x^{62} + x^{60} + x^{52} + x^{46} + x^{44} + x^{36} \\
& + x^{34} + x^{30} + x^{26} + x^{24} + x^{22} + x^{20} + x^{16} + x^{12} + x^6 + x^4 + x^2, \\
p_{12}(x) &= x^{90} + x^{88} + x^{74} + x^{72} + x^{66} + x^{64} + x^{42} + x^{40} + x^{18} + x^{16} + x^{10} + x^8 \\
& + x^2 + 1,
\end{aligned}$$

and the initial terms are $[1, 0, 0, 0, 1, 0, 1, 0]$.

For $\phi(W^e, 0, 1)$:

$$\begin{aligned}
p_0(x) &= x^{84} + x^{83} + x^{82} + x^{81} + x^{78} + x^{75} + x^{74} + x^{72} + x^{70} + x^{69} + x^{64} \\
& + x^{59} + x^{57} + x^{54} + x^{52} + x^{50} + x^{49} + x^{48} + x^{47} + x^{45} + x^{42} + x^{41} \\
& + x^{39} + x^{38} + x^{32} + x^{31} + x^{27}, \\
p_3(x) &= x^{75} + x^{74} + x^{72} + x^{70} + x^{68} + x^{67} + x^{65} + x^{64} + x^{63} + x^{59} + x^{58} \\
& + x^{56} + x^{55} + x^{51} + x^{50} + x^{48} + x^{47} + x^{39} + x^{38} + x^{36} + x^{34} + x^{33} \\
& + x^{27} + x^{26} + x^{24} + x^{22} + x^{20} + x^{18} + x^{17} + x^{15} + x^{14} + x^{12} + x^{10} \\
& + x^9, \\
p_6(x) &= x^{78} + x^{77} + x^{76} + x^{74} + x^{68} + x^{67} + x^{65} + x^{64} + x^{63} + x^{59} + x^{57} \\
& + x^{56} + x^{55} + x^{54} + x^{53} + x^{49} + x^{48} + x^{45} + x^{44} + x^{42} + x^{40} + x^{39} \\
& + x^{38} + x^{36} + x^{28} + x^{27} + x^{26} + x^{23} + x^{22} + x^{19} + x^{18} + x^{16} + x^{12} \\
& + x^{11} + x^9 + x^6, \\
p_9(x) &= x^{81} + x^{80} + x^{78} + x^{77} + x^{69} + x^{68} + x^{67} + x^{65} + x^{64} + x^{63} + x^{59} \\
& + x^{58} + x^{56} + x^{55} + x^{53} + x^{52} + x^{51} + x^{49} + x^{48} + x^{46} + x^{44} + x^{42} \\
& + x^{41} + x^{35} + x^{34} + x^{32} + x^{30} + x^{28} + x^{26} + x^{25} + x^{23} + x^{22} + x^{20} \\
& + x^{18} + x^{16} + x^{15} + x^{13} + x^{12} + x^{10} + x^8 + x^7 + x^5 + x^4 + x^3, \\
p_{12}(x) &= x^{84} + x^{83} + x^{82} + x^{79} + x^{78} + x^{75} + x^{74} + x^{72} + x^{68} + x^{67} + x^{66} \\
& + x^{64} + x^{36} + x^{35} + x^{34} + x^{31} + x^{30} + x^{27} + x^{26} + x^{24} + x^{16} + x^{15} \\
& + x^{14} + x^{11} + x^{10} + x^8 + x^4 + x^3 + x^2 + 1,
\end{aligned}$$

and the initial terms are $[0, 1, 1, 1, 1, 1, 1, 1]$.

For $\phi(W^e, 1, 0)$:

$$\begin{aligned}
p_0(x) &= x^{84} + x^{83} + x^{82} + x^{81} + x^{78} + x^{75} + x^{74} + x^{72} + x^{70} + x^{69} + x^{64} \\
& + x^{59} + x^{57} + x^{54} + x^{52} + x^{50} + x^{49} + x^{48} + x^{47} + x^{45} + x^{42} + x^{41} \\
& + x^{39} + x^{38} + x^{32} + x^{31} + x^{27},
\end{aligned}$$

$$\begin{aligned}
p_3(x) &= x^{75} + x^{74} + x^{72} + x^{70} + x^{68} + x^{67} + x^{65} + x^{64} + x^{63} + x^{59} + x^{58} \\
&\quad + x^{56} + x^{55} + x^{51} + x^{50} + x^{48} + x^{47} + x^{39} + x^{38} + x^{36} + x^{34} + x^{33} \\
&\quad + x^{27} + x^{26} + x^{24} + x^{22} + x^{20} + x^{18} + x^{17} + x^{15} + x^{14} + x^{12} + x^{10} \\
&\quad + x^9, \\
p_6(x) &= x^{78} + x^{77} + x^{76} + x^{74} + x^{68} + x^{67} + x^{65} + x^{64} + x^{63} + x^{59} + x^{57} \\
&\quad + x^{56} + x^{55} + x^{54} + x^{53} + x^{49} + x^{48} + x^{45} + x^{44} + x^{42} + x^{40} + x^{39} \\
&\quad + x^{38} + x^{36} + x^{28} + x^{27} + x^{26} + x^{23} + x^{22} + x^{19} + x^{18} + x^{16} + x^{12} \\
&\quad + x^{11} + x^9 + x^6, \\
p_9(x) &= x^{81} + x^{80} + x^{78} + x^{77} + x^{69} + x^{68} + x^{67} + x^{65} + x^{64} + x^{63} + x^{59} \\
&\quad + x^{58} + x^{56} + x^{55} + x^{53} + x^{52} + x^{51} + x^{49} + x^{48} + x^{46} + x^{44} + x^{42} \\
&\quad + x^{41} + x^{35} + x^{34} + x^{32} + x^{30} + x^{28} + x^{26} + x^{25} + x^{23} + x^{22} + x^{20} \\
&\quad + x^{18} + x^{16} + x^{15} + x^{13} + x^{12} + x^{10} + x^8 + x^7 + x^5 + x^4 + x^3, \\
p_{12}(x) &= x^{84} + x^{83} + x^{82} + x^{79} + x^{78} + x^{75} + x^{74} + x^{72} + x^{68} + x^{67} + x^{66} \\
&\quad + x^{64} + x^{36} + x^{35} + x^{34} + x^{31} + x^{30} + x^{27} + x^{26} + x^{24} + x^{16} + x^{15} \\
&\quad + x^{14} + x^{11} + x^{10} + x^8 + x^4 + x^3 + x^2 + 1,
\end{aligned}$$

and the initial terms are $[0, 1, 1, 1, 1, 1, 1, 1]$.

For $\phi(W^e, 1, 1)$:

$$\begin{aligned}
p_0(x) &= x^{78} + x^{76} + x^{72} + x^{64} + x^{62} + x^{60} + x^{56} + x^{52} + x^{50} + x^{48} + x^{46} \\
&\quad + x^{44} + x^{42}, \\
p_3(x) &= x^{76} + x^{74} + x^{70} + x^{66} + x^{56} + x^{54} + x^{50} + x^{48} + x^{46} + x^{42} + x^{40} \\
&\quad + x^{34} + x^{32} + x^{30} + x^{28} + x^{22} + x^{18} + x^{14} + x^{12} + x^{10} + x^8 + x^6, \\
p_6(x) &= x^{76} + x^{74} + x^{62} + x^{58} + x^{54} + x^{52} + x^{48} + x^{44} + x^{42} + x^{40} + x^{34} \\
&\quad + x^{32} + x^{24} + x^{22} + x^{18} + x^{14} + x^6 + 1, \\
p_9(x) &= x^{76} + x^{74} + x^{70} + x^{68} + x^{66} + x^{64} + x^{58} + x^{56} + x^{52} + x^{48} + x^{46} \\
&\quad + x^{44} + x^{42} + x^{34} + x^{20} + x^{14} + x^8 + x^6 + x^4 + x^2, \\
p_{12}(x) &= x^{78} + x^{74} + x^{72} + x^{70} + x^{66} + x^{64} + x^{30} + x^{26} + x^{24} + x^{22} + x^{18} \\
&\quad + x^{16} + x^{14} + x^{10} + x^8 + x^6 + x^2 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 1, 0, 1, 0, 0, 0]$.

For $\phi(M^o, 0, 0)$:

$$\begin{aligned}
p_0(x) &= x^{84} + x^{83} + x^{81} + x^{79} + x^{78} + x^{76} + x^{74} + x^{73} + x^{71} + x^{68} + x^{66} \\
&\quad + x^{62} + x^{61} + x^{60} + x^{57} + x^{56} + x^{55} + x^{51} + x^{48} + x^{47} + x^{45} + x^{44} \\
&\quad + x^{43} + x^{38} + x^{34} + x^{33} + x^{32} + x^{27} + x^{25} + x^{22} + x^{20} + x^{17} + x^{16} \\
&\quad + x^{15} + x^{12}, \\
p_3(x) &= x^{80} + x^{79} + x^{77} + x^{76} + x^{74} + x^{73} + x^{65} + x^{63} + x^{60} + x^{59} + x^{55} \\
&\quad + x^{54} + x^{52} + x^{51} + x^{50} + x^{49} + x^{46} + x^{44} + x^{43} + x^{42} + x^{40} + x^{39}
\end{aligned}$$

$$\begin{aligned}
& + x^{36} + x^{35} + x^{33} + x^{32} + x^{31} + x^{30} + x^{29} + x^{28} + x^{23} + x^{15} + x^{14} \\
& + x^{13} + x^{11} + x^8, \\
p_6(x) &= x^{82} + x^{81} + x^{76} + x^{75} + x^{73} + x^{72} + x^{69} + x^{67} + x^{66} + x^{63} + x^{62} \\
& + x^{57} + x^{55} + x^{53} + x^{52} + x^{51} + x^{50} + x^{48} + x^{47} + x^{46} + x^{45} + x^{43} \\
& + x^{40} + x^{38} + x^{37} + x^{34} + x^{32} + x^{30} + x^{29} + x^{26} + x^{22} + x^{21} + x^{20} \\
& + x^{19} + x^{18} + x^{17} + x^{14} + x^{12} + x^9 + x^8 + x^7 + x^3 + x^2 + 1, \\
p_9(x) &= x^{80} + x^{79} + x^{78} + x^{75} + x^{74} + x^{71} + x^{70} + x^{68} + x^{32} + x^{31} + x^{30} \\
& + x^{27} + x^{26} + x^{23} + x^{22} + x^{20} + x^{16} + x^{15} + x^{14} + x^{11} + x^{10} + x^7 \\
& + x^6 + x^4, \\
p_{12}(x) &= x^{84} + x^{83} + x^{82} + x^{80} + x^{72} + x^{71} + x^{70} + x^{67} + x^{66} + x^{64} + x^{36} \\
& + x^{35} + x^{34} + x^{32} + x^{24} + x^{23} + x^{22} + x^{20} + x^8 + x^7 + x^6 + x^3 + x^2 \\
& + 1,
\end{aligned}$$

and the initial terms are $[1, 0, 1, 0, 0, 0, 0, 1]$.

For $\phi(M^o, 0, 1)$:

$$\begin{aligned}
p_0(x) &= x^{92} + x^{89} + x^{88} + x^{86} + x^{84} + x^{82} + x^{81} + x^{79} + x^{75} + x^{73} + x^{71} \\
& + x^{69} + x^{68} + x^{67} + x^{65} + x^{62} + x^{61} + x^{60} + x^{59} + x^{56} + x^{55} + x^{54} \\
& + x^{52} + x^{50} + x^{49} + x^{46} + x^{44} + x^{42} + x^{41} + x^{38} + x^{36} + x^{33} + x^{29} \\
& + x^{27} + x^{26} + x^{24} + x^{21} + x^{18}, \\
p_3(x) &= x^{83} + x^{78} + x^{76} + x^{74} + x^{71} + x^{70} + x^{69} + x^{66} + x^{64} + x^{63} + x^{62} \\
& + x^{57} + x^{56} + x^{55} + x^{54} + x^{51} + x^{49} + x^{48} + x^{47} + x^{44} + x^{42} + x^{41} \\
& + x^{40} + x^{37} + x^{35} + x^{34} + x^{33} + x^{30} + x^{28} + x^{27} + x^{26} + x^{23} + x^{21} \\
& + x^{20} + x^{16} + x^{13} + x^{10} + x^9, \\
p_6(x) &= x^{86} + x^{78} + x^{76} + x^{72} + x^{64} + x^{54} + x^{52} + x^{48} + x^{42} + x^{40} + x^{32} \\
& + x^{20} + x^{18} + x^{16} + x^{12} + x^{10} + x^8 + x^6, \\
p_9(x) &= x^{89} + x^{84} + x^{82} + x^{80} + x^{79} + x^{77} + x^{76} + x^{74} + x^{73} + x^{71} + x^{70} \\
& + x^{69} + x^{68} + x^{67} + x^{41} + x^{36} + x^{34} + x^{32} + x^{31} + x^{29} + x^{28} + x^{26} \\
& + x^{23} + x^{22} + x^{21} + x^{19} + x^{18} + x^{16} + x^{15} + x^{13} + x^{12} + x^{10} + x^9 \\
& + x^7 + x^6 + x^5 + x^4 + x^3, \\
p_{12}(x) &= x^{92} + x^{84} + x^{76} + x^{72} + x^{64} + x^{44} + x^{36} + x^{24} + x^{20} + x^{16} + x^{12} + x^8 \\
& + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 1, 1, 0, 1, 1]$.

For $\phi(M^o, 1, 0)$:

$$\begin{aligned}
p_0(x) &= x^{84} + x^{81} + x^{77} + x^{71} + x^{68} + x^{66} + x^{64} + x^{62} + x^{60} + x^{57} + x^{55} \\
& + x^{54} + x^{53} + x^{51} + x^{50} + x^{48} + x^{47} + x^{46} + x^{45} + x^{44} + x^{43} + x^{42} \\
& + x^{41} + x^{40} + x^{38} + x^{36} + x^{35} + x^{34} + x^{33} + x^{30} + x^{27}, \\
p_3(x) &= x^{75} + x^{71} + x^{70} + x^{68} + x^{67} + x^{64} + x^{63} + x^{62} + x^{61} + x^{60} + x^{59}
\end{aligned}$$

$$\begin{aligned}
& + x^{58} + x^{57} + x^{56} + x^{55} + x^{54} + x^{53} + x^{52} + x^{51} + x^{50} + x^{49} + x^{48} \\
& + x^{47} + x^{46} + x^{45} + x^{44} + x^{42} + x^{41} + x^{40} + x^{37} + x^{34} + x^{33} + x^{27} \\
& + x^{23} + x^{22} + x^{20} + x^{16} + x^{13} + x^{10} + x^9, \\
p_6(x) &= x^{78} + x^{74} + x^{68} + x^{66} + x^{62} + x^{58} + x^{52} + x^{42} + x^{40} + x^{36} + x^{28} \\
& + x^{16} + x^{12} + x^{10} + x^8 + x^6, \\
p_9(x) &= x^{81} + x^{77} + x^{76} + x^{74} + x^{73} + x^{71} + x^{70} + x^{69} + x^{68} + x^{67} + x^{33} \\
& + x^{29} + x^{28} + x^{26} + x^{25} + x^{23} + x^{22} + x^{21} + x^{20} + x^{19} + x^{17} + x^{13} \\
& + x^{12} + x^{10} + x^9 + x^7 + x^6 + x^5 + x^4 + x^3, \\
p_{12}(x) &= x^{84} + x^{80} + x^{72} + x^{64} + x^{36} + x^{32} + x^{24} + x^{20} + x^8 + 1,
\end{aligned}$$

and the initial terms are $[0, 1, 1, 0, 0, 1, 1, 1]$.

For $\phi(M^o, 1, 1)$:

$$\begin{aligned}
p_0(x) &= x^{84} + x^{83} + x^{81} + x^{80} + x^{78} + x^{77} + x^{75} + x^{72} + x^{71} + x^{70} + x^{69} \\
& + x^{66} + x^{65} + x^{63} + x^{61} + x^{60} + x^{59} + x^{58} + x^{56} + x^{55} + x^{52} + x^{51} \\
& + x^{48} + x^{47} + x^{45} + x^{43} + x^{41} + x^{39} + x^{38} + x^{36} + x^{33}, \\
p_3(x) &= x^{80} + x^{79} + x^{77} + x^{75} + x^{74} + x^{72} + x^{71} + x^{70} + x^{69} + x^{64} + x^{63} \\
& + x^{62} + x^{57} + x^{55} + x^{50} + x^{44} + x^{43} + x^{42} + x^{37} + x^{33} + x^{31} + x^{30} \\
& + x^{29} + x^{27} + x^{25} + x^{24} + x^{22} + x^{21} + x^{20} + x^{19} + x^{14} + x^{12}, \\
p_6(x) &= x^{82} + x^{81} + x^{80} + x^{79} + x^{77} + x^{76} + x^{75} + x^{74} + x^{73} + x^{71} + x^{70} \\
& + x^{69} + x^{68} + x^{67} + x^{65} + x^{64} + x^{63} + x^{60} + x^{59} + x^{55} + x^{51} + x^{50} \\
& + x^{49} + x^{48} + x^{44} + x^{38} + x^{37} + x^{36} + x^{35} + x^{32} + x^{31} + x^{29} + x^{27} \\
& + x^{24} + x^{22} + x^{21} + x^{17} + x^{16} + x^{14} + x^{11} + x^4 + x^3 + x^2 + 1, \\
p_9(x) &= x^{80} + x^{79} + x^{78} + x^{75} + x^{74} + x^{71} + x^{70} + x^{68} + x^{32} + x^{31} + x^{30} \\
& + x^{27} + x^{26} + x^{23} + x^{22} + x^{20} + x^{16} + x^{15} + x^{14} + x^{11} + x^{10} + x^7 \\
& + x^6 + x^4, \\
p_{12}(x) &= x^{84} + x^{83} + x^{82} + x^{80} + x^{72} + x^{71} + x^{70} + x^{67} + x^{66} + x^{64} + x^{36} \\
& + x^{35} + x^{34} + x^{32} + x^{24} + x^{23} + x^{22} + x^{20} + x^8 + x^7 + x^6 + x^3 + x^2 \\
& + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 1, 0, 0, 1]$.

For $\phi(W^o, 0, 0)$:

$$\begin{aligned}
p_0(x) &= x^{84} + x^{83} + x^{81} + x^{79} + x^{78} + x^{76} + x^{74} + x^{73} + x^{71} + x^{68} + x^{66} \\
& + x^{62} + x^{61} + x^{60} + x^{57} + x^{56} + x^{55} + x^{51} + x^{48} + x^{47} + x^{45} + x^{44} \\
& + x^{43} + x^{38} + x^{34} + x^{33} + x^{32} + x^{27} + x^{25} + x^{22} + x^{20} + x^{17} + x^{16} \\
& + x^{15} + x^{12}, \\
p_3(x) &= x^{80} + x^{79} + x^{77} + x^{76} + x^{74} + x^{73} + x^{65} + x^{63} + x^{60} + x^{59} + x^{55} \\
& + x^{54} + x^{52} + x^{51} + x^{50} + x^{49} + x^{46} + x^{44} + x^{43} + x^{42} + x^{40} + x^{39}
\end{aligned}$$

$$\begin{aligned}
& + x^{36} + x^{35} + x^{33} + x^{32} + x^{31} + x^{30} + x^{29} + x^{28} + x^{23} + x^{15} + x^{14} \\
& + x^{13} + x^{11} + x^8, \\
p_6(x) &= x^{82} + x^{81} + x^{76} + x^{75} + x^{73} + x^{72} + x^{69} + x^{67} + x^{66} + x^{63} + x^{62} \\
& + x^{57} + x^{55} + x^{53} + x^{52} + x^{51} + x^{50} + x^{48} + x^{47} + x^{46} + x^{45} + x^{43} \\
& + x^{40} + x^{38} + x^{37} + x^{34} + x^{32} + x^{30} + x^{29} + x^{26} + x^{22} + x^{21} + x^{20} \\
& + x^{19} + x^{18} + x^{17} + x^{14} + x^{12} + x^9 + x^8 + x^7 + x^3 + x^2 + 1, \\
p_9(x) &= x^{80} + x^{79} + x^{78} + x^{75} + x^{74} + x^{71} + x^{70} + x^{68} + x^{32} + x^{31} + x^{30} \\
& + x^{27} + x^{26} + x^{23} + x^{22} + x^{20} + x^{16} + x^{15} + x^{14} + x^{11} + x^{10} + x^7 \\
& + x^6 + x^4, \\
p_{12}(x) &= x^{84} + x^{83} + x^{82} + x^{80} + x^{72} + x^{71} + x^{70} + x^{67} + x^{66} + x^{64} + x^{36} \\
& + x^{35} + x^{34} + x^{32} + x^{24} + x^{23} + x^{22} + x^{20} + x^8 + x^7 + x^6 + x^3 + x^2 \\
& + 1,
\end{aligned}$$

and the initial terms are $[1, 0, 1, 0, 0, 0, 0, 1]$.

For $\phi(W^o, 0, 1)$:

$$\begin{aligned}
p_0(x) &= x^{84} + x^{81} + x^{77} + x^{71} + x^{68} + x^{66} + x^{64} + x^{62} + x^{60} + x^{57} + x^{55} \\
& + x^{54} + x^{53} + x^{51} + x^{50} + x^{48} + x^{47} + x^{46} + x^{45} + x^{44} + x^{43} + x^{42} \\
& + x^{41} + x^{40} + x^{38} + x^{36} + x^{35} + x^{34} + x^{33} + x^{30} + x^{27}, \\
p_3(x) &= x^{75} + x^{71} + x^{70} + x^{68} + x^{67} + x^{64} + x^{63} + x^{62} + x^{61} + x^{60} + x^{59} \\
& + x^{58} + x^{57} + x^{56} + x^{55} + x^{54} + x^{53} + x^{52} + x^{51} + x^{50} + x^{49} + x^{48} \\
& + x^{47} + x^{46} + x^{45} + x^{44} + x^{42} + x^{41} + x^{40} + x^{37} + x^{34} + x^{33} + x^{27} \\
& + x^{23} + x^{22} + x^{20} + x^{16} + x^{13} + x^{10} + x^9, \\
p_6(x) &= x^{78} + x^{74} + x^{68} + x^{66} + x^{62} + x^{58} + x^{52} + x^{42} + x^{40} + x^{36} + x^{28} \\
& + x^{16} + x^{12} + x^{10} + x^8 + x^6, \\
p_9(x) &= x^{81} + x^{77} + x^{76} + x^{74} + x^{73} + x^{71} + x^{70} + x^{69} + x^{68} + x^{67} + x^{33} \\
& + x^{29} + x^{28} + x^{26} + x^{25} + x^{23} + x^{22} + x^{21} + x^{20} + x^{19} + x^{17} + x^{13} \\
& + x^{12} + x^{10} + x^9 + x^7 + x^6 + x^5 + x^4 + x^3, \\
p_{12}(x) &= x^{84} + x^{80} + x^{72} + x^{64} + x^{36} + x^{32} + x^{24} + x^{20} + x^8 + 1,
\end{aligned}$$

and the initial terms are $[0, 1, 1, 0, 0, 1, 1, 1]$.

For $\phi(W^o, 1, 0)$:

$$\begin{aligned}
p_0(x) &= x^{92} + x^{89} + x^{88} + x^{86} + x^{84} + x^{82} + x^{81} + x^{79} + x^{75} + x^{73} + x^{71} \\
& + x^{69} + x^{68} + x^{67} + x^{65} + x^{62} + x^{61} + x^{60} + x^{59} + x^{56} + x^{55} + x^{54} \\
& + x^{52} + x^{50} + x^{49} + x^{46} + x^{44} + x^{42} + x^{41} + x^{38} + x^{36} + x^{33} + x^{29} \\
& + x^{27} + x^{26} + x^{24} + x^{21} + x^{18}, \\
p_3(x) &= x^{83} + x^{78} + x^{76} + x^{74} + x^{71} + x^{70} + x^{69} + x^{66} + x^{64} + x^{63} + x^{62} \\
& + x^{57} + x^{56} + x^{55} + x^{54} + x^{51} + x^{49} + x^{48} + x^{47} + x^{44} + x^{42} + x^{41}
\end{aligned}$$

$$\begin{aligned}
& + x^{40} + x^{37} + x^{35} + x^{34} + x^{33} + x^{30} + x^{28} + x^{27} + x^{26} + x^{23} + x^{21} \\
& + x^{20} + x^{16} + x^{13} + x^{10} + x^9, \\
p_6(x) &= x^{86} + x^{78} + x^{76} + x^{72} + x^{64} + x^{54} + x^{52} + x^{48} + x^{42} + x^{40} + x^{32} \\
& + x^{20} + x^{18} + x^{16} + x^{12} + x^{10} + x^8 + x^6, \\
p_9(x) &= x^{89} + x^{84} + x^{82} + x^{80} + x^{79} + x^{77} + x^{76} + x^{74} + x^{73} + x^{71} + x^{70} \\
& + x^{69} + x^{68} + x^{67} + x^{41} + x^{36} + x^{34} + x^{32} + x^{31} + x^{29} + x^{28} + x^{26} \\
& + x^{23} + x^{22} + x^{21} + x^{19} + x^{18} + x^{16} + x^{15} + x^{13} + x^{12} + x^{10} + x^9 \\
& + x^7 + x^6 + x^5 + x^4 + x^3, \\
p_{12}(x) &= x^{92} + x^{84} + x^{76} + x^{72} + x^{64} + x^{44} + x^{36} + x^{24} + x^{20} + x^{16} + x^{12} + x^8 \\
& + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 1, 1, 0, 1, 1]$.

For $\phi(W^o, 1, 1)$:

$$\begin{aligned}
p_0(x) &= x^{84} + x^{83} + x^{81} + x^{80} + x^{78} + x^{77} + x^{75} + x^{72} + x^{71} + x^{70} + x^{69} \\
& + x^{66} + x^{65} + x^{63} + x^{61} + x^{60} + x^{59} + x^{58} + x^{56} + x^{55} + x^{52} + x^{51} \\
& + x^{48} + x^{47} + x^{45} + x^{43} + x^{41} + x^{39} + x^{38} + x^{36} + x^{33}, \\
p_3(x) &= x^{80} + x^{79} + x^{77} + x^{75} + x^{74} + x^{72} + x^{71} + x^{70} + x^{69} + x^{64} + x^{63} \\
& + x^{62} + x^{57} + x^{55} + x^{50} + x^{44} + x^{43} + x^{42} + x^{37} + x^{33} + x^{31} + x^{30} \\
& + x^{29} + x^{27} + x^{25} + x^{24} + x^{22} + x^{21} + x^{20} + x^{19} + x^{14} + x^{12}, \\
p_6(x) &= x^{82} + x^{81} + x^{80} + x^{79} + x^{77} + x^{76} + x^{75} + x^{74} + x^{73} + x^{71} + x^{70} \\
& + x^{69} + x^{68} + x^{67} + x^{65} + x^{64} + x^{63} + x^{60} + x^{59} + x^{55} + x^{51} + x^{50} \\
& + x^{49} + x^{48} + x^{44} + x^{38} + x^{37} + x^{36} + x^{35} + x^{32} + x^{31} + x^{29} + x^{27} \\
& + x^{24} + x^{22} + x^{21} + x^{17} + x^{16} + x^{14} + x^{11} + x^4 + x^3 + x^2 + 1, \\
p_9(x) &= x^{80} + x^{79} + x^{78} + x^{75} + x^{74} + x^{71} + x^{70} + x^{68} + x^{32} + x^{31} + x^{30} \\
& + x^{27} + x^{26} + x^{23} + x^{22} + x^{20} + x^{16} + x^{15} + x^{14} + x^{11} + x^{10} + x^7 \\
& + x^6 + x^4, \\
p_{12}(x) &= x^{84} + x^{83} + x^{82} + x^{80} + x^{72} + x^{71} + x^{70} + x^{67} + x^{66} + x^{64} + x^{36} \\
& + x^{35} + x^{34} + x^{32} + x^{24} + x^{23} + x^{22} + x^{20} + x^8 + x^7 + x^6 + x^3 + x^2 \\
& + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 1, 0, 0, 1]$.

Below is the transition function and output function of an 2-automaton that generates $T = M_{0,0}^o$:

Transition function $(n, j) \mapsto \delta(n, j)$ ($\Lambda(n) := [\delta(n, 0), \delta(n, 1)]$):

n	$\Lambda(n)$	n	$\Lambda(n)$	n	$\Lambda(n)$	n	$\Lambda(n)$	n	$\Lambda(n)$
0	[1, 2]	99	[171, 176]	198	[309, 310]	297	[325, 381]	396	[297, 25]
1	[3, 4]	100	[177, 178]	199	[311, 57]	298	[127, 317]	397	[82, 300]
2	[5, 6]	101	[179, 44]	200	[312, 313]	299	[376, 405]	398	[466, 456]
3	[7, 8]	102	[180, 32]	201	[125, 309]	300	[148, 174]	399	[302, 418]

4	[9, 10]	103	[181, 182]	202	[314, 315]	301	[406, 407]	400	[278, 464]
5	[11, 12]	104	[183, 184]	203	[189, 316]	302	[332, 402]	401	[467, 435]
6	[13, 14]	105	[185, 186]	204	[317, 318]	303	[354, 382]	402	[425, 17]
7	[15, 16]	106	[146, 11]	205	[319, 320]	304	[386, 406]	403	[451, 99]
8	[17, 18]	107	[176, 187]	206	[53, 46]	305	[326, 393]	404	[461, 261]
9	[19, 20]	108	[188, 189]	207	[321, 322]	306	[408, 235]	405	[283, 431]
10	[21, 22]	109	[190, 191]	208	[323, 181]	307	[28, 4]	406	[463, 468]
11	[23, 24]	110	[155, 192]	209	[324, 325]	308	[105, 409]	407	[429, 469]
12	[25, 26]	111	[184, 193]	210	[134, 326]	309	[209, 202]	408	[470, 344]
13	[27, 28]	112	[194, 195]	211	[327, 328]	310	[216, 410]	409	[156, 395]
14	[29, 30]	113	[196, 72]	212	[329, 330]	311	[411, 412]	410	[193, 471]
15	[31, 32]	114	[197, 198]	213	[331, 332]	312	[413, 282]	411	[390, 472]
16	[33, 34]	115	[199, 199]	214	[178, 321]	313	[231, 279]	412	[473, 311]
17	[35, 36]	116	[200, 201]	215	[333, 124]	314	[243, 414]	413	[474, 475]
18	[37, 38]	117	[202, 203]	216	[182, 334]	315	[415, 250]	414	[38, 383]
19	[39, 40]	118	[201, 197]	217	[139, 335]	316	[80, 288]	415	[142, 33]
20	[41, 42]	119	[204, 205]	218	[336, 337]	317	[260, 79]	416	[122, 380]
21	[10, 43]	120	[206, 207]	219	[316, 323]	318	[218, 206]	417	[49, 117]
22	[44, 45]	121	[208, 209]	220	[60, 338]	319	[416, 417]	418	[373, 476]
23	[46, 47]	122	[210, 211]	221	[152, 121]	320	[418, 275]	419	[404, 157]
24	[48, 49]	123	[212, 213]	222	[167, 339]	321	[419, 98]	420	[477, 478]
25	[50, 51]	124	[214, 215]	223	[340, 114]	322	[420, 16]	421	[479, 177]
26	[52, 53]	125	[111, 216]	224	[341, 342]	323	[299, 204]	422	[345, 480]
27	[54, 9]	126	[76, 217]	225	[343, 344]	324	[284, 292]	423	[481, 128]
28	[55, 56]	127	[26, 218]	226	[40, 345]	325	[421, 212]	424	[328, 35]
29	[57, 58]	128	[219, 220]	227	[346, 347]	326	[305, 200]	425	[482, 37]
30	[59, 60]	129	[20, 221]	228	[315, 348]	327	[422, 415]	426	[173, 387]
31	[61, 62]	130	[222, 223]	229	[318, 41]	328	[64, 71]	427	[363, 483]
32	[63, 64]	131	[91, 224]	230	[349, 350]	329	[228, 248]	428	[334, 143]
33	[65, 66]	132	[225, 226]	231	[351, 352]	330	[423, 251]	429	[400, 385]
34	[67, 68]	133	[227, 228]	232	[313, 353]	331	[424, 425]	430	[358, 163]
35	[69, 70]	134	[205, 229]	233	[322, 52]	332	[270, 95]	431	[476, 477]
36	[71, 72]	135	[230, 231]	234	[162, 354]	333	[426, 427]	432	[342, 351]
37	[73, 74]	136	[232, 233]	235	[355, 356]	334	[428, 429]	433	[403, 8]
38	[75, 76]	137	[234, 235]	236	[357, 358]	335	[237, 430]	434	[365, 481]
39	[16, 77]	138	[236, 237]	237	[359, 13]	336	[431, 274]	435	[484, 485]
40	[18, 78]	139	[238, 239]	238	[360, 194]	337	[281, 432]	436	[405, 388]
41	[79, 80]	140	[240, 241]	239	[361, 362]	338	[427, 69]	437	[131, 474]
42	[81, 82]	141	[62, 242]	240	[330, 363]	339	[272, 304]	438	[392, 378]
43	[83, 84]	142	[70, 243]	241	[160, 349]	340	[433, 65]	439	[486, 479]
44	[85, 5]	143	[77, 244]	242	[114, 364]	341	[22, 434]	440	[118, 340]
45	[86, 87]	144	[245, 246]	243	[129, 359]	342	[435, 436]	441	[310, 374]
46	[88, 89]	145	[247, 23]	244	[339, 365]	343	[66, 437]	442	[483, 50]
47	[90, 91]	146	[87, 110]	245	[366, 329]	344	[257, 438]	443	[485, 401]

48	[92, 93]	147	[248, 249]	246	[367, 183]	345	[289, 439]	444	[397, 59]
49	[94, 19]	148	[250, 27]	247	[368, 48]	346	[432, 230]	445	[472, 55]
50	[95, 63]	149	[251, 252]	248	[347, 135]	347	[440, 222]	446	[335, 368]
51	[96, 97]	150	[211, 253]	249	[350, 361]	348	[244, 441]	447	[487, 161]
52	[98, 88]	151	[254, 29]	250	[369, 175]	349	[263, 219]	448	[153, 119]
53	[99, 100]	152	[100, 96]	251	[370, 343]	350	[439, 423]	449	[169, 190]
54	[101, 21]	153	[255, 81]	252	[150, 147]	351	[6, 442]	450	[58, 327]
55	[102, 103]	154	[213, 256]	253	[371, 370]	352	[436, 443]	451	[394, 331]
56	[104, 105]	155	[242, 257]	254	[338, 314]	353	[444, 445]	452	[307, 170]
57	[106, 107]	156	[233, 258]	255	[372, 154]	354	[445, 419]	453	[14, 307]
58	[108, 109]	157	[259, 260]	256	[51, 373]	355	[258, 267]	454	[8, 39]
59	[110, 111]	158	[261, 262]	257	[374, 346]	356	[446, 90]	455	[362, 185]
60	[112, 67]	159	[203, 263]	258	[353, 172]	357	[414, 268]	456	[141, 188]
61	[113, 114]	160	[264, 265]	259	[375, 376]	358	[447, 448]	457	[320, 486]
62	[115, 116]	161	[262, 266]	260	[45, 145]	359	[220, 208]	458	[480, 482]
63	[117, 118]	162	[267, 62]	261	[348, 360]	360	[253, 413]	459	[391, 399]
64	[119, 120]	163	[268, 269]	262	[377, 369]	361	[449, 420]	460	[116, 54]
65	[121, 122]	164	[78, 270]	263	[378, 138]	362	[450, 416]	461	[478, 377]
66	[56, 123]	165	[221, 104]	264	[364, 168]	363	[84, 451]	462	[488, 166]
67	[124, 125]	166	[271, 272]	265	[379, 140]	364	[72, 452]	463	[475, 489]
68	[126, 127]	167	[273, 92]	266	[380, 341]	365	[430, 449]	464	[356, 151]
69	[128, 129]	168	[274, 112]	267	[381, 375]	366	[291, 450]	465	[149, 488]
70	[130, 131]	169	[275, 234]	268	[382, 324]	367	[442, 453]	466	[398, 487]
71	[132, 133]	170	[235, 276]	269	[383, 126]	368	[454, 94]	467	[489, 490]
72	[120, 134]	171	[277, 278]	270	[384, 372]	369	[455, 433]	468	[407, 484]
73	[135, 130]	172	[279, 280]	271	[385, 386]	370	[456, 457]	469	[471, 336]
74	[136, 137]	173	[266, 281]	272	[158, 371]	371	[217, 255]	470	[491, 465]
75	[138, 139]	174	[249, 73]	273	[123, 355]	372	[458, 298]	471	[293, 286]
76	[140, 136]	175	[282, 264]	274	[387, 180]	373	[459, 460]	472	[438, 227]
77	[32, 141]	176	[224, 283]	275	[388, 389]	374	[93, 440]	473	[412, 428]
78	[36, 142]	177	[246, 284]	276	[344, 390]	375	[103, 461]	474	[460, 290]
79	[143, 144]	178	[107, 285]	277	[191, 319]	376	[4, 454]	475	[276, 492]
80	[145, 146]	179	[74, 86]	278	[133, 366]	377	[437, 444]	476	[97, 426]
81	[147, 148]	180	[286, 287]	279	[192, 391]	378	[410, 424]	477	[223, 465]
82	[12, 149]	181	[288, 106]	280	[352, 367]	379	[207, 232]	478	[294, 455]
83	[43, 150]	182	[68, 289]	281	[337, 392]	380	[215, 458]	479	[452, 422]
84	[151, 152]	183	[290, 291]	282	[195, 379]	381	[453, 277]	480	[448, 225]
85	[34, 153]	184	[292, 293]	283	[393, 394]	382	[287, 273]	481	[441, 83]
86	[154, 155]	185	[285, 294]	284	[389, 2]	383	[269, 462]	482	[265, 75]
87	[137, 156]	186	[295, 214]	285	[395, 308]	384	[252, 446]	483	[24, 210]
88	[157, 158]	187	[296, 102]	286	[396, 397]	385	[301, 463]	484	[468, 467]
89	[159, 160]	188	[109, 297]	287	[47, 312]	386	[464, 245]	485	[469, 271]
90	[161, 162]	189	[298, 299]	288	[2, 398]	387	[280, 459]	486	[229, 447]
91	[163, 164]	190	[300, 295]	289	[399, 400]	388	[239, 247]	487	[434, 254]

92	[165, 132]	191	[241, 301]	290	[164, 179]	389	[303, 236]	488	[462, 421]
93	[166, 167]	192	[256, 302]	291	[401, 384]	390	[465, 259]	489	[443, 108]
94	[168, 169]	193	[226, 303]	292	[42, 396]	391	[457, 296]	490	[492, 411]
95	[170, 171]	194	[304, 305]	293	[144, 357]	392	[198, 466]	491	[1, 375]
96	[172, 173]	195	[89, 240]	294	[402, 403]	393	[417, 238]	492	[490, 473]
97	[174, 165]	196	[306, 307]	295	[186, 333]	394	[409, 85]		
98	[175, 159]	197	[308, 115]	296	[187, 404]	395	[30, 101]		

Output function $n \mapsto \tau(n)$:

n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$
0	0	71	1	142	0	213	0	284	1	426	0
1	0	72	0	143	1	214	1	285	0	427	0
2	1	73	1	144	0	215	0	286	1	428	0
3	0	74	0	145	0	216	1	287	1	429	0
4	1	75	0	146	1	217	1	288	1	430	1
5	1	76	1	147	0	218	0	289	0	431	0
6	0	77	1	148	1	219	0	290	1	432	0
7	0	78	1	149	0	220	0	291	0	433	1
8	0	79	1	150	0	221	1	292	0	434	1
9	1	80	0	151	0	222	0	293	0	435	0
10	1	81	0	152	1	223	1	294	0	436	1
11	1	82	1	153	0	224	1	295	0	437	1
12	1	83	0	154	0	225	1	296	0	438	1
13	0	84	0	155	0	226	1	297	1	439	0
14	1	85	1	156	1	227	0	298	0	440	0
15	0	86	0	157	1	228	0	299	1	441	1
16	1	87	1	158	1	229	0	300	1	442	1
17	0	88	1	159	0	230	1	301	0	443	0
18	1	89	0	160	0	231	0	302	0	444	1
19	1	90	1	161	1	232	0	303	1	445	1
20	1	91	1	162	1	233	1	304	0	446	0
21	1	92	1	163	1	234	1	305	0	447	1
22	1	93	0	164	1	235	1	306	0	448	0
23	1	94	1	165	1	236	0	307	1	449	1
24	1	95	1	166	0	237	0	308	0	450	1
25	1	96	0	167	0	238	1	309	1	451	1
26	0	97	0	168	1	239	1	310	1	452	1
27	0	98	0	169	1	240	1	311	0	453	1
28	1	99	0	170	1	241	0	312	1	454	0
29	1	100	1	171	0	242	0	313	0	455	1
30	0	101	0	172	0	243	1	314	1	456	0
31	0	102	1	173	0	244	1	315	0	457	0
32	1	103	1	174	0	245	0	316	0	458	0
33	1	104	1	175	0	246	1	317	0	459	0

34	1	105	0	176	1	247	0	318	0	389	1	460	1
35	0	106	1	177	1	248	0	319	1	390	0	461	0
36	1	107	1	178	1	249	0	320	0	391	0	462	1
37	1	108	1	179	0	250	1	321	0	392	1	463	0
38	0	109	1	180	1	251	0	322	1	393	1	464	0
39	1	110	0	181	1	252	0	323	1	394	1	465	0
40	1	111	0	182	1	253	1	324	1	395	0	466	1
41	1	112	0	183	1	254	0	325	1	396	1	467	0
42	0	113	0	184	0	255	0	326	0	397	1	468	0
43	0	114	0	185	0	256	0	327	0	398	1	469	0
44	1	115	0	186	0	257	0	328	0	399	0	470	0
45	0	116	1	187	0	258	1	329	0	400	0	471	0
46	1	117	1	188	1	259	1	330	1	401	0	472	1
47	1	118	0	189	0	260	0	331	0	402	0	473	1
48	1	119	0	190	1	261	1	332	0	403	1	474	1
49	1	120	0	191	0	262	1	333	0	404	0	475	0
50	1	121	1	192	0	263	1	334	0	405	1	476	0
51	0	122	1	193	1	264	0	335	0	406	0	477	1
52	0	123	0	194	0	265	0	336	0	407	0	478	0
53	0	124	1	195	0	266	0	337	1	408	0	479	1
54	0	125	0	196	0	267	1	338	0	409	1	480	0
55	1	126	1	197	0	268	1	339	1	410	1	481	1
56	1	127	0	198	1	269	1	340	1	411	0	482	0
57	1	128	0	199	0	270	0	341	1	412	1	483	1
58	1	129	1	200	1	271	0	342	0	413	1	484	0
59	0	130	0	201	0	272	1	343	1	414	0	485	0
60	0	131	1	202	1	273	0	344	0	415	0	486	0
61	0	132	1	203	0	274	1	345	0	416	1	487	1
62	0	133	0	204	0	275	1	346	0	417	1	488	1
63	1	134	1	205	1	276	0	347	0	418	0	489	0
64	0	135	1	206	0	277	0	348	1	419	0	490	0
65	1	136	0	207	0	278	0	349	1	420	1	491	0
66	1	137	1	208	1	279	0	350	0	421	1	492	0
67	1	138	0	209	1	280	1	351	0	422	0		
68	1	139	1	210	1	281	1	352	1	423	1		
69	0	140	1	211	0	282	0	353	1	424	0		
70	0	141	0	212	0	283	1	354	1	425	0		

UNIVERSITÉ DE STRASBOURG, CNRS, IRMA UMR 7501, F-67000 STRASBOURG, FRANCE
Email address: guoniu.han@unistra.fr

SCHOOL OF MATHEMATICS AND STATISTICS, HUAZHONG UNIVERSITY OF SCIENCE AND TECHNOLOGY, WUHAN, PR CHINA
Email address: huyining@hust.edu.cn