

PROOF OF THE CONTINUED FRACTION CONJECTURE FOR

$$(a, b) = (z^3 + 1, z^2 + 1)$$

GUO-NIU HAN* AND YINING HU*

ABSTRACT. The continued fraction conjecture claims that the continued fraction defined by the (a, b) -Thue-Morse sequence is algebraic over $\mathbb{F}_2(x)$ for all pairs of distinct elements (a, b) in $\mathbb{F}_2[x] \setminus \mathbb{F}_2$. In this paper we prove the conjecture for $(a, b) = (z^3 + 1, z^2 + 1)$.

1. INTRODUCTION

Let $\mathbb{F}_2$ be the finite field of cardinality 2. Given a sequence of polynomials $\mathbf{a} = (a_0(z), a_1(z), a_2(z), \dots)$ of elements from $\mathbb{F}_2[z] \setminus \mathbb{F}_2$, we define the infinite continued fraction

$$(1.1) \quad \text{CF}(\mathbf{a}(z)) = \frac{1}{a_0(z) + \frac{1}{a_1(z) + \frac{1}{a_2(z) + \frac{1}{\ddots}}}}$$

as the limit of the finite continued fractions

$$(1.2) \quad \text{CF}_n(\mathbf{a}(z)) = \frac{1}{a_0(z) + \frac{1}{a_1(z) + \frac{1}{\ddots + \frac{1}{a_n(z)}}}} \in \mathbb{F}_2((1/z)).$$

Conversely, each power series in $\mathbb{F}_2[[1/z]]$ without constant term

$$(1.3) \quad c_{-1}z^{-1} + c_{-2}z^{-2} + c_{-3}z^{-3} + \dots$$

admits a continued fraction expansion of form (1.1).

Let a and b be two distinct elements from $\mathbb{F}_2[z] \setminus \mathbb{F}_2$. We consider the continued fraction (1.1) associated with the (a, b) -Thue-Morse sequence

$$\mathbf{t} = (t_0(z), t_1(z), t_2(z), \dots) = (a, b, b, a, b, a, a, b, \dots)$$

that we shall denote

$$(1.4) \quad \text{CF}(a, b) := \text{CF}(\mathbf{t}).$$

Date: 9/04/2020 08:38:47 .

2010 Mathematics Subject Classification. 11B85, 11J70, 11B50, 11Y65, 05A15.

Key words and phrases. automatic sequence, continued fraction, Thue-Morse sequence.

* This paper was automatically generated from a computer program developed by the authors, with 10.9 minutes CPU time used.

Theorem 1.1. *When $(a, b) = (z^3 + 1, z^2 + 1)$, the two power series $\text{CF}(a, b)$ and $\text{CF}(b, a)$ are algebraic over $\mathbb{F}_2(z)$, with minimal polynomials of the form*

$$p_4(z)y^4 + p_3(z)y^3 + p_2(z)y^2 + p_1(z)y + p_0(z) = 0$$

where, for $\text{CF}(a, b)$,

$$\begin{aligned} p_0(z) &= z^{11} + z^9 + z^8 + z^7 + z^5 + z^3 + z^2 + 1, \\ p_1(z) &= z^{14} + z^{13} + z^{12} + z^4 + z^3 + z^2, \\ p_2(z) &= z^{16} + z^{10} + z^6 + 1, \\ p_3(z) &= z^{14} + z^{13} + z^{12} + z^4 + z^3 + z^2, \\ p_4(z) &= z^{12} + z^{11} + z^{10} + z^9 + z^7 + z^6 + z^5 + z^4 + z^3 + z^2 + 1, \end{aligned}$$

and for $\text{CF}(b, a)$,

$$\begin{aligned} p_0(z) &= z^{11} + z^9 + z^7 + z^5 + z^3 + z^2 + 1, \\ p_1(z) &= z^{14} + z^{13} + z^{12} + z^4 + z^3 + z^2, \\ p_2(z) &= z^{16} + z^{10} + z^6 + 1, \\ p_3(z) &= z^{14} + z^{13} + z^{12} + z^4 + z^3 + z^2, \\ p_4(z) &= z^{12} + z^{11} + z^{10} + z^9 + z^8 + z^7 + z^6 + z^5 + z^4 + z^3 + z^2 + 1. \end{aligned}$$

2. PROOF

Define the sequences of polynomials $(P_n(z))_{n \geq 0}$ and $(Q_n(z))_{n \geq 0}$ by

$$\begin{pmatrix} P_n(z) & Q_n(z) \\ P_{n-1}(z) & Q_{n-1}(z) \end{pmatrix} := \begin{pmatrix} t_n(z) & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} t_{n-1}(z) & 1 \\ 1 & 0 \end{pmatrix} \cdots \begin{pmatrix} t_0(z) & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}.$$

By the basic properties of continued fractions, we have

$$(2.1) \quad \text{CF}_n(a, b) = \frac{P_n(z)}{Q_n(z)} \in \mathbb{F}_2((1/z)).$$

As formula (2.1) is not efficient for calculating $\text{CF}_n(a, b)$, a fast method is to be derived and described as follows. Define

$$\begin{aligned} M_n(x) &= x^{\deg(t_{2^n-1})} \begin{pmatrix} t_{2^n-1}(1/x) & 1 \\ 1 & 0 \end{pmatrix} \times x^{\deg(t_{2^n-2})} \begin{pmatrix} t_{2^n-2}(1/x) & 1 \\ 1 & 0 \end{pmatrix} \times \\ &\quad \cdots \times x^{\deg(t_0)} \begin{pmatrix} t_0(1/x) & 1 \\ 1 & 0 \end{pmatrix}, \\ W_n(x) &= x^{\deg(\bar{t}_{2^n-1})} \begin{pmatrix} \bar{t}_{2^n-1}(1/x) & 1 \\ 1 & 0 \end{pmatrix} \times x^{\deg(\bar{t}_{2^n-2})} \begin{pmatrix} \bar{t}_{2^n-2}(1/x) & 1 \\ 1 & 0 \end{pmatrix} \times \\ &\quad \cdots \times x^{\deg(\bar{t}_0)} \begin{pmatrix} \bar{t}_0(1/x) & 1 \\ 1 & 0 \end{pmatrix}, \end{aligned}$$

where $\bar{t}$ is the (b, a) -Thue-Morse sequence. By the properties of the Thue-Morse sequence, we have for $n \geq 0$

$$\begin{aligned} M_{n+1}(x) &= W_n(x) \cdot M_n(x), \\ W_{n+1}(x) &= M_n(x) \cdot W_n(x). \end{aligned}$$

Let $x := 1/z$. For each non-zero polynomial $P(z)$, define $\tilde{P}(x)$ to be $P(1/x)$. Then,

$$\text{CF}_n(a, b) = \frac{P_n(z)}{Q_n(z)} = \frac{\tilde{P}_n(x)}{\tilde{Q}_n(x)} \in \mathbb{F}_2((x)) = \mathbb{F}_2((1/z)).$$

Comparing the definition of $M_n(x)$ with the definition of $P_n(x)$ and $Q_n(x)$, we see that

$$\begin{aligned} M_n(x)_{0,1} &= x^{d_n} \tilde{P}_{2^n-1}(x), \\ M_n(x)_{0,0} &= x^{d_n} \tilde{Q}_{2^n-1}(x), \end{aligned}$$

for some positive integer d_n . Hence,

$$(2.2) \quad \text{CF}_{2^{2^n}-1}(a, b) = \frac{\tilde{P}_{2^{2^n}-1}(x)}{\tilde{Q}_{2^{2^n}-1}(x)} = \frac{M_{2^n}(x)_{0,1}}{M_{2^n}(x)_{0,0}}.$$

Therefore, by the convergence theorem of continued fraction, the algebraicity of $\text{CF}(a, b)$ will be established if it is shown that both $M_{2^n}(x)_{0,1}$ and $M_{2^n}(x)_{0,0}$ converge to algebraic series in $\mathbb{F}_2[[x]]$.

Actually, we will prove that for all $0 \leq i, j \leq 1$ the four sequences $(M_{2^n}(x)_{i,j})_n$, $(M_{2^{n+1}}(x)_{i,j})_n$, $(W_{2^n}(x)_{i,j})_n$, and $(W_{2^{n+1}}(x)_{i,j})_n$ converge to algebraic series in $\mathbb{F}_2[[x]]$. For this purpose, we define four 2×2 matrices M^e, M^o, W^e, W^o as follows: For each $T \in \{M^e, M^o, W^e, W^o\}$ and $0 \leq i, j \leq 1$, $T_{i,j}$ is defined to be the unique solution in $\mathbb{F}_2[[x]]$ of the polynomial $\phi(T, i, j)$ under certain initial conditions; the polynomials $\phi(T, i, j)$ and initial conditions are given in Section 3. We will prove that these four matrices, whose components are algebraic by definition, are the limits of $(M_{2^n}(x))_n$, $(M_{2^{n+1}}(x))_n$, $(W_{2^n}(x))_n$, and $(W_{2^{n+1}}(x))_n$.

Let us explain how the polynomials $\phi(T, i, j)$ and initial conditions are found, and why the solutions exist and are unique. For $0 \leq i, j \leq 1$, the coefficients of the polynomial $\phi(M^e, i, j)$ (resp. $\phi(M^o, i, j)$, $\phi(W^e, i, j)$, and $\phi(W^o, i, j)$) are the Padé-Hermite approximants of type

$$(125, 125, 125, 125, 125)$$

of the vector

$$(1, T^3, T^6, T^9, T^{12}),$$

where $T = M_{12,i,j}$ (resp. $M_{11,i,j}$, $W_{12,i,j}$, and $W_{11,i,j}$), found by the Derksen algorithm. We take the first 8 terms of $M_{12,i,j}$ (resp. $M_{11,i,j}$, $W_{12,i,j}$, and $W_{11,i,j}$) as initial conditions for $\phi(M^e, i, j)$ (resp. $\phi(M^o, i, j)$, $\phi(W^e, i, j)$, and $\phi(W^o, i, j)$). The following fact will be used to ensure that the solution exists and is unique: let $P(x, y) \in \mathbb{F}_2[x, y]$ and for each series $f(x) = \sum_{n=0}^{\infty} a_n x^n \in \mathbb{F}_2[[x]]$ denote the partial sum $\sum_{j=0}^{n-1} a_j x^j$ by $f_n(x)$ for $n \geq 0$. If for some $n \geq 0$ and $a_0, a_1, \dots, a_{n-1} \in \mathbb{F}_2$ $P(x, \sum_{j=0}^{n-1} a_j x^j) = O(x^n)$ and $Q(x, y) := P(x, \sum_{j=0}^{n-1} a_j x^j + x^n y)$ can be written as $x^m \sum_{j=0}^{\infty} q_j(x) y^j$ where $q_j(x)$ are polynomials for $j \geq 0$, $q_1(0) = 1$, and $q_j(0) = 0$ for $j > 1$, then there exists a unique solution $f(x) \in \mathbb{F}_2[[x]]$ of $P(x, f(x)) = 0$ that satisfies the initial condition $f_n(x) = \sum_{j=0}^{n-1} a_j x^j$.

We state two lemmas concerning the four matrices M^e, M^o, W^e, W^o . The first one is about relations between them; the second, about the structure of the each matrix.

Lemma 2.1. *We have*

$$(2.3) \quad M^e = W^o \cdot M^o,$$

$$(2.4) \quad M^o = W^e \cdot M^e,$$

$$(2.5) \quad W^e = M^o \cdot W^o,$$

$$(2.6) \quad W^o = M^e \cdot W^e.$$

Proof. We give the proof of the identity

$$M_{0,0}^e = W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o,$$

the proofs of the others being similar.

First, we compute the minimal polynomials of $W_{0,0}^o M_{0,0}^o$ and $W_{0,1}^o M_{1,0}^o$. We know that

$$P(x, y) = \text{Res}_z (\phi(W^o, 0, 0)(x, z), z^{12} \cdot \phi(M^o, 0, 0)(x, y/z))$$

is an annihilating polynomial of $W_{0,0}^o M_{0,0}^o$. We use Padé-Hermite approximation to find a candidate for the minimal polynomial of $W_{0,0}^o M_{0,0}^o$, that will be called $\phi_0(x, y)$. To prove that $\phi_0(x, y)$ is indeed the minimal polynomial, it suffices to prove that it is an irreducible factor of $P(x, y)$ of multiplicity m and that $Q(x, y) := P(x, y)/\phi_0(x, y)^m$ is not an annihilating polynomial of $W_{0,0}^o M_{0,0}^o$. We verify the first point directly. For the second point, we truncate $W_{0,0}^o M_{0,0}^o$ to order 450 and substitute it for y in $Q(x, y)$. We get a series of valuation less than 450, which proves that $Q(x, y)$ is not an annihilating polynomial of $W_{0,0}^o M_{0,0}^o$. We find the minimal polynomial $\phi_1(x, y)$ of $W_{0,1}^o M_{1,0}^o$ in a similar way.

Now we prove that $\phi(M^e, 0, 0)$ is the minimal polynomial of $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$. We know that

$$S(x, y) = \text{Res}_z (\phi_0(x, z), \phi_1(x, y + z))$$

is an annihilating of $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$. We verify that $\phi(M^e, 0, 0)$ is an irreducible factor of $S(x, y)$ of multiplicity μ , and that $Q(x, y) := S(x, y)/\phi(M^e, 0, 0)^\mu$ is not an annihilating polynomial of $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$. To see the last point, we truncate $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$ to order 550 and substitute it for y in $Q(x, y)$. We get a series of valuation less than 550, and therefore $Q(x, y)$ is not an annihilating polynomial of $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$.

Finally, the first 16 terms of $M_{0,0}^e$ and $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$ coincide. As these first terms determine a unique solution of $\phi(M^e, 0, 0)$, we know that the two series are one and the same. $\square$

Define

$$R^e = \begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix} \quad \text{and} \quad R^o = \begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix}.$$

Lemma 2.2. *For all integers $k \geq 2$ and $u = 2^{2k-1}$, the following identities hold.*

$$\begin{aligned} M^e[:10u] &= M^e[:5u] + x^{2u} M^e[:3u] + x^{3u} M^e[:2u] + x^{4u} M^e[:u] + x^{2u} M^e[:5u] \\ &\quad + x^{4u} M^e[:3u] + x^{5u} M^e[:2u] + x^{6u} M^e[:u] + x^{3u} M^e[:5u] \\ &\quad + x^{5u} M^e[:3u] + x^{6u} M^e[:2u] + x^{7u} M^e[:u] + x^{5u} M^e[:5u] \\ &\quad + x^{7u} M^e[:3u] + x^{9u} M^e[:u] + (x^{5u} + x^{7u} + x^{8u}) R^e, \\ W^e[:10u] &= W^e[:5u] + x^{2u} W^e[:3u] + x^{3u} W^e[:2u] + x^{4u} W^e[:u] + x^{2u} W^e[:5u] \\ &\quad + x^{4u} W^e[:3u] + x^{5u} W^e[:2u] + x^{6u} W^e[:u] + x^{3u} W^e[:5u] \end{aligned}$$

$$\begin{aligned}
& + x^{5u}W^e[:3u] + x^{6u}W^e[:2u] + x^{7u}W^e[:u] + x^{5u}W^e[:5u] \\
& + x^{7u}W^e[:3u] + x^{9u}W^e[:u] + (x^{5u} + x^{7u} + x^{8u})R^e.
\end{aligned}$$

For all integers $k \geq 2$ and $u = 2^{2k}$, the following identities hold.

$$\begin{aligned}
M^o[:10u] &= M^o[:5u] + x^{2u}M^o[:3u] + x^{3u}M^o[:2u] + x^{4u}M^o[:u] + x^{2u}M^o[:5u] \\
& + x^{4u}M^o[:3u] + x^{5u}M^o[:2u] + x^{6u}M^o[:u] + x^{3u}M^o[:5u] \\
& + x^{5u}M^o[:3u] + x^{6u}M^o[:2u] + x^{7u}M^o[:u] + x^{5u}M^o[:5u] \\
& + x^{7u}M^o[:3u] + x^{9u}M^o[:u] + (x^{5u} + x^{7u} + x^{8u})R^o, \\
W^o[:10u] &= W^o[:5u] + x^{2u}W^o[:3u] + x^{3u}W^o[:2u] + x^{4u}W^o[:u] + x^{2u}W^o[:5u] \\
& + x^{4u}W^o[:3u] + x^{5u}W^o[:2u] + x^{6u}W^o[:u] + x^{3u}W^o[:5u] \\
& + x^{5u}W^o[:3u] + x^{6u}W^o[:2u] + x^{7u}W^o[:u] + x^{5u}W^o[:5u] \\
& + x^{7u}W^o[:3u] + x^{9u}W^o[:u] + (x^{5u} + x^{7u} + x^{8u})R^o.
\end{aligned}$$

Proof. To prove Lemma 2.2 we first construct an automaton for each sequence concerned, and then transform the conditions on infinitely many k 's into finitely many conditions on the states of the automaton. In the following, we will prove that for $T = M_{0,0}^o$, for all integer $k \geq 2$ and $u = 2^{2k}$,

$$\begin{aligned}
T[:10u] &= T[:5u] + x^{2u}T[:3u] + x^{3u}T[:2u] + x^{4u}T[:u] + x^{2u}T[:5u] + x^{4u}T[:3u] \\
& + x^{5u}T[:2u] + x^{6u}T[:u] + x^{3u}T[:5u] + x^{5u}T[:3u] + x^{6u}T[:2u] \\
& + x^{7u}T[:u] + x^{5u}T[:5u] + x^{7u}T[:3u] + x^{9u}T[:u] + (x^{5u} + x^{7u} + x^{8u}).
\end{aligned}$$

The proof of the other 15 cases are similar. We break down the above identity into 5 parts:

$$\begin{aligned}
x^5u &= x^{5u}T[:u] + x^{4u}T[u:2u] + x^{3u}T[2u:3u] + x^{2u}T[3u:4u] \\
& + T[5u:6u], \\
0 &= x^{5u}T[u:2u] + x^{4u}T[2u:3u] + x^{3u}T[3u:4u] + x^{2u}T[4u:5u] \\
& + T[6u:7u], \\
x^7u &= x^{6u}T[u:2u] + x^{3u}T[4u:5u] + T[7u:8u], \\
x^8u &= x^{7u}T[u:2u] + x^{5u}T[3u:4u] + T[8u:9u], \\
0 &= x^{9u}T[:u] + x^{7u}T[2u:3u] + x^{5u}T[4u:5u] + T[9u:10u],
\end{aligned}$$

which can be rewritten as

$$(2.7) \quad 0 = T[[w]_2] + T[[1w]_2] + T[[10w]_2] + T[[11w]_2] + T[[101w]_2],$$

$$(2.8) \quad 0 = T[[1w]_2] + T[[10w]_2] + T[[11w]_2] + T[[100w]_2] + T[[110w]_2],$$

$$(2.9) \quad 0 = T[[1w]_2] + T[[100w]_2] + T[[111w]_2],$$

$$(2.10) \quad 0 = T[[1w]_2] + T[[11w]_2] + T[[1000w]_2],$$

$$(2.11) \quad 0 = T[[w]_2] + T[[10w]_2] + T[[100w]_2] + T[[1001w]_2],$$

for all binary word w of length $2k$ and $w \neq 0^{2k}$, and

$$(2.12) \quad 1 = T[[w]_2] + T[[1w]_2] + T[[10w]_2] + T[[11w]_2] + T[[101w]_2],$$

$$(2.13) \quad 0 = T[[1w]_2] + T[[10w]_2] + T[[11w]_2] + T[[100w]_2] + T[[110w]_2],$$

$$(2.14) \quad 1 = T[[1w]_2] + T[[100w]_2] + T[[111w]_2],$$

$$(2.15) \quad 1 = T[[1w]_2] + T[[11w]_2] + T[[1000w]_2],$$

$$(2.16) \quad 0 = T[[w]_2] + T[[10w]_2] + T[[100w]_2] + T[[1001w]_2],$$

for $w = 0^{2k}$.

First we calculate an 2-automaton that generates T from its minimal polynomial and its first terms. This automaton has 368 states; its transition function and output function can be found in the annex. Let $A(s, w)$ denote the state reached after reading w from right to left starting from the state s , and τ the output function. Define

$$E_{2k} = \{A(i, w) : |w| = 2k, w \neq 0^{2k}\}.$$

Identities (2.7) through (2.16) can be written as

$$(2.17) \quad 0 = \tau(A(s, \epsilon)) + \tau(A(s, 1)) + \tau(A(s, 10)) + \tau(A(s, 11)) + \tau(A(s, 101)),$$

$$0 = \tau(A(s, 1)) + \tau(A(s, 10)) + \tau(A(s, 11)) + \tau(A(s, 100))$$

$$(2.18) \quad + \tau(A(s, 110)),$$

$$(2.19) \quad 0 = \tau(A(s, 1)) + \tau(A(s, 100)) + \tau(A(s, 111)),$$

$$(2.20) \quad 0 = \tau(A(s, 1)) + \tau(A(s, 11)) + \tau(A(s, 1000)),$$

$$(2.21) \quad 0 = \tau(A(s, \epsilon)) + \tau(A(s, 10)) + \tau(A(s, 100)) + \tau(A(s, 1001)),$$

for all $s \in E_{2k}$, and

$$(2.22) \quad 1 = \tau(A(s, \epsilon)) + \tau(A(s, 1)) + \tau(A(s, 10)) + \tau(A(s, 11)) + \tau(A(s, 101)),$$

$$0 = \tau(A(s, 1)) + \tau(A(s, 10)) + \tau(A(s, 11)) + \tau(A(s, 100))$$

$$(2.23) \quad + \tau(A(s, 110)),$$

$$(2.24) \quad 1 = \tau(A(s, 1)) + \tau(A(s, 100)) + \tau(A(s, 111)),$$

$$(2.25) \quad 1 = \tau(A(s, 1)) + \tau(A(s, 11)) + \tau(A(s, 1000)),$$

$$(2.26) \quad 0 = \tau(A(s, \epsilon)) + \tau(A(s, 10)) + \tau(A(s, 100)) + \tau(A(s, 1001)),$$

for $s = A(i, 0^{2k})$. We find that $(A(i, 0^{16}), E_{16}) = (A(i, 0^{14}), E_{14})$, so that we only have to verify that identities (2.17) through (2.26) hold for $2 \leq k \leq 8$, which turns out to be true. $\square$

In the following lemma, we express M_{2k} , M_{2k+1} , W_{2k} , and W_{2k+1} in terms of M^e , M^o , W^e , and W^o .

Lemma 2.3. *For all integer $k \geq 2$, and $u = 2^{2k-1}$,*

$$M_{2k} = M^e[:5u] + x^{2u}M^e[:3u] + x^{3u}M^e[:2u] + x^{4u}M^e[:u] + x^{5u}R^e,$$

$$W_{2k} = W^e[:5u] + x^{2u}W^e[:3u] + x^{3u}W^e[:2u] + x^{4u}W^e[:u] + x^{5u}R^e.$$

For all integer $k \geq 2$, and $u = 2^{2k}$,

$$M_{2k+1} = M^o[:5u] + x^{2u}M^o[:3u] + x^{3u}M^o[:2u] + x^{4u}M^o[:u] + x^{5u}R^o,$$

$$W_{2k+1} = W^o[:5u] + x^{2u}W^o[:3u] + x^{3u}W^o[:2u] + x^{4u}W^o[:u] + x^{5u}R^o.$$

Proof. We call the four identities in Lemma 2.3 also by the name M_{2k} , W_{2k} , M_{2k+1} , and W_{2k+1} . For $n = 2$, the identities can be verified directly. For $n \geq 2$, we claim that

$$M_{2k} \wedge W_{2k} \Rightarrow M_{2k+1} \wedge W_{2k+1},$$

$$M_{2k+1} \wedge W_{2k+1} \Rightarrow M_{2k+2} \wedge W_{2k+2}.$$

We give the proof of

$$M_{2k} \wedge W_{2k} \Rightarrow M_{2k+1},$$

the other ones being similar. Set $u = 2^{2k}$ and $v = 2^{2k-1}$. By definition and induction hypothesis, the left side of identity M_{2k+1} is equal to

$$(2.27) \quad W_{2k} M_{2k} = (W^e[:5v] + x^{2v} W^e[:3v] + x^{3v} W^e[:2v] + x^{4v} W^e[:v] + x^{5u} R^e) \times (M^e[:5v] + x^{2v} M^e[:3v] + x^{3v} M^e[:2v] + x^{4v} M^e[:v] + x^{5u} R^e);$$

Call this expression *lhs*. Note that both sides of identity M_{2k+1} have the same term of highest degree $x^{10v} R^o$. Therefore we only need to prove that their difference is $O(x^{10v})$. Using Lemma 2.1 it can be seen that the right side of identity M_{2k+1} is congruent, modulo x^{10v} , to

$$W^e[:10v] M^e[:10v] + x^{4v} W^e[:6v] M^e[:6v] + x^{6v} W^e[:4v] M^e[:4v] + x^{8v} W^e[:2v] M^e[:2v].$$

For all $n \leq 10$, replace the occurrences of $W^e[:n \cdot v]$ and $M^e[:n \cdot v]$ in the above expression by the reduction modulo $x^{n \cdot v}$ of the right side of the corresponding identity in Lemma 2.2 and get a new expression, which we call *rhs*. Define

$$\begin{aligned} X &:= x^v, \\ a_n &:= W^e[n \cdot v : (n+1) \cdot v] / X^n, \\ b_n &:= M^e[n \cdot v : (n+1) \cdot v] / X^n, \\ c &:= R^e. \end{aligned}$$

Using the notation introduced above, we can represent the expressions *lhs* (2.27) and *rhs* as polynomials in $\mathbb{F}_2[a_1, \dots, a_{10}, b_1, \dots, b_{10}, c][X]$. Note that it is not a problem that a_j commutes with b_k while W^e does not commute with M^e , because in the expressions concerned, the products of W^e -terms and M^e -terms are always in the same order. We let the computer do the simplification and check that the difference between these two polynomials is indeed $O(X^{10})$, which completes the proof. $\square$

Proof of Theorem 1.1. We prove the theorem for $\text{CF}(a, b)$; for $\text{CF}(b, a)$, the proof is similar. By Lemma 2.3, we have For all $0 \leq j, k \leq 1$,

$$\begin{aligned} \lim_{n \rightarrow \infty} M_{2n,j,k} &= M_{j,k}^e, \\ \lim_{n \rightarrow \infty} M_{2n+1,j,k} &= M_{j,k}^o, \\ \lim_{n \rightarrow \infty} W_{2n,j,k} &= W_{j,k}^e, \\ \lim_{n \rightarrow \infty} W_{2n+1,j,k} &= W_{j,k}^o. \end{aligned}$$

Let $z = 1/x$. By the convergence theorem and identity (2.2),

$$\text{CF}(a, b) = \frac{M_{0,1}^e(x)}{M_{0,0}^e(x)}.$$

By definition, $\phi(M^e, 0, 1)$ and $\phi(M^e, 0, 0)$ are minimal polynomials of $M_{0,1}^e$ and $M_{0,0}^e$. Therefore

$$P(x, y) = \text{Res}_t (\phi(M^e, 0, 1)(x, t), y^{12} \phi(M^e, 0, 1)(x, t/y))$$

is an annihilating polynomial of $f(x) = M_{0,1}^e / M_{0,0}^e$.

Define

$$Q(x, y) = q_4(x)y^4 + q_3(x)y^3 + q_2(x)y^2 + q_1(x)y + q_0(x),$$

where

$$\begin{aligned} q_0(x) &= x^{16} + x^{14} + x^{13} + x^{11} + x^9 + x^8 + x^7 + x^5, \\ q_1(x) &= x^{14} + x^{13} + x^{12} + x^4 + x^3 + x^2, \\ q_2(x) &= x^{16} + x^{10} + x^6 + 1, \\ q_3(x) &= x^{14} + x^{13} + x^{12} + x^4 + x^3 + x^2, \\ q_4(x) &= x^{16} + x^{14} + x^{13} + x^{12} + x^{11} + x^{10} + x^9 + x^7 + x^6 + x^5 + x^4. \end{aligned}$$

The polynomial $Q(x, y)$ is the candidate for the minimal polynomial of $f(x)$ found by Padé-Hermite approximation. To prove that it is indeed the minimal polynomial of $f(x)$, we only need to prove that it is an irreducible factor of $P(x, y)$ of multiplicity m and $R(x, y) := P(x, y)/Q(x, y)^m$ is not an annihilating polynomial of $f(x)$. We verify the first point directly. For the second point, we find that when we truncate $f(x)$ to order 156, and substitute it for y in $R(z, y)$, we get a series with valuation smaller than 156, which proves that $R(z, y)$ is not an annihilating polynomial of $f(x)$. Finally, $z^{12}Q(1/z, y)$ is the minimal polynomial of $\text{CF}(a, b) = f(1/z)$. $\square$

3. ANNEXE

All 16 polynomials are of the form

$$p_0(x) + p_3(x)y^3 + p_6(x)y^6 + p_9(x)y^9 + p_{12}(x)y^{12}.$$

The coefficients $p_j(x)$ and the 16 initial terms to determine the solutions uniquely are given below.

For $\phi(M^e, 0, 0)$:

$$\begin{aligned} p_0(x) &= x^{84} + x^{82} + x^{76} + x^{74} + x^{64} + x^{60} + x^{56} + x^{50} + x^{48} + x^{44} + x^{38} \\ &\quad + x^{34} + x^{32} + x^{30} + x^{24}, \\ p_3(x) &= x^{80} + x^{74} + x^{72} + x^{68} + x^{66} + x^{64} + x^{60} + x^{58} + x^{56} + x^{54} + x^{40} \\ &\quad + x^{34} + x^{32} + x^{28} + x^{26} + x^{24} + x^{20} + x^{18} + x^{16} + x^{14}, \\ p_6(x) &= x^{80} + x^{76} + x^{74} + x^{72} + x^{70} + x^{66} + x^{64} + x^{60} + x^{42} + x^{36} + x^{34} \\ &\quad + x^{32} + x^{30} + x^{26} + x^{24} + x^{20} + x^2 + 1, \\ p_9(x) &= x^{80} + x^{74} + x^{70} + x^{64} + x^{60} + x^{54} + x^{50} + x^{44} + x^{40} + x^{34} + x^{30} \\ &\quad + x^{24} + x^{20} + x^{14} + x^{10} + x^4, \\ p_{12}(x) &= x^{84} + x^{82} + x^{80} + x^4 + x^2 + 1, \end{aligned}$$

and the initial terms are $[1, 0, 0, 0, 1, 0, 0, 0, 0, 0, 1, 0, 1, 0, 1, 0]$.

For $\phi(M^e, 0, 1)$:

$$\begin{aligned} p_0(x) &= x^{87} + x^{84} + x^{81} + x^{80} + x^{78} + x^{76} + x^{75} + x^{74} + x^{73} + x^{69} + x^{68} \\ &\quad + x^{67} + x^{66} + x^{65} + x^{63} + x^{62} + x^{60} + x^{58} + x^{56} + x^{52} + x^{51} + x^{49} \\ &\quad + x^{47} + x^{45} + x^{43} + x^{42} + x^{40} + x^{38} + x^{37} + x^{36} + x^{35} + x^{30} + x^{29} \\ &\quad + x^{28} + x^{27}, \\ p_3(x) &= x^{69} + x^{68} + x^{67} + x^{66} + x^{63} + x^{62} + x^{57} + x^{56} + x^{55} + x^{54} + x^{53} \end{aligned}$$

$$\begin{aligned}
& + x^{52} + x^{51} + x^{50} + x^{49} + x^{48} + x^{47} + x^{46} + x^{45} + x^{44} + x^{43} + x^{42} \\
& + x^{41} + x^{40} + x^{39} + x^{38} + x^{37} + x^{36} + x^{35} + x^{34} + x^{33} + x^{32} + x^{31} \\
& + x^{30} + x^{25} + x^{24} + x^{21} + x^{20} + x^{19} + x^{18}, \\
p_6(x) &= x^{75} + x^{73} + x^{72} + x^{70} + x^{69} + x^{66} + x^{65} + x^{62} + x^{61} + x^{58} + x^{57} \\
& + x^{55} + x^{54} + x^{52} + x^{35} + x^{33} + x^{32} + x^{30} + x^{29} + x^{26} + x^{25} + x^{22} \\
& + x^{21} + x^{18} + x^{17} + x^{15} + x^{14} + x^{12}, \\
p_9(x) &= x^{81} + x^{80} + x^{73} + x^{72} + x^{71} + x^{70} + x^{69} + x^{68} + x^{65} + x^{64} + x^{63} \\
& + x^{62} + x^{59} + x^{58} + x^{57} + x^{56} + x^{55} + x^{54} + x^{47} + x^{46} + x^{41} + x^{40} \\
& + x^{33} + x^{32} + x^{31} + x^{30} + x^{29} + x^{28} + x^{25} + x^{24} + x^{23} + x^{22} + x^{19} \\
& + x^{18} + x^{17} + x^{16} + x^{15} + x^{14} + x^7 + x^6, \\
p_{12}(x) &= x^{87} + x^{84} + x^{79} + x^{76} + x^{75} + x^{72} + x^{71} + x^{68} + x^{67} + x^{64} + x^{63} \\
& + x^{60} + x^{59} + x^{56} + x^{55} + x^{52} + x^{51} + x^{48} + x^{47} + x^{44} + x^{43} + x^{40} \\
& + x^{39} + x^{36} + x^{35} + x^{32} + x^{31} + x^{28} + x^{27} + x^{24} + x^{23} + x^{20} + x^{19} \\
& + x^{16} + x^{15} + x^{12} + x^{11} + x^8 + x^3 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 1, 0, 0, 1, 1, 1, 1, 0, 0, 0, 1, 0, 0]$.

For $\phi(M^e, 1, 0)$:

$$\begin{aligned}
p_0(x) &= x^{87} + x^{84} + x^{81} + x^{80} + x^{78} + x^{76} + x^{75} + x^{74} + x^{73} + x^{69} + x^{68} \\
& + x^{67} + x^{66} + x^{65} + x^{63} + x^{62} + x^{60} + x^{58} + x^{56} + x^{52} + x^{51} + x^{49} \\
& + x^{47} + x^{45} + x^{43} + x^{42} + x^{40} + x^{38} + x^{37} + x^{36} + x^{35} + x^{30} + x^{29} \\
& + x^{28} + x^{27}, \\
p_3(x) &= x^{69} + x^{68} + x^{67} + x^{66} + x^{63} + x^{62} + x^{57} + x^{56} + x^{55} + x^{54} + x^{53} \\
& + x^{52} + x^{51} + x^{50} + x^{49} + x^{48} + x^{47} + x^{46} + x^{45} + x^{44} + x^{43} + x^{42} \\
& + x^{41} + x^{40} + x^{39} + x^{38} + x^{37} + x^{36} + x^{35} + x^{34} + x^{33} + x^{32} + x^{31} \\
& + x^{30} + x^{25} + x^{24} + x^{21} + x^{20} + x^{19} + x^{18}, \\
p_6(x) &= x^{75} + x^{73} + x^{72} + x^{70} + x^{69} + x^{66} + x^{65} + x^{62} + x^{61} + x^{58} + x^{57} \\
& + x^{55} + x^{54} + x^{52} + x^{35} + x^{33} + x^{32} + x^{30} + x^{29} + x^{26} + x^{25} + x^{22} \\
& + x^{21} + x^{18} + x^{17} + x^{15} + x^{14} + x^{12}, \\
p_9(x) &= x^{81} + x^{80} + x^{73} + x^{72} + x^{71} + x^{70} + x^{69} + x^{68} + x^{65} + x^{64} + x^{63} \\
& + x^{62} + x^{59} + x^{58} + x^{57} + x^{56} + x^{55} + x^{54} + x^{47} + x^{46} + x^{41} + x^{40} \\
& + x^{33} + x^{32} + x^{31} + x^{30} + x^{29} + x^{28} + x^{25} + x^{24} + x^{23} + x^{22} + x^{19} \\
& + x^{18} + x^{17} + x^{16} + x^{15} + x^{14} + x^7 + x^6, \\
p_{12}(x) &= x^{87} + x^{84} + x^{79} + x^{76} + x^{75} + x^{72} + x^{71} + x^{68} + x^{67} + x^{64} + x^{63} \\
& + x^{60} + x^{59} + x^{56} + x^{55} + x^{52} + x^{51} + x^{48} + x^{47} + x^{44} + x^{43} + x^{40} \\
& + x^{39} + x^{36} + x^{35} + x^{32} + x^{31} + x^{28} + x^{27} + x^{24} + x^{23} + x^{20} + x^{19} \\
& + x^{16} + x^{15} + x^{12} + x^{11} + x^8 + x^3 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 1, 0, 0, 1, 1, 1, 1, 0, 0, 0, 1, 0, 0]$.

For $\phi(M^e, 1, 1)$:

$$\begin{aligned}
p_0(x) &= x^{90} + x^{88} + x^{86} + x^{82} + x^{80} + x^{74} + x^{64} + x^{62} + x^{60} + x^{58} + x^{54} \\
&\quad + x^{52} + x^{50} + x^{48} + x^{46} + x^{40} + x^{34} + x^{32} + x^{30}, \\
p_3(x) &= x^{86} + x^{82} + x^{74} + x^{72} + x^{70} + x^{64} + x^{62} + x^{60} + x^{58} + x^{54} + x^{52} \\
&\quad + x^{46} + x^{42} + x^{34} + x^{32} + x^{30} + x^{24} + x^{22} + x^{20} + x^{18} + x^{14} + x^{12}, \\
p_6(x) &= x^{86} + x^{82} + x^{80} + x^{76} + x^{72} + x^{68} + x^{60} + x^{56} + x^{46} + x^{44} + x^{36} \\
&\quad + x^{32} + x^{28} + x^{20} + x^{16} + x^4 + x^2 + 1, \\
p_9(x) &= x^{86} + x^{82} + x^{76} + x^{72} + x^{70} + x^{60} + x^{58} + x^{54} + x^{48} + x^{46} + x^{44} \\
&\quad + x^{42} + x^{36} + x^{32} + x^{30} + x^{20} + x^{18} + x^{14} + x^8 + x^4, \\
p_{12}(x) &= x^{90} + x^{88} + x^{74} + x^{72} + x^{66} + x^{64} + x^{58} + x^{56} + x^{50} + x^{48} + x^{42} \\
&\quad + x^{40} + x^{34} + x^{32} + x^{26} + x^{24} + x^{18} + x^{16} + x^2 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 0, 1, 0, 0, 0, 1, 0, 1, 0, 0, 0]$.

For $\phi(W^e, 0, 0)$:

$$\begin{aligned}
p_0(x) &= x^{90} + x^{88} + x^{86} + x^{78} + x^{76} + x^{72} + x^{70} + x^{64} + x^{62} + x^{60} + x^{56} \\
&\quad + x^{54} + x^{52} + x^{48} + x^{46} + x^{44} + x^{40} + x^{28} + x^{24}, \\
p_3(x) &= x^{86} + x^{82} + x^{78} + x^{68} + x^{66} + x^{56} + x^{54} + x^{46} + x^{42} + x^{38} + x^{28} \\
&\quad + x^{26} + x^{16} + x^{14}, \\
p_6(x) &= x^{86} + x^{72} + x^{70} + x^{66} + x^{64} + x^{62} + x^{56} + x^{54} + x^{50} + x^{48} + x^{46} \\
&\quad + x^{44} + x^{42} + x^{40} + x^{32} + x^{30} + x^{26} + x^{24} + x^{22} + x^{16} + x^{14} + x^{10} \\
&\quad + x^8 + x^4 + x^2 + 1, \\
p_9(x) &= x^{86} + x^{82} + x^{76} + x^{72} + x^{70} + x^{60} + x^{58} + x^{54} + x^{48} + x^{46} + x^{44} \\
&\quad + x^{42} + x^{36} + x^{32} + x^{30} + x^{20} + x^{18} + x^{14} + x^8 + x^4, \\
p_{12}(x) &= x^{90} + x^{88} + x^{74} + x^{72} + x^{66} + x^{64} + x^{58} + x^{56} + x^{50} + x^{48} + x^{42} \\
&\quad + x^{40} + x^{34} + x^{32} + x^{26} + x^{24} + x^{18} + x^{16} + x^2 + 1,
\end{aligned}$$

and the initial terms are $[1, 0, 0, 0, 0, 0, 1, 0, 1, 0, 1, 0, 0, 0, 0, 0]$.

For $\phi(W^e, 0, 1)$:

$$\begin{aligned}
p_0(x) &= x^{87} + x^{84} + x^{81} + x^{80} + x^{78} + x^{76} + x^{74} + x^{72} + x^{70} + x^{68} + x^{65} \\
&\quad + x^{64} + x^{62} + x^{61} + x^{57} + x^{54} + x^{52} + x^{51} + x^{50} + x^{49} + x^{47} + x^{43} \\
&\quad + x^{40} + x^{38} + x^{37} + x^{35} + x^{29} + x^{28} + x^{27}, \\
p_3(x) &= x^{69} + x^{68} + x^{67} + x^{66} + x^{63} + x^{62} + x^{57} + x^{56} + x^{55} + x^{54} + x^{53} \\
&\quad + x^{52} + x^{51} + x^{50} + x^{49} + x^{48} + x^{47} + x^{46} + x^{45} + x^{44} + x^{43} + x^{42} \\
&\quad + x^{41} + x^{40} + x^{39} + x^{38} + x^{37} + x^{36} + x^{35} + x^{34} + x^{33} + x^{32} + x^{31} \\
&\quad + x^{30} + x^{25} + x^{24} + x^{21} + x^{20} + x^{19} + x^{18}, \\
p_6(x) &= x^{75} + x^{73} + x^{72} + x^{70} + x^{69} + x^{66} + x^{65} + x^{62} + x^{61} + x^{58} + x^{57} \\
&\quad + x^{55} + x^{54} + x^{52} + x^{35} + x^{33} + x^{32} + x^{30} + x^{29} + x^{26} + x^{25} + x^{22}
\end{aligned}$$

$$\begin{aligned}
& + x^{21} + x^{18} + x^{17} + x^{15} + x^{14} + x^{12}, \\
p_9(x) &= x^{81} + x^{80} + x^{73} + x^{72} + x^{71} + x^{70} + x^{69} + x^{68} + x^{65} + x^{64} + x^{63} \\
& + x^{62} + x^{59} + x^{58} + x^{57} + x^{56} + x^{55} + x^{54} + x^{47} + x^{46} + x^{41} + x^{40} \\
& + x^{33} + x^{32} + x^{31} + x^{30} + x^{29} + x^{28} + x^{25} + x^{24} + x^{23} + x^{22} + x^{19} \\
& + x^{18} + x^{17} + x^{16} + x^{15} + x^{14} + x^7 + x^6, \\
p_{12}(x) &= x^{87} + x^{84} + x^{79} + x^{76} + x^{75} + x^{72} + x^{71} + x^{68} + x^{67} + x^{64} + x^{63} \\
& + x^{60} + x^{59} + x^{56} + x^{55} + x^{52} + x^{51} + x^{48} + x^{47} + x^{44} + x^{43} + x^{40} \\
& + x^{39} + x^{36} + x^{35} + x^{32} + x^{31} + x^{28} + x^{27} + x^{24} + x^{23} + x^{20} + x^{19} \\
& + x^{16} + x^{15} + x^{12} + x^{11} + x^8 + x^3 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 1, 0, 1, 0, 1, 1, 0, 0, 0, 1, 1, 1, 0, 0]$.

For $\phi(W^e, 1, 0)$:

$$\begin{aligned}
p_0(x) &= x^{87} + x^{84} + x^{81} + x^{80} + x^{78} + x^{76} + x^{74} + x^{72} + x^{70} + x^{68} + x^{65} \\
& + x^{64} + x^{62} + x^{61} + x^{57} + x^{54} + x^{52} + x^{51} + x^{50} + x^{49} + x^{47} + x^{43} \\
& + x^{40} + x^{38} + x^{37} + x^{35} + x^{29} + x^{28} + x^{27}, \\
p_3(x) &= x^{69} + x^{68} + x^{67} + x^{66} + x^{63} + x^{62} + x^{57} + x^{56} + x^{55} + x^{54} + x^{53} \\
& + x^{52} + x^{51} + x^{50} + x^{49} + x^{48} + x^{47} + x^{46} + x^{45} + x^{44} + x^{43} + x^{42} \\
& + x^{41} + x^{40} + x^{39} + x^{38} + x^{37} + x^{36} + x^{35} + x^{34} + x^{33} + x^{32} + x^{31} \\
& + x^{30} + x^{25} + x^{24} + x^{21} + x^{20} + x^{19} + x^{18}, \\
p_6(x) &= x^{75} + x^{73} + x^{72} + x^{70} + x^{69} + x^{66} + x^{65} + x^{62} + x^{61} + x^{58} + x^{57} \\
& + x^{55} + x^{54} + x^{52} + x^{35} + x^{33} + x^{32} + x^{30} + x^{29} + x^{26} + x^{25} + x^{22} \\
& + x^{21} + x^{18} + x^{17} + x^{15} + x^{14} + x^{12}, \\
p_9(x) &= x^{81} + x^{80} + x^{73} + x^{72} + x^{71} + x^{70} + x^{69} + x^{68} + x^{65} + x^{64} + x^{63} \\
& + x^{62} + x^{59} + x^{58} + x^{57} + x^{56} + x^{55} + x^{54} + x^{47} + x^{46} + x^{41} + x^{40} \\
& + x^{33} + x^{32} + x^{31} + x^{30} + x^{29} + x^{28} + x^{25} + x^{24} + x^{23} + x^{22} + x^{19} \\
& + x^{18} + x^{17} + x^{16} + x^{15} + x^{14} + x^7 + x^6, \\
p_{12}(x) &= x^{87} + x^{84} + x^{79} + x^{76} + x^{75} + x^{72} + x^{71} + x^{68} + x^{67} + x^{64} + x^{63} \\
& + x^{60} + x^{59} + x^{56} + x^{55} + x^{52} + x^{51} + x^{48} + x^{47} + x^{44} + x^{43} + x^{40} \\
& + x^{39} + x^{36} + x^{35} + x^{32} + x^{31} + x^{28} + x^{27} + x^{24} + x^{23} + x^{20} + x^{19} \\
& + x^{16} + x^{15} + x^{12} + x^{11} + x^8 + x^3 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 1, 0, 1, 0, 1, 1, 0, 0, 0, 1, 1, 1, 0, 0]$.

For $\phi(W^e, 1, 1)$:

$$\begin{aligned}
p_0(x) &= x^{84} + x^{82} + x^{70} + x^{66} + x^{64} + x^{60} + x^{44} + x^{40} + x^{36} + x^{32} + x^{30}, \\
p_3(x) &= x^{80} + x^{74} + x^{68} + x^{66} + x^{58} + x^{56} + x^{54} + x^{52} + x^{40} + x^{34} + x^{28} \\
& + x^{26} + x^{18} + x^{16} + x^{14} + x^{12}, \\
p_6(x) &= x^{80} + x^{64} + x^{60} + x^{58} + x^{54} + x^{52} + x^{50} + x^{48} + x^{42} + x^{24} + x^{20}
\end{aligned}$$

$$\begin{aligned}
& + x^{18} + x^{14} + x^{12} + x^{10} + x^8 + x^2 + 1, \\
p_9(x) &= x^{80} + x^{74} + x^{70} + x^{64} + x^{60} + x^{54} + x^{50} + x^{44} + x^{40} + x^{34} + x^{30} \\
& + x^{24} + x^{20} + x^{14} + x^{10} + x^4, \\
p_{12}(x) &= x^{84} + x^{82} + x^{80} + x^4 + x^2 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 1, 0, 0, 0, 1, 0, 1, 0, 0, 0, 1, 0]$.

For $\phi(M^o, 0, 0)$:

$$\begin{aligned}
p_0(x) &= x^{87} + x^{84} + x^{83} + x^{81} + x^{80} + x^{79} + x^{78} + x^{75} + x^{74} + x^{71} + x^{69} \\
& + x^{66} + x^{63} + x^{62} + x^{60} + x^{58} + x^{55} + x^{54} + x^{52} + x^{49} + x^{48} + x^{47} \\
& + x^{45} + x^{44} + x^{43} + x^{42} + x^{41} + x^{37} + x^{33} + x^{32} + x^{30} + x^{28} + x^{27} \\
& + x^{26} + x^{24}, \\
p_3(x) &= x^{79} + x^{76} + x^{73} + x^{70} + x^{69} + x^{68} + x^{62} + x^{61} + x^{59} + x^{58} + x^{57} \\
& + x^{53} + x^{52} + x^{49} + x^{48} + x^{45} + x^{44} + x^{41} + x^{40} + x^{39} + x^{37} + x^{32} \\
& + x^{30} + x^{25} + x^{24} + x^{22} + x^{20} + x^{19} + x^{18} + x^{16}, \\
p_6(x) &= x^{83} + x^{81} + x^{80} + x^{78} + x^{76} + x^{74} + x^{72} + x^{69} + x^{68} + x^{66} + x^{62} \\
& + x^{60} + x^{56} + x^{53} + x^{50} + x^{49} + x^{48} + x^{45} + x^{44} + x^{40} + x^{38} + x^{37} \\
& + x^{34} + x^{33} + x^{26} + x^{25} + x^{24} + x^{22} + x^{21} + x^{17} + x^{12} + x^{10} + x^3 + 1, \\
p_9(x) &= x^{79} + x^{76} + x^{75} + x^{72} + x^{63} + x^{60} + x^{59} + x^{56} + x^{47} + x^{44} + x^{43} \\
& + x^{40} + x^{31} + x^{28} + x^{27} + x^{24} + x^{15} + x^{12} + x^{11} + x^8, \\
p_{12}(x) &= x^{87} + x^{84} + x^{71} + x^{68} + x^{67} + x^{64} + x^{55} + x^{52} + x^{51} + x^{48} + x^{39} \\
& + x^{36} + x^{35} + x^{32} + x^{23} + x^{20} + x^{19} + x^{16} + x^3 + 1,
\end{aligned}$$

and the initial terms are $[1, 0, 0, 0, 1, 1, 1, 1, 0, 0, 0, 0, 0, 1, 1, 0]$.

For $\phi(M^o, 0, 1)$:

$$\begin{aligned}
p_0(x) &= x^{90} + x^{88} + x^{84} + x^{83} + x^{82} + x^{81} + x^{77} + x^{76} + x^{74} + x^{72} + x^{69} \\
& + x^{68} + x^{67} + x^{66} + x^{65} + x^{64} + x^{62} + x^{59} + x^{58} + x^{57} + x^{56} + x^{50} \\
& + x^{49} + x^{46} + x^{45} + x^{43} + x^{39} + x^{34} + x^{32} + x^{31} + x^{30} + x^{28} + x^{27}, \\
p_3(x) &= x^{72} + x^{71} + x^{69} + x^{68} + x^{66} + x^{65} + x^{64} + x^{62} + x^{61} + x^{59} + x^{58} \\
& + x^{32} + x^{31} + x^{29} + x^{28} + x^{26} + x^{25} + x^{24} + x^{22} + x^{21} + x^{19} + x^{18}, \\
p_6(x) &= x^{78} + x^{74} + x^{72} + x^{68} + x^{66} + x^{60} + x^{54} + x^{50} + x^{48} + x^{46} + x^{44} \\
& + x^{42} + x^{40} + x^{36} + x^{30} + x^{24} + x^{22} + x^{18} + x^{16} + x^{12}, \\
p_9(x) &= x^{84} + x^{83} + x^{82} + x^{72} + x^{71} + x^{70} + x^{68} + x^{67} + x^{66} + x^{56} + x^{55} \\
& + x^{54} + x^{52} + x^{51} + x^{50} + x^{40} + x^{39} + x^{38} + x^{36} + x^{35} + x^{34} + x^{24} \\
& + x^{23} + x^{22} + x^{20} + x^{19} + x^{18} + x^8 + x^7 + x^6, \\
p_{12}(x) &= x^{90} + x^{88} + x^{82} + x^{80} + x^{78} + x^{76} + x^{62} + x^{60} + x^{46} + x^{44} + x^{30} \\
& + x^{28} + x^{14} + x^{12} + x^{10} + x^8 + x^2 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 1, 0, 0, 1, 1, 0, 0, 1, 1, 1, 0, 0, 0]$.

For $\phi(M^o, 1, 0)$:

$$\begin{aligned}
p_0(x) &= x^{88} + x^{84} + x^{82} + x^{81} + x^{80} + x^{78} + x^{77} + x^{76} + x^{75} + x^{74} + x^{72} \\
&\quad + x^{63} + x^{62} + x^{61} + x^{59} + x^{54} + x^{48} + x^{47} + x^{46} + x^{43} + x^{41} + x^{37} \\
&\quad + x^{36} + x^{35} + x^{31} + x^{30} + x^{29} + x^{28} + x^{27}, \\
p_3(x) &= x^{70} + x^{69} + x^{68} + x^{66} + x^{65} + x^{63} + x^{62} + x^{60} + x^{59} + x^{58} + x^{30} \\
&\quad + x^{29} + x^{28} + x^{26} + x^{25} + x^{23} + x^{22} + x^{20} + x^{19} + x^{18}, \\
p_6(x) &= x^{76} + x^{74} + x^{72} + x^{56} + x^{54} + x^{52} + x^{36} + x^{34} + x^{32} + x^{16} + x^{14} \\
&\quad + x^{12}, \\
p_9(x) &= x^{82} + x^{81} + x^{79} + x^{78} + x^{74} + x^{73} + x^{71} + x^{70} + x^{66} + x^{65} + x^{63} \\
&\quad + x^{62} + x^{58} + x^{57} + x^{55} + x^{54} + x^{50} + x^{49} + x^{47} + x^{46} + x^{42} + x^{41} \\
&\quad + x^{39} + x^{38} + x^{34} + x^{33} + x^{31} + x^{30} + x^{26} + x^{25} + x^{23} + x^{22} + x^{18} \\
&\quad + x^{17} + x^{15} + x^{14} + x^{10} + x^9 + x^7 + x^6, \\
p_{12}(x) &= x^{88} + x^{84} + x^{72} + x^{64} + x^{56} + x^{48} + x^{40} + x^{32} + x^{24} + x^{16} + x^4 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 1, 0, 1, 0, 0, 1, 0, 0, 0, 1, 1, 1, 0]$.

For $\phi(M^o, 1, 1)$:

$$\begin{aligned}
p_0(x) &= x^{87} + x^{84} + x^{83} + x^{81} + x^{80} + x^{78} + x^{76} + x^{75} + x^{74} + x^{73} + x^{71} \\
&\quad + x^{70} + x^{67} + x^{64} + x^{63} + x^{62} + x^{61} + x^{59} + x^{58} + x^{57} + x^{55} + x^{52} \\
&\quad + x^{51} + x^{50} + x^{49} + x^{48} + x^{47} + x^{46} + x^{44} + x^{43} + x^{41} + x^{40} + x^{39} \\
&\quad + x^{38} + x^{35} + x^{34} + x^{30}, \\
p_3(x) &= x^{79} + x^{76} + x^{73} + x^{71} + x^{70} + x^{69} + x^{67} + x^{65} + x^{64} + x^{63} + x^{61} \\
&\quad + x^{60} + x^{58} + x^{57} + x^{56} + x^{53} + x^{52} + x^{49} + x^{48} + x^{45} + x^{44} + x^{41} \\
&\quad + x^{40} + x^{39} + x^{37} + x^{32} + x^{31} + x^{30} + x^{28} + x^{27} + x^{23} + x^{18}, \\
p_6(x) &= x^{83} + x^{81} + x^{80} + x^{79} + x^{78} + x^{74} + x^{73} + x^{72} + x^{71} + x^{70} + x^{65} \\
&\quad + x^{61} + x^{60} + x^{59} + x^{58} + x^{57} + x^{54} + x^{53} + x^{51} + x^{50} + x^{49} + x^{45} \\
&\quad + x^{44} + x^{40} + x^{39} + x^{38} + x^{37} + x^{36} + x^{34} + x^{31} + x^{30} + x^{29} + x^{28} \\
&\quad + x^{24} + x^{19} + x^{18} + x^{16} + x^{14} + x^{12} + x^{11} + x^{10} + x^8 + x^3 + 1, \\
p_9(x) &= x^{79} + x^{76} + x^{75} + x^{72} + x^{63} + x^{60} + x^{59} + x^{56} + x^{47} + x^{44} + x^{43} \\
&\quad + x^{40} + x^{31} + x^{28} + x^{27} + x^{24} + x^{15} + x^{12} + x^{11} + x^8, \\
p_{12}(x) &= x^{87} + x^{84} + x^{71} + x^{68} + x^{67} + x^{64} + x^{55} + x^{52} + x^{51} + x^{48} + x^{39} \\
&\quad + x^{36} + x^{35} + x^{32} + x^{23} + x^{20} + x^{19} + x^{16} + x^3 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 1, 0, 1, 1, 0, 0, 0, 0, 1, 1, 0]$.

For $\phi(W^o, 0, 0)$:

$$\begin{aligned}
p_0(x) &= x^{87} + x^{84} + x^{83} + x^{81} + x^{80} + x^{79} + x^{78} + x^{75} + x^{74} + x^{71} + x^{69} \\
&\quad + x^{66} + x^{63} + x^{62} + x^{60} + x^{58} + x^{55} + x^{54} + x^{52} + x^{49} + x^{48} + x^{47} \\
&\quad + x^{45} + x^{44} + x^{43} + x^{42} + x^{41} + x^{37} + x^{33} + x^{32} + x^{30} + x^{28} + x^{27}
\end{aligned}$$

$$\begin{aligned}
& + x^{26} + x^{24}, \\
p_3(x) &= x^{79} + x^{76} + x^{73} + x^{70} + x^{69} + x^{68} + x^{62} + x^{61} + x^{59} + x^{58} + x^{57} \\
& + x^{53} + x^{52} + x^{49} + x^{48} + x^{45} + x^{44} + x^{41} + x^{40} + x^{39} + x^{37} + x^{32} \\
& + x^{30} + x^{25} + x^{24} + x^{22} + x^{20} + x^{19} + x^{18} + x^{16}, \\
p_6(x) &= x^{83} + x^{81} + x^{80} + x^{78} + x^{76} + x^{74} + x^{72} + x^{69} + x^{68} + x^{66} + x^{62} \\
& + x^{60} + x^{56} + x^{53} + x^{50} + x^{49} + x^{48} + x^{45} + x^{44} + x^{40} + x^{38} + x^{37} \\
& + x^{34} + x^{33} + x^{26} + x^{25} + x^{24} + x^{22} + x^{21} + x^{17} + x^{12} + x^{10} + x^3 + 1, \\
p_9(x) &= x^{79} + x^{76} + x^{75} + x^{72} + x^{63} + x^{60} + x^{59} + x^{56} + x^{47} + x^{44} + x^{43} \\
& + x^{40} + x^{31} + x^{28} + x^{27} + x^{24} + x^{15} + x^{12} + x^{11} + x^8, \\
p_{12}(x) &= x^{87} + x^{84} + x^{71} + x^{68} + x^{67} + x^{64} + x^{55} + x^{52} + x^{51} + x^{48} + x^{39} \\
& + x^{36} + x^{35} + x^{32} + x^{23} + x^{20} + x^{19} + x^{16} + x^3 + 1,
\end{aligned}$$

and the initial terms are $[1, 0, 0, 0, 1, 1, 1, 1, 0, 0, 0, 0, 1, 1, 0]$.

For $\phi(W^o, 0, 1)$:

$$\begin{aligned}
p_0(x) &= x^{88} + x^{84} + x^{82} + x^{81} + x^{80} + x^{78} + x^{77} + x^{76} + x^{75} + x^{74} + x^{72} \\
& + x^{63} + x^{62} + x^{61} + x^{59} + x^{54} + x^{48} + x^{47} + x^{46} + x^{43} + x^{41} + x^{37} \\
& + x^{36} + x^{35} + x^{31} + x^{30} + x^{29} + x^{28} + x^{27}, \\
p_3(x) &= x^{70} + x^{69} + x^{68} + x^{66} + x^{65} + x^{63} + x^{62} + x^{60} + x^{59} + x^{58} + x^{30} \\
& + x^{29} + x^{28} + x^{26} + x^{25} + x^{23} + x^{22} + x^{20} + x^{19} + x^{18}, \\
p_6(x) &= x^{76} + x^{74} + x^{72} + x^{56} + x^{54} + x^{52} + x^{36} + x^{34} + x^{32} + x^{16} + x^{14} \\
& + x^{12}, \\
p_9(x) &= x^{82} + x^{81} + x^{79} + x^{78} + x^{74} + x^{73} + x^{71} + x^{70} + x^{66} + x^{65} + x^{63} \\
& + x^{62} + x^{58} + x^{57} + x^{55} + x^{54} + x^{50} + x^{49} + x^{47} + x^{46} + x^{42} + x^{41} \\
& + x^{39} + x^{38} + x^{34} + x^{33} + x^{31} + x^{30} + x^{26} + x^{25} + x^{23} + x^{22} + x^{18} \\
& + x^{17} + x^{15} + x^{14} + x^{10} + x^9 + x^7 + x^6, \\
p_{12}(x) &= x^{88} + x^{84} + x^{72} + x^{64} + x^{56} + x^{48} + x^{40} + x^{32} + x^{24} + x^{16} + x^4 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 1, 0, 1, 0, 0, 0, 1, 0, 0, 0, 1, 1, 1, 0]$.

For $\phi(W^o, 1, 0)$:

$$\begin{aligned}
p_0(x) &= x^{90} + x^{88} + x^{84} + x^{83} + x^{82} + x^{81} + x^{77} + x^{76} + x^{74} + x^{72} + x^{69} \\
& + x^{68} + x^{67} + x^{66} + x^{65} + x^{64} + x^{62} + x^{59} + x^{58} + x^{57} + x^{56} + x^{50} \\
& + x^{49} + x^{46} + x^{45} + x^{43} + x^{39} + x^{34} + x^{32} + x^{31} + x^{30} + x^{28} + x^{27}, \\
p_3(x) &= x^{72} + x^{71} + x^{69} + x^{68} + x^{66} + x^{65} + x^{64} + x^{62} + x^{61} + x^{59} + x^{58} \\
& + x^{32} + x^{31} + x^{29} + x^{28} + x^{26} + x^{25} + x^{24} + x^{22} + x^{21} + x^{19} + x^{18}, \\
p_6(x) &= x^{78} + x^{74} + x^{72} + x^{68} + x^{66} + x^{60} + x^{54} + x^{50} + x^{48} + x^{46} + x^{44} \\
& + x^{42} + x^{40} + x^{36} + x^{30} + x^{24} + x^{22} + x^{18} + x^{16} + x^{12}, \\
p_9(x) &= x^{84} + x^{83} + x^{82} + x^{72} + x^{71} + x^{70} + x^{68} + x^{67} + x^{66} + x^{56} + x^{55}
\end{aligned}$$

$$\begin{aligned}
& + x^{54} + x^{52} + x^{51} + x^{50} + x^{40} + x^{39} + x^{38} + x^{36} + x^{35} + x^{34} + x^{24} \\
& + x^{23} + x^{22} + x^{20} + x^{19} + x^{18} + x^8 + x^7 + x^6, \\
p_{12}(x) = & x^{90} + x^{88} + x^{82} + x^{80} + x^{78} + x^{76} + x^{62} + x^{60} + x^{46} + x^{44} + x^{30} \\
& + x^{28} + x^{14} + x^{12} + x^{10} + x^8 + x^2 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 1, 0, 0, 1, 1, 0, 0, 1, 1, 1, 0, 0, 0]$.

For $\phi(W^o, 1, 1)$:

$$\begin{aligned}
p_0(x) = & x^{87} + x^{84} + x^{83} + x^{81} + x^{80} + x^{78} + x^{76} + x^{75} + x^{74} + x^{73} + x^{71} \\
& + x^{70} + x^{67} + x^{64} + x^{63} + x^{62} + x^{61} + x^{59} + x^{58} + x^{57} + x^{55} + x^{52} \\
& + x^{51} + x^{50} + x^{49} + x^{48} + x^{47} + x^{46} + x^{44} + x^{43} + x^{41} + x^{40} + x^{39} \\
& + x^{38} + x^{35} + x^{34} + x^{30}, \\
p_3(x) = & x^{79} + x^{76} + x^{73} + x^{71} + x^{70} + x^{69} + x^{67} + x^{65} + x^{64} + x^{63} + x^{61} \\
& + x^{60} + x^{58} + x^{57} + x^{56} + x^{53} + x^{52} + x^{49} + x^{48} + x^{45} + x^{44} + x^{41} \\
& + x^{40} + x^{39} + x^{37} + x^{32} + x^{31} + x^{30} + x^{28} + x^{27} + x^{23} + x^{18}, \\
p_6(x) = & x^{83} + x^{81} + x^{80} + x^{79} + x^{78} + x^{74} + x^{73} + x^{72} + x^{71} + x^{70} + x^{65} \\
& + x^{61} + x^{60} + x^{59} + x^{58} + x^{57} + x^{54} + x^{53} + x^{51} + x^{50} + x^{49} + x^{45} \\
& + x^{44} + x^{40} + x^{39} + x^{38} + x^{37} + x^{36} + x^{34} + x^{31} + x^{30} + x^{29} + x^{28} \\
& + x^{24} + x^{19} + x^{18} + x^{16} + x^{14} + x^{12} + x^{11} + x^{10} + x^8 + x^3 + 1, \\
p_9(x) = & x^{79} + x^{76} + x^{75} + x^{72} + x^{63} + x^{60} + x^{59} + x^{56} + x^{47} + x^{44} + x^{43} \\
& + x^{40} + x^{31} + x^{28} + x^{27} + x^{24} + x^{15} + x^{12} + x^{11} + x^8, \\
p_{12}(x) = & x^{87} + x^{84} + x^{71} + x^{68} + x^{67} + x^{64} + x^{55} + x^{52} + x^{51} + x^{48} + x^{39} \\
& + x^{36} + x^{35} + x^{32} + x^{23} + x^{20} + x^{19} + x^{16} + x^3 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 1, 0, 1, 1, 0, 0, 0, 0, 1, 1, 0]$.

Below is the transition function and output function of an 2-automaton that generates $T = M_{0,0}^o$:

Transition function $(n, j) \mapsto \delta(n, j)$ ($\Lambda(n) := [\delta(n, 0), \delta(n, 1)]$):

n	$\Lambda(n)$	n	$\Lambda(n)$	n	$\Lambda(n)$	n	$\Lambda(n)$	n	$\Lambda(n)$
0	[1, 2]	74	[125, 42]	148	[224, 225]	222	[58, 283]	296	[251, 279]
1	[3, 4]	75	[97, 106]	149	[52, 118]	223	[161, 55]	297	[114, 260]
2	[5, 6]	76	[126, 31]	150	[47, 109]	224	[284, 285]	298	[343, 238]
3	[1, 7]	77	[127, 128]	151	[120, 226]	225	[286, 188]	299	[345, 271]
4	[8, 9]	78	[129, 9]	152	[132, 41]	226	[73, 20]	300	[346, 278]
5	[10, 11]	79	[130, 131]	153	[227, 228]	227	[285, 287]	301	[271, 347]
6	[12, 13]	80	[103, 112]	154	[34, 221]	228	[288, 289]	302	[120, 262]
7	[14, 15]	81	[132, 125]	155	[229, 32]	229	[290, 152]	303	[227, 348]
8	[16, 17]	82	[133, 134]	156	[230, 231]	230	[291, 189]	304	[38, 221]
9	[18, 19]	83	[30, 135]	157	[223, 232]	231	[292, 62]	305	[230, 229]
10	[20, 21]	84	[127, 29]	158	[225, 36]	232	[23, 293]	306	[136, 256]
11	[22, 23]	85	[136, 137]	159	[233, 35]	233	[167, 294]	307	[7, 244]

12	[24, 25]	86	[138, 139]	160	[234, 49]	234	[295, 288]	308	[245, 140]
13	[26, 27]	87	[140, 141]	161	[111, 235]	235	[182, 296]	309	[349, 34]
14	[28, 29]	88	[41, 142]	162	[236, 44]	236	[297, 298]	310	[276, 350]
15	[30, 31]	89	[118, 50]	163	[237, 124]	237	[299, 300]	311	[351, 258]
16	[32, 33]	90	[51, 46]	164	[124, 238]	238	[300, 301]	312	[275, 242]
17	[34, 12]	91	[122, 143]	165	[105, 98]	239	[180, 191]	313	[345, 258]
18	[35, 36]	92	[144, 140]	166	[45, 47]	240	[298, 302]	314	[273, 253]
19	[37, 38]	93	[113, 122]	167	[104, 97]	241	[303, 18]	315	[277, 232]
20	[39, 40]	94	[145, 146]	168	[12, 239]	242	[304, 305]	316	[225, 227]
21	[41, 42]	95	[147, 148]	169	[231, 32]	243	[215, 306]	317	[224, 35]
22	[43, 44]	96	[149, 88]	170	[231, 240]	244	[4, 307]	318	[266, 352]
23	[45, 46]	97	[150, 151]	171	[10, 222]	245	[308, 185]	319	[222, 277]
24	[47, 48]	98	[152, 149]	172	[241, 242]	246	[169, 309]	320	[353, 231]
25	[49, 50]	99	[153, 154]	173	[243, 244]	247	[310, 292]	321	[140, 131]
26	[51, 47]	100	[155, 156]	174	[245, 130]	248	[289, 150]	322	[112, 105]
27	[52, 49]	101	[157, 158]	175	[241, 246]	249	[311, 312]	323	[133, 251]
28	[53, 54]	102	[146, 159]	176	[247, 137]	250	[313, 314]	324	[99, 352]
29	[55, 56]	103	[160, 161]	177	[248, 104]	251	[207, 197]	325	[267, 350]
30	[57, 58]	104	[162, 163]	178	[46, 48]	252	[315, 170]	326	[240, 33]
31	[59, 60]	105	[164, 165]	179	[121, 114]	253	[316, 317]	327	[36, 348]
32	[61, 59]	106	[166, 167]	180	[249, 41]	254	[60, 72]	328	[353, 229]
33	[62, 63]	107	[168, 169]	181	[250, 251]	255	[190, 57]	329	[277, 354]
34	[54, 26]	108	[170, 171]	182	[252, 253]	256	[151, 318]	330	[11, 223]
35	[64, 65]	109	[172, 173]	183	[254, 255]	257	[319, 320]	331	[35, 227]
36	[66, 67]	110	[174, 175]	184	[247, 256]	258	[15, 214]	332	[355, 11]
37	[68, 69]	111	[176, 76]	185	[8, 257]	259	[312, 321]	333	[144, 130]
38	[70, 71]	112	[177, 166]	186	[258, 259]	260	[294, 208]	334	[252, 274]
39	[72, 73]	113	[91, 178]	187	[260, 261]	261	[71, 322]	335	[221, 239]
40	[74, 75]	114	[179, 180]	188	[249, 125]	262	[283, 297]	336	[355, 222]
41	[76, 77]	115	[181, 182]	189	[116, 235]	263	[323, 172]	337	[240, 356]
42	[78, 79]	116	[183, 184]	190	[40, 262]	264	[324, 286]	338	[32, 356]
43	[63, 74]	117	[185, 186]	191	[104, 105]	265	[287, 325]	339	[40, 226]
44	[80, 81]	118	[69, 187]	192	[96, 234]	266	[326, 319]	340	[258, 347]
45	[82, 4]	119	[188, 68]	193	[115, 111]	267	[327, 212]	341	[130, 141]
46	[83, 83]	120	[189, 190]	194	[263, 243]	268	[158, 328]	342	[206, 357]
47	[84, 85]	121	[191, 66]	195	[129, 257]	269	[329, 330]	343	[306, 14]
48	[86, 87]	122	[192, 193]	196	[264, 265]	270	[331, 332]	344	[293, 282]
49	[88, 89]	123	[194, 195]	197	[28, 128]	271	[77, 84]	345	[358, 359]
50	[90, 91]	124	[196, 183]	198	[266, 100]	272	[333, 334]	346	[360, 304]
51	[92, 86]	125	[197, 196]	199	[267, 268]	273	[335, 316]	347	[359, 361]
52	[93, 90]	126	[198, 93]	200	[269, 270]	274	[330, 336]	348	[65, 291]
53	[94, 95]	127	[199, 162]	201	[36, 228]	275	[337, 155]	349	[27, 24]
54	[96, 52]	128	[89, 200]	202	[271, 259]	276	[338, 331]	350	[211, 362]
55	[97, 98]	129	[201, 147]	203	[272, 7]	277	[21, 339]	351	[363, 78]

56	[99, 100]	130	[85, 176]	204	[273, 274]	278	[17, 5]	352	[154, 364]
57	[101, 102]	131	[175, 202]	205	[263, 7]	279	[314, 340]	353	[322, 61]
58	[103, 104]	132	[203, 204]	206	[275, 246]	280	[334, 341]	354	[81, 284]
59	[105, 106]	133	[205, 206]	207	[254, 265]	281	[193, 160]	355	[365, 164]
60	[107, 108]	134	[184, 207]	208	[49, 119]	282	[124, 342]	356	[163, 365]
61	[46, 109]	135	[208, 209]	209	[276, 268]	283	[237, 343]	357	[134, 366]
62	[110, 111]	136	[200, 179]	210	[143, 261]	284	[122, 260]	358	[272, 243]
63	[112, 97]	137	[210, 53]	211	[11, 277]	285	[125, 142]	359	[346, 139]
64	[113, 114]	138	[6, 211]	212	[229, 240]	286	[344, 234]	360	[223, 354]
65	[115, 116]	139	[212, 213]	213	[233, 225]	287	[260, 281]	361	[243, 280]
66	[116, 117]	140	[214, 215]	214	[264, 255]	288	[123, 343]	362	[37, 34]
67	[118, 119]	141	[216, 217]	215	[126, 135]	289	[39, 120]	363	[351, 271]
68	[44, 120]	142	[204, 218]	216	[138, 278]	290	[236, 39]	364	[349, 38]
69	[111, 117]	143	[178, 219]	217	[134, 279]	291	[114, 143]	365	[234, 118]
70	[121, 122]	144	[220, 216]	218	[7, 280]	292	[248, 112]	366	[195, 367]
71	[123, 124]	145	[221, 13]	219	[143, 281]	293	[44, 40]	367	[251, 366]
72	[43, 39]	146	[38, 12]	220	[250, 134]	294	[343, 342]		
73	[110, 116]	147	[222, 223]	221	[282, 210]	295	[344, 52]		

Output function $n \mapsto \tau(n)$:

n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$
0	0	53	1	106	1	159	1	212	1	265	0	318	0
1	0	54	0	107	0	160	1	213	1	266	0	319	1
2	0	55	0	108	0	161	1	214	1	267	0	320	0
3	0	56	1	109	1	162	1	215	0	268	1	321	1
4	1	57	1	110	1	163	0	216	0	269	0	322	0
5	0	58	1	111	1	164	1	217	1	270	0	323	1
6	0	59	1	112	0	165	1	218	1	271	0	324	1
7	1	60	0	113	0	166	1	219	1	272	0	325	0
8	1	61	1	114	1	167	1	220	0	273	1	326	0
9	0	62	1	115	0	168	0	221	1	274	0	327	0
10	0	63	0	116	0	169	0	222	1	275	0	328	0
11	0	64	0	117	1	170	0	223	1	276	1	329	0
12	0	65	0	118	1	171	0	224	0	277	0	330	0
13	0	66	0	119	1	172	1	225	1	278	0	331	0
14	1	67	1	120	0	173	0	226	1	279	1	332	1
15	1	68	1	121	1	174	1	227	1	280	0	333	0
16	1	69	1	122	0	175	1	228	1	281	0	334	0
17	0	70	1	123	1	176	1	229	1	282	1	335	1
18	0	71	1	124	1	177	0	230	1	283	0	336	1
19	1	72	0	125	1	178	1	231	0	284	0	337	0
20	0	73	1	126	0	179	1	232	1	285	1	338	1
21	0	74	1	127	0	180	1	233	1	286	1	339	1
22	0	75	0	128	1	181	0	234	1	287	0	340	1

23	1	76	0	129	0	182	0	235	0	288	1	341	0
24	0	77	0	130	0	183	0	236	1	289	0	342	0
25	0	78	0	131	1	184	1	237	0	290	1	343	0
26	0	79	0	132	0	185	1	238	1	291	1	344	1
27	0	80	1	133	1	186	1	239	1	292	0	345	0
28	1	81	0	134	1	187	0	240	0	293	1	346	1
29	0	82	1	135	0	188	1	241	1	294	0	347	1
30	1	83	1	136	0	189	0	242	1	295	1	348	0
31	1	84	0	137	1	190	1	243	0	296	0	349	0
32	1	85	0	138	0	191	1	244	1	297	1	350	0
33	1	86	0	139	1	192	0	245	1	298	0	351	1
34	0	87	1	140	1	193	0	246	0	299	0	352	0
35	0	88	0	141	0	194	1	247	1	300	1	353	0
36	0	89	1	142	1	195	0	248	0	301	0	354	0
37	1	90	0	143	1	196	1	249	1	302	0	355	1
38	1	91	0	144	0	197	1	250	0	303	1	356	0
39	0	92	0	145	1	198	0	251	0	304	1	357	1
40	1	93	0	146	1	199	0	252	0	305	1	358	0
41	0	94	1	147	1	200	0	253	1	306	0	359	1
42	0	95	1	148	0	201	0	254	0	307	1	360	1
43	0	96	0	149	0	202	0	255	1	308	1	361	0
44	1	97	0	150	0	203	0	256	0	309	0	362	1
45	1	98	0	151	0	204	1	257	1	310	1	363	1
46	1	99	1	152	0	205	1	258	1	311	1	364	0
47	0	100	1	153	1	206	0	259	0	312	0	365	1
48	0	101	1	154	0	207	0	260	0	313	0	366	0
49	0	102	1	155	1	208	0	261	1	314	1	367	0
50	0	103	1	156	1	209	1	262	0	315	0		
51	0	104	1	157	1	210	1	263	1	316	1		
52	0	105	1	158	1	211	0	264	1	317	0		

UNIVERSITÉ DE STRASBOURG, CNRS, IRMA UMR 7501, F-67000 STRASBOURG, FRANCE
Email address: `guoniu.han@unistra.fr`

SCHOOL OF MATHEMATICS AND STATISTICS, HUAZHONG UNIVERSITY OF SCIENCE AND TECHNOLOGY, WUHAN, PR CHINA
Email address: `huyining@hust.edu.cn`