

PROOF OF THE CONTINUED FRACTION CONJECTURE FOR

$$(a, b) = (z^2 + z + 1, z^3 + z^2 + z + 1)$$

GUO-NIU HAN* AND YINING HU*

ABSTRACT. The continued fraction conjecture claims that the continued fraction defined by the (a, b) -Thue-Morse sequence is algebraic over $\mathbb{F}_2(x)$ for all pairs of distinct elements (a, b) in $\mathbb{F}_2[x] \setminus \mathbb{F}_2$. In this paper we prove the conjecture for $(a, b) = (z^2 + z + 1, z^3 + z^2 + z + 1)$.

1. INTRODUCTION

Let $\mathbb{F}_2$ be the finite field of cardinality 2. Given a sequence of polynomials $\mathbf{a} = (a_0(z), a_1(z), a_2(z), \dots)$ of elements from $\mathbb{F}_2[z] \setminus \mathbb{F}_2$, we define the infinite continued fraction

$$(1.1) \quad \text{CF}(\mathbf{a}(z)) = \frac{1}{a_0(z) + \frac{1}{a_1(z) + \frac{1}{a_2(z) + \frac{1}{\ddots}}}}$$

as the limit of the finite continued fractions

$$(1.2) \quad \text{CF}_n(\mathbf{a}(z)) = \frac{1}{a_0(z) + \frac{1}{a_1(z) + \frac{1}{\ddots + \frac{1}{a_n(z)}}}} \in \mathbb{F}_2((1/z)).$$

Conversely, each power series in $\mathbb{F}_2[[1/z]]$ without constant term

$$(1.3) \quad c_{-1}z^{-1} + c_{-2}z^{-2} + c_{-3}z^{-3} + \dots$$

admits a continued fraction expansion of form (1.1).

Let a and b be two distinct elements from $\mathbb{F}_2[z] \setminus \mathbb{F}_2$. We consider the continued fraction (1.1) associated with the (a, b) -Thue-Morse sequence

$$\mathbf{t} = (t_0(z), t_1(z), t_2(z), \dots) = (a, b, b, a, b, a, a, b, \dots)$$

that we shall denote

$$(1.4) \quad \text{CF}(a, b) := \text{CF}(\mathbf{t}).$$

Date: 9/04/2020 08:53:08 .

2010 Mathematics Subject Classification. 11B85, 11J70, 11B50, 11Y65, 05A15.

Key words and phrases. automatic sequence, continued fraction, Thue-Morse sequence.

* This paper was automatically generated from a computer program developed by the authors, with 10.7 minutes CPU time used.

Theorem 1.1. *When $(a, b) = (z^2 + z + 1, z^3 + z^2 + z + 1)$, the two power series $\text{CF}(a, b)$ and $\text{CF}(b, a)$ are algebraic over $\mathbb{F}_2(z)$, with minimal polynomials of the form*

$$p_4(z)y^4 + p_3(z)y^3 + p_2(z)y^2 + p_1(z)y + p_0(z) = 0$$

where, for $\text{CF}(a, b)$,

$$\begin{aligned} p_0(z) &= z^{15} + z^{13} + z^9 + z^8 + z^7 + z^5 + z^4 + z^3 + z^2 + 1, \\ p_1(z) &= z^{18} + z^{16} + z^{15} + z^{13} + z^{12} + z^{10} + z^9 + z^8 + z^7 + z^6 + z^5 + z^3, \\ p_2(z) &= z^{20} + z^{16} + z^{10} + z^8 + z^6 + 1, \\ p_3(z) &= z^{18} + z^{16} + z^{15} + z^{13} + z^{12} + z^{10} + z^9 + z^8 + z^7 + z^6 + z^5 + z^3, \\ p_4(z) &= z^{16} + z^{15} + z^{13} + z^{10} + z^9 + z^8 + z^7 + z^6 + z^5 + z^4 + z^3 + z^2 + 1, \end{aligned}$$

and for $\text{CF}(b, a)$,

$$\begin{aligned} p_0(z) &= z^{15} + z^{13} + z^{12} + z^9 + z^8 + z^7 + z^5 + z^4 + z^3 + z^2 + 1, \\ p_1(z) &= z^{18} + z^{16} + z^{15} + z^{13} + z^{12} + z^{10} + z^9 + z^8 + z^7 + z^6 + z^5 + z^3, \\ p_2(z) &= z^{20} + z^{16} + z^{10} + z^8 + z^6 + 1, \\ p_3(z) &= z^{18} + z^{16} + z^{15} + z^{13} + z^{12} + z^{10} + z^9 + z^8 + z^7 + z^6 + z^5 + z^3, \\ p_4(z) &= z^{16} + z^{15} + z^{13} + z^{12} + z^{10} + z^9 + z^8 + z^7 + z^6 + z^5 + z^4 + z^3 + z^2 + 1. \end{aligned}$$

2. PROOF

Define the sequences of polynomials $(P_n(z))_{n \geq 0}$ and $(Q_n(z))_{n \geq 0}$ by

$$\begin{pmatrix} P_n(z) & Q_n(z) \\ P_{n-1}(z) & Q_{n-1}(z) \end{pmatrix} := \begin{pmatrix} t_n(z) & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} t_{n-1}(z) & 1 \\ 1 & 0 \end{pmatrix} \cdots \begin{pmatrix} t_0(z) & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}.$$

By the basic properties of continued fractions, we have

$$(2.1) \quad \text{CF}_n(a, b) = \frac{P_n(z)}{Q_n(z)} \in \mathbb{F}_2((1/z)).$$

As formula (2.1) is not efficient for calculating $\text{CF}_n(a, b)$, a fast method is to be derived and described as follows. Define

$$\begin{aligned} M_n(x) &= x^{\deg(t_{2^n-1})} \begin{pmatrix} t_{2^n-1}(1/x) & 1 \\ 1 & 0 \end{pmatrix} \times x^{\deg(t_{2^n-2})} \begin{pmatrix} t_{2^n-2}(1/x) & 1 \\ 1 & 0 \end{pmatrix} \times \\ &\quad \cdots \times x^{\deg(t_0)} \begin{pmatrix} t_0(1/x) & 1 \\ 1 & 0 \end{pmatrix}, \\ W_n(x) &= x^{\deg(\bar{t}_{2^n-1})} \begin{pmatrix} \bar{t}_{2^n-1}(1/x) & 1 \\ 1 & 0 \end{pmatrix} \times x^{\deg(\bar{t}_{2^n-2})} \begin{pmatrix} \bar{t}_{2^n-2}(1/x) & 1 \\ 1 & 0 \end{pmatrix} \times \\ &\quad \cdots \times x^{\deg(\bar{t}_0)} \begin{pmatrix} \bar{t}_0(1/x) & 1 \\ 1 & 0 \end{pmatrix}, \end{aligned}$$

where $\bar{\mathbf{t}}$ is the (b, a) -Thue-Morse sequence. By the properties of the Thue-Morse sequence, we have for $n \geq 0$

$$\begin{aligned} M_{n+1}(x) &= W_n(x) \cdot M_n(x), \\ W_{n+1}(x) &= M_n(x) \cdot W_n(x). \end{aligned}$$

Let $x := 1/z$. For each non-zero polynomial $P(z)$, define $\tilde{P}(x)$ to be $P(1/x)$. Then,

$$\text{CF}_n(a, b) = \frac{P_n(z)}{Q_n(z)} = \frac{\tilde{P}_n(x)}{\tilde{Q}_n(x)} \in \mathbb{F}_2((x)) = \mathbb{F}_2((1/z)).$$

Comparing the definition of $M_n(x)$ with the definition of $P_n(x)$ and $Q_n(x)$, we see that

$$\begin{aligned} M_n(x)_{0,1} &= x^{d_n} \tilde{P}_{2^n-1}(x), \\ M_n(x)_{0,0} &= x^{d_n} \tilde{Q}_{2^n-1}(x), \end{aligned}$$

for some positive integer d_n . Hence,

$$(2.2) \quad \text{CF}_{2^{2^n}-1}(a, b) = \frac{\tilde{P}_{2^{2^n}-1}(x)}{\tilde{Q}_{2^{2^n}-1}(x)} = \frac{M_{2^n}(x)_{0,1}}{M_{2^n}(x)_{0,0}}.$$

Therefore, by the convergence theorem of continued fraction, the algebraicity of $\text{CF}(a, b)$ will be established if it is shown that both $M_{2^n}(x)_{0,1}$ and $M_{2^n}(x)_{0,0}$ converge to algebraic series in $\mathbb{F}_2[[x]]$.

Actually, we will prove that for all $0 \leq i, j \leq 1$ the four sequences $(M_{2^n}(x)_{i,j})_n$, $(M_{2^{n+1}}(x)_{i,j})_n$, $(W_{2^n}(x)_{i,j})_n$, and $(W_{2^{n+1}}(x)_{i,j})_n$ converge to algebraic series in $\mathbb{F}_2[[x]]$. For this purpose, we define four 2×2 matrices M^e, M^o, W^e, W^o as follows: For each $T \in \{M^e, M^o, W^e, W^o\}$ and $0 \leq i, j \leq 1$, $T_{i,j}$ is defined to be the unique solution in $\mathbb{F}_2[[x]]$ of the polynomial $\phi(T, i, j)$ under certain initial conditions; the polynomials $\phi(T, i, j)$ and initial conditions are given in Section 3. We will prove that these four matrices, whose components are algebraic by definition, are the limits of $(M_{2^n}(x))_n$, $(M_{2^{n+1}}(x))_n$, $(W_{2^n}(x))_n$, and $(W_{2^{n+1}}(x))_n$.

Let us explain how the polynomials $\phi(T, i, j)$ and initial conditions are found, and why the solutions exist and are unique. For $0 \leq i, j \leq 1$, the coefficients of the polynomial $\phi(M^e, i, j)$ (resp. $\phi(M^o, i, j)$, $\phi(W^e, i, j)$, and $\phi(W^o, i, j)$) are the Padé-Hermite approximants of type

$$(125, 125, 125, 125, 125)$$

of the vector

$$(1, T^3, T^6, T^9, T^{12}),$$

where $T = M_{12,i,j}$ (resp. $M_{11,i,j}$, $W_{12,i,j}$, and $W_{11,i,j}$), found by the Derksen algorithm. We take the first 8 terms of $M_{12,i,j}$ (resp. $M_{11,i,j}$, $W_{12,i,j}$, and $W_{11,i,j}$) as initial conditions for $\phi(M^e, i, j)$ (resp. $\phi(M^o, i, j)$, $\phi(W^e, i, j)$, and $\phi(W^o, i, j)$). The following fact will be used to ensure that the solution exists and is unique: let $P(x, y) \in \mathbb{F}_2[x, y]$ and for each series $f(x) = \sum_{n=0}^{\infty} a_n x^n \in \mathbb{F}_2[[x]]$ denote the partial sum $\sum_{j=0}^{n-1} a_j x^j$ by $f_n(x)$ for $n \geq 0$. If for some $n \geq 0$ and $a_0, a_1, \dots, a_{n-1} \in \mathbb{F}_2$ $P(x, \sum_{j=0}^{n-1} a_j x^j) = O(x^n)$ and $Q(x, y) := P(x, \sum_{j=0}^{n-1} a_j x^j + x^n y)$ can be written as $x^m \sum_{j=0}^{\infty} q_j(x) y^j$ where $q_j(x)$ are polynomials for $j \geq 0$, $q_1(0) = 1$, and $q_j(0) = 0$ for $j > 1$, then there exists a unique solution $f(x) \in \mathbb{F}_2[[x]]$ of $P(x, f(x)) = 0$ that satisfies the initial condition $f_n(x) = \sum_{j=0}^{n-1} a_j x^j$.

We state two lemmas concerning the four matrices M^e, M^o, W^e, W^o . The first one is about relations between them; the second, about the structure of the each matrix.

Lemma 2.1. *We have*

$$(2.3) \quad M^e = W^o \cdot M^o,$$

$$(2.4) \quad M^o = W^e \cdot M^e,$$

$$(2.5) \quad W^e = M^o \cdot W^o,$$

$$(2.6) \quad W^o = M^e \cdot W^e.$$

Proof. We give the proof of the identity

$$M_{0,0}^e = W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o,$$

the proofs of the others being similar.

First, we compute the minimal polynomials of $W_{0,0}^o M_{0,0}^o$ and $W_{0,1}^o M_{1,0}^o$. We know that

$$P(x, y) = \text{Res}_z (\phi(W^o, 0, 0)(x, z), z^{12} \cdot \phi(M^o, 0, 0)(x, y/z))$$

is an annihilating polynomial of $W_{0,0}^o M_{0,0}^o$. We use Padé-Hermite approximation to find a candidate for the minimal polynomial of $W_{0,0}^o M_{0,0}^o$, that will be called $\phi_0(x, y)$. To prove that $\phi_0(x, y)$ is indeed the minimal polynomial, it suffices to prove that it is an irreducible factor of $P(x, y)$ of multiplicity m and that $Q(x, y) := P(x, y)/\phi_0(x, y)^m$ is not an annihilating polynomial of $W_{0,0}^o M_{0,0}^o$. We verify the first point directly. For the second point, we truncate $W_{0,0}^o M_{0,0}^o$ to order 450 and substitute it for y in $Q(x, y)$. We get a series of valuation less than 450, which proves that $Q(x, y)$ is not an annihilating polynomial of $W_{0,0}^o M_{0,0}^o$. We find the minimal polynomial $\phi_1(x, y)$ of $W_{0,1}^o M_{1,0}^o$ in a similar way.

Now we prove that $\phi(M^e, 0, 0)$ is the minimal polynomial of $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$. We know that

$$S(x, y) = \text{Res}_z (\phi_0(x, z), \phi_1(x, y + z))$$

is an annihilating of $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$. We verify that $\phi(M^e, 0, 0)$ is an irreducible factor of $S(x, y)$ of multiplicity μ , and that $Q(x, y) := S(x, y)/\phi(M^e, 0, 0)^\mu$ is not an annihilating polynomial of $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$. To see the last point, we truncate $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$ to order 550 and substitute it for y in $Q(x, y)$. We get a series of valuation less than 550, and therefore $Q(x, y)$ is not an annihilating polynomial of $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$.

Finally, the first 16 terms of $M_{0,0}^e$ and $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$ coincide. As these first terms determine a unique solution of $\phi(M^e, 0, 0)$, we know that the two series are one and the same. $\square$

Define

$$R^e = \begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix} \quad \text{and} \quad R^o = \begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix}.$$

Lemma 2.2. *For all integers $k \geq 2$ and $u = 2^{2k-1}$, the following identities hold.*

$$\begin{aligned} M^e[:10u] &= M^e[:5u] + x^{2u} M^e[:3u] + x^{3u} M^e[:2u] + x^{4u} M^e[:u] + x^{2u} M^e[:5u] \\ &\quad + x^{4u} M^e[:3u] + x^{5u} M^e[:2u] + x^{6u} M^e[:u] + x^{3u} M^e[:5u] \\ &\quad + x^{5u} M^e[:3u] + x^{6u} M^e[:2u] + x^{7u} M^e[:u] + x^{5u} M^e[:5u] \\ &\quad + x^{7u} M^e[:3u] + x^{6u} M^e[:4u] + x^{9u} M^e[:u] + (x^{5u} + x^{7u} + x^{8u}) R^e, \\ W^e[:10u] &= W^e[:5u] + x^{2u} W^e[:3u] + x^{3u} W^e[:2u] + x^{4u} W^e[:u] + x^{2u} W^e[:5u] \\ &\quad + x^{4u} W^e[:3u] + x^{5u} W^e[:2u] + x^{6u} W^e[:u] + x^{3u} W^e[:5u] \end{aligned}$$

$$\begin{aligned}
& + x^{5u}W^e[:3u] + x^{6u}W^e[:2u] + x^{7u}W^e[:u] + x^{5u}W^e[:5u] \\
& + x^{7u}W^e[:3u] + x^{6u}W^e[:4u] + x^{9u}W^e[:u] + (x^{5u} + x^{7u} + x^{8u})R^e.
\end{aligned}$$

For all integers $k \geq 2$ and $u = 2^{2k}$, the following identities hold.

$$\begin{aligned}
M^o[:10u] &= M^o[:5u] + x^{2u}M^o[:3u] + x^{3u}M^o[:2u] + x^{4u}M^o[:u] + x^{2u}M^o[:5u] \\
& + x^{4u}M^o[:3u] + x^{5u}M^o[:2u] + x^{6u}M^o[:u] + x^{3u}M^o[:5u] \\
& + x^{5u}M^o[:3u] + x^{6u}M^o[:2u] + x^{7u}M^o[:u] + x^{5u}M^o[:5u] \\
& + x^{7u}M^o[:3u] + x^{6u}M^o[:4u] + x^{9u}M^o[:u] + (x^{5u} + x^{7u} + x^{8u})R^o, \\
W^o[:10u] &= W^o[:5u] + x^{2u}W^o[:3u] + x^{3u}W^o[:2u] + x^{4u}W^o[:u] + x^{2u}W^o[:5u] \\
& + x^{4u}W^o[:3u] + x^{5u}W^o[:2u] + x^{6u}W^o[:u] + x^{3u}W^o[:5u] \\
& + x^{5u}W^o[:3u] + x^{6u}W^o[:2u] + x^{7u}W^o[:u] + x^{5u}W^o[:5u] \\
& + x^{7u}W^o[:3u] + x^{6u}W^o[:4u] + x^{9u}W^o[:u] + (x^{5u} + x^{7u} + x^{8u})R^o.
\end{aligned}$$

Proof. To prove Lemma 2.2 we first construct an automaton for each sequence concerned, and then transform the conditions on infinitely many k 's into finitely many conditions on the states of the automaton. In the following, we will prove that for $T = M_{0,0}^o$, for all integer $k \geq 2$ and $u = 2^{2k}$,

$$\begin{aligned}
T[:10u] &= T[:5u] + x^{2u}T[:3u] + x^{3u}T[:2u] + x^{4u}T[:u] + x^{2u}T[:5u] + x^{4u}T[:3u] \\
& + x^{5u}T[:2u] + x^{6u}T[:u] + x^{3u}T[:5u] + x^{5u}T[:3u] + x^{6u}T[:2u] \\
& + x^{7u}T[:u] + x^{5u}T[:5u] + x^{7u}T[:3u] + x^{6u}T[:4u] + x^{9u}T[:u] \\
& + (x^{5u} + x^{7u} + x^{8u}).
\end{aligned}$$

The proof of the other 15 cases are similar. We break down the above identity into 5 parts:

$$\begin{aligned}
x^5u &= x^{5u}T[:u] + x^{4u}T[u:2u] + x^{3u}T[2u:3u] + x^{2u}T[3u:4u] \\
& + T[5u:6u], \\
0 &= x^{6u}T[:u] + x^{5u}T[u:2u] + x^{4u}T[2u:3u] + x^{3u}T[3u:4u] \\
& + x^{2u}T[4u:5u] + T[6u:7u], \\
x^7u &= x^{3u}T[4u:5u] + T[7u:8u], \\
x^8u &= x^{7u}T[u:2u] + x^{6u}T[2u:3u] + x^{5u}T[3u:4u] + T[8u:9u], \\
0 &= x^{9u}T[:u] + x^{7u}T[2u:3u] + x^{6u}T[3u:4u] + x^{5u}T[4u:5u] \\
& + T[9u:10u],
\end{aligned}$$

which can be rewritten as

$$(2.7) \quad 0 = T[[w]_2] + T[[1w]_2] + T[[10w]_2] + T[[11w]_2] + T[[101w]_2],$$

$$0 = T[[w]_2] + T[[1w]_2] + T[[10w]_2] + T[[11w]_2] + T[[100w]_2]$$

$$(2.8) \quad + T[[110w]_2],$$

$$(2.9) \quad 0 = T[[100w]_2] + T[[111w]_2],$$

$$(2.10) \quad 0 = T[[1w]_2] + T[[10w]_2] + T[[11w]_2] + T[[1000w]_2],$$

$$(2.11) \quad 0 = T[[w]_2] + T[[10w]_2] + T[[11w]_2] + T[[100w]_2] + T[[1001w]_2],$$

for all binary word w of length $2k$ and $w \neq 0^{2k}$, and

$$(2.12) \quad 1 = T[[w]_2] + T[[1w]_2] + T[[10w]_2] + T[[11w]_2] + T[[101w]_2],$$

$$0 = T[[w]_2] + T[[1w]_2] + T[[10w]_2] + T[[11w]_2] + T[[100w]_2]$$

$$(2.13) \quad + T[[110w]_2],$$

$$(2.14) \quad 1 = T[[100w]_2] + T[[111w]_2],$$

$$(2.15) \quad 1 = T[[1w]_2] + T[[10w]_2] + T[[11w]_2] + T[[1000w]_2],$$

$$(2.16) \quad 0 = T[[w]_2] + T[[10w]_2] + T[[11w]_2] + T[[100w]_2] + T[[1001w]_2],$$

for $w = 0^{2k}$.

First we calculate an 2-automaton that generates T from its minimal polynomial and its first terms. This automaton has 389 states; its transition function and output function can be found in the annex. Let $A(s, w)$ denote the state reached after reading w from right to left starting from the state s , and τ the output function. Define

$$E_{2k} = \{A(i, w) : |w| = 2k, w \neq 0^{2k}\}.$$

Identities (2.7) through (2.16) can be written as

$$(2.17) \quad 0 = \tau(A(s, \epsilon)) + \tau(A(s, 1)) + \tau(A(s, 10)) + \tau(A(s, 11)) + \tau(A(s, 101)),$$

$$0 = \tau(A(s, \epsilon)) + \tau(A(s, 1)) + \tau(A(s, 10)) + \tau(A(s, 11)) + \tau(A(s, 100))$$

$$(2.18) \quad + \tau(A(s, 110)),$$

$$(2.19) \quad 0 = \tau(A(s, 100)) + \tau(A(s, 111)),$$

$$(2.20) \quad 0 = \tau(A(s, 1)) + \tau(A(s, 10)) + \tau(A(s, 11)) + \tau(A(s, 1000)),$$

$$0 = \tau(A(s, \epsilon)) + \tau(A(s, 10)) + \tau(A(s, 11)) + \tau(A(s, 100))$$

$$(2.21) \quad + \tau(A(s, 1001)),$$

for all $s \in E_{2k}$, and

$$(2.22) \quad 1 = \tau(A(s, \epsilon)) + \tau(A(s, 1)) + \tau(A(s, 10)) + \tau(A(s, 11)) + \tau(A(s, 101)),$$

$$0 = \tau(A(s, \epsilon)) + \tau(A(s, 1)) + \tau(A(s, 10)) + \tau(A(s, 11)) + \tau(A(s, 100))$$

$$(2.23) \quad + \tau(A(s, 110)),$$

$$(2.24) \quad 1 = \tau(A(s, 100)) + \tau(A(s, 111)),$$

$$(2.25) \quad 1 = \tau(A(s, 1)) + \tau(A(s, 10)) + \tau(A(s, 11)) + \tau(A(s, 1000)),$$

$$0 = \tau(A(s, \epsilon)) + \tau(A(s, 10)) + \tau(A(s, 11)) + \tau(A(s, 100))$$

$$(2.26) \quad + \tau(A(s, 1001)),$$

for $s = A(i, 0^{2k})$. We find that $(A(i, 0^{20}), E_{20}) = (A(i, 0^{18}), E_{18})$, so that we only have to verify that identities (2.17) through (2.26) hold for $2 \leq k \leq 10$, which turns out to be true. $\square$

In the following lemma, we express M_{2k} , M_{2k+1} , W_{2k} , and W_{2k+1} in terms of M^e , M^o , W^e , and W^o .

Lemma 2.3. *For all integer $k \geq 2$, and $u = 2^{2k-1}$,*

$$M_{2k} = M^e[:5u] + x^{2u}M^e[:3u] + x^{3u}M^e[:2u] + x^{4u}M^e[:u] + x^{5u}R^e,$$

$$W_{2k} = W^e[:5u] + x^{2u}W^e[:3u] + x^{3u}W^e[:2u] + x^{4u}W^e[:u] + x^{5u}R^e.$$

For all integer $k \geq 2$, and $u = 2^{2k}$,

$$\begin{aligned} M_{2k+1} &= M^o[:5u] + x^{2u}M^o[:3u] + x^{3u}M^o[:2u] + x^{4u}M^o[:u] + x^{5u}R^o, \\ W_{2k+1} &= W^o[:5u] + x^{2u}W^o[:3u] + x^{3u}W^o[:2u] + x^{4u}W^o[:u] + x^{5u}R^o. \end{aligned}$$

Proof. We call the four identities in Lemma 2.3 also by the name M_{2k} , W_{2k} , M_{2k+1} , and W_{2k+1} . For $n = 2$, the identities can be verified directly. For $n \geq 2$, we claim that

$$\begin{aligned} M_{2k} \wedge W_{2k} &\Rightarrow M_{2k+1} \wedge W_{2k+1}, \\ M_{2k+1} \wedge W_{2k+1} &\Rightarrow M_{2k+2} \wedge W_{2k+2}. \end{aligned}$$

We give the proof of

$$M_{2k} \wedge W_{2k} \Rightarrow M_{2k+1},$$

the other ones being similar. Set $u = 2^{2k}$ and $v = 2^{2k-1}$. By definition and induction hypothesis, the left side of identity M_{2k+1} is equal to

$$(2.27) \quad \begin{aligned} W_{2k}M_{2k} &= (W^e[:5v] + x^{2v}W^e[:3v] + x^{3v}W^e[:2v] + x^{4v}W^e[:v] + x^{5v}R^e) \times (\\ &M^e[:5v] + x^{2v}M^e[:3v] + x^{3v}M^e[:2v] + x^{4v}M^e[:v] + x^{5v}R^e); \end{aligned}$$

Call this expression *lhs*. Note that both sides of identity M_{2k+1} have the same term of highest degree $x^{10v}R^o$. Therefore we only need to prove that their difference is $O(x^{10v})$. Using Lemma 2.1 it can be seen that the right side of identity M_{2k+1} is congruent, modulo x^{10v} , to

$$\begin{aligned} &W^e[:10v]M^e[:10v] + x^{4v}W^e[:6v]M^e[:6v] + x^{6v}W^e[:4v]M^e[:4v] \\ &+ x^{8v}W^e[:2v]M^e[:2v]. \end{aligned}$$

For all $n \leq 10$, replace the occurrences of $W^e[:n \cdot v]$ and $M^e[:n \cdot v]$ in the above expression by the reduction modulo $x^{n \cdot v}$ of the right side of the corresponding identity in Lemma 2.2 and get a new expression, which we call *rhs*. Define

$$\begin{aligned} X &:= x^v, \\ a_n &:= W^e[n \cdot v : (n+1) \cdot v] / X^n, \\ b_n &:= M^e[n \cdot v : (n+1) \cdot v] / X^n, \\ c &:= R^e. \end{aligned}$$

Using the notation introduced above, we can represent the expressions *lhs* (2.27) and *rhs* as polynomials in $\mathbb{F}_2[a_1, \dots, a_{10}, b_1, \dots, b_{10}, c][X]$. Note that it is not a problem that a_j commutes with b_k while W^e does not commute with M^e , because in the expressions concerned, the products of W^e -terms and M^e -terms are always in the same order. We let the computer do the simplification and check that the difference between these two polynomials is indeed $O(X^{10})$, which completes the proof. $\square$

Proof of Theorem 1.1. We prove the theorem for $\text{CF}(a, b)$; for $\text{CF}(b, a)$, the proof is similar. By Lemma 2.3, we have For all $0 \leq j, k \leq 1$,

$$\begin{aligned} \lim_{n \rightarrow \infty} M_{2n,j,k} &= M_{j,k}^e, \\ \lim_{n \rightarrow \infty} M_{2n+1,j,k} &= M_{j,k}^o, \\ \lim_{n \rightarrow \infty} W_{2n,j,k} &= W_{j,k}^e, \\ \lim_{n \rightarrow \infty} W_{2n+1,j,k} &= W_{j,k}^o. \end{aligned}$$

Let $z = 1/x$. By the convergence theorem and identity (2.2),

$$\text{CF}(a, b) = \frac{M_{0,1}^e(x)}{M_{0,0}^e(x)}.$$

By definition, $\phi(M^e, 0, 1)$ and $\phi(M^e, 0, 0)$ are minimal polynomials of $M_{0,1}^e$ and $M_{0,0}^e$. Therefore

$$P(x, y) = \text{Res}_t (\phi(M^e, 0, 1)(x, t), y^{12}\phi(M^e, 0, 1)(x, t/y))$$

is an annihilating polynomial of $f(x) = M_{0,1}^e/M_{0,0}^e$.

Define

$$Q(x, y) = q_4(x)y^4 + q_3(x)y^3 + q_2(x)y^2 + q_1(x)y + q_0(x),$$

where

$$\begin{aligned} q_0(x) &= x^{20} + x^{18} + x^{17} + x^{16} + x^{15} + x^{13} + x^{12} + x^{11} + x^7 + x^5, \\ q_1(x) &= x^{17} + x^{15} + x^{14} + x^{13} + x^{12} + x^{11} + x^{10} + x^8 + x^7 + x^5 + x^4 + x^2, \\ q_2(x) &= x^{20} + x^{14} + x^{12} + x^{10} + x^4 + 1, \\ q_3(x) &= x^{17} + x^{15} + x^{14} + x^{13} + x^{12} + x^{11} + x^{10} + x^8 + x^7 + x^5 + x^4 + x^2, \\ q_4(x) &= x^{20} + x^{18} + x^{17} + x^{16} + x^{15} + x^{14} + x^{13} + x^{12} + x^{11} + x^{10} + x^7 + x^5 \\ &\quad + x^4. \end{aligned}$$

The polynomial $Q(x, y)$ is the candidate for the minimal polynomial of $f(x)$ found by Padé-Hermite approximation. To prove that it is indeed the minimal polynomial of $f(x)$, we only need to prove that it is an irreducible factor of $P(x, y)$ of multiplicity m and $R(x, y) := P(x, y)/Q(x, y)^m$ is not an annihilating polynomial of $f(x)$. We verify the first point directly. For the second point, we find that when we truncate $f(x)$ to order 156, and substitute it for y in $R(z, y)$, we get a series with valuation smaller than 156, which proves that $R(z, y)$ is not an annihilating polynomial of $f(x)$. Finally, $z^{12}Q(1/z, y)$ is the minimal polynomial of $\text{CF}(a, b) = f(1/z)$. $\square$

3. ANNEXE

All 16 polynomials are of the form

$$p_0(x) + p_3(x)y^3 + p_6(x)y^6 + p_9(x)y^9 + p_{12}(x)y^{12}.$$

The coefficients $p_j(x)$ and the 16 initial terms to determine the solutions uniquely are given below.

For $\phi(M^e, 0, 0)$:

$$\begin{aligned} p_0(x) &= x^{108} + x^{106} + x^{100} + x^{98} + x^{92} + x^{90} + x^{86} + x^{84} + x^{82} + x^{78} + x^{74} \\ &\quad + x^{70} + x^{66} + x^{64} + x^{60} + x^{54} + x^{50} + x^{48} + x^{40} + x^{38} + x^{34} + x^{28} \\ &\quad + x^{24}, \\ p_3(x) &= x^{102} + x^{100} + x^{98} + x^{90} + x^{88} + x^{84} + x^{80} + x^{78} + x^{76} + x^{68} + x^{66} \\ &\quad + x^{62} + x^{60} + x^{56} + x^{52} + x^{50} + x^{44} + x^{42} + x^{40} + x^{36} + x^{34} + x^{32} \\ &\quad + x^{30} + x^{28} + x^{24} + x^{22} + x^{20} + x^{14}, \\ p_6(x) &= x^{104} + x^{100} + x^{94} + x^{92} + x^{80} + x^{70} + x^{66} + x^{64} + x^{58} + x^{56} + x^{54} \\ &\quad + x^{52} + x^{44} + x^{34} + x^{32} + x^{28} + x^{22} + x^{18} + x^{14} + x^{12} + x^{10} + x^6 \end{aligned}$$

$$\begin{aligned}
& + x^2 + 1, \\
p_9(x) &= x^{102} + x^{100} + x^{96} + x^{94} + x^{92} + x^{90} + x^{88} + x^{86} + x^{84} + x^{82} + x^{78} \\
& + x^{74} + x^{72} + x^{68} + x^{66} + x^{62} + x^{60} + x^{58} + x^{48} + x^{46} + x^{38} + x^{36} \\
& + x^{34} + x^{32} + x^{30} + x^{24} + x^{20} + x^{18} + x^{16} + x^{14} + x^{10} + x^8 + x^6 + x^4, \\
p_{12}(x) &= x^{108} + x^{106} + x^{104} + x^{100} + x^{98} + x^{96} + x^{92} + x^{90} + x^{88} + x^{84} + x^{82} \\
& + x^{80} + x^{76} + x^{74} + x^{72} + x^{68} + x^{66} + x^{64} + x^{60} + x^{58} + x^{56} + x^{52} \\
& + x^{50} + x^{48} + x^{28} + x^{26} + x^{24} + x^{20} + x^{18} + x^{16} + x^{12} + x^{10} + x^8 \\
& + x^4 + x^2 + 1,
\end{aligned}$$

and the initial terms are $[1, 0, 0, 0, 0, 0, 1, 0, 0, 0, 1, 0, 0, 0, 1, 0]$.

For $\phi(M^e, 0, 1)$:

$$\begin{aligned}
p_0(x) &= x^{105} + x^{103} + x^{102} + x^{101} + x^{100} + x^{99} + x^{98} + x^{95} + x^{91} + x^{90} + x^{89} \\
& + x^{87} + x^{85} + x^{84} + x^{83} + x^{79} + x^{78} + x^{75} + x^{74} + x^{72} + x^{71} + x^{69} \\
& + x^{66} + x^{65} + x^{64} + x^{63} + x^{62} + x^{61} + x^{58} + x^{56} + x^{52} + x^{50} + x^{46} \\
& + x^{45} + x^{43} + x^{40} + x^{39} + x^{38} + x^{35} + x^{33} + x^{31} + x^{30} + x^{29} + x^{28} \\
& + x^{27}, \\
p_3(x) &= x^{78} + x^{72} + x^{70} + x^{68} + x^{58} + x^{56} + x^{54} + x^{52} + x^{46} + x^{42} + x^{38} \\
& + x^{32} + x^{30} + x^{28} + x^{22} + x^{18}, \\
p_6(x) &= x^{87} + x^{85} + x^{84} + x^{83} + x^{82} + x^{80} + x^{67} + x^{65} + x^{64} + x^{63} + x^{62} \\
& + x^{60} + x^{59} + x^{57} + x^{56} + x^{54} + x^{53} + x^{50} + x^{49} + x^{46} + x^{45} + x^{42} \\
& + x^{41} + x^{39} + x^{38} + x^{36} + x^{31} + x^{29} + x^{28} + x^{26} + x^{25} + x^{23} + x^{22} \\
& + x^{20} + x^{19} + x^{17} + x^{16} + x^{15} + x^{14} + x^{12}, \\
p_9(x) &= x^{96} + x^{92} + x^{86} + x^{84} + x^{80} + x^{76} + x^{74} + x^{72} + x^{68} + x^{66} + x^{64} \\
& + x^{60} + x^{58} + x^{56} + x^{50} + x^{48} + x^{40} + x^{38} + x^{36} + x^{34} + x^{32} + x^{26} \\
& + x^{24} + x^{20} + x^{16} + x^{14} + x^{12} + x^6, \\
p_{12}(x) &= x^{105} + x^{103} + x^{102} + x^{100} + x^{97} + x^{95} + x^{94} + x^{93} + x^{92} + x^{91} + x^{90} \\
& + x^{88} + x^{85} + x^{83} + x^{82} + x^{80} + x^{73} + x^{71} + x^{70} + x^{68} + x^{65} + x^{63} \\
& + x^{62} + x^{61} + x^{60} + x^{59} + x^{58} + x^{56} + x^{53} + x^{51} + x^{50} + x^{48} + x^{25} \\
& + x^{23} + x^{22} + x^{20} + x^{17} + x^{15} + x^{14} + x^{13} + x^{12} + x^{11} + x^{10} + x^8 \\
& + x^5 + x^3 + x^2 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 1, 1, 0, 1, 1, 1, 1, 1, 1, 1, 0, 0, 1]$.

For $\phi(M^e, 1, 0)$:

$$\begin{aligned}
p_0(x) &= x^{105} + x^{103} + x^{102} + x^{101} + x^{100} + x^{99} + x^{98} + x^{95} + x^{91} + x^{90} + x^{89} \\
& + x^{87} + x^{85} + x^{84} + x^{83} + x^{79} + x^{78} + x^{75} + x^{74} + x^{72} + x^{71} + x^{69} \\
& + x^{66} + x^{65} + x^{64} + x^{63} + x^{62} + x^{61} + x^{58} + x^{56} + x^{52} + x^{50} + x^{46} \\
& + x^{45} + x^{43} + x^{40} + x^{39} + x^{38} + x^{35} + x^{33} + x^{31} + x^{30} + x^{29} + x^{28}
\end{aligned}$$

$$\begin{aligned}
& + x^{27}, \\
p_3(x) &= x^{78} + x^{72} + x^{70} + x^{68} + x^{58} + x^{56} + x^{54} + x^{52} + x^{46} + x^{42} + x^{38} \\
& + x^{32} + x^{30} + x^{28} + x^{22} + x^{18}, \\
p_6(x) &= x^{87} + x^{85} + x^{84} + x^{83} + x^{82} + x^{80} + x^{67} + x^{65} + x^{64} + x^{63} + x^{62} \\
& + x^{60} + x^{59} + x^{57} + x^{56} + x^{54} + x^{53} + x^{50} + x^{49} + x^{46} + x^{45} + x^{42} \\
& + x^{41} + x^{39} + x^{38} + x^{36} + x^{31} + x^{29} + x^{28} + x^{26} + x^{25} + x^{23} + x^{22} \\
& + x^{20} + x^{19} + x^{17} + x^{16} + x^{15} + x^{14} + x^{12}, \\
p_9(x) &= x^{96} + x^{92} + x^{86} + x^{84} + x^{80} + x^{76} + x^{74} + x^{72} + x^{68} + x^{66} + x^{64} \\
& + x^{60} + x^{58} + x^{56} + x^{50} + x^{48} + x^{40} + x^{38} + x^{36} + x^{34} + x^{32} + x^{26} \\
& + x^{24} + x^{20} + x^{16} + x^{14} + x^{12} + x^6, \\
p_{12}(x) &= x^{105} + x^{103} + x^{102} + x^{100} + x^{97} + x^{95} + x^{94} + x^{93} + x^{92} + x^{91} + x^{90} \\
& + x^{88} + x^{85} + x^{83} + x^{82} + x^{80} + x^{73} + x^{71} + x^{70} + x^{68} + x^{65} + x^{63} \\
& + x^{62} + x^{61} + x^{60} + x^{59} + x^{58} + x^{56} + x^{53} + x^{51} + x^{50} + x^{48} + x^{25} \\
& + x^{23} + x^{22} + x^{20} + x^{17} + x^{15} + x^{14} + x^{13} + x^{12} + x^{11} + x^{10} + x^8 \\
& + x^5 + x^3 + x^2 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 1, 1, 0, 1, 1, 1, 1, 1, 1, 1, 1, 0, 0, 1]$.

For $\phi(M^e, 1, 1)$:

$$\begin{aligned}
p_0(x) &= x^{102} + x^{100} + x^{96} + x^{94} + x^{86} + x^{78} + x^{76} + x^{68} + x^{66} + x^{62} + x^{58} \\
& + x^{54} + x^{52} + x^{50} + x^{42} + x^{40} + x^{38} + x^{32} + x^{30}, \\
p_3(x) &= x^{96} + x^{94} + x^{92} + x^{90} + x^{78} + x^{76} + x^{72} + x^{66} + x^{62} + x^{54} + x^{48} \\
& + x^{46} + x^{42} + x^{40} + x^{36} + x^{34} + x^{32} + x^{26} + x^{24} + x^{22} + x^{18} + x^{12}, \\
p_6(x) &= x^{98} + x^{94} + x^{92} + x^{88} + x^{84} + x^{82} + x^{78} + x^{76} + x^{74} + x^{68} + x^{66} \\
& + x^{64} + x^{62} + x^{60} + x^{58} + x^{54} + x^{48} + x^{46} + x^{36} + x^{22} + x^{20} + x^{18} \\
& + x^{16} + x^{12} + x^{10} + x^8 + x^2 + 1, \\
p_9(x) &= x^{96} + x^{94} + x^{88} + x^{84} + x^{82} + x^{78} + x^{76} + x^{74} + x^{70} + x^{68} + x^{64} \\
& + x^{62} + x^{60} + x^{58} + x^{54} + x^{50} + x^{42} + x^{40} + x^{38} + x^{36} + x^{34} + x^{28} \\
& + x^{26} + x^{22} + x^{14} + x^8 + x^6 + x^4, \\
p_{12}(x) &= x^{102} + x^{100} + x^{98} + x^{96} + x^{94} + x^{92} + x^{90} + x^{88} + x^{86} + x^{84} + x^{82} \\
& + x^{80} + x^{70} + x^{68} + x^{66} + x^{64} + x^{62} + x^{60} + x^{58} + x^{56} + x^{54} + x^{52} \\
& + x^{50} + x^{48} + x^{22} + x^{20} + x^{18} + x^{16} + x^{14} + x^{12} + x^{10} + x^8 + x^6 + x^4 \\
& + x^2 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 1, 0, 1, 0, 0, 0, 0, 0, 0, 0, 0, 0]$.

For $\phi(W^e, 0, 0)$:

$$\begin{aligned}
p_0(x) &= x^{102} + x^{100} + x^{96} + x^{94} + x^{90} + x^{88} + x^{84} + x^{82} + x^{80} + x^{78} + x^{76} \\
& + x^{72} + x^{66} + x^{64} + x^{62} + x^{60} + x^{58} + x^{52} + x^{50} + x^{44} + x^{42} + x^{34}
\end{aligned}$$

$$\begin{aligned}
& + x^{30} + x^{28} + x^{24}, \\
p_3(x) &= x^{96} + x^{94} + x^{92} + x^{90} + x^{84} + x^{82} + x^{72} + x^{68} + x^{58} + x^{52} + x^{48} \\
& + x^{46} + x^{42} + x^{40} + x^{36} + x^{34} + x^{32} + x^{26} + x^{24} + x^{22} + x^{20} + x^{14}, \\
p_6(x) &= x^{98} + x^{94} + x^{92} + x^{90} + x^{86} + x^{84} + x^{78} + x^{76} + x^{74} + x^{70} + x^{68} \\
& + x^{66} + x^{60} + x^{58} + x^{56} + x^{54} + x^{50} + x^{46} + x^{36} + x^{26} + x^{24} + x^{20} \\
& + x^{16} + x^{12} + x^2 + 1, \\
p_9(x) &= x^{96} + x^{94} + x^{88} + x^{84} + x^{82} + x^{78} + x^{76} + x^{74} + x^{70} + x^{68} + x^{64} \\
& + x^{62} + x^{60} + x^{58} + x^{54} + x^{50} + x^{42} + x^{40} + x^{38} + x^{36} + x^{34} + x^{28} \\
& + x^{26} + x^{22} + x^{14} + x^8 + x^6 + x^4, \\
p_{12}(x) &= x^{102} + x^{100} + x^{98} + x^{96} + x^{94} + x^{92} + x^{90} + x^{88} + x^{86} + x^{84} + x^{82} \\
& + x^{80} + x^{70} + x^{68} + x^{66} + x^{64} + x^{62} + x^{60} + x^{58} + x^{56} + x^{54} + x^{52} \\
& + x^{50} + x^{48} + x^{22} + x^{20} + x^{18} + x^{16} + x^{14} + x^{12} + x^{10} + x^8 + x^6 + x^4 \\
& + x^2 + 1,
\end{aligned}$$

and the initial terms are $[1, 0, 0, 0, 1, 0, 1, 0, 1, 0, 0, 0, 0, 0, 0, 0]$.

For $\phi(W^e, 0, 1)$:

$$\begin{aligned}
p_0(x) &= x^{105} + x^{103} + x^{102} + x^{101} + x^{100} + x^{99} + x^{98} + x^{95} + x^{91} + x^{90} + x^{89} \\
& + x^{82} + x^{80} + x^{78} + x^{77} + x^{76} + x^{75} + x^{73} + x^{72} + x^{70} + x^{69} + x^{68} \\
& + x^{66} + x^{64} + x^{63} + x^{60} + x^{57} + x^{53} + x^{51} + x^{50} + x^{45} + x^{43} + x^{42} \\
& + x^{41} + x^{39} + x^{37} + x^{36} + x^{35} + x^{33} + x^{31} + x^{29} + x^{28} + x^{27}, \\
p_3(x) &= x^{78} + x^{72} + x^{70} + x^{68} + x^{58} + x^{56} + x^{54} + x^{52} + x^{46} + x^{42} + x^{38} \\
& + x^{32} + x^{30} + x^{28} + x^{22} + x^{18}, \\
p_6(x) &= x^{87} + x^{85} + x^{84} + x^{83} + x^{82} + x^{80} + x^{67} + x^{65} + x^{64} + x^{63} + x^{62} \\
& + x^{60} + x^{59} + x^{57} + x^{56} + x^{54} + x^{53} + x^{50} + x^{49} + x^{46} + x^{45} + x^{42} \\
& + x^{41} + x^{39} + x^{38} + x^{36} + x^{31} + x^{29} + x^{28} + x^{26} + x^{25} + x^{23} + x^{22} \\
& + x^{20} + x^{19} + x^{17} + x^{16} + x^{15} + x^{14} + x^{12}, \\
p_9(x) &= x^{96} + x^{92} + x^{86} + x^{84} + x^{80} + x^{76} + x^{74} + x^{72} + x^{68} + x^{66} + x^{64} \\
& + x^{60} + x^{58} + x^{56} + x^{50} + x^{48} + x^{40} + x^{38} + x^{36} + x^{34} + x^{32} + x^{26} \\
& + x^{24} + x^{20} + x^{16} + x^{14} + x^{12} + x^6, \\
p_{12}(x) &= x^{105} + x^{103} + x^{102} + x^{100} + x^{97} + x^{95} + x^{94} + x^{93} + x^{92} + x^{91} + x^{90} \\
& + x^{88} + x^{85} + x^{83} + x^{82} + x^{80} + x^{73} + x^{71} + x^{70} + x^{68} + x^{65} + x^{63} \\
& + x^{62} + x^{61} + x^{60} + x^{59} + x^{58} + x^{56} + x^{53} + x^{51} + x^{50} + x^{48} + x^{25} \\
& + x^{23} + x^{22} + x^{20} + x^{17} + x^{15} + x^{14} + x^{13} + x^{12} + x^{11} + x^{10} + x^8 \\
& + x^5 + x^3 + x^2 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 1, 1, 0, 0, 0, 1, 0, 0, 1, 0, 1, 0, 0]$.

For $\phi(W^e, 1, 0)$:

$$\begin{aligned}
p_0(x) &= x^{105} + x^{103} + x^{102} + x^{101} + x^{100} + x^{99} + x^{98} + x^{95} + x^{91} + x^{90} + x^{89} \\
&\quad + x^{82} + x^{80} + x^{78} + x^{77} + x^{76} + x^{75} + x^{73} + x^{72} + x^{70} + x^{69} + x^{68} \\
&\quad + x^{66} + x^{64} + x^{63} + x^{60} + x^{57} + x^{53} + x^{51} + x^{50} + x^{45} + x^{43} + x^{42} \\
&\quad + x^{41} + x^{39} + x^{37} + x^{36} + x^{35} + x^{33} + x^{31} + x^{29} + x^{28} + x^{27}, \\
p_3(x) &= x^{78} + x^{72} + x^{70} + x^{68} + x^{58} + x^{56} + x^{54} + x^{52} + x^{46} + x^{42} + x^{38} \\
&\quad + x^{32} + x^{30} + x^{28} + x^{22} + x^{18}, \\
p_6(x) &= x^{87} + x^{85} + x^{84} + x^{83} + x^{82} + x^{80} + x^{67} + x^{65} + x^{64} + x^{63} + x^{62} \\
&\quad + x^{60} + x^{59} + x^{57} + x^{56} + x^{54} + x^{53} + x^{50} + x^{49} + x^{46} + x^{45} + x^{42} \\
&\quad + x^{41} + x^{39} + x^{38} + x^{36} + x^{31} + x^{29} + x^{28} + x^{26} + x^{25} + x^{23} + x^{22} \\
&\quad + x^{20} + x^{19} + x^{17} + x^{16} + x^{15} + x^{14} + x^{12}, \\
p_9(x) &= x^{96} + x^{92} + x^{86} + x^{84} + x^{80} + x^{76} + x^{74} + x^{72} + x^{68} + x^{66} + x^{64} \\
&\quad + x^{60} + x^{58} + x^{56} + x^{50} + x^{48} + x^{40} + x^{38} + x^{36} + x^{34} + x^{32} + x^{26} \\
&\quad + x^{24} + x^{20} + x^{16} + x^{14} + x^{12} + x^6, \\
p_{12}(x) &= x^{105} + x^{103} + x^{102} + x^{100} + x^{97} + x^{95} + x^{94} + x^{93} + x^{92} + x^{91} + x^{90} \\
&\quad + x^{88} + x^{85} + x^{83} + x^{82} + x^{80} + x^{73} + x^{71} + x^{70} + x^{68} + x^{65} + x^{63} \\
&\quad + x^{62} + x^{61} + x^{60} + x^{59} + x^{58} + x^{56} + x^{53} + x^{51} + x^{50} + x^{48} + x^{25} \\
&\quad + x^{23} + x^{22} + x^{20} + x^{17} + x^{15} + x^{14} + x^{13} + x^{12} + x^{11} + x^{10} + x^8 \\
&\quad + x^5 + x^3 + x^2 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 1, 1, 0, 0, 0, 1, 0, 0, 1, 0, 1, 0, 0]$.

For $\phi(W^e, 1, 1)$:

$$\begin{aligned}
p_0(x) &= x^{108} + x^{106} + x^{100} + x^{98} + x^{96} + x^{94} + x^{90} + x^{88} + x^{84} + x^{78} + x^{74} \\
&\quad + x^{70} + x^{64} + x^{62} + x^{58} + x^{50} + x^{48} + x^{44} + x^{40} + x^{32} + x^{30}, \\
p_3(x) &= x^{102} + x^{100} + x^{98} + x^{84} + x^{82} + x^{70} + x^{66} + x^{62} + x^{60} + x^{58} + x^{52} \\
&\quad + x^{50} + x^{46} + x^{36} + x^{26} + x^{12}, \\
p_6(x) &= x^{104} + x^{100} + x^{96} + x^{90} + x^{88} + x^{84} + x^{64} + x^{62} + x^{60} + x^{56} + x^{52} \\
&\quad + x^{50} + x^{36} + x^{34} + x^{32} + x^{30} + x^{26} + x^{22} + x^{18} + x^{12} + x^8 + x^6 + x^2 \\
&\quad + 1, \\
p_9(x) &= x^{102} + x^{100} + x^{96} + x^{94} + x^{92} + x^{90} + x^{88} + x^{86} + x^{84} + x^{82} + x^{78} \\
&\quad + x^{74} + x^{72} + x^{68} + x^{66} + x^{62} + x^{60} + x^{58} + x^{48} + x^{46} + x^{38} + x^{36} \\
&\quad + x^{34} + x^{32} + x^{30} + x^{24} + x^{20} + x^{18} + x^{16} + x^{14} + x^{10} + x^8 + x^6 + x^4, \\
p_{12}(x) &= x^{108} + x^{106} + x^{104} + x^{100} + x^{98} + x^{96} + x^{92} + x^{90} + x^{88} + x^{84} + x^{82} \\
&\quad + x^{80} + x^{76} + x^{74} + x^{72} + x^{68} + x^{66} + x^{64} + x^{60} + x^{58} + x^{56} + x^{52} \\
&\quad + x^{50} + x^{48} + x^{28} + x^{26} + x^{24} + x^{20} + x^{18} + x^{16} + x^{12} + x^{10} + x^8 \\
&\quad + x^4 + x^2 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 0, 1, 0, 1, 0, 1, 0, 0, 0, 1, 0]$.

For $\phi(M^o, 0, 0)$:

$$\begin{aligned}
p_0(x) &= x^{105} + x^{103} + x^{102} + x^{101} + x^{100} + x^{98} + x^{93} + x^{89} + x^{86} + x^{85} + x^{84} \\
&\quad + x^{83} + x^{79} + x^{76} + x^{71} + x^{70} + x^{67} + x^{65} + x^{63} + x^{61} + x^{60} + x^{59} \\
&\quad + x^{57} + x^{55} + x^{54} + x^{53} + x^{51} + x^{50} + x^{48} + x^{47} + x^{45} + x^{43} + x^{41} \\
&\quad + x^{40} + x^{39} + x^{38} + x^{37} + x^{36} + x^{31} + x^{29} + x^{28} + x^{27} + x^{24}, \\
p_3(x) &= x^{93} + x^{91} + x^{90} + x^{89} + x^{88} + x^{86} + x^{81} + x^{77} + x^{74} + x^{73} + x^{72} \\
&\quad + x^{71} + x^{67} + x^{64} + x^{61} + x^{56} + x^{55} + x^{54} + x^{53} + x^{50} + x^{46} + x^{45} \\
&\quad + x^{43} + x^{41} + x^{40} + x^{35} + x^{31} + x^{30} + x^{27} + x^{25} + x^{24} + x^{21} + x^{19} \\
&\quad + x^{16}, \\
p_6(x) &= x^{101} + x^{98} + x^{93} + x^{91} + x^{89} + x^{88} + x^{86} + x^{84} + x^{83} + x^{79} + x^{77} \\
&\quad + x^{75} + x^{73} + x^{70} + x^{69} + x^{68} + x^{67} + x^{64} + x^{61} + x^{59} + x^{58} + x^{55} \\
&\quad + x^{54} + x^{53} + x^{52} + x^{51} + x^{49} + x^{47} + x^{46} + x^{44} + x^{43} + x^{38} + x^{36} \\
&\quad + x^{34} + x^{32} + x^{29} + x^{27} + x^{23} + x^{19} + x^{16} + x^{15} + x^{14} + x^{12} + x^{10} \\
&\quad + x^5 + x^3 + x^2 + 1, \\
p_9(x) &= x^{93} + x^{91} + x^{90} + x^{88} + x^{61} + x^{59} + x^{58} + x^{56} + x^{13} + x^{11} + x^{10} + x^8, \\
p_{12}(x) &= x^{105} + x^{103} + x^{102} + x^{100} + x^{97} + x^{95} + x^{94} + x^{92} + x^{85} + x^{83} + x^{82} \\
&\quad + x^{80} + x^{73} + x^{71} + x^{70} + x^{68} + x^{65} + x^{63} + x^{62} + x^{60} + x^{53} + x^{51} \\
&\quad + x^{50} + x^{48} + x^{25} + x^{23} + x^{22} + x^{20} + x^{17} + x^{15} + x^{14} + x^{12} + x^5 \\
&\quad + x^3 + x^2 + 1,
\end{aligned}$$

and the initial terms are $[1, 0, 0, 0, 1, 1, 0, 1, 0, 0, 1, 1, 1, 0, 1, 0]$.

For $\phi(M^o, 0, 1)$:

$$\begin{aligned}
p_0(x) &= x^{108} + x^{99} + x^{98} + x^{97} + x^{95} + x^{94} + x^{92} + x^{91} + x^{90} + x^{86} + x^{83} \\
&\quad + x^{78} + x^{77} + x^{76} + x^{74} + x^{73} + x^{72} + x^{71} + x^{70} + x^{69} + x^{68} + x^{64} \\
&\quad + x^{63} + x^{62} + x^{61} + x^{59} + x^{58} + x^{57} + x^{56} + x^{55} + x^{53} + x^{52} + x^{50} \\
&\quad + x^{49} + x^{48} + x^{47} + x^{46} + x^{45} + x^{43} + x^{42} + x^{41} + x^{39} + x^{37} + x^{36} \\
&\quad + x^{34} + x^{33} + x^{32} + x^{30} + x^{28} + x^{27}, \\
p_3(x) &= x^{81} + x^{79} + x^{78} + x^{76} + x^{75} + x^{73} + x^{72} + x^{70} + x^{69} + x^{67} + x^{66} \\
&\quad + x^{65} + x^{64} + x^{62} + x^{61} + x^{59} + x^{58} + x^{57} + x^{56} + x^{54} + x^{53} + x^{51} \\
&\quad + x^{50} + x^{48} + x^{47} + x^{45} + x^{44} + x^{42} + x^{41} + x^{39} + x^{38} + x^{36} + x^{35} \\
&\quad + x^{33} + x^{32} + x^{30} + x^{29} + x^{27} + x^{26} + x^{24} + x^{23} + x^{21} + x^{20} + x^{18}, \\
p_6(x) &= x^{90} + x^{84} + x^{82} + x^{78} + x^{76} + x^{74} + x^{72} + x^{68} + x^{66} + x^{62} + x^{60} \\
&\quad + x^{58} + x^{56} + x^{52} + x^{50} + x^{44} + x^{38} + x^{32} + x^{18} + x^{12}, \\
p_9(x) &= x^{99} + x^{97} + x^{96} + x^{95} + x^{94} + x^{93} + x^{92} + x^{91} + x^{90} + x^{89} + x^{88} \\
&\quad + x^{86} + x^{67} + x^{65} + x^{64} + x^{63} + x^{62} + x^{61} + x^{60} + x^{59} + x^{58} + x^{57}
\end{aligned}$$

$$\begin{aligned}
& + x^{56} + x^{54} + x^{19} + x^{17} + x^{16} + x^{15} + x^{14} + x^{13} + x^{12} + x^{11} + x^{10} \\
& + x^9 + x^8 + x^6, \\
p_{12}(x) = & x^{108} + x^{104} + x^{96} + x^{92} + x^{88} + x^{84} + x^{80} + x^{76} + x^{72} + x^{64} + x^{60} \\
& + x^{56} + x^{52} + x^{48} + x^{28} + x^{24} + x^{16} + x^{12} + x^8 + x^4 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 1, 1, 0, 1, 0, 1, 1, 0, 0, 0, 0, 1, 0, 1]$.

For $\phi(M^o, 1, 0)$:

$$\begin{aligned}
p_0(x) = & x^{112} + x^{103} + x^{102} + x^{101} + x^{100} + x^{99} + x^{98} + x^{95} + x^{91} + x^{89} + x^{88} \\
& + x^{87} + x^{85} + x^{84} + x^{81} + x^{80} + x^{78} + x^{76} + x^{74} + x^{71} + x^{69} + x^{68} \\
& + x^{65} + x^{64} + x^{62} + x^{60} + x^{57} + x^{54} + x^{52} + x^{50} + x^{48} + x^{46} + x^{42} \\
& + x^{40} + x^{38} + x^{37} + x^{35} + x^{28} + x^{27}, \\
p_3(x) = & x^{85} + x^{83} + x^{82} + x^{80} + x^{79} + x^{77} + x^{76} + x^{74} + x^{53} + x^{51} + x^{50} \\
& + x^{48} + x^{47} + x^{44} + x^{43} + x^{40} + x^{39} + x^{37} + x^{36} + x^{34} + x^{29} + x^{27} \\
& + x^{26} + x^{24} + x^{23} + x^{21} + x^{20} + x^{18}, \\
p_6(x) = & x^{94} + x^{88} + x^{86} + x^{80} + x^{74} + x^{68} + x^{62} + x^{56} + x^{46} + x^{42} + x^{40} \\
& + x^{38} + x^{36} + x^{34} + x^{32} + x^{30} + x^{28} + x^{24} + x^{18} + x^{12}, \\
p_9(x) = & x^{103} + x^{101} + x^{100} + x^{99} + x^{98} + x^{97} + x^{96} + x^{95} + x^{94} + x^{93} + x^{92} \\
& + x^{91} + x^{90} + x^{89} + x^{88} + x^{86} + x^{71} + x^{69} + x^{68} + x^{67} + x^{66} + x^{65} \\
& + x^{64} + x^{63} + x^{62} + x^{61} + x^{60} + x^{59} + x^{58} + x^{57} + x^{56} + x^{54} + x^{23} \\
& + x^{21} + x^{20} + x^{19} + x^{18} + x^{17} + x^{16} + x^{15} + x^{14} + x^{13} + x^{12} + x^{11} \\
& + x^{10} + x^9 + x^8 + x^6, \\
p_{12}(x) = & x^{112} + x^{108} + x^{96} + x^{88} + x^{84} + x^{76} + x^{64} + x^{56} + x^{52} + x^{48} + x^{32} \\
& + x^{28} + x^{16} + x^8 + x^4 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 1, 1, 0, 0, 0, 0, 0, 0, 1, 0, 1, 0, 1]$.

For $\phi(M^o, 1, 1)$:

$$\begin{aligned}
p_0(x) = & x^{105} + x^{103} + x^{102} + x^{101} + x^{100} + x^{98} + x^{91} + x^{90} + x^{89} + x^{88} + x^{87} \\
& + x^{86} + x^{85} + x^{83} + x^{77} + x^{74} + x^{70} + x^{68} + x^{67} + x^{66} + x^{65} + x^{61} \\
& + x^{60} + x^{57} + x^{54} + x^{51} + x^{47} + x^{45} + x^{43} + x^{42} + x^{41} + x^{38} + x^{36} \\
& + x^{35} + x^{34} + x^{32} + x^{30}, \\
p_3(x) = & x^{93} + x^{91} + x^{90} + x^{89} + x^{88} + x^{86} + x^{79} + x^{78} + x^{77} + x^{76} + x^{75} \\
& + x^{74} + x^{73} + x^{71} + x^{69} + x^{67} + x^{66} + x^{64} + x^{58} + x^{56} + x^{55} + x^{54} \\
& + x^{49} + x^{47} + x^{44} + x^{42} + x^{39} + x^{38} + x^{36} + x^{32} + x^{31} + x^{30} + x^{29} \\
& + x^{27} + x^{26} + x^{24} + x^{23} + x^{22} + x^{20} + x^{18}, \\
p_6(x) = & x^{101} + x^{98} + x^{90} + x^{89} + x^{87} + x^{86} + x^{85} + x^{82} + x^{81} + x^{80} + x^{78} \\
& + x^{77} + x^{76} + x^{75} + x^{71} + x^{69} + x^{61} + x^{58} + x^{56} + x^{54} + x^{51} + x^{50} \\
& + x^{47} + x^{44} + x^{40} + x^{38} + x^{36} + x^{34} + x^{33} + x^{32} + x^{30} + x^{27} + x^{26}
\end{aligned}$$

$$\begin{aligned}
& + x^{25} + x^{22} + x^{21} + x^{20} + x^{19} + x^{18} + x^{17} + x^{16} + x^{13} + x^{11} + x^8 \\
& + x^5 + x^3 + x^2 + 1, \\
p_9(x) &= x^{93} + x^{91} + x^{90} + x^{88} + x^{61} + x^{59} + x^{58} + x^{56} + x^{13} + x^{11} + x^{10} + x^8, \\
p_{12}(x) &= x^{105} + x^{103} + x^{102} + x^{100} + x^{97} + x^{95} + x^{94} + x^{92} + x^{85} + x^{83} + x^{82} \\
& + x^{80} + x^{73} + x^{71} + x^{70} + x^{68} + x^{65} + x^{63} + x^{62} + x^{60} + x^{53} + x^{51} \\
& + x^{50} + x^{48} + x^{25} + x^{23} + x^{22} + x^{20} + x^{17} + x^{15} + x^{14} + x^{12} + x^5 \\
& + x^3 + x^2 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 1, 0, 1, 1, 0, 0, 1, 0, 0, 0, 0]$.

For $\phi(W^o, 0, 0)$:

$$\begin{aligned}
p_0(x) &= x^{105} + x^{103} + x^{102} + x^{101} + x^{100} + x^{98} + x^{93} + x^{89} + x^{86} + x^{85} + x^{84} \\
& + x^{83} + x^{79} + x^{76} + x^{71} + x^{70} + x^{67} + x^{65} + x^{63} + x^{61} + x^{60} + x^{59} \\
& + x^{57} + x^{55} + x^{54} + x^{53} + x^{51} + x^{50} + x^{48} + x^{47} + x^{45} + x^{43} + x^{41} \\
& + x^{40} + x^{39} + x^{38} + x^{37} + x^{36} + x^{31} + x^{29} + x^{28} + x^{27} + x^{24}, \\
p_3(x) &= x^{93} + x^{91} + x^{90} + x^{89} + x^{88} + x^{86} + x^{81} + x^{77} + x^{74} + x^{73} + x^{72} \\
& + x^{71} + x^{67} + x^{64} + x^{61} + x^{56} + x^{55} + x^{54} + x^{53} + x^{50} + x^{46} + x^{45} \\
& + x^{43} + x^{41} + x^{40} + x^{35} + x^{31} + x^{30} + x^{27} + x^{25} + x^{24} + x^{21} + x^{19} \\
& + x^{16}, \\
p_6(x) &= x^{101} + x^{98} + x^{93} + x^{91} + x^{89} + x^{88} + x^{86} + x^{84} + x^{83} + x^{79} + x^{77} \\
& + x^{75} + x^{73} + x^{70} + x^{69} + x^{68} + x^{67} + x^{64} + x^{61} + x^{59} + x^{58} + x^{55} \\
& + x^{54} + x^{53} + x^{52} + x^{51} + x^{49} + x^{47} + x^{46} + x^{44} + x^{43} + x^{38} + x^{36} \\
& + x^{34} + x^{32} + x^{29} + x^{27} + x^{23} + x^{19} + x^{16} + x^{15} + x^{14} + x^{12} + x^{10} \\
& + x^5 + x^3 + x^2 + 1, \\
p_9(x) &= x^{93} + x^{91} + x^{90} + x^{88} + x^{61} + x^{59} + x^{58} + x^{56} + x^{13} + x^{11} + x^{10} + x^8, \\
p_{12}(x) &= x^{105} + x^{103} + x^{102} + x^{100} + x^{97} + x^{95} + x^{94} + x^{92} + x^{85} + x^{83} + x^{82} \\
& + x^{80} + x^{73} + x^{71} + x^{70} + x^{68} + x^{65} + x^{63} + x^{62} + x^{60} + x^{53} + x^{51} \\
& + x^{50} + x^{48} + x^{25} + x^{23} + x^{22} + x^{20} + x^{17} + x^{15} + x^{14} + x^{12} + x^5 \\
& + x^3 + x^2 + 1,
\end{aligned}$$

and the initial terms are $[1, 0, 0, 0, 1, 1, 0, 1, 0, 0, 1, 1, 1, 0, 1, 0]$.

For $\phi(W^o, 0, 1)$:

$$\begin{aligned}
p_0(x) &= x^{112} + x^{103} + x^{102} + x^{101} + x^{100} + x^{99} + x^{98} + x^{95} + x^{91} + x^{89} + x^{88} \\
& + x^{87} + x^{85} + x^{84} + x^{81} + x^{80} + x^{78} + x^{76} + x^{74} + x^{71} + x^{69} + x^{68} \\
& + x^{65} + x^{64} + x^{62} + x^{60} + x^{57} + x^{54} + x^{52} + x^{50} + x^{48} + x^{46} + x^{42} \\
& + x^{40} + x^{38} + x^{37} + x^{35} + x^{28} + x^{27}, \\
p_3(x) &= x^{85} + x^{83} + x^{82} + x^{80} + x^{79} + x^{77} + x^{76} + x^{74} + x^{53} + x^{51} + x^{50} \\
& + x^{48} + x^{47} + x^{44} + x^{43} + x^{40} + x^{39} + x^{37} + x^{36} + x^{34} + x^{29} + x^{27}
\end{aligned}$$

$$\begin{aligned}
& + x^{26} + x^{24} + x^{23} + x^{21} + x^{20} + x^{18}, \\
p_6(x) &= x^{94} + x^{88} + x^{86} + x^{80} + x^{74} + x^{68} + x^{62} + x^{56} + x^{46} + x^{42} + x^{40} \\
& + x^{38} + x^{36} + x^{34} + x^{32} + x^{30} + x^{28} + x^{24} + x^{18} + x^{12}, \\
p_9(x) &= x^{103} + x^{101} + x^{100} + x^{99} + x^{98} + x^{97} + x^{96} + x^{95} + x^{94} + x^{93} + x^{92} \\
& + x^{91} + x^{90} + x^{89} + x^{88} + x^{86} + x^{71} + x^{69} + x^{68} + x^{67} + x^{66} + x^{65} \\
& + x^{64} + x^{63} + x^{62} + x^{61} + x^{60} + x^{59} + x^{58} + x^{57} + x^{56} + x^{54} + x^{23} \\
& + x^{21} + x^{20} + x^{19} + x^{18} + x^{17} + x^{16} + x^{15} + x^{14} + x^{13} + x^{12} + x^{11} \\
& + x^{10} + x^9 + x^8 + x^6, \\
p_{12}(x) &= x^{112} + x^{108} + x^{96} + x^{88} + x^{84} + x^{76} + x^{64} + x^{56} + x^{52} + x^{48} + x^{32} \\
& + x^{28} + x^{16} + x^8 + x^4 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 1, 1, 0, 0, 0, 0, 0, 0, 1, 0, 1, 0, 1]$.

For $\phi(W^o, 1, 0)$:

$$\begin{aligned}
p_0(x) &= x^{108} + x^{99} + x^{98} + x^{97} + x^{95} + x^{94} + x^{92} + x^{91} + x^{90} + x^{86} + x^{83} \\
& + x^{78} + x^{77} + x^{76} + x^{74} + x^{73} + x^{72} + x^{71} + x^{70} + x^{69} + x^{68} + x^{64} \\
& + x^{63} + x^{62} + x^{61} + x^{59} + x^{58} + x^{57} + x^{56} + x^{55} + x^{53} + x^{52} + x^{50} \\
& + x^{49} + x^{48} + x^{47} + x^{46} + x^{45} + x^{43} + x^{42} + x^{41} + x^{39} + x^{37} + x^{36} \\
& + x^{34} + x^{33} + x^{32} + x^{30} + x^{28} + x^{27}, \\
p_3(x) &= x^{81} + x^{79} + x^{78} + x^{76} + x^{75} + x^{73} + x^{72} + x^{70} + x^{69} + x^{67} + x^{66} \\
& + x^{65} + x^{64} + x^{62} + x^{61} + x^{59} + x^{58} + x^{57} + x^{56} + x^{54} + x^{53} + x^{51} \\
& + x^{50} + x^{48} + x^{47} + x^{45} + x^{44} + x^{42} + x^{41} + x^{39} + x^{38} + x^{36} + x^{35} \\
& + x^{33} + x^{32} + x^{30} + x^{29} + x^{27} + x^{26} + x^{24} + x^{23} + x^{21} + x^{20} + x^{18}, \\
p_6(x) &= x^{90} + x^{84} + x^{82} + x^{78} + x^{76} + x^{74} + x^{72} + x^{68} + x^{66} + x^{62} + x^{60} \\
& + x^{58} + x^{56} + x^{52} + x^{50} + x^{44} + x^{38} + x^{32} + x^{18} + x^{12}, \\
p_9(x) &= x^{99} + x^{97} + x^{96} + x^{95} + x^{94} + x^{93} + x^{92} + x^{91} + x^{90} + x^{89} + x^{88} \\
& + x^{86} + x^{67} + x^{65} + x^{64} + x^{63} + x^{62} + x^{61} + x^{60} + x^{59} + x^{58} + x^{57} \\
& + x^{56} + x^{54} + x^{19} + x^{17} + x^{16} + x^{15} + x^{14} + x^{13} + x^{12} + x^{11} + x^{10} \\
& + x^9 + x^8 + x^6, \\
p_{12}(x) &= x^{108} + x^{104} + x^{96} + x^{92} + x^{88} + x^{84} + x^{80} + x^{76} + x^{72} + x^{64} + x^{60} \\
& + x^{56} + x^{52} + x^{48} + x^{28} + x^{24} + x^{16} + x^{12} + x^8 + x^4 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 1, 1, 0, 1, 0, 1, 1, 0, 0, 0, 0, 1, 0, 1]$.

For $\phi(W^o, 1, 1)$:

$$\begin{aligned}
p_0(x) &= x^{105} + x^{103} + x^{102} + x^{101} + x^{100} + x^{98} + x^{91} + x^{90} + x^{89} + x^{88} + x^{87} \\
& + x^{86} + x^{85} + x^{83} + x^{77} + x^{74} + x^{70} + x^{68} + x^{67} + x^{66} + x^{65} + x^{61} \\
& + x^{60} + x^{57} + x^{54} + x^{51} + x^{47} + x^{45} + x^{43} + x^{42} + x^{41} + x^{38} + x^{36} \\
& + x^{35} + x^{34} + x^{32} + x^{30},
\end{aligned}$$

$$\begin{aligned}
p_3(x) &= x^{93} + x^{91} + x^{90} + x^{89} + x^{88} + x^{86} + x^{79} + x^{78} + x^{77} + x^{76} + x^{75} \\
&\quad + x^{74} + x^{73} + x^{71} + x^{69} + x^{67} + x^{66} + x^{64} + x^{58} + x^{56} + x^{55} + x^{54} \\
&\quad + x^{49} + x^{47} + x^{44} + x^{42} + x^{39} + x^{38} + x^{36} + x^{32} + x^{31} + x^{30} + x^{29} \\
&\quad + x^{27} + x^{26} + x^{24} + x^{23} + x^{22} + x^{20} + x^{18}, \\
p_6(x) &= x^{101} + x^{98} + x^{90} + x^{89} + x^{87} + x^{86} + x^{85} + x^{82} + x^{81} + x^{80} + x^{78} \\
&\quad + x^{77} + x^{76} + x^{75} + x^{71} + x^{69} + x^{61} + x^{58} + x^{56} + x^{54} + x^{51} + x^{50} \\
&\quad + x^{47} + x^{44} + x^{40} + x^{38} + x^{36} + x^{34} + x^{33} + x^{32} + x^{30} + x^{27} + x^{26} \\
&\quad + x^{25} + x^{22} + x^{21} + x^{20} + x^{19} + x^{18} + x^{17} + x^{16} + x^{13} + x^{11} + x^8 \\
&\quad + x^5 + x^3 + x^2 + 1, \\
p_9(x) &= x^{93} + x^{91} + x^{90} + x^{88} + x^{61} + x^{59} + x^{58} + x^{56} + x^{13} + x^{11} + x^{10} + x^8, \\
p_{12}(x) &= x^{105} + x^{103} + x^{102} + x^{100} + x^{97} + x^{95} + x^{94} + x^{92} + x^{85} + x^{83} + x^{82} \\
&\quad + x^{80} + x^{73} + x^{71} + x^{70} + x^{68} + x^{65} + x^{63} + x^{62} + x^{60} + x^{53} + x^{51} \\
&\quad + x^{50} + x^{48} + x^{25} + x^{23} + x^{22} + x^{20} + x^{17} + x^{15} + x^{14} + x^{12} + x^5 \\
&\quad + x^3 + x^2 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 1, 0, 1, 1, 0, 0, 1, 0, 0, 0, 0]$.

Below is the transition function and output function of an 2-automaton that generates $T = M_{0,0}^o$:

Transition function $(n, j) \mapsto \delta(n, j)$ ($\Lambda(n) := [\delta(n, 0), \delta(n, 1)]$):

n	$\Lambda(n)$	n	$\Lambda(n)$	n	$\Lambda(n)$	n	$\Lambda(n)$	n	$\Lambda(n)$
0	[1, 2]	78	[129, 42]	156	[59, 71]	234	[45, 45]	312	[151, 288]
1	[3, 4]	79	[130, 129]	157	[235, 189]	235	[300, 136]	313	[31, 269]
2	[5, 6]	80	[98, 131]	158	[236, 237]	236	[211, 164]	314	[300, 145]
3	[7, 8]	81	[132, 53]	159	[238, 109]	237	[223, 158]	315	[342, 345]
4	[9, 10]	82	[133, 134]	160	[110, 239]	238	[97, 191]	316	[156, 346]
5	[11, 12]	83	[135, 136]	161	[240, 112]	239	[301, 193]	317	[140, 347]
6	[13, 14]	84	[137, 138]	162	[241, 242]	240	[302, 303]	318	[348, 126]
7	[15, 16]	85	[139, 137]	163	[243, 244]	241	[218, 198]	319	[247, 102]
8	[17, 18]	86	[140, 98]	164	[245, 122]	242	[304, 202]	320	[349, 350]
9	[19, 20]	87	[141, 142]	165	[246, 123]	243	[60, 305]	321	[269, 267]
10	[21, 22]	88	[13, 106]	166	[247, 248]	244	[306, 214]	322	[281, 266]
11	[23, 24]	89	[143, 144]	167	[154, 249]	245	[177, 307]	323	[351, 154]
12	[18, 25]	90	[145, 146]	168	[250, 251]	246	[210, 163]	324	[338, 352]
13	[26, 27]	91	[147, 13]	169	[113, 240]	247	[90, 177]	325	[295, 353]
14	[28, 29]	92	[30, 148]	170	[252, 253]	248	[226, 308]	326	[354, 47]
15	[7, 30]	93	[149, 150]	171	[123, 101]	249	[309, 77]	327	[52, 30]
16	[31, 32]	94	[151, 152]	172	[147, 254]	250	[162, 310]	328	[355, 253]
17	[33, 34]	95	[153, 112]	173	[236, 255]	251	[74, 163]	329	[349, 356]
18	[35, 36]	96	[154, 155]	174	[136, 146]	252	[97, 311]	330	[357, 340]
19	[35, 37]	97	[156, 157]	175	[256, 114]	253	[312, 181]	331	[243, 139]
20	[38, 39]	98	[158, 159]	176	[257, 258]	254	[191, 313]	332	[358, 273]

21	[40, 41]	99	[160, 161]	177	[114, 259]	255	[314, 194]	333	[272, 359]
22	[42, 43]	100	[162, 158]	178	[260, 9]	256	[77, 315]	334	[360, 345]
23	[44, 45]	101	[163, 164]	179	[125, 261]	257	[316, 6]	335	[290, 294]
24	[46, 47]	102	[165, 166]	180	[153, 262]	258	[310, 159]	336	[248, 245]
25	[48, 49]	103	[167, 92]	181	[262, 113]	259	[315, 76]	337	[347, 99]
26	[50, 8]	104	[168, 169]	182	[123, 263]	260	[78, 21]	338	[18, 361]
27	[51, 52]	105	[165, 170]	183	[46, 264]	261	[317, 318]	339	[362, 235]
28	[53, 54]	106	[170, 171]	184	[43, 129]	262	[169, 221]	340	[71, 301]
29	[55, 36]	107	[172, 67]	185	[12, 155]	263	[319, 225]	341	[26, 363]
30	[56, 57]	108	[173, 174]	186	[121, 35]	264	[176, 91]	342	[224, 73]
31	[58, 59]	109	[175, 176]	187	[265, 246]	265	[222, 162]	343	[364, 212]
32	[60, 61]	110	[177, 178]	188	[9, 266]	266	[320, 79]	344	[220, 90]
33	[62, 63]	111	[179, 70]	189	[267, 157]	267	[80, 69]	345	[365, 223]
34	[64, 65]	112	[180, 95]	190	[120, 268]	268	[58, 70]	346	[366, 62]
35	[66, 67]	113	[181, 65]	191	[269, 156]	269	[321, 235]	347	[316, 364]
36	[68, 69]	114	[182, 87]	192	[121, 55]	270	[88, 322]	348	[182, 367]
37	[70, 71]	115	[183, 184]	193	[270, 248]	271	[216, 161]	349	[17, 219]
38	[19, 72]	116	[6, 185]	194	[8, 148]	272	[188, 323]	350	[368, 27]
39	[73, 74]	117	[186, 187]	195	[204, 271]	273	[198, 166]	351	[199, 369]
40	[28, 75]	118	[188, 189]	196	[272, 273]	274	[180, 312]	352	[370, 301]
41	[63, 61]	119	[190, 18]	197	[274, 252]	275	[225, 210]	353	[171, 318]
42	[23, 76]	120	[191, 96]	198	[275, 39]	276	[76, 234]	354	[59, 370]
43	[77, 23]	121	[192, 193]	199	[128, 130]	277	[324, 187]	355	[173, 209]
44	[78, 79]	122	[166, 171]	200	[44, 44]	278	[167, 325]	356	[311, 313]
45	[79, 22]	123	[4, 194]	201	[130, 276]	279	[91, 307]	357	[201, 371]
46	[80, 81]	124	[195, 196]	202	[276, 128]	280	[79, 326]	358	[213, 372]
47	[82, 83]	125	[71, 192]	203	[250, 277]	281	[225, 74]	359	[373, 308]
48	[84, 85]	126	[197, 198]	204	[204, 204]	282	[158, 327]	360	[211, 374]
49	[86, 87]	127	[95, 64]	205	[278, 111]	283	[192, 328]	361	[268, 55]
50	[88, 83]	128	[199, 184]	206	[30, 279]	284	[329, 23]	362	[344, 375]
51	[85, 89]	129	[200, 77]	207	[254, 14]	285	[321, 188]	363	[353, 242]
52	[90, 91]	130	[201, 202]	208	[280, 281]	286	[195, 169]	364	[133, 297]
53	[92, 93]	131	[203, 204]	209	[282, 258]	287	[66, 330]	365	[257, 376]
54	[94, 95]	132	[181, 205]	210	[102, 245]	288	[233, 215]	366	[292, 50]
55	[96, 97]	133	[206, 207]	211	[53, 144]	289	[331, 29]	367	[244, 137]
56	[98, 99]	134	[208, 209]	212	[283, 269]	290	[302, 332]	368	[108, 377]
57	[100, 101]	135	[210, 211]	213	[241, 284]	291	[313, 97]	369	[378, 41]
58	[102, 103]	136	[6, 212]	214	[116, 246]	292	[306, 56]	370	[274, 355]
59	[33, 104]	137	[213, 62]	215	[285, 244]	293	[333, 303]	371	[379, 356]
60	[50, 30]	138	[214, 57]	216	[262, 286]	294	[206, 230]	372	[256, 340]
61	[105, 13]	139	[85, 215]	217	[287, 288]	295	[76, 334]	373	[299, 380]
62	[106, 107]	140	[216, 217]	218	[103, 289]	296	[62, 60]	374	[381, 355]
63	[108, 109]	141	[218, 165]	219	[290, 33]	297	[231, 311]	375	[372, 63]
64	[110, 111]	142	[219, 25]	220	[106, 291]	298	[335, 20]	376	[5, 364]

65	[112, 113]	143	[220, 176]	221	[271, 204]	299	[171, 336]	377	[325, 93]
66	[114, 115]	144	[196, 221]	222	[145, 236]	300	[186, 337]	378	[170, 317]
67	[14, 107]	145	[222, 223]	223	[292, 52]	301	[338, 49]	379	[56, 382]
68	[116, 117]	146	[224, 225]	224	[252, 293]	302	[113, 153]	380	[383, 233]
69	[118, 119]	147	[226, 227]	225	[137, 119]	303	[286, 240]	381	[169, 384]
70	[120, 121]	148	[16, 92]	226	[139, 118]	304	[40, 339]	382	[385, 347]
71	[103, 122]	149	[92, 228]	227	[294, 104]	305	[340, 259]	383	[124, 143]
72	[117, 123]	150	[229, 230]	228	[32, 156]	306	[246, 100]	384	[386, 277]
73	[124, 53]	151	[96, 231]	229	[295, 241]	307	[146, 237]	385	[204, 203]
74	[125, 126]	152	[232, 214]	230	[296, 297]	308	[341, 121]	386	[4, 387]
75	[127, 33]	153	[95, 181]	231	[11, 154]	309	[342, 343]	387	[388, 353]
76	[128, 43]	154	[84, 233]	232	[275, 298]	310	[280, 9]	388	[200, 329]
77	[129, 128]	155	[24, 234]	233	[299, 142]	311	[344, 150]		

Output function $n \mapsto \tau(n)$:

n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$
0	0	56	0	112	0	168	1	224	1	280	1	336	1
1	0	57	1	113	1	169	1	225	1	281	1	337	1
2	0	58	0	114	0	170	1	226	1	282	0	338	0
3	0	59	1	115	0	171	0	227	0	283	1	339	1
4	0	60	1	116	1	172	1	228	1	284	1	340	1
5	0	61	0	117	1	173	0	229	1	285	0	341	1
6	1	62	1	118	0	174	1	230	1	286	0	342	1
7	0	63	0	119	0	175	0	231	0	287	0	343	0
8	1	64	0	120	0	176	1	232	1	288	0	344	1
9	0	65	0	121	1	177	0	233	0	289	1	345	1
10	0	66	0	122	0	178	0	234	1	290	1	346	0
11	0	67	0	123	0	179	1	235	1	291	0	347	1
12	0	68	1	124	0	180	0	236	0	292	0	348	0
13	1	69	0	125	1	181	1	237	0	293	0	349	1
14	0	70	0	126	0	182	0	238	1	294	0	350	0
15	0	71	1	127	0	183	0	239	0	295	1	351	1
16	0	72	1	128	1	184	0	240	1	296	1	352	0
17	1	73	0	129	0	185	0	241	1	297	0	353	0
18	0	74	1	130	1	186	1	242	0	298	1	354	1
19	0	75	0	131	1	187	0	243	1	299	0	355	0
20	0	76	1	132	1	188	0	244	0	300	1	356	1
21	0	77	0	133	0	189	0	245	0	301	0	357	1
22	0	78	0	134	1	190	0	246	0	302	1	358	1
23	0	79	1	135	0	191	0	247	0	303	0	359	0
24	0	80	0	136	1	192	1	248	1	304	0	360	0
25	1	81	1	137	1	193	1	249	1	305	1	361	0
26	1	82	0	138	1	194	1	250	1	306	0	362	1
27	1	83	0	139	1	195	0	251	1	307	1	363	0

28	0	84	1	140	1	196	0	252	1	308	1	364	0
29	1	85	1	141	1	197	0	253	1	309	1	365	1
30	0	86	1	142	1	198	1	254	0	310	1	366	0
31	0	87	1	143	1	199	1	255	1	311	1	367	0
32	1	88	1	144	0	200	0	256	0	312	1	368	0
33	1	89	1	145	0	201	1	257	1	313	0	369	1
34	0	90	0	146	1	202	1	258	1	314	1	370	0
35	0	91	1	147	1	203	1	259	1	315	1	371	0
36	1	92	0	148	0	204	0	260	0	316	1	372	0
37	0	93	0	149	0	205	1	261	1	317	1	373	0
38	0	94	1	150	1	206	0	262	1	318	0	374	1
39	0	95	0	151	1	207	0	263	0	319	0	375	0
40	0	96	1	152	1	208	1	264	1	320	1	376	0
41	0	97	1	153	0	209	0	265	0	321	0	377	1
42	0	98	0	154	1	210	0	266	1	322	1	378	1
43	0	99	0	155	0	211	0	267	0	323	1	379	0
44	0	100	1	156	1	212	1	268	0	324	0	380	0
45	1	101	1	157	1	213	1	269	0	325	1	381	1
46	0	102	0	158	0	214	1	270	1	326	1	382	0
47	0	103	1	159	1	215	0	271	1	327	0	383	0
48	1	104	1	160	0	216	1	272	0	328	0	384	0
49	1	105	0	161	1	217	0	273	1	329	1	385	0
50	1	106	1	162	1	218	1	274	0	330	1	386	0
51	1	107	1	163	1	219	1	275	1	331	1	387	0
52	0	108	0	164	0	220	1	276	1	332	1	388	0
53	0	109	0	165	0	221	1	277	0	333	0		
54	1	110	0	166	0	222	0	278	1	334	0		
55	1	111	1	167	1	223	0	279	1	335	1		

UNIVERSITÉ DE STRASBOURG, CNRS, IRMA UMR 7501, F-67000 STRASBOURG, FRANCE

Email address: `guoniu.han@unistra.fr`

SCHOOL OF MATHEMATICS AND STATISTICS, HUAZHONG UNIVERSITY OF SCIENCE AND TECHNOLOGY, WUHAN, PR CHINA

Email address: `huyining@hust.edu.cn`