

PROOF OF THE CONTINUED FRACTION CONJECTURE FOR

$$(a, b) = (z^5 + z^4 + z^3, z)$$

GUO-NIU HAN* AND YINING HU*

ABSTRACT. The continued fraction conjecture claims that the continued fraction defined by the (a, b) -Thue-Morse sequence is algebraic over $\mathbb{F}_2(x)$ for all pairs of distinct elements (a, b) in $\mathbb{F}_2[x] \setminus \mathbb{F}_2$. In this paper we prove the conjecture for $(a, b) = (z^5 + z^4 + z^3, z)$.

1. INTRODUCTION

Let $\mathbb{F}_2$ be the finite field of cardinality 2. Given a sequence of polynomials $\mathbf{a} = (a_0(z), a_1(z), a_2(z), \dots)$ of elements from $\mathbb{F}_2[z] \setminus \mathbb{F}_2$, we define the infinite continued fraction

$$(1.1) \quad \text{CF}(\mathbf{a}(z)) = \frac{1}{a_0(z) + \frac{1}{a_1(z) + \frac{1}{a_2(z) + \frac{1}{\ddots}}}}$$

as the limit of the finite continued fractions

$$(1.2) \quad \text{CF}_n(\mathbf{a}(z)) = \frac{1}{a_0(z) + \frac{1}{a_1(z) + \frac{1}{\ddots + \frac{1}{a_n(z)}}}} \in \mathbb{F}_2((1/z)).$$

Conversely, each power series in $\mathbb{F}_2[[1/z]]$ without constant term

$$(1.3) \quad c_{-1}z^{-1} + c_{-2}z^{-2} + c_{-3}z^{-3} + \dots$$

admits a continued fraction expansion of form (1.1).

Let a and b be two distinct elements from $\mathbb{F}_2[z] \setminus \mathbb{F}_2$. We consider the continued fraction (1.1) associated with the (a, b) -Thue-Morse sequence

$$\mathbf{t} = (t_0(z), t_1(z), t_2(z), \dots) = (a, b, b, a, b, a, a, b, \dots)$$

that we shall denote

$$(1.4) \quad \text{CF}(a, b) := \text{CF}(\mathbf{t}).$$

Date: 9/04/2020 09:39:27 .

2010 Mathematics Subject Classification. 11B85, 11J70, 11B50, 11Y65, 05A15.

Key words and phrases. automatic sequence, continued fraction, Thue-Morse sequence.

* This paper was automatically generated from a computer program developed by the authors, with 10.7 minutes CPU time used.

Theorem 1.1. *When $(a, b) = (z^5 + z^4 + z^3, z)$, the two power series $\text{CF}(a, b)$ and $\text{CF}(b, a)$ are algebraic over $\mathbb{F}_2(z)$, with minimal polynomials of the form*

$$p_4(z)y^4 + p_3(z)y^3 + p_2(z)y^2 + p_1(z)y + p_0(z) = 0$$

where, for $\text{CF}(a, b)$,

$$\begin{aligned} p_0(z) &= z^{14} + z^{13} + z^{12} + z^7 + z^3 + z^2 + 1, \\ p_1(z) &= z^{19} + z^{17} + z^{14} + z^{11} + z^9 + z^8 + z^4 + z^3, \\ p_2(z) &= z^{20} + z^{18} + z^{16} + z^{14} + z^{12} + z^8 + z^6, \\ p_3(z) &= z^{19} + z^{17} + z^{14} + z^{11} + z^9 + z^8 + z^4 + z^3, \\ p_4(z) &= z^{18} + z^{16} + z^{13} + z^7 + z^6 + z^3 + z^2, \end{aligned}$$

and for $\text{CF}(b, a)$,

$$\begin{aligned} p_0(z) &= z^{16} + z^{14} + z^{13} + z^8 + z^7 + z^3 + z^2, \\ p_1(z) &= z^{19} + z^{17} + z^{14} + z^{11} + z^9 + z^8 + z^4 + z^3, \\ p_2(z) &= z^{20} + z^{18} + z^{16} + z^{14} + z^{12} + z^8 + z^6, \\ p_3(z) &= z^{19} + z^{17} + z^{14} + z^{11} + z^9 + z^8 + z^4 + z^3, \\ p_4(z) &= z^{18} + z^{13} + z^{12} + z^8 + z^7 + z^6 + z^3 + z^2 + 1. \end{aligned}$$

2. PROOF

Define the sequences of polynomials $(P_n(z))_{n \geq 0}$ and $(Q_n(z))_{n \geq 0}$ by

$$\begin{pmatrix} P_n(z) & Q_n(z) \\ P_{n-1}(z) & Q_{n-1}(z) \end{pmatrix} := \begin{pmatrix} t_n(z) & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} t_{n-1}(z) & 1 \\ 1 & 0 \end{pmatrix} \cdots \begin{pmatrix} t_0(z) & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}.$$

By the basic properties of continued fractions, we have

$$(2.1) \quad \text{CF}_n(a, b) = \frac{P_n(z)}{Q_n(z)} \in \mathbb{F}_2((1/z)).$$

As formula (2.1) is not efficient for calculating $\text{CF}_n(a, b)$, a fast method is to be derived and described as follows. Define

$$\begin{aligned} M_n(x) &= x^{\deg(t_{2^n-1})} \begin{pmatrix} t_{2^n-1}(1/x) & 1 \\ 1 & 0 \end{pmatrix} \times x^{\deg(t_{2^n-2})} \begin{pmatrix} t_{2^n-2}(1/x) & 1 \\ 1 & 0 \end{pmatrix} \times \\ &\quad \cdots \times x^{\deg(t_0)} \begin{pmatrix} t_0(1/x) & 1 \\ 1 & 0 \end{pmatrix}, \\ W_n(x) &= x^{\deg(\bar{t}_{2^n-1})} \begin{pmatrix} \bar{t}_{2^n-1}(1/x) & 1 \\ 1 & 0 \end{pmatrix} \times x^{\deg(\bar{t}_{2^n-2})} \begin{pmatrix} \bar{t}_{2^n-2}(1/x) & 1 \\ 1 & 0 \end{pmatrix} \times \\ &\quad \cdots \times x^{\deg(\bar{t}_0)} \begin{pmatrix} \bar{t}_0(1/x) & 1 \\ 1 & 0 \end{pmatrix}, \end{aligned}$$

where $\bar{t}$ is the (b, a) -Thue-Morse sequence. By the properties of the Thue-Morse sequence, we have for $n \geq 0$

$$\begin{aligned} M_{n+1}(x) &= W_n(x) \cdot M_n(x), \\ W_{n+1}(x) &= M_n(x) \cdot W_n(x). \end{aligned}$$

Let $x := 1/z$. For each non-zero polynomial $P(z)$, define $\tilde{P}(x)$ to be $P(1/x)$. Then,

$$\text{CF}_n(a, b) = \frac{P_n(z)}{Q_n(z)} = \frac{\tilde{P}_n(x)}{\tilde{Q}_n(x)} \in \mathbb{F}_2((x)) = \mathbb{F}_2((1/z)).$$

Comparing the definition of $M_n(x)$ with the definition of $P_n(x)$ and $Q_n(x)$, we see that

$$\begin{aligned} M_n(x)_{0,1} &= x^{d_n} \tilde{P}_{2^n-1}(x), \\ M_n(x)_{0,0} &= x^{d_n} \tilde{Q}_{2^n-1}(x), \end{aligned}$$

for some positive integer d_n . Hence,

$$(2.2) \quad \text{CF}_{2^{2^n}-1}(a, b) = \frac{\tilde{P}_{2^{2^n}-1}(x)}{\tilde{Q}_{2^{2^n}-1}(x)} = \frac{M_{2^n}(x)_{0,1}}{M_{2^n}(x)_{0,0}}.$$

Therefore, by the convergence theorem of continued fraction, the algebraicity of $\text{CF}(a, b)$ will be established if it is shown that both $M_{2^n}(x)_{0,1}$ and $M_{2^n}(x)_{0,0}$ converge to algebraic series in $\mathbb{F}_2[[x]]$.

Actually, we will prove that for all $0 \leq i, j \leq 1$ the four sequences $(M_{2^n}(x)_{i,j})_n$, $(M_{2^{n+1}}(x)_{i,j})_n$, $(W_{2^n}(x)_{i,j})_n$, and $(W_{2^{n+1}}(x)_{i,j})_n$ converge to algebraic series in $\mathbb{F}_2[[x]]$. For this purpose, we define four 2×2 matrices M^e, M^o, W^e, W^o as follows: For each $T \in \{M^e, M^o, W^e, W^o\}$ and $0 \leq i, j \leq 1$, $T_{i,j}$ is defined to be the unique solution in $\mathbb{F}_2[[x]]$ of the polynomial $\phi(T, i, j)$ under certain initial conditions; the polynomials $\phi(T, i, j)$ and initial conditions are given in Section 3. We will prove that these four matrices, whose components are algebraic by definition, are the limits of $(M_{2^n}(x))_n$, $(M_{2^{n+1}}(x))_n$, $(W_{2^n}(x))_n$, and $(W_{2^{n+1}}(x))_n$.

Let us explain how the polynomials $\phi(T, i, j)$ and initial conditions are found, and why the solutions exist and are unique. For $0 \leq i, j \leq 1$, the coefficients of the polynomial $\phi(M^e, i, j)$ (resp. $\phi(M^o, i, j)$, $\phi(W^e, i, j)$, and $\phi(W^o, i, j)$) are the Padé-Hermite approximants of type

$$(150, 150, 150, 150, 150)$$

of the vector

$$(1, T^3, T^6, T^9, T^{12}),$$

where $T = M_{12,i,j}$ (resp. $M_{11,i,j}$, $W_{12,i,j}$, and $W_{11,i,j}$), found by the Derksen algorithm. We take the first 8 terms of $M_{12,i,j}$ (resp. $M_{11,i,j}$, $W_{12,i,j}$, and $W_{11,i,j}$) as initial conditions for $\phi(M^e, i, j)$ (resp. $\phi(M^o, i, j)$, $\phi(W^e, i, j)$, and $\phi(W^o, i, j)$). The following fact will be used to ensure that the solution exists and is unique: let $P(x, y) \in \mathbb{F}_2[x, y]$ and for each series $f(x) = \sum_{n=0}^{\infty} a_n x^n \in \mathbb{F}_2[[x]]$ denote the partial sum $\sum_{j=0}^{n-1} a_j x^j$ by $f_n(x)$ for $n \geq 0$. If for some $n \geq 0$ and $a_0, a_1, \dots, a_{n-1} \in \mathbb{F}_2$ $P(x, \sum_{j=0}^{n-1} a_j x^j) = O(x^n)$ and $Q(x, y) := P(x, \sum_{j=0}^{n-1} a_j x^j + x^n y)$ can be written as $x^m \sum_{j=0}^{\infty} q_j(x) y^j$ where $q_j(x)$ are polynomials for $j \geq 0$, $q_1(0) = 1$, and $q_j(0) = 0$ for $j > 1$, then there exists a unique solution $f(x) \in \mathbb{F}_2[[x]]$ of $P(x, f(x)) = 0$ that satisfies the initial condition $f_n(x) = \sum_{j=0}^{n-1} a_j x^j$.

We state two lemmas concerning the four matrices M^e, M^o, W^e, W^o . The first one is about relations between them; the second, about the structure of the each matrix.

Lemma 2.1. *We have*

$$(2.3) \quad M^e = W^o \cdot M^o,$$

$$(2.4) \quad M^o = W^e \cdot M^e,$$

$$(2.5) \quad W^e = M^o \cdot W^o,$$

$$(2.6) \quad W^o = M^e \cdot W^e.$$

Proof. We give the proof of the identity

$$M_{0,0}^e = W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o,$$

the proofs of the others being similar.

First, we compute the minimal polynomials of $W_{0,0}^o M_{0,0}^o$ and $W_{0,1}^o M_{1,0}^o$. We know that

$$P(x, y) = \text{Res}_z (\phi(W^o, 0, 0)(x, z), z^{12} \cdot \phi(M^o, 0, 0)(x, y/z))$$

is an annihilating polynomial of $W_{0,0}^o M_{0,0}^o$. We use Padé-Hermite approximation to find a candidate for the minimal polynomial of $W_{0,0}^o M_{0,0}^o$, that will be called $\phi_0(x, y)$. To prove that $\phi_0(x, y)$ is indeed the minimal polynomial, it suffices to prove that it is an irreducible factor of $P(x, y)$ of multiplicity m and that $Q(x, y) := P(x, y)/\phi_0(x, y)^m$ is not an annihilating polynomial of $W_{0,0}^o M_{0,0}^o$. We verify the first point directly. For the second point, we truncate $W_{0,0}^o M_{0,0}^o$ to order 540 and substitute it for y in $Q(x, y)$. We get a series of valuation less than 540, which proves that $Q(x, y)$ is not an annihilating polynomial of $W_{0,0}^o M_{0,0}^o$. We find the minimal polynomial $\phi_1(x, y)$ of $W_{0,1}^o M_{1,0}^o$ in a similar way.

Now we prove that $\phi(M^e, 0, 0)$ is the minimal polynomial of $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$. We know that

$$S(x, y) = \text{Res}_z (\phi_0(x, z), \phi_1(x, y + z))$$

is an annihilating of $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$. We verify that $\phi(M^e, 0, 0)$ is an irreducible factor of $S(x, y)$ of multiplicity μ , and that $Q(x, y) := S(x, y)/\phi(M^e, 0, 0)^\mu$ is not an annihilating polynomial of $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$. To see the last point, we truncate $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$ to order 660 and substitute it for y in $Q(x, y)$. We get a series of valuation less than 660, and therefore $Q(x, y)$ is not an annihilating polynomial of $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$.

Finally, the first 16 terms of $M_{0,0}^e$ and $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$ coincide. As these first terms determine a unique solution of $\phi(M^e, 0, 0)$, we know that the two series are one and the same. $\square$

Define

$$R^e = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \quad \text{and} \quad R^o = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}.$$

Lemma 2.2. *For all integers $k \geq 2$ and $u = 2^{2k-1}$, the following identities hold.*

$$\begin{aligned} M^e[:12u] &= M^e[:6u] + x^u M^e[:5u] + x^{3u} M^e[:3u] + x^{5u} M^e[:u] + x^u M^e[:6u] \\ &\quad + x^{2u} M^e[:5u] + x^{4u} M^e[:3u] + x^{6u} M^e[:u] + x^{2u} M^e[:6u] \\ &\quad + x^{3u} M^e[:5u] + x^{5u} M^e[:3u] + x^{7u} M^e[:u] + x^{4u} M^e[:6u] \\ &\quad + x^{5u} M^e[:5u] + x^{7u} M^e[:3u] + x^{9u} M^e[:u] + x^{5u} M^e[:6u] \\ &\quad + x^{6u} M^e[:5u] + x^{8u} M^e[:3u] + x^{10u} M^e[:u] + x^{7u} M^e[:5u] \\ &\quad + x^{8u} M^e[:4u] + x^{9u} M^e[:3u] + x^{10u} M^e[:2u] + x^{11u} M^e[:u] \end{aligned}$$

$$\begin{aligned}
& + (x^{6u} + x^{7u} + x^{8u} + x^{10u} + x^{11u})R^e, \\
W^e[:12u] &= W^e[:6u] + x^u W^e[:5u] + x^{3u} W^e[:3u] + x^{5u} W^e[:u] + x^u W^e[:6u] \\
& + x^{2u} W^e[:5u] + x^{4u} W^e[:3u] + x^{6u} W^e[:u] + x^{2u} W^e[:6u] \\
& + x^{3u} W^e[:5u] + x^{5u} W^e[:3u] + x^{7u} W^e[:u] + x^{4u} W^e[:6u] \\
& + x^{5u} W^e[:5u] + x^{7u} W^e[:3u] + x^{9u} W^e[:u] + x^{5u} W^e[:6u] \\
& + x^{6u} W^e[:5u] + x^{8u} W^e[:3u] + x^{10u} W^e[:u] + x^{7u} W^e[:5u] \\
& + x^{8u} W^e[:4u] + x^{9u} W^e[:3u] + x^{10u} W^e[:2u] + x^{11u} W^e[:u] \\
& + (x^{6u} + x^{7u} + x^{8u} + x^{10u} + x^{11u})R^e.
\end{aligned}$$

For all integers $k \geq 2$ and $u = 2^{2k}$, the following identities hold.

$$\begin{aligned}
M^o[:12u] &= M^o[:6u] + x^u M^o[:5u] + x^{3u} M^o[:3u] + x^{5u} M^o[:u] + x^u M^o[:6u] \\
& + x^{2u} M^o[:5u] + x^{4u} M^o[:3u] + x^{6u} M^o[:u] + x^{2u} M^o[:6u] \\
& + x^{3u} M^o[:5u] + x^{5u} M^o[:3u] + x^{7u} M^o[:u] + x^{4u} M^o[:6u] \\
& + x^{5u} M^o[:5u] + x^{7u} M^o[:3u] + x^{9u} M^o[:u] + x^{5u} M^o[:6u] \\
& + x^{6u} M^o[:5u] + x^{8u} M^o[:3u] + x^{10u} M^o[:u] + x^{7u} M^o[:5u] \\
& + x^{8u} M^o[:4u] + x^{9u} M^o[:3u] + x^{10u} M^o[:2u] + x^{11u} M^o[:u] \\
& + (x^{6u} + x^{7u} + x^{8u} + x^{10u} + x^{11u})R^o, \\
W^o[:12u] &= W^o[:6u] + x^u W^o[:5u] + x^{3u} W^o[:3u] + x^{5u} W^o[:u] + x^u W^o[:6u] \\
& + x^{2u} W^o[:5u] + x^{4u} W^o[:3u] + x^{6u} W^o[:u] + x^{2u} W^o[:6u] \\
& + x^{3u} W^o[:5u] + x^{5u} W^o[:3u] + x^{7u} W^o[:u] + x^{4u} W^o[:6u] \\
& + x^{5u} W^o[:5u] + x^{7u} W^o[:3u] + x^{9u} W^o[:u] + x^{5u} W^o[:6u] \\
& + x^{6u} W^o[:5u] + x^{8u} W^o[:3u] + x^{10u} W^o[:u] + x^{7u} W^o[:5u] \\
& + x^{8u} W^o[:4u] + x^{9u} W^o[:3u] + x^{10u} W^o[:2u] + x^{11u} W^o[:u] \\
& + (x^{6u} + x^{7u} + x^{8u} + x^{10u} + x^{11u})R^o.
\end{aligned}$$

Proof. To prove Lemma 2.2 we first construct an automaton for each sequence concerned, and then transform the conditions on infinitely many k 's into finitely many conditions on the states of the automaton. In the following, we will prove that for $T = M_{0,0}^o$, for all integer $k \geq 2$ and $u = 2^{2k}$,

$$\begin{aligned}
T[:12u] &= T[:6u] + x^u T[:5u] + x^{3u} T[:3u] + x^{5u} T[:u] + x^u T[:6u] + x^{2u} T[:5u] \\
& + x^{4u} T[:3u] + x^{6u} T[:u] + x^{2u} T[:6u] + x^{3u} T[:5u] + x^{5u} T[:3u] \\
& + x^{7u} T[:u] + x^{4u} T[:6u] + x^{5u} T[:5u] + x^{7u} T[:3u] + x^{9u} T[:u] \\
& + x^{5u} T[:6u] + x^{6u} T[:5u] + x^{8u} T[:3u] + x^{10u} T[:u] + x^{7u} T[:5u] \\
& + x^{8u} T[:4u] + x^{9u} T[:3u] + x^{10u} T[:2u] + x^{11u} T[:u].
\end{aligned}$$

The proof of the other 15 cases are similar. We break down the above identity into 6 parts:

$$\begin{aligned}
0 &= x^{5u} T[u:2u] + x^{3u} T[3u:4u] + x^u T[5u:6u] + T[6u:7u], \\
0 &= x^{7u} T[:u] + x^{6u} T[u:2u] + x^{5u} T[2u:3u] + x^{4u} T[3u:4u]
\end{aligned}$$

$$\begin{aligned}
& + x^{3u}T[4u:5u] + x^{2u}T[5u:6u] + T[7u:8u], \\
0 & = x^{6u}T[2u:3u] + x^{4u}T[4u:5u] + T[8u:9u], \\
0 & = x^{6u}T[3u:4u] + x^{4u}T[5u:6u] + T[9u:10u], \\
0 & = x^{9u}T[u:2u] + x^{7u}T[3u:4u] + x^{6u}T[4u:5u] + x^{5u}T[5u:6u] \\
& \quad + T[10u:11u], \\
0 & = x^{11u}T[:u] + x^{10u}T[u:2u] + x^{9u}T[2u:3u] + x^{8u}T[3u:4u] \\
& \quad + x^{7u}T[4u:5u] + T[11u:12u],
\end{aligned}$$

which can be rewritten as

$$\begin{aligned}
(2.7) \quad 0 & = T[[1w]_2] + T[[11w]_2] + T[[101w]_2] + T[[110w]_2], \\
0 & = T[[w]_2] + T[[1w]_2] + T[[10w]_2] + T[[11w]_2] + T[[100w]_2] \\
(2.8) \quad & + T[[101w]_2] + T[[111w]_2], \\
(2.9) \quad 0 & = T[[10w]_2] + T[[100w]_2] + T[[1000w]_2], \\
(2.10) \quad 0 & = T[[11w]_2] + T[[101w]_2] + T[[1001w]_2], \\
(2.11) \quad 0 & = T[[1w]_2] + T[[11w]_2] + T[[100w]_2] + T[[101w]_2] + T[[1010w]_2], \\
0 & = T[[w]_2] + T[[1w]_2] + T[[10w]_2] + T[[11w]_2] + T[[100w]_2] \\
(2.12) \quad & + T[[1011w]_2],
\end{aligned}$$

for all binary word w of length $2k$ and $w \neq 0^{2k}$, and

$$\begin{aligned}
(2.13) \quad 0 & = T[[1w]_2] + T[[11w]_2] + T[[101w]_2] + T[[110w]_2], \\
0 & = T[[w]_2] + T[[1w]_2] + T[[10w]_2] + T[[11w]_2] + T[[100w]_2] \\
(2.14) \quad & + T[[101w]_2] + T[[111w]_2], \\
(2.15) \quad 0 & = T[[10w]_2] + T[[100w]_2] + T[[1000w]_2], \\
(2.16) \quad 0 & = T[[11w]_2] + T[[101w]_2] + T[[1001w]_2], \\
(2.17) \quad 0 & = T[[1w]_2] + T[[11w]_2] + T[[100w]_2] + T[[101w]_2] + T[[1010w]_2], \\
0 & = T[[w]_2] + T[[1w]_2] + T[[10w]_2] + T[[11w]_2] + T[[100w]_2] \\
(2.18) \quad & + T[[1011w]_2],
\end{aligned}$$

for $w = 0^{2k}$.

First we calculate an 2-automaton that generates T from its minimal polynomial and its first terms. This automaton has 319 states; its transition function and output function can be found in the annex. Let $A(s, w)$ denote the state reached after reading w from right to left starting from the state s , and τ the output function. Define

$$E_{2k} = \{A(i, w) : |w| = 2k, w \neq 0^{2k}\}.$$

Identities (2.7) through (2.18) can be written as

$$\begin{aligned}
(2.19) \quad 0 & = \tau(A(s, 1)) + \tau(A(s, 11)) + \tau(A(s, 101)) + \tau(A(s, 110)), \\
0 & = \tau(A(s, \epsilon)) + \tau(A(s, 1)) + \tau(A(s, 10)) + \tau(A(s, 11)) + \tau(A(s, 100)) \\
(2.20) \quad & + \tau(A(s, 101)) + \tau(A(s, 111)), \\
(2.21) \quad 0 & = \tau(A(s, 10)) + \tau(A(s, 100)) + \tau(A(s, 1000)), \\
(2.22) \quad 0 & = \tau(A(s, 11)) + \tau(A(s, 101)) + \tau(A(s, 1001)),
\end{aligned}$$

$$(2.23) \quad \begin{aligned} 0 &= \tau(A(s, 1)) + \tau(A(s, 11)) + \tau(A(s, 100)) + \tau(A(s, 101)) \\ &\quad + \tau(A(s, 1010)), \end{aligned}$$

$$(2.24) \quad \begin{aligned} 0 &= \tau(A(s, \epsilon)) + \tau(A(s, 1)) + \tau(A(s, 10)) + \tau(A(s, 11)) + \tau(A(s, 100)) \\ &\quad + \tau(A(s, 1011)), \end{aligned}$$

for all $s \in E_{2k}$, and

$$(2.25) \quad 0 = \tau(A(s, 1)) + \tau(A(s, 11)) + \tau(A(s, 101)) + \tau(A(s, 110)),$$

$$(2.26) \quad \begin{aligned} 0 &= \tau(A(s, \epsilon)) + \tau(A(s, 1)) + \tau(A(s, 10)) + \tau(A(s, 11)) + \tau(A(s, 100)) \\ &\quad + \tau(A(s, 101)) + \tau(A(s, 111)), \end{aligned}$$

$$(2.27) \quad 0 = \tau(A(s, 10)) + \tau(A(s, 100)) + \tau(A(s, 1000)),$$

$$(2.28) \quad 0 = \tau(A(s, 11)) + \tau(A(s, 101)) + \tau(A(s, 1001)),$$

$$(2.29) \quad \begin{aligned} 0 &= \tau(A(s, 1)) + \tau(A(s, 11)) + \tau(A(s, 100)) + \tau(A(s, 101)) \\ &\quad + \tau(A(s, 1010)), \end{aligned}$$

$$(2.30) \quad \begin{aligned} 0 &= \tau(A(s, \epsilon)) + \tau(A(s, 1)) + \tau(A(s, 10)) + \tau(A(s, 11)) + \tau(A(s, 100)) \\ &\quad + \tau(A(s, 1011)), \end{aligned}$$

for $s = A(i, 0^{2k})$. We find that $(A(i, 0^{22}), E_{22}) = (A(i, 0^{18}), E_{18})$, so that we only have to verify that identities (2.19) through (2.30) hold for $2 \leq k \leq 11$, which turns out to be true. $\square$

In the following lemma, we express M_{2k} , M_{2k+1} , W_{2k} , and W_{2k+1} in terms of M^e , M^o , W^e , and W^o .

Lemma 2.3. *For all integer $k \geq 2$, and $u = 2^{2k-1}$,*

$$M_{2k} = M^e[:6u] + x^u M^e[:5u] + x^{3u} M^e[:3u] + x^{5u} M^e[:u] + x^{6u} R^e,$$

$$W_{2k} = W^e[:6u] + x^u W^e[:5u] + x^{3u} W^e[:3u] + x^{5u} W^e[:u] + x^{6u} R^e.$$

For all integer $k \geq 2$, and $u = 2^{2k}$,

$$M_{2k+1} = M^o[:6u] + x^u M^o[:5u] + x^{3u} M^o[:3u] + x^{5u} M^o[:u] + x^{6u} R^o,$$

$$W_{2k+1} = W^o[:6u] + x^u W^o[:5u] + x^{3u} W^o[:3u] + x^{5u} W^o[:u] + x^{6u} R^o.$$

Proof. We call the four identities in Lemma 2.3 also by the name M_{2k} , W_{2k} , M_{2k+1} , and W_{2k+1} . For $n = 2$, the identities can be verified directly. For $n \geq 2$, we claim that

$$\begin{aligned} M_{2k} \wedge W_{2k} &\Rightarrow M_{2k+1} \wedge W_{2k+1}, \\ M_{2k+1} \wedge W_{2k+1} &\Rightarrow M_{2k+2} \wedge W_{2k+2}. \end{aligned}$$

We give the proof of

$$M_{2k} \wedge W_{2k} \Rightarrow M_{2k+1},$$

the other ones being similar. Set $u = 2^{2k}$ and $v = 2^{2k-1}$. By definition and induction hypothesis, the left side of identity M_{2k+1} is equal to

$$(2.31) \quad \begin{aligned} W_{2k} M_{2k} &= (W^e[:6v] + x^v W^e[:5v] + x^{3v} W^e[:3v] + x^{5v} W^e[:v] + x^{6v} R^e) \times (\\ &\quad M^e[:6v] + x^v M^e[:5v] + x^{3v} M^e[:3v] + x^{5v} M^e[:v] + x^{6v} R^e); \end{aligned}$$

Call this expression lhs . Note that both sides of identity M_{2k+1} have the same term of highest degree $x^{12v} R^o$. Therefore we only need to prove that their difference is

$O(x^{12v})$. Using Lemma 2.1 it can be seen that the right side of identity M_{2k+1} is congruent, modulo x^{12v} , to

$$W^e[:12v]M^e[:12v] + x^{2v}W^e[:10v]M^e[:10v] + x^{6v}W^e[:6v]M^e[:6v] + x^{10v}W^e[:2v]M^e[:2v].$$

For all $n \leq 12$, replace the occurrences of $W^e[:n \cdot v]$ and $M^e[:n \cdot v]$ in the above expression by the reduction modulo $x^{n \cdot v}$ of the right side of the corresponding identity in Lemma 2.2 and get a new expression, which we call *rhs*. Define

$$\begin{aligned} X &:= x^v, \\ a_n &:= W^e[n \cdot v : (n+1) \cdot v] / X^n, \\ b_n &:= M^e[n \cdot v : (n+1) \cdot v] / X^n, \\ c &:= R^e. \end{aligned}$$

Using the notation introduced above, we can represent the expressions *lhs* (2.31) and *rhs* as polynomials in $\mathbb{F}_2[a_1, \dots, a_{12}, b_1, \dots, b_{12}, c][X]$. Note that it is not a problem that a_j commutes with b_k while W^e does not commute with M^e , because in the expressions concerned, the products of W^e -terms and M^e -terms are always in the same order. We let the computer do the simplification and check that the difference between these two polynomials is indeed $O(X^{12})$, which completes the proof. $\square$

Proof of Theorem 1.1. We prove the theorem for $\text{CF}(a, b)$; for $\text{CF}(b, a)$, the proof is similar. By Lemma 2.3, we have For all $0 \leq j, k \leq 1$,

$$\begin{aligned} \lim_{n \rightarrow \infty} M_{2n,j,k} &= M_{j,k}^e, \\ \lim_{n \rightarrow \infty} M_{2n+1,j,k} &= M_{j,k}^o, \\ \lim_{n \rightarrow \infty} W_{2n,j,k} &= W_{j,k}^e, \\ \lim_{n \rightarrow \infty} W_{2n+1,j,k} &= W_{j,k}^o. \end{aligned}$$

Let $z = 1/x$. By the convergence theorem and identity (2.2),

$$\text{CF}(a, b) = \frac{M_{0,1}^e(x)}{M_{0,0}^e(x)}.$$

By definition, $\phi(M^e, 0, 1)$ and $\phi(M^e, 0, 0)$ are minimal polynomials of $M_{0,1}^e$ and $M_{0,0}^e$. Therefore

$$P(x, y) = \text{Res}_t (\phi(M^e, 0, 1)(x, t), y^{12}\phi(M^e, 0, 1)(x, t/y))$$

is an annihilating polynomial of $f(x) = M_{0,1}^e/M_{0,0}^e$.

Define

$$Q(x, y) = q_4(x)y^4 + q_3(x)y^3 + q_2(x)y^2 + q_1(x)y + q_0(x),$$

where

$$\begin{aligned} q_0(x) &= x^{20} + x^{18} + x^{17} + x^{13} + x^8 + x^7 + x^6, \\ q_1(x) &= x^{17} + x^{16} + x^{12} + x^{11} + x^9 + x^6 + x^3 + x, \\ q_2(x) &= x^{14} + x^{12} + x^8 + x^6 + x^4 + x^2 + 1, \\ q_3(x) &= x^{17} + x^{16} + x^{12} + x^{11} + x^9 + x^6 + x^3 + x, \\ q_4(x) &= x^{18} + x^{17} + x^{14} + x^{13} + x^7 + x^4 + x^2. \end{aligned}$$

The polynomial $Q(x, y)$ is the candidate for the minimal polynomial of $f(x)$ found by Padé-Hermite approximation. To prove that it is indeed the minimal polynomial of $f(x)$, we only need to prove that it is an irreducible factor of $P(x, y)$ of multiplicity m and $R(x, y) := P(x, y)/Q(x, y)^m$ is not an annihilating polynomial of $f(x)$. We verify the first point directly. For the second point, we find that when we truncate $f(x)$ to order 186, and substitute it for y in $R(z, y)$, we get a series with valuation smaller than 186, which proves that $R(z, y)$ is not an annihilating polynomial of $f(x)$. Finally, $z^{12}Q(1/z, y)$ is the minimal polynomial of $\text{CF}(a, b) = f(1/z)$. $\square$

3. ANNEXE

All 16 polynomials are of the form

$$p_0(x) + p_3(x)y^3 + p_6(x)y^6 + p_9(x)y^9 + p_{12}(x)y^{12}.$$

The coefficients $p_j(x)$ and the 16 initial terms to determine the solutions uniquely are given below.

For $\phi(M^e, 0, 0)$:

$$\begin{aligned} p_0(x) &= x^{96} + x^{92} + x^{90} + x^{86} + x^{74} + x^{68} + x^{64} + x^{62} + x^{58} + x^{56} + x^{54} \\ &\quad + x^{50} + x^{48} + x^{46} + x^{42} + x^{38} + x^{34} + x^{32} + x^{30} + x^{28} + x^{26} + x^{24} \\ &\quad + x^{22} + x^{20} + x^{18} + x^{14} + x^{12}, \\ p_3(x) &= x^{86} + x^{80} + x^{74} + x^{72} + x^{70} + x^{68} + x^{64} + x^{62} + x^{58} + x^{52} + x^{48} \\ &\quad + x^{34} + x^{28} + x^{22} + x^{16} + x^8, \\ p_6(x) &= x^{86} + x^{70} + x^{68} + x^{56} + x^{46} + x^{44} + x^{36} + x^{24} + x^{16} + x^{12} + x^4 + 1, \\ p_9(x) &= x^{84} + x^{82} + x^{80} + x^{78} + x^{70} + x^{68} + x^{66} + x^{64} + x^{62} + x^{56} + x^{54} \\ &\quad + x^{52} + x^{48} + x^{46} + x^{44} + x^{38} + x^{36} + x^{32} + x^{26} + x^{16} + x^{12} + x^8 \\ &\quad + x^6 + x^2, \\ p_{12}(x) &= x^{84} + x^{82} + x^{80} + x^{52} + x^{50} + x^{48} + x^4 + x^2 + 1, \end{aligned}$$

and the initial terms are $[1, 0, 1, 0, 1, 0, 1, 0, 0, 0, 0, 0, 1, 0, 1, 0]$.

For $\phi(M^e, 0, 1)$:

$$\begin{aligned} p_0(x) &= x^{108} + x^{103} + x^{98} + x^{95} + x^{94} + x^{92} + x^{91} + x^{89} + x^{88} + x^{87} + x^{84} \\ &\quad + x^{83} + x^{82} + x^{81} + x^{79} + x^{76} + x^{74} + x^{73} + x^{72} + x^{71} + x^{66} + x^{63} \\ &\quad + x^{60} + x^{59} + x^{55} + x^{54} + x^{53} + x^{52} + x^{51} + x^{47} + x^{46} + x^{45} + x^{44} \\ &\quad + x^{43} + x^{40} + x^{39} + x^{36} + x^{34} + x^{32} + x^{31} + x^{30} + x^{29} + x^{28} + x^{26} \\ &\quad + x^{24}, \\ p_3(x) &= x^{99} + x^{96} + x^{94} + x^{90} + x^{89} + x^{88} + x^{87} + x^{86} + x^{85} + x^{84} + x^{82} \\ &\quad + x^{81} + x^{75} + x^{72} + x^{70} + x^{66} + x^{65} + x^{64} + x^{63} + x^{62} + x^{61} + x^{60} \\ &\quad + x^{58} + x^{57} + x^{51} + x^{48} + x^{46} + x^{43} + x^{42} + x^{41} + x^{39} + x^{37} + x^{36} \\ &\quad + x^{35} + x^{31} + x^{29} + x^{28} + x^{27} + x^{23} + x^{21} + x^{20} + x^{16} + x^{15} + x^{14} \\ &\quad + x^{13} + x^{12} + x^{10} + x^9, \\ p_6(x) &= x^{96} + x^{95} + x^{93} + x^{92} + x^{88} + x^{87} + x^{86} + x^{82} + x^{81} + x^{79} + x^{77} \end{aligned}$$

$$\begin{aligned}
& + x^{75} + x^{74} + x^{72} + x^{71} + x^{69} + x^{67} + x^{66} + x^{62} + x^{61} + x^{60} + x^{58} \\
& + x^{57} + x^{55} + x^{54} + x^{52} + x^{51} + x^{49} + x^{48} + x^{46} + x^{45} + x^{43} + x^{42} \\
& + x^{36} + x^{35} + x^{34} + x^{32} + x^{31} + x^{29} + x^{27} + x^{25} + x^{24} + x^{22} + x^{21} \\
& + x^{19} + x^{17} + x^{16} + x^8 + x^7 + x^6, \\
p_9(x) &= x^{93} + x^{91} + x^{90} + x^{86} + x^{84} + x^{81} + x^{80} + x^{77} + x^{76} + x^{72} + x^{71} \\
& + x^{70} + x^{68} + x^{64} + x^{62} + x^{59} + x^{58} + x^{57} + x^{53} + x^{51} + x^{50} + x^{46} \\
& + x^{44} + x^{43} + x^{39} + x^{36} + x^{35} + x^{34} + x^{33} + x^{32} + x^{31} + x^{30} + x^{28} \\
& + x^{27} + x^{25} + x^{22} + x^{20} + x^{17} + x^{16} + x^{15} + x^9 + x^6 + x^4 + x^3, \\
p_{12}(x) &= x^{90} + x^{89} + x^{88} + x^{86} + x^{85} + x^{84} + x^{82} + x^{81} + x^{80} + x^{58} + x^{57} \\
& + x^{56} + x^{54} + x^{53} + x^{52} + x^{50} + x^{49} + x^{48} + x^{10} + x^9 + x^8 + x^6 + x^5 \\
& + x^4 + x^2 + x + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 1, 1, 1, 0, 0, 0, 0, 1, 1, 1, 0]$.

For $\phi(M^e, 1, 0)$:

$$\begin{aligned}
p_0(x) &= x^{108} + x^{103} + x^{98} + x^{95} + x^{94} + x^{92} + x^{91} + x^{89} + x^{88} + x^{87} + x^{84} \\
& + x^{83} + x^{82} + x^{81} + x^{79} + x^{76} + x^{74} + x^{73} + x^{72} + x^{71} + x^{66} + x^{63} \\
& + x^{60} + x^{59} + x^{55} + x^{54} + x^{53} + x^{52} + x^{51} + x^{47} + x^{46} + x^{45} + x^{44} \\
& + x^{43} + x^{40} + x^{39} + x^{36} + x^{34} + x^{32} + x^{31} + x^{30} + x^{29} + x^{28} + x^{26} \\
& + x^{24}, \\
p_3(x) &= x^{99} + x^{96} + x^{94} + x^{90} + x^{89} + x^{88} + x^{87} + x^{86} + x^{85} + x^{84} + x^{82} \\
& + x^{81} + x^{75} + x^{72} + x^{70} + x^{66} + x^{65} + x^{64} + x^{63} + x^{62} + x^{61} + x^{60} \\
& + x^{58} + x^{57} + x^{51} + x^{48} + x^{46} + x^{43} + x^{42} + x^{41} + x^{39} + x^{37} + x^{36} \\
& + x^{35} + x^{31} + x^{29} + x^{28} + x^{27} + x^{23} + x^{21} + x^{20} + x^{16} + x^{15} + x^{14} \\
& + x^{13} + x^{12} + x^{10} + x^9, \\
p_6(x) &= x^{96} + x^{95} + x^{93} + x^{92} + x^{88} + x^{87} + x^{86} + x^{82} + x^{81} + x^{79} + x^{77} \\
& + x^{75} + x^{74} + x^{72} + x^{71} + x^{69} + x^{67} + x^{66} + x^{62} + x^{61} + x^{60} + x^{58} \\
& + x^{57} + x^{55} + x^{54} + x^{52} + x^{51} + x^{49} + x^{48} + x^{46} + x^{45} + x^{43} + x^{42} \\
& + x^{36} + x^{35} + x^{34} + x^{32} + x^{31} + x^{29} + x^{27} + x^{25} + x^{24} + x^{22} + x^{21} \\
& + x^{19} + x^{17} + x^{16} + x^8 + x^7 + x^6, \\
p_9(x) &= x^{93} + x^{91} + x^{90} + x^{86} + x^{84} + x^{81} + x^{80} + x^{77} + x^{76} + x^{72} + x^{71} \\
& + x^{70} + x^{68} + x^{64} + x^{62} + x^{59} + x^{58} + x^{57} + x^{53} + x^{51} + x^{50} + x^{46} \\
& + x^{44} + x^{43} + x^{39} + x^{36} + x^{35} + x^{34} + x^{33} + x^{32} + x^{31} + x^{30} + x^{28} \\
& + x^{27} + x^{25} + x^{22} + x^{20} + x^{17} + x^{16} + x^{15} + x^9 + x^6 + x^4 + x^3, \\
p_{12}(x) &= x^{90} + x^{89} + x^{88} + x^{86} + x^{85} + x^{84} + x^{82} + x^{81} + x^{80} + x^{58} + x^{57} \\
& + x^{56} + x^{54} + x^{53} + x^{52} + x^{50} + x^{49} + x^{48} + x^{10} + x^9 + x^8 + x^6 + x^5 \\
& + x^4 + x^2 + x + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 1, 1, 1, 0, 0, 0, 0, 1, 1, 1, 0]$.

For $\phi(M^e, 1, 1)$:

$$\begin{aligned}
p_0(x) &= x^{120} + x^{116} + x^{114} + x^{112} + x^{104} + x^{102} + x^{98} + x^{96} + x^{82} + x^{80} \\
&\quad + x^{78} + x^{72} + x^{68} + x^{64} + x^{62} + x^{60} + x^{54} + x^{50} + x^{48} + x^{46} + x^{42} \\
&\quad + x^{38} + x^{36}, \\
p_3(x) &= x^{102} + x^{100} + x^{92} + x^{86} + x^{82} + x^{78} + x^{76} + x^{68} + x^{62} + x^{58} + x^{52} \\
&\quad + x^{50} + x^{48} + x^{38} + x^{32} + x^{30} + x^{26} + x^{22} + x^{20} + x^{18} + x^8 + x^6, \\
p_6(x) &= x^{102} + x^{100} + x^{96} + x^{90} + x^{88} + x^{86} + x^{84} + x^{82} + x^{68} + x^{64} + x^{60} \\
&\quad + x^{56} + x^{54} + x^{52} + x^{50} + x^{42} + x^{40} + x^{38} + x^{30} + x^{28} + x^{24} + x^{22} \\
&\quad + x^{18} + x^{16} + x^{14} + x^{10} + x^8 + x^6 + x^2 + 1, \\
p_9(x) &= x^{96} + x^{90} + x^{84} + x^{82} + x^{78} + x^{76} + x^{74} + x^{72} + x^{70} + x^{62} + x^{50} \\
&\quad + x^{46} + x^{44} + x^{40} + x^{34} + x^{24} + x^{20} + x^{14} + x^{12} + x^{10} + x^8 + x^6 + x^4 \\
&\quad + x^2, \\
p_{12}(x) &= x^{96} + x^{88} + x^{80} + x^{64} + x^{56} + x^{48} + x^{16} + x^8 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 1, 0, 0, 0, 0, 0]$.

For $\phi(W^e, 0, 0)$:

$$\begin{aligned}
p_0(x) &= x^{120} + x^{116} + x^{114} + x^{112} + x^{108} + x^{102} + x^{100} + x^{98} + x^{92} + x^{90} \\
&\quad + x^{88} + x^{82} + x^{78} + x^{76} + x^{74} + x^{70} + x^{68} + x^{58} + x^{56} + x^{54} + x^{46} \\
&\quad + x^{34} + x^{24} + x^{22} + x^{12}, \\
p_3(x) &= x^{102} + x^{100} + x^{92} + x^{90} + x^{88} + x^{86} + x^{78} + x^{66} + x^{54} + x^{44} + x^{42} \\
&\quad + x^{40} + x^{36} + x^{34} + x^{28} + x^{26} + x^{18} + x^{16} + x^{14} + x^8, \\
p_6(x) &= x^{102} + x^{100} + x^{96} + x^{86} + x^{80} + x^{72} + x^{68} + x^{66} + x^{64} + x^{62} + x^{58} \\
&\quad + x^{44} + x^{40} + x^{34} + x^{28} + x^{20} + x^{18} + x^{10} + x^8 + x^4 + x^2 + 1, \\
p_9(x) &= x^{96} + x^{90} + x^{84} + x^{82} + x^{78} + x^{76} + x^{74} + x^{72} + x^{70} + x^{62} + x^{50} \\
&\quad + x^{46} + x^{44} + x^{40} + x^{34} + x^{24} + x^{20} + x^{14} + x^{12} + x^{10} + x^8 + x^6 + x^4 \\
&\quad + x^2, \\
p_{12}(x) &= x^{96} + x^{88} + x^{80} + x^{64} + x^{56} + x^{48} + x^{16} + x^8 + 1,
\end{aligned}$$

and the initial terms are $[1, 0, 0, 0, 1, 0, 0, 0, 0, 0, 1, 0, 0, 0, 0, 0]$.

For $\phi(W^e, 0, 1)$:

$$\begin{aligned}
p_0(x) &= x^{108} + x^{104} + x^{102} + x^{99} + x^{97} + x^{96} + x^{95} + x^{93} + x^{90} + x^{89} + x^{83} \\
&\quad + x^{82} + x^{80} + x^{77} + x^{74} + x^{73} + x^{71} + x^{66} + x^{63} + x^{62} + x^{61} + x^{60} \\
&\quad + x^{59} + x^{57} + x^{53} + x^{52} + x^{49} + x^{47} + x^{44} + x^{42} + x^{40} + x^{39} + x^{36} \\
&\quad + x^{35} + x^{33} + x^{27} + x^{24} + x^{19} + x^{18}, \\
p_3(x) &= x^{99} + x^{96} + x^{94} + x^{90} + x^{89} + x^{88} + x^{87} + x^{86} + x^{85} + x^{84} + x^{82} \\
&\quad + x^{81} + x^{75} + x^{72} + x^{70} + x^{66} + x^{65} + x^{64} + x^{63} + x^{62} + x^{61} + x^{60} \\
&\quad + x^{58} + x^{57} + x^{51} + x^{48} + x^{46} + x^{43} + x^{42} + x^{41} + x^{39} + x^{37} + x^{36}
\end{aligned}$$

$$\begin{aligned}
& + x^{35} + x^{31} + x^{29} + x^{28} + x^{27} + x^{23} + x^{21} + x^{20} + x^{16} + x^{15} + x^{14} \\
& + x^{13} + x^{12} + x^{10} + x^9, \\
p_6(x) &= x^{96} + x^{95} + x^{93} + x^{92} + x^{88} + x^{87} + x^{86} + x^{82} + x^{81} + x^{79} + x^{77} \\
& + x^{75} + x^{74} + x^{72} + x^{71} + x^{69} + x^{67} + x^{66} + x^{62} + x^{61} + x^{60} + x^{58} \\
& + x^{57} + x^{55} + x^{54} + x^{52} + x^{51} + x^{49} + x^{48} + x^{46} + x^{45} + x^{43} + x^{42} \\
& + x^{36} + x^{35} + x^{34} + x^{32} + x^{31} + x^{29} + x^{27} + x^{25} + x^{24} + x^{22} + x^{21} \\
& + x^{19} + x^{17} + x^{16} + x^8 + x^7 + x^6, \\
p_9(x) &= x^{93} + x^{91} + x^{90} + x^{86} + x^{84} + x^{81} + x^{80} + x^{77} + x^{76} + x^{72} + x^{71} \\
& + x^{70} + x^{68} + x^{64} + x^{62} + x^{59} + x^{58} + x^{57} + x^{53} + x^{51} + x^{50} + x^{46} \\
& + x^{44} + x^{43} + x^{39} + x^{36} + x^{35} + x^{34} + x^{33} + x^{32} + x^{31} + x^{30} + x^{28} \\
& + x^{27} + x^{25} + x^{22} + x^{20} + x^{17} + x^{16} + x^{15} + x^9 + x^6 + x^4 + x^3, \\
p_{12}(x) &= x^{90} + x^{89} + x^{88} + x^{86} + x^{85} + x^{84} + x^{82} + x^{81} + x^{80} + x^{58} + x^{57} \\
& + x^{56} + x^{54} + x^{53} + x^{52} + x^{50} + x^{49} + x^{48} + x^{10} + x^9 + x^8 + x^6 + x^5 \\
& + x^4 + x^2 + x + 1,
\end{aligned}$$

and the initial terms are $[0, 1, 0, 0, 0, 1, 0, 1, 1, 0, 1, 1, 1, 0, 0, 0]$.

For $\phi(W^e, 1, 0)$:

$$\begin{aligned}
p_0(x) &= x^{108} + x^{104} + x^{102} + x^{99} + x^{97} + x^{96} + x^{95} + x^{93} + x^{90} + x^{89} + x^{83} \\
& + x^{82} + x^{80} + x^{77} + x^{74} + x^{73} + x^{71} + x^{66} + x^{63} + x^{62} + x^{61} + x^{60} \\
& + x^{59} + x^{57} + x^{53} + x^{52} + x^{49} + x^{47} + x^{44} + x^{42} + x^{40} + x^{39} + x^{36} \\
& + x^{35} + x^{33} + x^{27} + x^{24} + x^{19} + x^{18}, \\
p_3(x) &= x^{99} + x^{96} + x^{94} + x^{90} + x^{89} + x^{88} + x^{87} + x^{86} + x^{85} + x^{84} + x^{82} \\
& + x^{81} + x^{75} + x^{72} + x^{70} + x^{66} + x^{65} + x^{64} + x^{63} + x^{62} + x^{61} + x^{60} \\
& + x^{58} + x^{57} + x^{51} + x^{48} + x^{46} + x^{43} + x^{42} + x^{41} + x^{39} + x^{37} + x^{36} \\
& + x^{35} + x^{31} + x^{29} + x^{28} + x^{27} + x^{23} + x^{21} + x^{20} + x^{16} + x^{15} + x^{14} \\
& + x^{13} + x^{12} + x^{10} + x^9, \\
p_6(x) &= x^{96} + x^{95} + x^{93} + x^{92} + x^{88} + x^{87} + x^{86} + x^{82} + x^{81} + x^{79} + x^{77} \\
& + x^{75} + x^{74} + x^{72} + x^{71} + x^{69} + x^{67} + x^{66} + x^{62} + x^{61} + x^{60} + x^{58} \\
& + x^{57} + x^{55} + x^{54} + x^{52} + x^{51} + x^{49} + x^{48} + x^{46} + x^{45} + x^{43} + x^{42} \\
& + x^{36} + x^{35} + x^{34} + x^{32} + x^{31} + x^{29} + x^{27} + x^{25} + x^{24} + x^{22} + x^{21} \\
& + x^{19} + x^{17} + x^{16} + x^8 + x^7 + x^6, \\
p_9(x) &= x^{93} + x^{91} + x^{90} + x^{86} + x^{84} + x^{81} + x^{80} + x^{77} + x^{76} + x^{72} + x^{71} \\
& + x^{70} + x^{68} + x^{64} + x^{62} + x^{59} + x^{58} + x^{57} + x^{53} + x^{51} + x^{50} + x^{46} \\
& + x^{44} + x^{43} + x^{39} + x^{36} + x^{35} + x^{34} + x^{33} + x^{32} + x^{31} + x^{30} + x^{28} \\
& + x^{27} + x^{25} + x^{22} + x^{20} + x^{17} + x^{16} + x^{15} + x^9 + x^6 + x^4 + x^3, \\
p_{12}(x) &= x^{90} + x^{89} + x^{88} + x^{86} + x^{85} + x^{84} + x^{82} + x^{81} + x^{80} + x^{58} + x^{57}
\end{aligned}$$

$$\begin{aligned}
& + x^{56} + x^{54} + x^{53} + x^{52} + x^{50} + x^{49} + x^{48} + x^{10} + x^9 + x^8 + x^6 + x^5 \\
& + x^4 + x^2 + x + 1,
\end{aligned}$$

and the initial terms are $[0, 1, 0, 0, 0, 1, 0, 1, 1, 0, 1, 1, 1, 0, 0, 0]$.

For $\phi(W^e, 1, 1)$:

$$\begin{aligned}
p_0(x) &= x^{96} + x^{92} + x^{90} + x^{88} + x^{84} + x^{80} + x^{72} + x^{68} + x^{66} + x^{64} + x^{60} \\
&\quad + x^{58} + x^{56} + x^{54} + x^{52} + x^{50} + x^{48} + x^{46} + x^{44} + x^{42} + x^{38} + x^{36} \\
&\quad + x^{34} + x^{26} + x^{24}, \\
p_3(x) &= x^{86} + x^{80} + x^{78} + x^{74} + x^{70} + x^{66} + x^{64} + x^{62} + x^{42} + x^{40} + x^{38} \\
&\quad + x^{30} + x^{26} + x^{18} + x^{10} + x^6, \\
p_6(x) &= x^{86} + x^{78} + x^{64} + x^{58} + x^{56} + x^{50} + x^{48} + x^{42} + x^{34} + x^{30} + x^{22} \\
&\quad + x^{18} + x^{16} + x^{14} + x^{10} + 1, \\
p_9(x) &= x^{84} + x^{82} + x^{80} + x^{78} + x^{70} + x^{68} + x^{66} + x^{64} + x^{62} + x^{56} + x^{54} \\
&\quad + x^{52} + x^{48} + x^{46} + x^{44} + x^{38} + x^{36} + x^{32} + x^{26} + x^{16} + x^{12} + x^8 \\
&\quad + x^6 + x^2, \\
p_{12}(x) &= x^{84} + x^{82} + x^{80} + x^{52} + x^{50} + x^{48} + x^4 + x^2 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 1, 0, 0, 0, 1, 0, 0, 0, 0, 0, 1, 0, 1, 0]$.

For $\phi(M^o, 0, 0)$:

$$\begin{aligned}
p_0(x) &= x^{108} + x^{103} + x^{102} + x^{99} + x^{98} + x^{95} + x^{94} + x^{93} + x^{92} + x^{91} + x^{87} \\
&\quad + x^{84} + x^{82} + x^{78} + x^{74} + x^{72} + x^{71} + x^{70} + x^{68} + x^{59} + x^{58} + x^{57} \\
&\quad + x^{56} + x^{54} + x^{53} + x^{51} + x^{49} + x^{48} + x^{47} + x^{46} + x^{42} + x^{41} + x^{40} \\
&\quad + x^{38} + x^{37} + x^{33} + x^{32} + x^{31} + x^{28} + x^{26} + x^{25} + x^{24} + x^{22} + x^{21} \\
&\quad + x^{18} + x^{16} + x^{15} + x^{13} + x^{12}, \\
p_3(x) &= x^{102} + x^{101} + x^{98} + x^{96} + x^{95} + x^{92} + x^{90} + x^{87} + x^{86} + x^{85} + x^{84} \\
&\quad + x^{82} + x^{81} + x^{79} + x^{78} + x^{76} + x^{74} + x^{72} + x^{71} + x^{68} + x^{65} + x^{62} \\
&\quad + x^{51} + x^{50} + x^{49} + x^{47} + x^{46} + x^{45} + x^{44} + x^{42} + x^{41} + x^{40} + x^{39} \\
&\quad + x^{38} + x^{36} + x^{35} + x^{33} + x^{24} + x^{22} + x^{21} + x^{18} + x^{15} + x^{12} + x^{11} \\
&\quad + x^9 + x^8, \\
p_6(x) &= x^{102} + x^{101} + x^{98} + x^{96} + x^{95} + x^{92} + x^{90} + x^{88} + x^{87} + x^{85} + x^{84} \\
&\quad + x^{83} + x^{82} + x^{80} + x^{75} + x^{74} + x^{73} + x^{70} + x^{69} + x^{67} + x^{66} + x^{61} \\
&\quad + x^{60} + x^{59} + x^{58} + x^{55} + x^{54} + x^{53} + x^{52} + x^{50} + x^{47} + x^{46} + x^{44} \\
&\quad + x^{43} + x^{42} + x^{40} + x^{38} + x^{35} + x^{34} + x^{33} + x^{30} + x^{29} + x^{28} + x^{25} \\
&\quad + x^{24} + x^{23} + x^{22} + x^{17} + x^{15} + x^{14} + x^{12} + x^{10} + x^8 + x^7 + x^6 + x^2 \\
&\quad + x + 1, \\
p_9(x) &= x^{102} + x^{101} + x^{100} + x^{94} + x^{93} + x^{92} + x^{90} + x^{89} + x^{88} + x^{86} + x^{85} \\
&\quad + x^{84} + x^{70} + x^{69} + x^{68} + x^{62} + x^{61} + x^{60} + x^{58} + x^{57} + x^{56} + x^{54}
\end{aligned}$$

$$\begin{aligned}
& + x^{53} + x^{52} + x^{22} + x^{21} + x^{20} + x^{14} + x^{13} + x^{12} + x^{10} + x^9 + x^8 + x^6 \\
& + x^5 + x^4, \\
p_{12}(x) = & x^{102} + x^{101} + x^{100} + x^{94} + x^{93} + x^{92} + x^{82} + x^{81} + x^{80} + x^{70} + x^{69} \\
& + x^{68} + x^{62} + x^{61} + x^{60} + x^{50} + x^{49} + x^{48} + x^{22} + x^{21} + x^{20} + x^{14} \\
& + x^{13} + x^{12} + x^2 + x + 1,
\end{aligned}$$

and the initial terms are $[1, 0, 1, 0, 0, 0, 1, 1, 0, 0, 1, 1, 0, 1, 1, 1]$.

For $\phi(M^o, 0, 1)$:

$$\begin{aligned}
p_0(x) = & x^{120} + x^{116} + x^{112} + x^{111} + x^{110} + x^{109} + x^{108} + x^{107} + x^{105} + x^{104} \\
& + x^{98} + x^{97} + x^{96} + x^{93} + x^{91} + x^{90} + x^{86} + x^{85} + x^{84} + x^{82} + x^{80} \\
& + x^{79} + x^{78} + x^{76} + x^{73} + x^{71} + x^{70} + x^{59} + x^{57} + x^{56} + x^{55} + x^{54} \\
& + x^{53} + x^{51} + x^{48} + x^{46} + x^{45} + x^{44} + x^{42} + x^{41} + x^{39} + x^{38} + x^{35} \\
& + x^{34} + x^{32} + x^{30} + x^{27} + x^{25} + x^{24}, \\
p_3(x) = & x^{99} + x^{98} + x^{97} + x^{96} + x^{95} + x^{93} + x^{92} + x^{91} + x^{88} + x^{86} + x^{85} \\
& + x^{81} + x^{75} + x^{74} + x^{73} + x^{72} + x^{71} + x^{69} + x^{68} + x^{67} + x^{64} + x^{62} \\
& + x^{61} + x^{57} + x^{51} + x^{50} + x^{49} + x^{48} + x^{47} + x^{45} + x^{44} + x^{42} + x^{41} \\
& + x^{39} + x^{38} + x^{36} + x^{34} + x^{31} + x^{30} + x^{28} + x^{26} + x^{23} + x^{22} + x^{20} \\
& + x^{19} + x^{17} + x^{16} + x^{14} + x^{13} + x^9, \\
p_6(x) = & x^{102} + x^{100} + x^{98} + x^{96} + x^{92} + x^{90} + x^{88} + x^{78} + x^{74} + x^{72} + x^{66} \\
& + x^{62} + x^{46} + x^{44} + x^{36} + x^{34} + x^{28} + x^{24} + x^{22} + x^{18} + x^{16} + x^{14} \\
& + x^{10} + x^6, \\
p_9(x) = & x^{105} + x^{104} + x^{101} + x^{98} + x^{97} + x^{96} + x^{94} + x^{91} + x^{90} + x^{88} + x^{87} \\
& + x^{83} + x^{73} + x^{72} + x^{69} + x^{66} + x^{65} + x^{64} + x^{62} + x^{59} + x^{58} + x^{56} \\
& + x^{55} + x^{51} + x^{25} + x^{24} + x^{21} + x^{18} + x^{17} + x^{16} + x^{14} + x^{11} + x^{10} \\
& + x^8 + x^7 + x^3, \\
p_{12}(x) = & x^{108} + x^{104} + x^{96} + x^{92} + x^{88} + x^{84} + x^{80} + x^{76} + x^{72} + x^{64} + x^{60} \\
& + x^{56} + x^{52} + x^{48} + x^{28} + x^{24} + x^{16} + x^{12} + x^8 + x^4 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 1, 1, 1, 0, 1, 1, 0, 1, 1, 1, 0]$.

For $\phi(M^o, 1, 0)$:

$$\begin{aligned}
p_0(x) = & x^{100} + x^{98} + x^{96} + x^{95} + x^{94} + x^{93} + x^{91} + x^{90} + x^{89} + x^{88} + x^{87} \\
& + x^{84} + x^{83} + x^{82} + x^{78} + x^{75} + x^{73} + x^{72} + x^{71} + x^{70} + x^{69} + x^{68} \\
& + x^{66} + x^{63} + x^{60} + x^{57} + x^{56} + x^{54} + x^{48} + x^{46} + x^{43} + x^{42} + x^{41} \\
& + x^{40} + x^{38} + x^{37} + x^{36} + x^{35} + x^{34} + x^{33} + x^{32} + x^{28} + x^{24} + x^{21} \\
& + x^{18}, \\
p_3(x) = & x^{91} + x^{90} + x^{89} + x^{88} + x^{86} + x^{81} + x^{67} + x^{66} + x^{65} + x^{64} + x^{62} \\
& + x^{57} + x^{43} + x^{42} + x^{41} + x^{40} + x^{38} + x^{35} + x^{34} + x^{32} + x^{30} + x^{27}
\end{aligned}$$

$$\begin{aligned}
& + x^{26} + x^{24} + x^{22} + x^{19} + x^{18} + x^{16} + x^{14} + x^9, \\
p_6(x) &= x^{94} + x^{92} + x^{86} + x^{80} + x^{78} + x^{76} + x^{74} + x^{70} + x^{68} + x^{66} + x^{60} \\
& + x^{56} + x^{54} + x^{50} + x^{48} + x^{44} + x^{42} + x^{34} + x^{30} + x^{28} + x^{26} + x^{24} \\
& + x^{20} + x^{18} + x^{16} + x^6, \\
p_9(x) &= x^{97} + x^{96} + x^{92} + x^{90} + x^{88} + x^{83} + x^{65} + x^{64} + x^{60} + x^{58} + x^{56} \\
& + x^{51} + x^{17} + x^{16} + x^{12} + x^{10} + x^8 + x^3, \\
p_{12}(x) &= x^{100} + x^{92} + x^{80} + x^{68} + x^{60} + x^{48} + x^{20} + x^{12} + 1,
\end{aligned}$$

and the initial terms are $[0, 1, 0, 1, 0, 0, 0, 0, 0, 1, 0, 0, 0, 0, 0, 1]$.

For $\phi(M^o, 1, 1)$:

$$\begin{aligned}
p_0(x) &= x^{108} + x^{104} + x^{98} + x^{96} + x^{94} + x^{93} + x^{92} + x^{90} + x^{89} + x^{86} + x^{81} \\
& + x^{78} + x^{74} + x^{72} + x^{70} + x^{68} + x^{64} + x^{63} + x^{62} + x^{60} + x^{59} + x^{58} \\
& + x^{53} + x^{52} + x^{51} + x^{45} + x^{43} + x^{42} + x^{40} + x^{39} + x^{38} + x^{34} + x^{33} \\
& + x^{32} + x^{30}, \\
p_3(x) &= x^{102} + x^{101} + x^{98} + x^{93} + x^{89} + x^{86} + x^{84} + x^{83} + x^{82} + x^{80} + x^{79} \\
& + x^{78} + x^{76} + x^{74} + x^{69} + x^{66} + x^{63} + x^{62} + x^{61} + x^{59} + x^{57} + x^{56} \\
& + x^{51} + x^{50} + x^{49} + x^{48} + x^{46} + x^{39} + x^{38} + x^{34} + x^{33} + x^{28} + x^{27} \\
& + x^{26} + x^{24} + x^{22} + x^{21} + x^{20} + x^{19} + x^{16} + x^{13} + x^{12}, \\
p_6(x) &= x^{102} + x^{101} + x^{98} + x^{93} + x^{89} + x^{88} + x^{83} + x^{82} + x^{79} + x^{78} + x^{76} \\
& + x^{73} + x^{72} + x^{71} + x^{70} + x^{61} + x^{60} + x^{59} + x^{57} + x^{53} + x^{51} + x^{50} \\
& + x^{49} + x^{48} + x^{47} + x^{45} + x^{44} + x^{40} + x^{38} + x^{36} + x^{34} + x^{31} + x^{30} \\
& + x^{28} + x^{27} + x^{26} + x^{25} + x^{24} + x^{23} + x^{21} + x^{19} + x^{18} + x^{17} + x^{15} \\
& + x^{13} + x^{10} + x^6 + x^5 + x^4 + x^2 + x + 1, \\
p_9(x) &= x^{102} + x^{101} + x^{100} + x^{94} + x^{93} + x^{92} + x^{90} + x^{89} + x^{88} + x^{86} + x^{85} \\
& + x^{84} + x^{70} + x^{69} + x^{68} + x^{62} + x^{61} + x^{60} + x^{58} + x^{57} + x^{56} + x^{54} \\
& + x^{53} + x^{52} + x^{22} + x^{21} + x^{20} + x^{14} + x^{13} + x^{12} + x^{10} + x^9 + x^8 + x^6 \\
& + x^5 + x^4, \\
p_{12}(x) &= x^{102} + x^{101} + x^{100} + x^{94} + x^{93} + x^{92} + x^{82} + x^{81} + x^{80} + x^{70} + x^{69} \\
& + x^{68} + x^{62} + x^{61} + x^{60} + x^{50} + x^{49} + x^{48} + x^{22} + x^{21} + x^{20} + x^{14} \\
& + x^{13} + x^{12} + x^2 + x + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 0, 1, 1, 1, 0, 1, 1, 1, 1, 1, 1]$.

For $\phi(W^o, 0, 0)$:

$$\begin{aligned}
p_0(x) &= x^{108} + x^{103} + x^{102} + x^{99} + x^{98} + x^{95} + x^{94} + x^{93} + x^{92} + x^{91} + x^{87} \\
& + x^{84} + x^{82} + x^{78} + x^{74} + x^{72} + x^{71} + x^{70} + x^{68} + x^{59} + x^{58} + x^{57} \\
& + x^{56} + x^{54} + x^{53} + x^{51} + x^{49} + x^{48} + x^{47} + x^{46} + x^{42} + x^{41} + x^{40} \\
& + x^{38} + x^{37} + x^{33} + x^{32} + x^{31} + x^{28} + x^{26} + x^{25} + x^{24} + x^{22} + x^{21}
\end{aligned}$$

$$\begin{aligned}
& + x^{18} + x^{16} + x^{15} + x^{13} + x^{12}, \\
p_3(x) &= x^{102} + x^{101} + x^{98} + x^{96} + x^{95} + x^{92} + x^{90} + x^{87} + x^{86} + x^{85} + x^{84} \\
& + x^{82} + x^{81} + x^{79} + x^{78} + x^{76} + x^{74} + x^{72} + x^{71} + x^{68} + x^{65} + x^{62} \\
& + x^{51} + x^{50} + x^{49} + x^{47} + x^{46} + x^{45} + x^{44} + x^{42} + x^{41} + x^{40} + x^{39} \\
& + x^{38} + x^{36} + x^{35} + x^{33} + x^{24} + x^{22} + x^{21} + x^{18} + x^{15} + x^{12} + x^{11} \\
& + x^9 + x^8, \\
p_6(x) &= x^{102} + x^{101} + x^{98} + x^{96} + x^{95} + x^{92} + x^{90} + x^{88} + x^{87} + x^{85} + x^{84} \\
& + x^{83} + x^{82} + x^{80} + x^{75} + x^{74} + x^{73} + x^{70} + x^{69} + x^{67} + x^{66} + x^{61} \\
& + x^{60} + x^{59} + x^{58} + x^{55} + x^{54} + x^{53} + x^{52} + x^{50} + x^{47} + x^{46} + x^{44} \\
& + x^{43} + x^{42} + x^{40} + x^{38} + x^{35} + x^{34} + x^{33} + x^{30} + x^{29} + x^{28} + x^{25} \\
& + x^{24} + x^{23} + x^{22} + x^{17} + x^{15} + x^{14} + x^{12} + x^{10} + x^8 + x^7 + x^6 + x^2 \\
& + x + 1, \\
p_9(x) &= x^{102} + x^{101} + x^{100} + x^{94} + x^{93} + x^{92} + x^{90} + x^{89} + x^{88} + x^{86} + x^{85} \\
& + x^{84} + x^{70} + x^{69} + x^{68} + x^{62} + x^{61} + x^{60} + x^{58} + x^{57} + x^{56} + x^{54} \\
& + x^{53} + x^{52} + x^{22} + x^{21} + x^{20} + x^{14} + x^{13} + x^{12} + x^{10} + x^9 + x^8 + x^6 \\
& + x^5 + x^4, \\
p_{12}(x) &= x^{102} + x^{101} + x^{100} + x^{94} + x^{93} + x^{92} + x^{82} + x^{81} + x^{80} + x^{70} + x^{69} \\
& + x^{68} + x^{62} + x^{61} + x^{60} + x^{50} + x^{49} + x^{48} + x^{22} + x^{21} + x^{20} + x^{14} \\
& + x^{13} + x^{12} + x^2 + x + 1,
\end{aligned}$$

and the initial terms are $[1, 0, 1, 0, 0, 0, 1, 1, 0, 0, 1, 1, 0, 1, 1, 1]$.

For $\phi(W^o, 0, 1)$:

$$\begin{aligned}
p_0(x) &= x^{100} + x^{98} + x^{96} + x^{95} + x^{94} + x^{93} + x^{91} + x^{90} + x^{89} + x^{88} + x^{87} \\
& + x^{84} + x^{83} + x^{82} + x^{78} + x^{75} + x^{73} + x^{72} + x^{71} + x^{70} + x^{69} + x^{68} \\
& + x^{66} + x^{63} + x^{60} + x^{57} + x^{56} + x^{54} + x^{48} + x^{46} + x^{43} + x^{42} + x^{41} \\
& + x^{40} + x^{38} + x^{37} + x^{36} + x^{35} + x^{34} + x^{33} + x^{32} + x^{28} + x^{24} + x^{21} \\
& + x^{18}, \\
p_3(x) &= x^{91} + x^{90} + x^{89} + x^{88} + x^{86} + x^{81} + x^{67} + x^{66} + x^{65} + x^{64} + x^{62} \\
& + x^{57} + x^{43} + x^{42} + x^{41} + x^{40} + x^{38} + x^{35} + x^{34} + x^{32} + x^{30} + x^{27} \\
& + x^{26} + x^{24} + x^{22} + x^{19} + x^{18} + x^{16} + x^{14} + x^9, \\
p_6(x) &= x^{94} + x^{92} + x^{86} + x^{80} + x^{78} + x^{76} + x^{74} + x^{70} + x^{68} + x^{66} + x^{60} \\
& + x^{56} + x^{54} + x^{50} + x^{48} + x^{44} + x^{42} + x^{34} + x^{30} + x^{28} + x^{26} + x^{24} \\
& + x^{20} + x^{18} + x^{16} + x^6, \\
p_9(x) &= x^{97} + x^{96} + x^{92} + x^{90} + x^{88} + x^{83} + x^{65} + x^{64} + x^{60} + x^{58} + x^{56} \\
& + x^{51} + x^{17} + x^{16} + x^{12} + x^{10} + x^8 + x^3, \\
p_{12}(x) &= x^{100} + x^{92} + x^{80} + x^{68} + x^{60} + x^{48} + x^{20} + x^{12} + 1,
\end{aligned}$$

and the initial terms are $[0, 1, 0, 1, 0, 0, 0, 0, 0, 1, 0, 0, 0, 0, 1]$.

For $\phi(W^o, 1, 0)$:

$$\begin{aligned}
p_0(x) &= x^{120} + x^{116} + x^{112} + x^{111} + x^{110} + x^{109} + x^{108} + x^{107} + x^{105} + x^{104} \\
&\quad + x^{98} + x^{97} + x^{96} + x^{93} + x^{91} + x^{90} + x^{86} + x^{85} + x^{84} + x^{82} + x^{80} \\
&\quad + x^{79} + x^{78} + x^{76} + x^{73} + x^{71} + x^{70} + x^{59} + x^{57} + x^{56} + x^{55} + x^{54} \\
&\quad + x^{53} + x^{51} + x^{48} + x^{46} + x^{45} + x^{44} + x^{42} + x^{41} + x^{39} + x^{38} + x^{35} \\
&\quad + x^{34} + x^{32} + x^{30} + x^{27} + x^{25} + x^{24}, \\
p_3(x) &= x^{99} + x^{98} + x^{97} + x^{96} + x^{95} + x^{93} + x^{92} + x^{91} + x^{88} + x^{86} + x^{85} \\
&\quad + x^{81} + x^{75} + x^{74} + x^{73} + x^{72} + x^{71} + x^{69} + x^{68} + x^{67} + x^{64} + x^{62} \\
&\quad + x^{61} + x^{57} + x^{51} + x^{50} + x^{49} + x^{48} + x^{47} + x^{45} + x^{44} + x^{42} + x^{41} \\
&\quad + x^{39} + x^{38} + x^{36} + x^{34} + x^{31} + x^{30} + x^{28} + x^{26} + x^{23} + x^{22} + x^{20} \\
&\quad + x^{19} + x^{17} + x^{16} + x^{14} + x^{13} + x^9, \\
p_6(x) &= x^{102} + x^{100} + x^{98} + x^{96} + x^{92} + x^{90} + x^{88} + x^{78} + x^{74} + x^{72} + x^{66} \\
&\quad + x^{62} + x^{46} + x^{44} + x^{36} + x^{34} + x^{28} + x^{24} + x^{22} + x^{18} + x^{16} + x^{14} \\
&\quad + x^{10} + x^6, \\
p_9(x) &= x^{105} + x^{104} + x^{101} + x^{98} + x^{97} + x^{96} + x^{94} + x^{91} + x^{90} + x^{88} + x^{87} \\
&\quad + x^{83} + x^{73} + x^{72} + x^{69} + x^{66} + x^{65} + x^{64} + x^{62} + x^{59} + x^{58} + x^{56} \\
&\quad + x^{55} + x^{51} + x^{25} + x^{24} + x^{21} + x^{18} + x^{17} + x^{16} + x^{14} + x^{11} + x^{10} \\
&\quad + x^8 + x^7 + x^3, \\
p_{12}(x) &= x^{108} + x^{104} + x^{96} + x^{92} + x^{88} + x^{84} + x^{80} + x^{76} + x^{72} + x^{64} + x^{60} \\
&\quad + x^{56} + x^{52} + x^{48} + x^{28} + x^{24} + x^{16} + x^{12} + x^8 + x^4 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 1, 1, 1, 0, 1, 1, 0, 1, 1, 1, 0]$.

For $\phi(W^o, 1, 1)$:

$$\begin{aligned}
p_0(x) &= x^{108} + x^{104} + x^{98} + x^{96} + x^{94} + x^{93} + x^{92} + x^{90} + x^{89} + x^{86} + x^{81} \\
&\quad + x^{78} + x^{74} + x^{72} + x^{70} + x^{68} + x^{64} + x^{63} + x^{62} + x^{60} + x^{59} + x^{58} \\
&\quad + x^{53} + x^{52} + x^{51} + x^{45} + x^{43} + x^{42} + x^{40} + x^{39} + x^{38} + x^{34} + x^{33} \\
&\quad + x^{32} + x^{30}, \\
p_3(x) &= x^{102} + x^{101} + x^{98} + x^{93} + x^{89} + x^{86} + x^{84} + x^{83} + x^{82} + x^{80} + x^{79} \\
&\quad + x^{78} + x^{76} + x^{74} + x^{69} + x^{66} + x^{63} + x^{62} + x^{61} + x^{59} + x^{57} + x^{56} \\
&\quad + x^{51} + x^{50} + x^{49} + x^{48} + x^{46} + x^{39} + x^{38} + x^{34} + x^{33} + x^{28} + x^{27} \\
&\quad + x^{26} + x^{24} + x^{22} + x^{21} + x^{20} + x^{19} + x^{16} + x^{13} + x^{12}, \\
p_6(x) &= x^{102} + x^{101} + x^{98} + x^{93} + x^{89} + x^{88} + x^{83} + x^{82} + x^{79} + x^{78} + x^{76} \\
&\quad + x^{73} + x^{72} + x^{71} + x^{70} + x^{61} + x^{60} + x^{59} + x^{57} + x^{53} + x^{51} + x^{50} \\
&\quad + x^{49} + x^{48} + x^{47} + x^{45} + x^{44} + x^{40} + x^{38} + x^{36} + x^{34} + x^{31} + x^{30} \\
&\quad + x^{28} + x^{27} + x^{26} + x^{25} + x^{24} + x^{23} + x^{21} + x^{19} + x^{18} + x^{17} + x^{15}
\end{aligned}$$

$$\begin{aligned}
& + x^{13} + x^{10} + x^6 + x^5 + x^4 + x^2 + x + 1, \\
p_9(x) = & x^{102} + x^{101} + x^{100} + x^{94} + x^{93} + x^{92} + x^{90} + x^{89} + x^{88} + x^{86} + x^{85} \\
& + x^{84} + x^{70} + x^{69} + x^{68} + x^{62} + x^{61} + x^{60} + x^{58} + x^{57} + x^{56} + x^{54} \\
& + x^{53} + x^{52} + x^{22} + x^{21} + x^{20} + x^{14} + x^{13} + x^{12} + x^{10} + x^9 + x^8 + x^6 \\
& + x^5 + x^4, \\
p_{12}(x) = & x^{102} + x^{101} + x^{100} + x^{94} + x^{93} + x^{92} + x^{82} + x^{81} + x^{80} + x^{70} + x^{69} \\
& + x^{68} + x^{62} + x^{61} + x^{60} + x^{50} + x^{49} + x^{48} + x^{22} + x^{21} + x^{20} + x^{14} \\
& + x^{13} + x^{12} + x^2 + x + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 0, 1, 1, 1, 0, 1, 1, 1, 1, 1, 1]$.

Below is the transition function and output function of an 2-automaton that generates $T = M_{0,0}^o$:

Transition function $(n, j) \mapsto \delta(n, j)$ ($\Lambda(n) := [\delta(n, 0), \delta(n, 1)]$):

n	$\Lambda(n)$	n	$\Lambda(n)$	n	$\Lambda(n)$	n	$\Lambda(n)$	n	$\Lambda(n)$
0	[1, 2]	64	[105, 8]	128	[146, 159]	192	[101, 197]	256	[97, 102]
1	[3, 4]	65	[106, 107]	129	[184, 179]	193	[228, 45]	257	[118, 54]
2	[5, 6]	66	[108, 38]	130	[185, 186]	194	[213, 202]	258	[215, 114]
3	[7, 8]	67	[108, 109]	131	[187, 152]	195	[203, 2]	259	[279, 198]
4	[9, 10]	68	[110, 111]	132	[188, 151]	196	[229, 160]	260	[208, 34]
5	[11, 12]	69	[112, 113]	133	[189, 190]	197	[62, 230]	261	[280, 49]
6	[13, 14]	70	[105, 114]	134	[191, 168]	198	[6, 186]	262	[273, 202]
7	[15, 16]	71	[115, 53]	135	[192, 139]	199	[231, 232]	263	[221, 37]
8	[17, 18]	72	[102, 116]	136	[193, 194]	200	[233, 234]	264	[223, 137]
9	[19, 20]	73	[117, 118]	137	[195, 155]	201	[235, 15]	265	[281, 38]
10	[21, 22]	74	[31, 119]	138	[107, 106]	202	[236, 237]	266	[276, 32]
11	[23, 24]	75	[120, 121]	139	[136, 196]	203	[171, 238]	267	[282, 38]
12	[25, 26]	76	[114, 31]	140	[197, 198]	204	[239, 232]	268	[283, 207]
13	[27, 28]	77	[122, 53]	141	[199, 102]	205	[240, 67]	269	[284, 99]
14	[29, 30]	78	[96, 100]	142	[200, 36]	206	[234, 241]	270	[285, 58]
15	[8, 31]	79	[34, 123]	143	[130, 198]	207	[242, 153]	271	[286, 287]
16	[10, 32]	80	[124, 96]	144	[201, 202]	208	[243, 73]	272	[158, 240]
17	[33, 34]	81	[125, 118]	145	[203, 204]	209	[244, 70]	273	[140, 148]
18	[35, 36]	82	[98, 126]	146	[109, 95]	210	[245, 178]	274	[288, 92]
19	[37, 38]	83	[54, 42]	147	[197, 116]	211	[246, 247]	275	[156, 289]
20	[39, 40]	84	[127, 128]	148	[14, 47]	212	[248, 249]	276	[72, 26]
21	[41, 42]	85	[39, 129]	149	[122, 205]	213	[250, 231]	277	[290, 291]
22	[43, 12]	86	[130, 116]	150	[34, 42]	214	[251, 252]	278	[292, 293]
23	[44, 45]	87	[131, 113]	151	[206, 207]	215	[253, 254]	279	[294, 169]
24	[46, 47]	88	[120, 100]	152	[36, 102]	216	[255, 256]	280	[182, 295]
25	[48, 49]	89	[132, 10]	153	[117, 208]	217	[257, 258]	281	[296, 146]
26	[50, 51]	90	[133, 134]	154	[209, 96]	218	[259, 260]	282	[297, 182]
27	[52, 31]	91	[114, 44]	155	[95, 37]	219	[254, 261]	283	[298, 299]
28	[49, 53]	92	[135, 14]	156	[38, 128]	220	[262, 258]	284	[175, 90]

29	[33, 54]	93	[128, 37]	157	[95, 108]	221	[263, 67]	285	[11, 278]
30	[13, 46]	94	[136, 137]	158	[128, 108]	222	[150, 257]	286	[300, 108]
31	[16, 55]	95	[66, 138]	159	[109, 128]	223	[186, 237]	287	[301, 129]
32	[20, 56]	96	[139, 140]	160	[37, 109]	224	[264, 175]	288	[222, 211]
33	[57, 58]	97	[141, 142]	161	[210, 121]	225	[265, 155]	289	[283, 274]
34	[59, 60]	98	[143, 144]	162	[50, 211]	226	[266, 79]	290	[41, 123]
35	[61, 62]	99	[145, 146]	163	[208, 54]	227	[267, 268]	291	[216, 211]
36	[63, 64]	100	[70, 147]	164	[132, 212]	228	[269, 147]	292	[271, 98]
37	[65, 66]	101	[148, 6]	165	[213, 214]	229	[219, 270]	293	[201, 214]
38	[67, 19]	102	[149, 150]	166	[124, 47]	230	[102, 198]	294	[284, 126]
39	[28, 68]	103	[151, 65]	167	[215, 8]	231	[118, 34]	295	[302, 204]
40	[69, 70]	104	[152, 153]	168	[216, 51]	232	[125, 208]	296	[281, 109]
41	[71, 72]	105	[154, 77]	169	[44, 119]	233	[52, 44]	297	[303, 128]
42	[73, 74]	106	[106, 106]	170	[217, 34]	234	[111, 99]	298	[277, 137]
43	[75, 76]	107	[155, 156]	171	[212, 103]	235	[226, 123]	299	[304, 136]
44	[77, 78]	108	[157, 158]	172	[199, 197]	236	[271, 111]	300	[305, 93]
45	[18, 79]	109	[159, 157]	173	[135, 46]	237	[131, 134]	301	[306, 307]
46	[80, 81]	110	[160, 145]	174	[115, 205]	238	[272, 49]	302	[308, 309]
47	[82, 83]	111	[161, 162]	175	[54, 123]	239	[273, 214]	303	[310, 56]
48	[84, 85]	112	[78, 163]	176	[210, 100]	240	[206, 274]	304	[93, 229]
49	[86, 87]	113	[164, 165]	177	[218, 12]	241	[275, 111]	305	[311, 106]
50	[88, 57]	114	[166, 167]	178	[96, 121]	242	[112, 134]	306	[212, 32]
51	[89, 90]	115	[74, 168]	179	[200, 97]	243	[104, 31]	307	[312, 136]
52	[91, 92]	116	[64, 169]	180	[209, 47]	244	[36, 197]	308	[49, 205]
53	[85, 93]	117	[170, 171]	181	[219, 220]	245	[276, 103]	309	[313, 10]
54	[94, 23]	118	[172, 173]	182	[107, 107]	246	[227, 277]	310	[311, 107]
55	[31, 45]	119	[153, 72]	183	[221, 108]	247	[43, 278]	311	[314, 160]
56	[38, 95]	120	[174, 74]	184	[222, 51]	248	[228, 119]	312	[138, 315]
57	[14, 96]	121	[30, 175]	185	[223, 196]	249	[133, 113]	313	[56, 316]
58	[35, 97]	122	[176, 177]	186	[47, 121]	250	[224, 45]	314	[303, 95]
59	[98, 99]	123	[58, 178]	187	[224, 119]	251	[48, 122]	315	[301, 40]
60	[47, 100]	124	[76, 179]	188	[225, 106]	252	[218, 278]	316	[317, 270]
61	[101, 102]	125	[180, 59]	189	[226, 42]	253	[217, 54]	317	[318, 4]
62	[10, 103]	126	[181, 182]	190	[46, 96]	254	[277, 196]	318	[111, 126]
63	[104, 44]	127	[183, 65]	191	[227, 136]	255	[279, 116]		

Output function $n \mapsto \tau(n)$:

n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$
0	0	46	1	92	0	138	1	184	0	230	1	276	1
1	0	47	0	93	0	139	1	185	0	231	1	277	0
2	1	48	0	94	1	140	0	186	0	232	0	278	1
3	0	49	0	95	1	141	1	187	0	233	1	279	1
4	0	50	0	96	1	142	1	188	0	234	1	280	1
5	1	51	0	97	1	143	0	189	1	235	1	281	0

6	1	52	1	98	0	144	1	190	1	236	1	282	1
7	0	53	0	99	1	145	1	191	1	237	0	283	0
8	0	54	1	100	0	146	0	192	0	238	0	284	1
9	0	55	0	101	0	147	0	193	1	239	0	285	1
10	0	56	1	102	1	148	0	194	0	240	1	286	1
11	1	57	0	103	1	149	1	195	1	241	1	287	1
12	0	58	0	104	0	150	0	196	0	242	1	288	0
13	1	59	0	105	0	151	1	197	0	243	0	289	0
14	0	60	0	106	0	152	0	198	1	244	0	290	0
15	0	61	0	107	1	153	1	199	1	245	1	291	1
16	0	62	0	108	1	154	0	200	1	246	1	292	1
17	0	63	0	109	0	155	1	201	1	247	0	293	1
18	0	64	0	110	0	156	1	202	1	248	1	294	1
19	0	65	0	111	1	157	1	203	1	249	1	295	0
20	0	66	1	112	1	158	0	204	0	250	0	296	0
21	0	67	1	113	0	159	0	205	1	251	0	297	1
22	0	68	0	114	1	160	0	206	1	252	1	298	0
23	1	69	1	115	0	161	1	207	1	253	1	299	0
24	1	70	0	116	0	162	0	208	0	254	0	300	1
25	0	71	0	117	1	163	0	209	0	255	1	301	1
26	0	72	1	118	1	164	0	210	1	256	1	302	0
27	1	73	1	119	1	165	0	211	1	257	1	303	1
28	0	74	0	120	0	166	1	212	1	258	1	304	0
29	0	75	0	121	1	167	1	213	0	259	1	305	1
30	1	76	1	122	1	168	1	214	0	260	0	306	1
31	0	77	1	123	0	169	1	215	1	261	1	307	1
32	0	78	1	124	1	170	1	216	1	262	0	308	0
33	0	79	0	125	0	171	1	217	1	263	0	309	1
34	0	80	1	126	0	172	1	218	1	264	0	310	1
35	0	81	0	127	0	173	0	219	0	265	0	311	1
36	0	82	0	128	0	174	0	220	0	266	1	312	1
37	0	83	1	129	0	175	1	221	0	267	1	313	1
38	1	84	0	130	0	176	1	222	0	268	0	314	1
39	0	85	0	131	0	177	1	223	0	269	1	315	1
40	1	86	0	132	0	178	1	224	0	270	1	316	1
41	0	87	0	133	1	179	1	225	0	271	1	317	1
42	1	88	0	134	1	180	0	226	1	272	0	318	1
43	0	89	0	135	0	181	0	227	1	273	0		
44	1	90	1	136	1	182	1	228	1	274	0		
45	0	91	1	137	1	183	0	229	0	275	1		

UNIVERSITÉ DE STRASBOURG, CNRS, IRMA UMR 7501, F-67000 STRASBOURG, FRANCE
Email address: `guoniu.han@unistra.fr`

SCHOOL OF MATHEMATICS AND STATISTICS, HUAZHONG UNIVERSITY OF SCIENCE AND TECHNOLOGY, WUHAN, PR CHINA
Email address: `huyining@hust.edu.cn`