

PROOF OF THE CONTINUED FRACTION CONJECTURE FOR

$$(a, b) = (z, z^5 + z^4 + z^2 + z)$$

GUO-NIU HAN* AND YINING HU*

ABSTRACT. The continued fraction conjecture claims that the continued fraction defined by the (a, b) -Thue-Morse sequence is algebraic over $\mathbb{F}_2(x)$ for all pairs of distinct elements (a, b) in $\mathbb{F}_2[x] \setminus \mathbb{F}_2$. In this paper we prove the conjecture for $(a, b) = (z, z^5 + z^4 + z^2 + z)$.

1. INTRODUCTION

Let $\mathbb{F}_2$ be the finite field of cardinality 2. Given a sequence of polynomials $\mathbf{a} = (a_0(z), a_1(z), a_2(z), \dots)$ of elements from $\mathbb{F}_2[z] \setminus \mathbb{F}_2$, we define the infinite continued fraction

$$(1.1) \quad \text{CF}(\mathbf{a}(z)) = \frac{1}{a_0(z) + \frac{1}{a_1(z) + \frac{1}{a_2(z) + \frac{1}{\ddots}}}}$$

as the limit of the finite continued fractions

$$(1.2) \quad \text{CF}_n(\mathbf{a}(z)) = \frac{1}{a_0(z) + \frac{1}{a_1(z) + \frac{1}{\ddots + \frac{1}{a_n(z)}}}} \in \mathbb{F}_2((1/z)).$$

Conversely, each power series in $\mathbb{F}_2[[1/z]]$ without constant term

$$(1.3) \quad c_{-1}z^{-1} + c_{-2}z^{-2} + c_{-3}z^{-3} + \dots$$

admits a continued fraction expansion of form (1.1).

Let a and b be two distinct elements from $\mathbb{F}_2[z] \setminus \mathbb{F}_2$. We consider the continued fraction (1.1) associated with the (a, b) -Thue-Morse sequence

$$\mathbf{t} = (t_0(z), t_1(z), t_2(z), \dots) = (a, b, b, a, b, a, a, b, \dots)$$

that we shall denote

$$(1.4) \quad \text{CF}(a, b) := \text{CF}(\mathbf{t}).$$

Date: 9/04/2020 15:46:58 .

2010 Mathematics Subject Classification. 11B85, 11J70, 11B50, 11Y65, 05A15.

Key words and phrases. automatic sequence, continued fraction, Thue-Morse sequence.

* This paper was automatically generated from a computer program developed by the authors, with 6.5 minutes CPU time used.

Theorem 1.1. *When $(a, b) = (z, z^5 + z^4 + z^2 + z)$, the two power series $\text{CF}(a, b)$ and $\text{CF}(b, a)$ are algebraic over $\mathbb{F}_2(z)$, with minimal polynomials of the form*

$$p_4(z)y^4 + p_3(z)y^3 + p_2(z)y^2 + p_1(z)y + p_0(z) = 0$$

where, for $\text{CF}(a, b)$,

$$\begin{aligned} p_0(z) &= z^{14} + z^{12} + z^{11} + z^{10} + z^9 + z^7 + z^3 + 1, \\ p_1(z) &= z^{17} + z^{15} + z^{12} + z^{11} + z^{10} + z^8 + z^5 + z^4, \\ p_2(z) &= z^{18} + z^{16} + z^{14} + z^{12} + z^8 + z^4, \\ p_3(z) &= z^{17} + z^{15} + z^{12} + z^{11} + z^{10} + z^8 + z^5 + z^4, \\ p_4(z) &= z^{16} + z^{11} + z^9 + z^8 + z^7 + z^6 + z^4 + z^3 + 1, \end{aligned}$$

and for $\text{CF}(b, a)$,

$$\begin{aligned} p_0(z) &= z^{12} + z^{11} + z^9 + z^7 + z^3 + z^2 + 1, \\ p_1(z) &= z^{17} + z^{15} + z^{12} + z^{11} + z^{10} + z^8 + z^5 + z^4, \\ p_2(z) &= z^{18} + z^{16} + z^{14} + z^{12} + z^8 + z^4, \\ p_3(z) &= z^{17} + z^{15} + z^{12} + z^{11} + z^{10} + z^8 + z^5 + z^4, \\ p_4(z) &= z^{16} + z^{14} + z^{11} + z^{10} + z^9 + z^8 + z^7 + z^6 + z^4 + z^3 + z^2 + 1. \end{aligned}$$

2. PROOF

Define the sequences of polynomials $(P_n(z))_{n \geq 0}$ and $(Q_n(z))_{n \geq 0}$ by

$$\begin{pmatrix} P_n(z) & Q_n(z) \\ P_{n-1}(z) & Q_{n-1}(z) \end{pmatrix} := \begin{pmatrix} t_n(z) & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} t_{n-1}(z) & 1 \\ 1 & 0 \end{pmatrix} \cdots \begin{pmatrix} t_0(z) & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}.$$

By the basic properties of continued fractions, we have

$$(2.1) \quad \text{CF}_n(a, b) = \frac{P_n(z)}{Q_n(z)} \in \mathbb{F}_2((1/z)).$$

As formula (2.1) is not efficient for calculating $\text{CF}_n(a, b)$, a fast method is to be derived and described as follows. Define

$$\begin{aligned} M_n(x) &= x^{\deg(t_{2^n-1})} \begin{pmatrix} t_{2^n-1}(1/x) & 1 \\ 1 & 0 \end{pmatrix} \times x^{\deg(t_{2^n-2})} \begin{pmatrix} t_{2^n-2}(1/x) & 1 \\ 1 & 0 \end{pmatrix} \times \\ &\quad \cdots \times x^{\deg(t_0)} \begin{pmatrix} t_0(1/x) & 1 \\ 1 & 0 \end{pmatrix}, \\ W_n(x) &= x^{\deg(\bar{t}_{2^n-1})} \begin{pmatrix} \bar{t}_{2^n-1}(1/x) & 1 \\ 1 & 0 \end{pmatrix} \times x^{\deg(\bar{t}_{2^n-2})} \begin{pmatrix} \bar{t}_{2^n-2}(1/x) & 1 \\ 1 & 0 \end{pmatrix} \times \\ &\quad \cdots \times x^{\deg(\bar{t}_0)} \begin{pmatrix} \bar{t}_0(1/x) & 1 \\ 1 & 0 \end{pmatrix}, \end{aligned}$$

where $\bar{t}$ is the (b, a) -Thue-Morse sequence. By the properties of the Thue-Morse sequence, we have for $n \geq 0$

$$\begin{aligned} M_{n+1}(x) &= W_n(x) \cdot M_n(x), \\ W_{n+1}(x) &= M_n(x) \cdot W_n(x). \end{aligned}$$

Let $x := 1/z$. For each non-zero polynomial $P(z)$, define $\tilde{P}(x)$ to be $P(1/x)$. Then,

$$\text{CF}_n(a, b) = \frac{P_n(z)}{Q_n(z)} = \frac{\tilde{P}_n(x)}{\tilde{Q}_n(x)} \in \mathbb{F}_2((x)) = \mathbb{F}_2((1/z)).$$

Comparing the definition of $M_n(x)$ with the definition of $P_n(x)$ and $Q_n(x)$, we see that

$$\begin{aligned} M_n(x)_{0,1} &= x^{d_n} \tilde{P}_{2^n-1}(x), \\ M_n(x)_{0,0} &= x^{d_n} \tilde{Q}_{2^n-1}(x), \end{aligned}$$

for some positive integer d_n . Hence,

$$(2.2) \quad \text{CF}_{2^{2^n}-1}(a, b) = \frac{\tilde{P}_{2^{2^n}-1}(x)}{\tilde{Q}_{2^{2^n}-1}(x)} = \frac{M_{2^n}(x)_{0,1}}{M_{2^n}(x)_{0,0}}.$$

Therefore, by the convergence theorem of continued fraction, the algebraicity of $\text{CF}(a, b)$ will be established if it is shown that both $M_{2^n}(x)_{0,1}$ and $M_{2^n}(x)_{0,0}$ converge to algebraic series in $\mathbb{F}_2[[x]]$.

Actually, we will prove that for all $0 \leq i, j \leq 1$ the four sequences $(M_{2^n}(x)_{i,j})_n$, $(M_{2^{n+1}}(x)_{i,j})_n$, $(W_{2^n}(x)_{i,j})_n$, and $(W_{2^{n+1}}(x)_{i,j})_n$ converge to algebraic series in $\mathbb{F}_2[[x]]$. For this purpose, we define four 2×2 matrices M^e, M^o, W^e, W^o as follows: For each $T \in \{M^e, M^o, W^e, W^o\}$ and $0 \leq i, j \leq 1$, $T_{i,j}$ is defined to be the unique solution in $\mathbb{F}_2[[x]]$ of the polynomial $\phi(T, i, j)$ under certain initial conditions; the polynomials $\phi(T, i, j)$ and initial conditions are given in Section 3. We will prove that these four matrices, whose components are algebraic by definition, are the limits of $(M_{2^n}(x))_n$, $(M_{2^{n+1}}(x))_n$, $(W_{2^n}(x))_n$, and $(W_{2^{n+1}}(x))_n$.

Let us explain how the polynomials $\phi(T, i, j)$ and initial conditions are found, and why the solutions exist and are unique. For $0 \leq i, j \leq 1$, the coefficients of the polynomial $\phi(M^e, i, j)$ (resp. $\phi(M^o, i, j)$, $\phi(W^e, i, j)$, and $\phi(W^o, i, j)$) are the Padé-Hermite approximants of type

$$(150, 150, 150, 150, 150)$$

of the vector

$$(1, T^3, T^6, T^9, T^{12}),$$

where $T = M_{12,i,j}$ (resp. $M_{11,i,j}$, $W_{12,i,j}$, and $W_{11,i,j}$), found by the Derksen algorithm. We take the first 8 terms of $M_{12,i,j}$ (resp. $M_{11,i,j}$, $W_{12,i,j}$, and $W_{11,i,j}$) as initial conditions for $\phi(M^e, i, j)$ (resp. $\phi(M^o, i, j)$, $\phi(W^e, i, j)$, and $\phi(W^o, i, j)$). The following fact will be used to ensure that the solution exists and is unique: let $P(x, y) \in \mathbb{F}_2[x, y]$ and for each series $f(x) = \sum_{n=0}^{\infty} a_n x^n \in \mathbb{F}_2[[x]]$ denote the partial sum $\sum_{j=0}^{n-1} a_j x^j$ by $f_n(x)$ for $n \geq 0$. If for some $n \geq 0$ and $a_0, a_1, \dots, a_{n-1} \in \mathbb{F}_2$ $P(x, \sum_{j=0}^{n-1} a_j x^j) = O(x^n)$ and $Q(x, y) := P(x, \sum_{j=0}^{n-1} a_j x^j + x^n y)$ can be written as $x^m \sum_{j=0}^{\infty} q_j(x) y^j$ where $q_j(x)$ are polynomials for $j \geq 0$, $q_1(0) = 1$, and $q_j(0) = 0$ for $j > 1$, then there exists a unique solution $f(x) \in \mathbb{F}_2[[x]]$ of $P(x, f(x)) = 0$ that satisfies the initial condition $f_n(x) = \sum_{j=0}^{n-1} a_j x^j$.

We state two lemmas concerning the four matrices M^e, M^o, W^e, W^o . The first one is about relations between them; the second, about the structure of the each matrix.

Lemma 2.1. *We have*

$$(2.3) \quad M^e = W^o \cdot M^o,$$

$$(2.4) \quad M^o = W^e \cdot M^e,$$

$$(2.5) \quad W^e = M^o \cdot W^o,$$

$$(2.6) \quad W^o = M^e \cdot W^e.$$

Proof. We give the proof of the identity

$$M_{0,0}^e = W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o,$$

the proofs of the others being similar.

First, we compute the minimal polynomials of $W_{0,0}^o M_{0,0}^o$ and $W_{0,1}^o M_{1,0}^o$. We know that

$$P(x, y) = \text{Res}_z (\phi(W^o, 0, 0)(x, z), z^{12} \cdot \phi(M^o, 0, 0)(x, y/z))$$

is an annihilating polynomial of $W_{0,0}^o M_{0,0}^o$. We use Padé-Hermite approximation to find a candidate for the minimal polynomial of $W_{0,0}^o M_{0,0}^o$, that will be called $\phi_0(x, y)$. To prove that $\phi_0(x, y)$ is indeed the minimal polynomial, it suffices to prove that it is an irreducible factor of $P(x, y)$ of multiplicity m and that $Q(x, y) := P(x, y)/\phi_0(x, y)^m$ is not an annihilating polynomial of $W_{0,0}^o M_{0,0}^o$. We verify the first point directly. For the second point, we truncate $W_{0,0}^o M_{0,0}^o$ to order 540 and substitute it for y in $Q(x, y)$. We get a series of valuation less than 540, which proves that $Q(x, y)$ is not an annihilating polynomial of $W_{0,0}^o M_{0,0}^o$. We find the minimal polynomial $\phi_1(x, y)$ of $W_{0,1}^o M_{1,0}^o$ in a similar way.

Now we prove that $\phi(M^e, 0, 0)$ is the minimal polynomial of $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$. We know that

$$S(x, y) = \text{Res}_z (\phi_0(x, z), \phi_1(x, y + z))$$

is an annihilating of $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$. We verify that $\phi(M^e, 0, 0)$ is an irreducible factor of $S(x, y)$ of multiplicity μ , and that $Q(x, y) := S(x, y)/\phi(M^e, 0, 0)^\mu$ is not an annihilating polynomial of $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$. To see the last point, we truncate $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$ to order 660 and substitute it for y in $Q(x, y)$. We get a series of valuation less than 660, and therefore $Q(x, y)$ is not an annihilating polynomial of $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$.

Finally, the first 16 terms of $M_{0,0}^e$ and $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$ coincide. As these first terms determine a unique solution of $\phi(M^e, 0, 0)$, we know that the two series are one and the same. $\square$

Define

$$R^e = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \quad \text{and} \quad R^o = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}.$$

Lemma 2.2. *For all integers $k \geq 2$ and $u = 2^{2k-1}$, the following identities hold.*

$$\begin{aligned} M^e[:12u] &= M^e[:6u] + x^u M^e[:5u] + x^{2u} M^e[:4u] + x^u M^e[:6u] + x^{2u} M^e[:5u] \\ &\quad + x^{3u} M^e[:4u] + x^{3u} M^e[:6u] + x^{4u} M^e[:5u] + x^{5u} M^e[:4u] \\ &\quad + x^{4u} M^e[:6u] + x^{5u} M^e[:5u] + x^{6u} M^e[:4u] + x^{6u} M^e[:6u] \\ &\quad + x^{8u} M^e[:4u] + x^{10u} M^e[:2u] + (x^{6u} + x^{7u} + x^{9u} + x^{10u}) R^e, \\ W^e[:12u] &= W^e[:6u] + x^u W^e[:5u] + x^{2u} W^e[:4u] + x^u W^e[:6u] + x^{2u} W^e[:5u] \\ &\quad + x^{3u} W^e[:4u] + x^{3u} W^e[:6u] + x^{4u} W^e[:5u] + x^{5u} W^e[:4u] \end{aligned}$$

$$\begin{aligned}
& + x^{4u}W^e[:6u] + x^{5u}W^e[:5u] + x^{6u}W^e[:4u] + x^{6u}W^e[:6u] \\
& + x^{8u}W^e[:4u] + x^{10u}W^e[:2u] + (x^{6u} + x^{7u} + x^{9u} + x^{10u})R^e.
\end{aligned}$$

For all integers $k \geq 2$ and $u = 2^{2k}$, the following identities hold.

$$\begin{aligned}
M^o[:12u] &= M^o[:6u] + x^u M^o[:5u] + x^{2u} M^o[:4u] + x^u M^o[:6u] + x^{2u} M^o[:5u] \\
&+ x^{3u} M^o[:4u] + x^{3u} M^o[:6u] + x^{4u} M^o[:5u] + x^{5u} M^o[:4u] \\
&+ x^{4u} M^o[:6u] + x^{5u} M^o[:5u] + x^{6u} M^o[:4u] + x^{6u} M^o[:6u] \\
&+ x^{8u} M^o[:4u] + x^{10u} M^o[:2u] + (x^{6u} + x^{7u} + x^{9u} + x^{10u})R^o, \\
W^o[:12u] &= W^o[:6u] + x^u W^o[:5u] + x^{2u} W^o[:4u] + x^u W^o[:6u] + x^{2u} W^o[:5u] \\
&+ x^{3u} W^o[:4u] + x^{3u} W^o[:6u] + x^{4u} W^o[:5u] + x^{5u} W^o[:4u] \\
&+ x^{4u} W^o[:6u] + x^{5u} W^o[:5u] + x^{6u} W^o[:4u] + x^{6u} W^o[:6u] \\
&+ x^{8u} W^o[:4u] + x^{10u} W^o[:2u] + (x^{6u} + x^{7u} + x^{9u} + x^{10u})R^o.
\end{aligned}$$

Proof. To prove Lemma 2.2 we first construct an automaton for each sequence concerned, and then transform the conditions on infinitely many k 's into finitely many conditions on the states of the automaton. In the following, we will prove that for $T = M_{0,0}^o$, for all integer $k \geq 2$ and $u = 2^{2k}$,

$$\begin{aligned}
T[:12u] &= T[:6u] + x^u T[:5u] + x^{2u} T[:4u] + x^u T[:6u] + x^{2u} T[:5u] + x^{3u} T[:4u] \\
&+ x^{3u} T[:6u] + x^{4u} T[:5u] + x^{5u} T[:4u] + x^{4u} T[:6u] + x^{5u} T[:5u] \\
&+ x^{6u} T[:4u] + x^{6u} T[:6u] + x^{8u} T[:4u] + x^{10u} T[:2u].
\end{aligned}$$

The proof of the other 15 cases are similar. We break down the above identity into 6 parts:

$$\begin{aligned}
0 &= x^{2u}T[4u:5u] + x^u T[5u:6u] + T[6u:7u], \\
0 &= x^{3u}T[4u:5u] + T[7u:8u], \\
0 &= x^{8u}T[:u] + x^{3u}T[5u:6u] + T[8u:9u], \\
0 &= x^{8u}T[u:2u] + x^{5u}T[4u:5u] + x^{4u}T[5u:6u] + T[9u:10u], \\
0 &= x^{10u}T[:u] + x^{8u}T[2u:3u] + x^{6u}T[4u:5u] + T[10u:11u], \\
0 &= x^{10u}T[u:2u] + x^{8u}T[3u:4u] + x^{6u}T[5u:6u] + T[11u:12u],
\end{aligned}$$

which can be rewritten as

$$(2.7) \quad 0 = T[[100w]_2] + T[[101w]_2] + T[[110w]_2],$$

$$(2.8) \quad 0 = T[[100w]_2] + T[[111w]_2],$$

$$(2.9) \quad 0 = T[[w]_2] + T[[101w]_2] + T[[1000w]_2],$$

$$(2.10) \quad 0 = T[[1w]_2] + T[[100w]_2] + T[[101w]_2] + T[[1001w]_2],$$

$$(2.11) \quad 0 = T[[w]_2] + T[[10w]_2] + T[[100w]_2] + T[[1010w]_2],$$

$$(2.12) \quad 0 = T[[1w]_2] + T[[11w]_2] + T[[101w]_2] + T[[1011w]_2],$$

for all binary word w of length $2k$ and $w \neq 0^{2k}$, and

$$(2.13) \quad 0 = T[[100w]_2] + T[[101w]_2] + T[[110w]_2],$$

$$(2.14) \quad 0 = T[[100w]_2] + T[[111w]_2],$$

$$(2.15) \quad 0 = T[[w]_2] + T[[101w]_2] + T[[1000w]_2],$$

$$(2.16) \quad 0 = T[[1w]_2] + T[[100w]_2] + T[[101w]_2] + T[[1001w]_2],$$

$$(2.17) \quad 0 = T[[w]_2] + T[[10w]_2] + T[[100w]_2] + T[[1010w]_2],$$

$$(2.18) \quad 0 = T[[1w]_2] + T[[11w]_2] + T[[101w]_2] + T[[1011w]_2],$$

for $w = 0^{2k}$.

First we calculate an 2-automaton that generates T from its minimal polynomial and its first terms. This automaton has 280 states; its transition function and output function can be found in the annex. Let $A(s, w)$ denote the state reached after reading w from right to left starting from the state s , and τ the output function. Define

$$E_{2k} = \{A(i, w) : |w| = 2k, w \neq 0^{2k}\}.$$

Identities (2.7) through (2.18) can be written as

$$(2.19) \quad 0 = \tau(A(s, 100)) + \tau(A(s, 101)) + \tau(A(s, 110)),$$

$$(2.20) \quad 0 = \tau(A(s, 100)) + \tau(A(s, 111)),$$

$$(2.21) \quad 0 = \tau(A(s, \epsilon)) + \tau(A(s, 101)) + \tau(A(s, 1000)),$$

$$(2.22) \quad 0 = \tau(A(s, 1)) + \tau(A(s, 100)) + \tau(A(s, 101)) + \tau(A(s, 1001)),$$

$$(2.23) \quad 0 = \tau(A(s, \epsilon)) + \tau(A(s, 10)) + \tau(A(s, 100)) + \tau(A(s, 1010)),$$

$$(2.24) \quad 0 = \tau(A(s, 1)) + \tau(A(s, 11)) + \tau(A(s, 101)) + \tau(A(s, 1011)),$$

for all $s \in E_{2k}$, and

$$(2.25) \quad 0 = \tau(A(s, 100)) + \tau(A(s, 101)) + \tau(A(s, 110)),$$

$$(2.26) \quad 0 = \tau(A(s, 100)) + \tau(A(s, 111)),$$

$$(2.27) \quad 0 = \tau(A(s, \epsilon)) + \tau(A(s, 101)) + \tau(A(s, 1000)),$$

$$(2.28) \quad 0 = \tau(A(s, 1)) + \tau(A(s, 100)) + \tau(A(s, 101)) + \tau(A(s, 1001)),$$

$$(2.29) \quad 0 = \tau(A(s, \epsilon)) + \tau(A(s, 10)) + \tau(A(s, 100)) + \tau(A(s, 1010)),$$

$$(2.30) \quad 0 = \tau(A(s, 1)) + \tau(A(s, 11)) + \tau(A(s, 101)) + \tau(A(s, 1011)),$$

for $s = A(i, 0^{2k})$. We find that $(A(i, 0^{18}), E_{18}) = (A(i, 0^{16}), E_{16})$, so that we only have to verify that identities (2.19) through (2.30) hold for $2 \leq k \leq 9$, which turns out to be true. $\square$

In the following lemma, we express M_{2k} , M_{2k+1} , W_{2k} , and W_{2k+1} in terms of M^e , M^o , W^e , and W^o .

Lemma 2.3. *For all integer $k \geq 2$, and $u = 2^{2k-1}$,*

$$M_{2k} = M^e[:6u] + x^u M^e[:5u] + x^{2u} M^e[:4u] + x^{6u} R^e,$$

$$W_{2k} = W^e[:6u] + x^u W^e[:5u] + x^{2u} W^e[:4u] + x^{6u} R^e.$$

For all integer $k \geq 2$, and $u = 2^{2k}$,

$$M_{2k+1} = M^o[:6u] + x^u M^o[:5u] + x^{2u} M^o[:4u] + x^{6u} R^o,$$

$$W_{2k+1} = W^o[:6u] + x^u W^o[:5u] + x^{2u} W^o[:4u] + x^{6u} R^o.$$

Proof. We call the four identities in Lemma 2.3 also by the name M_{2k} , W_{2k} , M_{2k+1} , and W_{2k+1} . For $n = 2$, the identities can be verified directly. For $n \geq 2$, we claim that

$$\begin{aligned} M_{2k} \wedge W_{2k} &\Rightarrow M_{2k+1} \wedge W_{2k+1}, \\ M_{2k+1} \wedge W_{2k+1} &\Rightarrow M_{2k+2} \wedge W_{2k+2}. \end{aligned}$$

We give the proof of

$$M_{2k} \wedge W_{2k} \Rightarrow M_{2k+1},$$

the other ones being similar. Set $u = 2^{2k}$ and $v = 2^{2k-1}$. By definition and induction hypothesis, the left side of identity M_{2k+1} is equal to

$$\begin{aligned} (2.31) \quad W_{2k} M_{2k} &= (W^e[:6v] + x^v W^e[:5v] + x^{2v} W^e[:4v] + x^{6u} R^e) \times (M^e[:6v] \\ &\quad + x^v M^e[:5v] + x^{2v} M^e[:4v] + x^{6u} R^e); \end{aligned}$$

Call this expression *lhs*. Note that both sides of identity M_{2k+1} have the same term of highest degree $x^{12v} R^o$. Therefore we only need to prove that their difference is $O(x^{12v})$. Using Lemma 2.1 it can be seen that the right side of identity M_{2k+1} is congruent, modulo x^{12v} , to

$$W^e[:12v] M^e[:12v] + x^{2v} W^e[:10v] M^e[:10v] + x^{4v} W^e[:8v] M^e[:8v].$$

For all $n \leq 12$, replace the occurrences of $W^e[:n \cdot v]$ and $M^e[:n \cdot v]$ in the above expression by the reduction modulo $x^{n \cdot v}$ of the right side of the corresponding identity in Lemma 2.2 and get a new expression, which we call *rhs*. Define

$$\begin{aligned} X &:= x^v, \\ a_n &:= W^e[n \cdot v : (n+1) \cdot v] / X^n, \\ b_n &:= M^e[n \cdot v : (n+1) \cdot v] / X^n, \\ c &:= R^e. \end{aligned}$$

Using the notation introduced above, we can represent the expressions *lhs* (2.31) and *rhs* as polynomials in $\mathbb{F}_2[a_1, \dots, a_{12}, b_1, \dots, b_{12}, c][X]$. Note that it is not a problem that a_j commutes with b_k while W^e does not commute with M^e , because in the expressions concerned, the products of W^e -terms and M^e -terms are always in the same order. We let the computer do the simplification and check that the difference between these two polynomials is indeed $O(X^{12})$, which completes the proof. $\square$

Proof of Theorem 1.1. We prove the theorem for $\text{CF}(a, b)$; for $\text{CF}(b, a)$, the proof is similar. By Lemma 2.3, we have For all $0 \leq j, k \leq 1$,

$$\begin{aligned} \lim_{n \rightarrow \infty} M_{2n,j,k} &= M_{j,k}^e, \\ \lim_{n \rightarrow \infty} M_{2n+1,j,k} &= M_{j,k}^o, \\ \lim_{n \rightarrow \infty} W_{2n,j,k} &= W_{j,k}^e, \\ \lim_{n \rightarrow \infty} W_{2n+1,j,k} &= W_{j,k}^o. \end{aligned}$$

Let $z = 1/x$. By the convergence theorem and identity (2.2),

$$\text{CF}(a, b) = \frac{M_{0,1}^e(x)}{M_{0,0}^e(x)}.$$

By definition, $\phi(M^e, 0, 1)$ and $\phi(M^e, 0, 0)$ are minimal polynomials of $M_{0,1}^e$ and $M_{0,0}^e$. Therefore

$$P(x, y) = \text{Res}_t (\phi(M^e, 0, 1)(x, t), y^{12}\phi(M^e, 0, 1)(x, t/y))$$

is an annihilating polynomial of $f(x) = M_{0,1}^e/M_{0,0}^e$.

Define

$$Q(x, y) = q_4(x)y^4 + q_3(x)y^3 + q_2(x)y^2 + q_1(x)y + q_0(x),$$

where

$$q_0(x) = x^{18} + x^{15} + x^{11} + x^9 + x^8 + x^7 + x^6 + x^4,$$

$$q_1(x) = x^{14} + x^{13} + x^{10} + x^8 + x^7 + x^6 + x^3 + x,$$

$$q_2(x) = x^{14} + x^{10} + x^6 + x^4 + x^2 + 1,$$

$$q_3(x) = x^{14} + x^{13} + x^{10} + x^8 + x^7 + x^6 + x^3 + x,$$

$$q_4(x) = x^{18} + x^{15} + x^{14} + x^{12} + x^{11} + x^{10} + x^9 + x^7 + x^2.$$

The polynomial $Q(x, y)$ is the candidate for the minimal polynomial of $f(x)$ found by Padé-Hermite approximation. To prove that it is indeed the minimal polynomial of $f(x)$, we only need to prove that it is an irreducible factor of $P(x, y)$ of multiplicity m and $R(x, y) := P(x, y)/Q(x, y)^m$ is not an annihilating polynomial of $f(x)$. We verify the first point directly. For the second point, we find that when we truncate $f(x)$ to order 186, and substitute it for y in $R(z, y)$, we get a series with valuation smaller than 186, which proves that $R(z, y)$ is not an annihilating polynomial of $f(x)$. Finally, $z^{12}Q(1/z, y)$ is the minimal polynomial of $\text{CF}(a, b) = f(1/z)$. $\square$

3. ANNEXE

All 16 polynomials are of the form

$$p_0(x) + p_3(x)y^3 + p_6(x)y^6 + p_9(x)y^9 + p_{12}(x)y^{12}.$$

The coefficients $p_j(x)$ and the 16 initial terms to determine the solutions uniquely are given below.

For $\phi(M^e, 0, 0)$:

$$\begin{aligned} p_0(x) = & x^{108} + x^{102} + x^{100} + x^{94} + x^{88} + x^{84} + x^{82} + x^{80} + x^{78} + x^{74} + x^{72} \\ & + x^{70} + x^{68} + x^{66} + x^{64} + x^{62} + x^{60} + x^{46} + x^{42} + x^{40} + x^{38} + x^{36} \\ & + x^{30} + x^{28} + x^{26} + x^{22} + x^{12}, \end{aligned}$$

$$\begin{aligned} p_3(x) = & x^{92} + x^{90} + x^{84} + x^{82} + x^{80} + x^{78} + x^{72} + x^{68} + x^{66} + x^{60} + x^{58} \\ & + x^{56} + x^{50} + x^{48} + x^{46} + x^{34} + x^{30} + x^{28} + x^{20} + x^{16} + x^{14} + x^8, \end{aligned}$$

$$\begin{aligned} p_6(x) = & x^{90} + x^{82} + x^{76} + x^{72} + x^{68} + x^{66} + x^{62} + x^{60} + x^{52} + x^{50} + x^{48} \\ & + x^{44} + x^{42} + x^{38} + x^{36} + x^{34} + x^{26} + x^{24} + x^{22} + x^{20} + x^{18} + x^{16} \\ & + x^{14} + x^{10} + x^8 + x^4 + x^2 + 1, \end{aligned}$$

$$\begin{aligned} p_9(x) = & x^{82} + x^{80} + x^{70} + x^{66} + x^{62} + x^{60} + x^{58} + x^{54} + x^{50} + x^{46} + x^{44} \\ & + x^{40} + x^{38} + x^{34} + x^{28} + x^{20} + x^{18} + x^{16} + x^{12} + x^{10} + x^8 + x^6 + x^4 \\ & + x^2, \end{aligned}$$

$$p_{12}(x) = x^{80} + x^{72} + x^{64} + x^{48} + x^{40} + x^{24} + x^8 + 1,$$

and the initial terms are $[1, 0, 0, 0, 1, 0, 0, 0, 0, 0, 1, 0, 0, 0, 1, 0]$.

For $\phi(M^e, 0, 1)$:

$$\begin{aligned}
p_0(x) &= x^{96} + x^{95} + x^{93} + x^{92} + x^{89} + x^{88} + x^{87} + x^{86} + x^{85} + x^{84} + x^{83} \\
&\quad + x^{82} + x^{81} + x^{79} + x^{76} + x^{73} + x^{72} + x^{71} + x^{70} + x^{65} + x^{63} + x^{62} \\
&\quad + x^{60} + x^{58} + x^{57} + x^{56} + x^{55} + x^{54} + x^{53} + x^{51} + x^{50} + x^{48} + x^{47} \\
&\quad + x^{46} + x^{42} + x^{41} + x^{39} + x^{36} + x^{35} + x^{33} + x^{31} + x^{30} + x^{29} + x^{28} \\
&\quad + x^{27} + x^{24} + x^{22} + x^{21} + x^{20} + x^{19} + x^{18}, \\
p_3(x) &= x^{74} + x^{72} + x^{71} + x^{70} + x^{68} + x^{67} + x^{64} + x^{62} + x^{61} + x^{60} + x^{58} \\
&\quad + x^{57} + x^{56} + x^{54} + x^{53} + x^{52} + x^{50} + x^{49} + x^{48} + x^{46} + x^{45} + x^{44} \\
&\quad + x^{42} + x^{41} + x^{40} + x^{38} + x^{37} + x^{36} + x^{34} + x^{33} + x^{32} + x^{30} + x^{29} \\
&\quad + x^{28} + x^{26} + x^{25} + x^{24} + x^{22} + x^{21} + x^{20} + x^{17} + x^{15} + x^{13} + x^{11} \\
&\quad + x^{10} + x^9, \\
p_6(x) &= x^{72} + x^{71} + x^{70} + x^{68} + x^{67} + x^{66} + x^{64} + x^{63} + x^{61} + x^{60} + x^{54} \\
&\quad + x^{53} + x^{51} + x^{50} + x^{48} + x^{47} + x^{46} + x^{40} + x^{39} + x^{37} + x^{36} + x^{34} \\
&\quad + x^{33} + x^{32} + x^{26} + x^{25} + x^{23} + x^{22} + x^{20} + x^{19} + x^{18} + x^{16} + x^{15} \\
&\quad + x^{14} + x^{10} + x^9 + x^7 + x^6, \\
p_9(x) &= x^{70} + x^{67} + x^{65} + x^{63} + x^{61} + x^{59} + x^{57} + x^{56} + x^{55} + x^{50} + x^{47} \\
&\quad + x^{46} + x^{45} + x^{36} + x^{34} + x^{33} + x^{32} + x^{30} + x^{29} + x^{26} + x^{23} + x^{22} \\
&\quad + x^{21} + x^{20} + x^{18} + x^{17} + x^{16} + x^{13} + x^{12} + x^{11} + x^8 + x^5 + x^4 + x^3, \\
p_{12}(x) &= x^{68} + x^{67} + x^{65} + x^{63} + x^{61} + x^{60} + x^{56} + x^{55} + x^{53} + x^{52} + x^{48} \\
&\quad + x^{47} + x^{45} + x^{44} + x^{40} + x^{39} + x^{37} + x^{35} + x^{33} + x^{32} + x^{20} + x^{19} \\
&\quad + x^{17} + x^{15} + x^{13} + x^{12} + x^8 + x^7 + x^5 + x^3 + x + 1,
\end{aligned}$$

and the initial terms are $[0, 1, 0, 0, 0, 1, 0, 1, 1, 1, 0, 0, 1, 1, 1, 1]$.

For $\phi(M^e, 1, 0)$:

$$\begin{aligned}
p_0(x) &= x^{96} + x^{95} + x^{93} + x^{92} + x^{89} + x^{88} + x^{87} + x^{86} + x^{85} + x^{84} + x^{83} \\
&\quad + x^{82} + x^{81} + x^{79} + x^{76} + x^{73} + x^{72} + x^{71} + x^{70} + x^{65} + x^{63} + x^{62} \\
&\quad + x^{60} + x^{58} + x^{57} + x^{56} + x^{55} + x^{54} + x^{53} + x^{51} + x^{50} + x^{48} + x^{47} \\
&\quad + x^{46} + x^{42} + x^{41} + x^{39} + x^{36} + x^{35} + x^{33} + x^{31} + x^{30} + x^{29} + x^{28} \\
&\quad + x^{27} + x^{24} + x^{22} + x^{21} + x^{20} + x^{19} + x^{18}, \\
p_3(x) &= x^{74} + x^{72} + x^{71} + x^{70} + x^{68} + x^{67} + x^{64} + x^{62} + x^{61} + x^{60} + x^{58} \\
&\quad + x^{57} + x^{56} + x^{54} + x^{53} + x^{52} + x^{50} + x^{49} + x^{48} + x^{46} + x^{45} + x^{44} \\
&\quad + x^{42} + x^{41} + x^{40} + x^{38} + x^{37} + x^{36} + x^{34} + x^{33} + x^{32} + x^{30} + x^{29} \\
&\quad + x^{28} + x^{26} + x^{25} + x^{24} + x^{22} + x^{21} + x^{20} + x^{17} + x^{15} + x^{13} + x^{11} \\
&\quad + x^{10} + x^9, \\
p_6(x) &= x^{72} + x^{71} + x^{70} + x^{68} + x^{67} + x^{66} + x^{64} + x^{63} + x^{61} + x^{60} + x^{54}
\end{aligned}$$

$$\begin{aligned}
& + x^{53} + x^{51} + x^{50} + x^{48} + x^{47} + x^{46} + x^{40} + x^{39} + x^{37} + x^{36} + x^{34} \\
& + x^{33} + x^{32} + x^{26} + x^{25} + x^{23} + x^{22} + x^{20} + x^{19} + x^{18} + x^{16} + x^{15} \\
& + x^{14} + x^{10} + x^9 + x^7 + x^6, \\
p_9(x) &= x^{70} + x^{67} + x^{65} + x^{63} + x^{61} + x^{59} + x^{57} + x^{56} + x^{55} + x^{50} + x^{47} \\
& + x^{46} + x^{45} + x^{36} + x^{34} + x^{33} + x^{32} + x^{30} + x^{29} + x^{26} + x^{23} + x^{22} \\
& + x^{21} + x^{20} + x^{18} + x^{17} + x^{16} + x^{13} + x^{12} + x^{11} + x^8 + x^5 + x^4 + x^3, \\
p_{12}(x) &= x^{68} + x^{67} + x^{65} + x^{63} + x^{61} + x^{60} + x^{56} + x^{55} + x^{53} + x^{52} + x^{48} \\
& + x^{47} + x^{45} + x^{44} + x^{40} + x^{39} + x^{37} + x^{35} + x^{33} + x^{32} + x^{20} + x^{19} \\
& + x^{17} + x^{15} + x^{13} + x^{12} + x^8 + x^7 + x^5 + x^3 + x + 1,
\end{aligned}$$

and the initial terms are $[0, 1, 0, 0, 0, 1, 0, 1, 1, 1, 0, 0, 1, 1, 1, 1]$.

For $\phi(M^e, 1, 1)$:

$$\begin{aligned}
p_0(x) &= x^{84} + x^{82} + x^{76} + x^{74} + x^{70} + x^{64} + x^{60} + x^{54} + x^{52} + x^{48} + x^{44} \\
& + x^{42} + x^{40} + x^{34} + x^{32} + x^{30} + x^{28} + x^{26} + x^{24}, \\
p_3(x) &= x^{68} + x^{64} + x^{62} + x^{60} + x^{58} + x^{56} + x^{54} + x^{44} + x^{40} + x^{26} + x^{22} \\
& + x^{18} + x^{14} + x^6, \\
p_6(x) &= x^{66} + x^{64} + x^{60} + x^{56} + x^{54} + x^{52} + x^{50} + x^{44} + x^{42} + x^{32} + x^{28} \\
& + x^{26} + x^{24} + x^{22} + x^{14} + x^{10} + x^4 + 1, \\
p_9(x) &= x^{58} + x^{54} + x^{52} + x^{50} + x^{48} + x^{40} + x^{36} + x^{32} + x^{28} + x^{22} + x^{20} \\
& + x^{18} + x^{14} + x^{10} + x^8 + x^2, \\
p_{12}(x) &= x^{56} + x^{54} + x^{50} + x^{48} + x^{40} + x^{38} + x^{34} + x^{32} + x^8 + x^6 + x^2 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 1, 0, 0, 0, 1, 0, 0, 0, 0, 0, 1, 0, 1, 0]$.

For $\phi(W^e, 0, 0)$:

$$\begin{aligned}
p_0(x) &= x^{84} + x^{82} + x^{80} + x^{78} + x^{76} + x^{70} + x^{68} + x^{66} + x^{64} + x^{60} + x^{56} \\
& + x^{52} + x^{48} + x^{46} + x^{32} + x^{28} + x^{24} + x^{20} + x^{16} + x^{14} + x^{12}, \\
p_3(x) &= x^{68} + x^{62} + x^{60} + x^{48} + x^{42} + x^{34} + x^{30} + x^{28} + x^{22} + x^{20} + x^{18} \\
& + x^{16} + x^{12} + x^8, \\
p_6(x) &= x^{66} + x^{64} + x^{62} + x^{58} + x^{56} + x^{54} + x^{50} + x^{46} + x^{42} + x^{40} + x^{38} \\
& + x^{36} + x^{34} + x^{32} + x^{30} + x^{28} + x^{26} + x^{22} + x^{18} + x^{16} + x^{14} + x^{12} \\
& + x^8 + 1, \\
p_9(x) &= x^{58} + x^{54} + x^{52} + x^{50} + x^{48} + x^{40} + x^{36} + x^{32} + x^{28} + x^{22} + x^{20} \\
& + x^{18} + x^{14} + x^{10} + x^8 + x^2, \\
p_{12}(x) &= x^{56} + x^{54} + x^{50} + x^{48} + x^{40} + x^{38} + x^{34} + x^{32} + x^8 + x^6 + x^2 + 1,
\end{aligned}$$

and the initial terms are $[1, 0, 1, 0, 1, 0, 1, 0, 0, 0, 0, 0, 1, 0, 1, 0]$.

For $\phi(W^e, 0, 1)$:

$$\begin{aligned}
p_0(x) &= x^{96} + x^{95} + x^{93} + x^{91} + x^{90} + x^{89} + x^{88} + x^{87} + x^{83} + x^{82} + x^{79} \\
& + x^{78} + x^{77} + x^{76} + x^{75} + x^{74} + x^{73} + x^{71} + x^{70} + x^{66} + x^{65} + x^{64}
\end{aligned}$$

$$\begin{aligned}
& + x^{63} + x^{62} + x^{60} + x^{58} + x^{56} + x^{54} + x^{53} + x^{48} + x^{46} + x^{45} + x^{40} \\
& + x^{39} + x^{38} + x^{36} + x^{34} + x^{33} + x^{32} + x^{30} + x^{29} + x^{26} + x^{24}, \\
p_3(x) &= x^{74} + x^{72} + x^{71} + x^{70} + x^{68} + x^{67} + x^{64} + x^{62} + x^{61} + x^{60} + x^{58} \\
& + x^{57} + x^{56} + x^{54} + x^{53} + x^{52} + x^{50} + x^{49} + x^{48} + x^{46} + x^{45} + x^{44} \\
& + x^{42} + x^{41} + x^{40} + x^{38} + x^{37} + x^{36} + x^{34} + x^{33} + x^{32} + x^{30} + x^{29} \\
& + x^{28} + x^{26} + x^{25} + x^{24} + x^{22} + x^{21} + x^{20} + x^{17} + x^{15} + x^{13} + x^{11} \\
& + x^{10} + x^9, \\
p_6(x) &= x^{72} + x^{71} + x^{70} + x^{68} + x^{67} + x^{66} + x^{64} + x^{63} + x^{61} + x^{60} + x^{54} \\
& + x^{53} + x^{51} + x^{50} + x^{48} + x^{47} + x^{46} + x^{40} + x^{39} + x^{37} + x^{36} + x^{34} \\
& + x^{33} + x^{32} + x^{26} + x^{25} + x^{23} + x^{22} + x^{20} + x^{19} + x^{18} + x^{16} + x^{15} \\
& + x^{14} + x^{10} + x^9 + x^7 + x^6, \\
p_9(x) &= x^{70} + x^{67} + x^{65} + x^{63} + x^{61} + x^{59} + x^{57} + x^{56} + x^{55} + x^{50} + x^{47} \\
& + x^{46} + x^{45} + x^{36} + x^{34} + x^{33} + x^{32} + x^{30} + x^{29} + x^{26} + x^{23} + x^{22} \\
& + x^{21} + x^{20} + x^{18} + x^{17} + x^{16} + x^{13} + x^{12} + x^{11} + x^8 + x^5 + x^4 + x^3, \\
p_{12}(x) &= x^{68} + x^{67} + x^{65} + x^{63} + x^{61} + x^{60} + x^{56} + x^{55} + x^{53} + x^{52} + x^{48} \\
& + x^{47} + x^{45} + x^{44} + x^{40} + x^{39} + x^{37} + x^{35} + x^{33} + x^{32} + x^{20} + x^{19} \\
& + x^{17} + x^{15} + x^{13} + x^{12} + x^8 + x^7 + x^5 + x^3 + x + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 1, 1, 0, 1, 0, 1, 0, 0, 0, 1, 0]$.

For $\phi(W^e, 1, 0)$:

$$\begin{aligned}
p_0(x) &= x^{96} + x^{95} + x^{93} + x^{91} + x^{90} + x^{89} + x^{88} + x^{87} + x^{83} + x^{82} + x^{79} \\
& + x^{78} + x^{77} + x^{76} + x^{75} + x^{74} + x^{73} + x^{71} + x^{70} + x^{66} + x^{65} + x^{64} \\
& + x^{63} + x^{62} + x^{60} + x^{58} + x^{56} + x^{54} + x^{53} + x^{48} + x^{46} + x^{45} + x^{40} \\
& + x^{39} + x^{38} + x^{36} + x^{34} + x^{33} + x^{32} + x^{30} + x^{29} + x^{26} + x^{24}, \\
p_3(x) &= x^{74} + x^{72} + x^{71} + x^{70} + x^{68} + x^{67} + x^{64} + x^{62} + x^{61} + x^{60} + x^{58} \\
& + x^{57} + x^{56} + x^{54} + x^{53} + x^{52} + x^{50} + x^{49} + x^{48} + x^{46} + x^{45} + x^{44} \\
& + x^{42} + x^{41} + x^{40} + x^{38} + x^{37} + x^{36} + x^{34} + x^{33} + x^{32} + x^{30} + x^{29} \\
& + x^{28} + x^{26} + x^{25} + x^{24} + x^{22} + x^{21} + x^{20} + x^{17} + x^{15} + x^{13} + x^{11} \\
& + x^{10} + x^9, \\
p_6(x) &= x^{72} + x^{71} + x^{70} + x^{68} + x^{67} + x^{66} + x^{64} + x^{63} + x^{61} + x^{60} + x^{54} \\
& + x^{53} + x^{51} + x^{50} + x^{48} + x^{47} + x^{46} + x^{40} + x^{39} + x^{37} + x^{36} + x^{34} \\
& + x^{33} + x^{32} + x^{26} + x^{25} + x^{23} + x^{22} + x^{20} + x^{19} + x^{18} + x^{16} + x^{15} \\
& + x^{14} + x^{10} + x^9 + x^7 + x^6, \\
p_9(x) &= x^{70} + x^{67} + x^{65} + x^{63} + x^{61} + x^{59} + x^{57} + x^{56} + x^{55} + x^{50} + x^{47} \\
& + x^{46} + x^{45} + x^{36} + x^{34} + x^{33} + x^{32} + x^{30} + x^{29} + x^{26} + x^{23} + x^{22} \\
& + x^{21} + x^{20} + x^{18} + x^{17} + x^{16} + x^{13} + x^{12} + x^{11} + x^8 + x^5 + x^4 + x^3,
\end{aligned}$$

$$\begin{aligned}
p_{12}(x) = & x^{68} + x^{67} + x^{65} + x^{63} + x^{61} + x^{60} + x^{56} + x^{55} + x^{53} + x^{52} + x^{48} \\
& + x^{47} + x^{45} + x^{44} + x^{40} + x^{39} + x^{37} + x^{35} + x^{33} + x^{32} + x^{20} + x^{19} \\
& + x^{17} + x^{15} + x^{13} + x^{12} + x^8 + x^7 + x^5 + x^3 + x + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 1, 1, 0, 1, 0, 1, 0, 0, 0, 1, 0]$.

For $\phi(W^e, 1, 1)$:

$$\begin{aligned}
p_0(x) = & x^{108} + x^{104} + x^{102} + x^{100} + x^{92} + x^{84} + x^{80} + x^{68} + x^{64} + x^{62} + x^{58} \\
& + x^{48} + x^{40} + x^{38} + x^{36}, \\
p_3(x) = & x^{92} + x^{90} + x^{88} + x^{86} + x^{78} + x^{74} + x^{72} + x^{70} + x^{64} + x^{62} + x^{60} \\
& + x^{56} + x^{54} + x^{52} + x^{48} + x^{46} + x^{44} + x^{40} + x^{38} + x^{36} + x^{34} + x^{32} \\
& + x^{30} + x^{22} + x^{20} + x^{18} + x^8 + x^6, \\
p_6(x) = & x^{90} + x^{86} + x^{74} + x^{70} + x^{60} + x^{58} + x^{56} + x^{54} + x^{48} + x^{44} + x^{42} \\
& + x^{40} + x^{36} + x^{34} + x^{32} + x^{28} + x^{26} + x^{10} + x^8 + x^6 + x^2 + 1, \\
p_9(x) = & x^{82} + x^{80} + x^{70} + x^{66} + x^{62} + x^{60} + x^{58} + x^{54} + x^{50} + x^{46} + x^{44} \\
& + x^{40} + x^{38} + x^{34} + x^{28} + x^{20} + x^{18} + x^{16} + x^{12} + x^{10} + x^8 + x^6 + x^4 \\
& + x^2, \\
p_{12}(x) = & x^{80} + x^{72} + x^{64} + x^{48} + x^{40} + x^{24} + x^8 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 1, 0, 0, 0, 1, 0]$.

For $\phi(M^o, 0, 0)$:

$$\begin{aligned}
p_0(x) = & x^{96} + x^{95} + x^{94} + x^{92} + x^{88} + x^{84} + x^{83} + x^{82} + x^{80} + x^{79} + x^{78} \\
& + x^{77} + x^{71} + x^{70} + x^{68} + x^{67} + x^{66} + x^{65} + x^{63} + x^{61} + x^{57} + x^{55} \\
& + x^{51} + x^{50} + x^{46} + x^{45} + x^{44} + x^{41} + x^{37} + x^{33} + x^{32} + x^{30} + x^{25} \\
& + x^{23} + x^{22} + x^{20} + x^{17} + x^{16} + x^{14} + x^{13} + x^{12}, \\
p_3(x) = & x^{80} + x^{79} + x^{78} + x^{75} + x^{73} + x^{70} + x^{69} + x^{67} + x^{66} + x^{65} + x^{63} \\
& + x^{58} + x^{56} + x^{55} + x^{54} + x^{53} + x^{51} + x^{46} + x^{41} + x^{40} + x^{37} + x^{36} \\
& + x^{33} + x^{31} + x^{30} + x^{29} + x^{28} + x^{26} + x^{24} + x^{23} + x^{22} + x^{21} + x^{18} \\
& + x^{15} + x^{14} + x^{13} + x^{12} + x^{10} + x^9 + x^8, \\
p_6(x) = & x^{76} + x^{75} + x^{74} + x^{72} + x^{70} + x^{69} + x^{67} + x^{65} + x^{64} + x^{62} + x^{60} \\
& + x^{59} + x^{57} + x^{56} + x^{55} + x^{54} + x^{53} + x^{52} + x^{47} + x^{45} + x^{44} + x^{43} \\
& + x^{42} + x^{40} + x^{37} + x^{35} + x^{34} + x^{33} + x^{32} + x^{28} + x^{25} + x^{23} + x^{22} \\
& + x^{20} + x^{11} + x^8 + x^7 + x^6 + x^4 + x^3 + x + 1, \\
p_9(x) = & x^{68} + x^{67} + x^{65} + x^{64} + x^{60} + x^{59} + x^{57} + x^{55} + x^{53} + x^{51} + x^{49} \\
& + x^{48} + x^{44} + x^{43} + x^{41} + x^{39} + x^{37} + x^{36} + x^{20} + x^{19} + x^{17} + x^{16} \\
& + x^{12} + x^{11} + x^9 + x^7 + x^5 + x^4, \\
p_{12}(x) = & x^{64} + x^{63} + x^{61} + x^{59} + x^{57} + x^{56} + x^{52} + x^{51} + x^{49} + x^{47} + x^{45} \\
& + x^{43} + x^{41} + x^{40} + x^{36} + x^{35} + x^{33} + x^{32} + x^{16} + x^{15} + x^{13} + x^{11}
\end{aligned}$$

$$+ x^9 + x^8 + x^4 + x^3 + x + 1,$$

and the initial terms are $[1, 0, 1, 0, 0, 0, 1, 1, 1, 1, 1, 0, 1, 1, 1, 0]$.

For $\phi(M^o, 0, 1)$:

$$\begin{aligned} p_0(x) = & x^{92} + x^{90} + x^{88} + x^{87} + x^{85} + x^{81} + x^{76} + x^{75} + x^{74} + x^{71} + x^{70} \\ & + x^{69} + x^{65} + x^{64} + x^{59} + x^{58} + x^{57} + x^{55} + x^{54} + x^{52} + x^{51} + x^{48} \\ & + x^{47} + x^{44} + x^{42} + x^{41} + x^{38} + x^{37} + x^{36} + x^{35} + x^{30} + x^{29} + x^{28} \\ & + x^{27} + x^{25} + x^{24} + x^{23} + x^{22} + x^{21} + x^{18}, \end{aligned}$$

$$\begin{aligned} p_3(x) = & x^{72} + x^{71} + x^{68} + x^{66} + x^{65} + x^{64} + x^{62} + x^{60} + x^{58} + x^{56} + x^{54} \\ & + x^{52} + x^{50} + x^{48} + x^{46} + x^{44} + x^{42} + x^{40} + x^{38} + x^{36} + x^{34} + x^{32} \\ & + x^{30} + x^{28} + x^{26} + x^{24} + x^{22} + x^{20} + x^{18} + x^{15} + x^{14} + x^9, \end{aligned}$$

$$\begin{aligned} p_6(x) = & x^{68} + x^{66} + x^{60} + x^{50} + x^{44} + x^{42} + x^{40} + x^{38} + x^{34} + x^{32} + x^{22} \\ & + x^{16} + x^{14} + x^6, \end{aligned}$$

$$\begin{aligned} p_9(x) = & x^{64} + x^{63} + x^{62} + x^{61} + x^{60} + x^{57} + x^{56} + x^{51} + x^{48} + x^{47} + x^{46} \\ & + x^{45} + x^{44} + x^{41} + x^{40} + x^{35} + x^{16} + x^{15} + x^{14} + x^{13} + x^{12} + x^9 \\ & + x^8 + x^3, \end{aligned}$$

$$p_{12}(x) = x^{60} + x^{56} + x^{48} + x^{44} + x^{40} + x^{32} + x^{12} + x^8 + 1,$$

and the initial terms are $[0, 1, 0, 1, 0, 0, 0, 0, 0, 1, 0, 0, 0, 0, 0, 0]$.

For $\phi(M^o, 1, 0)$:

$$\begin{aligned} p_0(x) = & x^{108} + x^{106} + x^{103} + x^{101} + x^{99} + x^{93} + x^{91} + x^{90} + x^{88} + x^{86} + x^{85} \\ & + x^{84} + x^{83} + x^{81} + x^{79} + x^{78} + x^{75} + x^{70} + x^{69} + x^{67} + x^{66} + x^{64} \\ & + x^{63} + x^{62} + x^{61} + x^{60} + x^{59} + x^{58} + x^{57} + x^{56} + x^{54} + x^{53} + x^{49} \\ & + x^{46} + x^{44} + x^{38} + x^{36} + x^{35} + x^{33} + x^{29} + x^{26} + x^{25} + x^{24}, \end{aligned}$$

$$\begin{aligned} p_3(x) = & x^{88} + x^{87} + x^{83} + x^{82} + x^{81} + x^{77} + x^{76} + x^{75} + x^{71} + x^{70} + x^{69} \\ & + x^{65} + x^{32} + x^{31} + x^{27} + x^{26} + x^{25} + x^{21} + x^{20} + x^{19} + x^{15} + x^{14} \\ & + x^{13} + x^9, \end{aligned}$$

$$\begin{aligned} p_6(x) = & x^{84} + x^{82} + x^{80} + x^{78} + x^{76} + x^{70} + x^{68} + x^{64} + x^{62} + x^{58} + x^{54} \\ & + x^{52} + x^{50} + x^{44} + x^{40} + x^{38} + x^{36} + x^{30} + x^{28} + x^{20} + x^{16} + x^{14} \\ & + x^{10} + x^6, \end{aligned}$$

$$\begin{aligned} p_9(x) = & x^{80} + x^{79} + x^{78} + x^{77} + x^{75} + x^{74} + x^{69} + x^{66} + x^{65} + x^{64} + x^{63} \\ & + x^{61} + x^{59} + x^{58} + x^{57} + x^{56} + x^{55} + x^{53} + x^{51} + x^{50} + x^{49} + x^{46} \\ & + x^{41} + x^{40} + x^{39} + x^{35} + x^{32} + x^{31} + x^{30} + x^{29} + x^{27} + x^{26} + x^{21} \\ & + x^{18} + x^{17} + x^{14} + x^9 + x^8 + x^7 + x^3, \end{aligned}$$

$$\begin{aligned} p_{12}(x) = & x^{76} + x^{68} + x^{56} + x^{48} + x^{44} + x^{40} + x^{36} + x^{32} + x^{28} + x^{20} + x^{12} + x^8 \\ & + x^4 + 1, \end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 1, 1, 0, 1, 1, 0, 1, 1, 0, 0, 1]$.

For $\phi(M^o, 1, 1)$:

$$\begin{aligned}
p_0(x) &= x^{96} + x^{95} + x^{94} + x^{92} + x^{90} + x^{89} + x^{88} + x^{87} + x^{85} + x^{83} + x^{82} \\
&\quad + x^{80} + x^{78} + x^{76} + x^{74} + x^{73} + x^{71} + x^{66} + x^{63} + x^{62} + x^{60} + x^{57} \\
&\quad + x^{55} + x^{53} + x^{51} + x^{50} + x^{48} + x^{46} + x^{43} + x^{42} + x^{39} + x^{36} + x^{35} \\
&\quad + x^{34} + x^{33} + x^{32} + x^{30}, \\
p_3(x) &= x^{80} + x^{79} + x^{78} + x^{75} + x^{74} + x^{71} + x^{70} + x^{65} + x^{64} + x^{61} + x^{59} \\
&\quad + x^{58} + x^{57} + x^{56} + x^{54} + x^{49} + x^{47} + x^{46} + x^{45} + x^{43} + x^{40} + x^{39} \\
&\quad + x^{36} + x^{35} + x^{30} + x^{28} + x^{27} + x^{26} + x^{25} + x^{24} + x^{22} + x^{19} + x^{15} \\
&\quad + x^{14} + x^{13} + x^{12}, \\
p_6(x) &= x^{76} + x^{75} + x^{74} + x^{72} + x^{66} + x^{65} + x^{64} + x^{62} + x^{55} + x^{50} + x^{49} \\
&\quad + x^{48} + x^{47} + x^{46} + x^{44} + x^{42} + x^{41} + x^{40} + x^{39} + x^{36} + x^{35} + x^{32} \\
&\quad + x^{31} + x^{29} + x^{26} + x^{24} + x^{23} + x^{21} + x^{18} + x^{17} + x^{15} + x^{14} + x^{11} \\
&\quad + x^{10} + x^9 + x^7 + x^5 + x^3 + x + 1, \\
p_9(x) &= x^{68} + x^{67} + x^{65} + x^{64} + x^{60} + x^{59} + x^{57} + x^{55} + x^{53} + x^{51} + x^{49} \\
&\quad + x^{48} + x^{44} + x^{43} + x^{41} + x^{39} + x^{37} + x^{36} + x^{20} + x^{19} + x^{17} + x^{16} \\
&\quad + x^{12} + x^{11} + x^9 + x^7 + x^5 + x^4, \\
p_{12}(x) &= x^{64} + x^{63} + x^{61} + x^{59} + x^{57} + x^{56} + x^{52} + x^{51} + x^{49} + x^{47} + x^{45} \\
&\quad + x^{43} + x^{41} + x^{40} + x^{36} + x^{35} + x^{33} + x^{32} + x^{16} + x^{15} + x^{13} + x^{11} \\
&\quad + x^9 + x^8 + x^4 + x^3 + x + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 0, 0, 1, 1, 0, 1, 1, 0, 0, 1, 0, 0]$.

For $\phi(W^o, 0, 0)$:

$$\begin{aligned}
p_0(x) &= x^{96} + x^{95} + x^{94} + x^{92} + x^{88} + x^{84} + x^{83} + x^{82} + x^{80} + x^{79} + x^{78} \\
&\quad + x^{77} + x^{71} + x^{70} + x^{68} + x^{67} + x^{66} + x^{65} + x^{63} + x^{61} + x^{57} + x^{55} \\
&\quad + x^{51} + x^{50} + x^{46} + x^{45} + x^{44} + x^{41} + x^{37} + x^{33} + x^{32} + x^{30} + x^{25} \\
&\quad + x^{23} + x^{22} + x^{20} + x^{17} + x^{16} + x^{14} + x^{13} + x^{12}, \\
p_3(x) &= x^{80} + x^{79} + x^{78} + x^{75} + x^{73} + x^{70} + x^{69} + x^{67} + x^{66} + x^{65} + x^{63} \\
&\quad + x^{58} + x^{56} + x^{55} + x^{54} + x^{53} + x^{51} + x^{46} + x^{41} + x^{40} + x^{37} + x^{36} \\
&\quad + x^{33} + x^{31} + x^{30} + x^{29} + x^{28} + x^{26} + x^{24} + x^{23} + x^{22} + x^{21} + x^{18} \\
&\quad + x^{15} + x^{14} + x^{13} + x^{12} + x^{10} + x^9 + x^8, \\
p_6(x) &= x^{76} + x^{75} + x^{74} + x^{72} + x^{70} + x^{69} + x^{67} + x^{65} + x^{64} + x^{62} + x^{60} \\
&\quad + x^{59} + x^{57} + x^{56} + x^{55} + x^{54} + x^{53} + x^{52} + x^{47} + x^{45} + x^{44} + x^{43} \\
&\quad + x^{42} + x^{40} + x^{37} + x^{35} + x^{34} + x^{33} + x^{32} + x^{28} + x^{25} + x^{23} + x^{22} \\
&\quad + x^{20} + x^{11} + x^8 + x^7 + x^6 + x^4 + x^3 + x + 1, \\
p_9(x) &= x^{68} + x^{67} + x^{65} + x^{64} + x^{60} + x^{59} + x^{57} + x^{55} + x^{53} + x^{51} + x^{49} \\
&\quad + x^{48} + x^{44} + x^{43} + x^{41} + x^{39} + x^{37} + x^{36} + x^{20} + x^{19} + x^{17} + x^{16}
\end{aligned}$$

$$\begin{aligned}
& + x^{12} + x^{11} + x^9 + x^7 + x^5 + x^4, \\
p_{12}(x) = & x^{64} + x^{63} + x^{61} + x^{59} + x^{57} + x^{56} + x^{52} + x^{51} + x^{49} + x^{47} + x^{45} \\
& + x^{43} + x^{41} + x^{40} + x^{36} + x^{35} + x^{33} + x^{32} + x^{16} + x^{15} + x^{13} + x^{11} \\
& + x^9 + x^8 + x^4 + x^3 + x + 1,
\end{aligned}$$

and the initial terms are $[1, 0, 1, 0, 0, 0, 1, 1, 1, 1, 1, 0, 1, 1, 1, 0]$.

For $\phi(W^o, 0, 1)$:

$$\begin{aligned}
p_0(x) = & x^{108} + x^{106} + x^{103} + x^{101} + x^{99} + x^{93} + x^{91} + x^{90} + x^{88} + x^{86} + x^{85} \\
& + x^{84} + x^{83} + x^{81} + x^{79} + x^{78} + x^{75} + x^{70} + x^{69} + x^{67} + x^{66} + x^{64} \\
& + x^{63} + x^{62} + x^{61} + x^{60} + x^{59} + x^{58} + x^{57} + x^{56} + x^{54} + x^{53} + x^{49} \\
& + x^{46} + x^{44} + x^{38} + x^{36} + x^{35} + x^{33} + x^{29} + x^{26} + x^{25} + x^{24}, \\
p_3(x) = & x^{88} + x^{87} + x^{83} + x^{82} + x^{81} + x^{77} + x^{76} + x^{75} + x^{71} + x^{70} + x^{69} \\
& + x^{65} + x^{32} + x^{31} + x^{27} + x^{26} + x^{25} + x^{21} + x^{20} + x^{19} + x^{15} + x^{14} \\
& + x^{13} + x^9, \\
p_6(x) = & x^{84} + x^{82} + x^{80} + x^{78} + x^{76} + x^{70} + x^{68} + x^{64} + x^{62} + x^{58} + x^{54} \\
& + x^{52} + x^{50} + x^{44} + x^{40} + x^{38} + x^{36} + x^{30} + x^{28} + x^{20} + x^{16} + x^{14} \\
& + x^{10} + x^6, \\
p_9(x) = & x^{80} + x^{79} + x^{78} + x^{77} + x^{75} + x^{74} + x^{69} + x^{66} + x^{65} + x^{64} + x^{63} \\
& + x^{61} + x^{59} + x^{58} + x^{57} + x^{56} + x^{55} + x^{53} + x^{51} + x^{50} + x^{49} + x^{46} \\
& + x^{41} + x^{40} + x^{39} + x^{35} + x^{32} + x^{31} + x^{30} + x^{29} + x^{27} + x^{26} + x^{21} \\
& + x^{18} + x^{17} + x^{14} + x^9 + x^8 + x^7 + x^3, \\
p_{12}(x) = & x^{76} + x^{68} + x^{56} + x^{48} + x^{44} + x^{40} + x^{36} + x^{32} + x^{28} + x^{20} + x^{12} + x^8 \\
& + x^4 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 1, 1, 0, 1, 1, 0, 1, 1, 0, 0, 1]$.

For $\phi(W^o, 1, 0)$:

$$\begin{aligned}
p_0(x) = & x^{92} + x^{90} + x^{88} + x^{87} + x^{85} + x^{81} + x^{76} + x^{75} + x^{74} + x^{71} + x^{70} \\
& + x^{69} + x^{65} + x^{64} + x^{59} + x^{58} + x^{57} + x^{55} + x^{54} + x^{52} + x^{51} + x^{48} \\
& + x^{47} + x^{44} + x^{42} + x^{41} + x^{38} + x^{37} + x^{36} + x^{35} + x^{30} + x^{29} + x^{28} \\
& + x^{27} + x^{25} + x^{24} + x^{23} + x^{22} + x^{21} + x^{18}, \\
p_3(x) = & x^{72} + x^{71} + x^{68} + x^{66} + x^{65} + x^{64} + x^{62} + x^{60} + x^{58} + x^{56} + x^{54} \\
& + x^{52} + x^{50} + x^{48} + x^{46} + x^{44} + x^{42} + x^{40} + x^{38} + x^{36} + x^{34} + x^{32} \\
& + x^{30} + x^{28} + x^{26} + x^{24} + x^{22} + x^{20} + x^{18} + x^{15} + x^{14} + x^9, \\
p_6(x) = & x^{68} + x^{66} + x^{60} + x^{50} + x^{44} + x^{42} + x^{40} + x^{38} + x^{34} + x^{32} + x^{22} \\
& + x^{16} + x^{14} + x^6, \\
p_9(x) = & x^{64} + x^{63} + x^{62} + x^{61} + x^{60} + x^{57} + x^{56} + x^{51} + x^{48} + x^{47} + x^{46} \\
& + x^{45} + x^{44} + x^{41} + x^{40} + x^{35} + x^{16} + x^{15} + x^{14} + x^{13} + x^{12} + x^9
\end{aligned}$$

$$+ x^8 + x^3,$$

$$p_{12}(x) = x^{60} + x^{56} + x^{48} + x^{44} + x^{40} + x^{32} + x^{12} + x^8 + 1,$$

and the initial terms are $[0, 1, 0, 1, 0, 0, 0, 0, 0, 1, 0, 0, 0, 0, 0, 0]$.

For $\phi(W^o, 1, 1)$:

$$p_0(x) = x^{96} + x^{95} + x^{94} + x^{92} + x^{90} + x^{89} + x^{88} + x^{87} + x^{85} + x^{83} + x^{82}$$

$$+ x^{80} + x^{78} + x^{76} + x^{74} + x^{73} + x^{71} + x^{66} + x^{63} + x^{62} + x^{60} + x^{57}$$

$$+ x^{55} + x^{53} + x^{51} + x^{50} + x^{48} + x^{46} + x^{43} + x^{42} + x^{39} + x^{36} + x^{35}$$

$$+ x^{34} + x^{33} + x^{32} + x^{30},$$

$$p_3(x) = x^{80} + x^{79} + x^{78} + x^{75} + x^{74} + x^{71} + x^{70} + x^{65} + x^{64} + x^{61} + x^{59}$$

$$+ x^{58} + x^{57} + x^{56} + x^{54} + x^{49} + x^{47} + x^{46} + x^{45} + x^{43} + x^{40} + x^{39}$$

$$+ x^{36} + x^{35} + x^{30} + x^{28} + x^{27} + x^{26} + x^{25} + x^{24} + x^{22} + x^{19} + x^{15}$$

$$+ x^{14} + x^{13} + x^{12},$$

$$p_6(x) = x^{76} + x^{75} + x^{74} + x^{72} + x^{66} + x^{65} + x^{64} + x^{62} + x^{55} + x^{50} + x^{49}$$

$$+ x^{48} + x^{47} + x^{46} + x^{44} + x^{42} + x^{41} + x^{40} + x^{39} + x^{36} + x^{35} + x^{32}$$

$$+ x^{31} + x^{29} + x^{26} + x^{24} + x^{23} + x^{21} + x^{18} + x^{17} + x^{15} + x^{14} + x^{11}$$

$$+ x^{10} + x^9 + x^7 + x^5 + x^3 + x + 1,$$

$$p_9(x) = x^{68} + x^{67} + x^{65} + x^{64} + x^{60} + x^{59} + x^{57} + x^{55} + x^{53} + x^{51} + x^{49}$$

$$+ x^{48} + x^{44} + x^{43} + x^{41} + x^{39} + x^{37} + x^{36} + x^{20} + x^{19} + x^{17} + x^{16}$$

$$+ x^{12} + x^{11} + x^9 + x^7 + x^5 + x^4,$$

$$p_{12}(x) = x^{64} + x^{63} + x^{61} + x^{59} + x^{57} + x^{56} + x^{52} + x^{51} + x^{49} + x^{47} + x^{45}$$

$$+ x^{43} + x^{41} + x^{40} + x^{36} + x^{35} + x^{33} + x^{32} + x^{16} + x^{15} + x^{13} + x^{11}$$

$$+ x^9 + x^8 + x^4 + x^3 + x + 1,$$

and the initial terms are $[0, 0, 0, 0, 0, 0, 1, 1, 0, 1, 1, 0, 0, 1, 0, 0]$.

Below is the transition function and output function of an 2-automaton that generates $T = M_{0,0}^o$:

Transition function $(n, j) \mapsto \delta(n, j)$ ($\Lambda(n) := [\delta(n, 0), \delta(n, 1)]$):

n	$\Lambda(n)$	n	$\Lambda(n)$	n	$\Lambda(n)$	n	$\Lambda(n)$	n	$\Lambda(n)$
0	[1, 2]	57	[101, 102]	114	[162, 167]	171	[221, 42]	228	[251, 252]
1	[3, 4]	58	[70, 103]	115	[168, 169]	172	[129, 137]	229	[253, 254]
2	[5, 6]	59	[104, 105]	116	[167, 170]	173	[50, 111]	230	[255, 189]
3	[7, 8]	60	[106, 55]	117	[171, 18]	174	[53, 51]	231	[256, 97]
4	[9, 10]	61	[56, 107]	118	[172, 173]	175	[222, 223]	232	[257, 252]
5	[11, 12]	62	[108, 109]	119	[174, 25]	176	[120, 109]	233	[258, 201]
6	[13, 14]	63	[110, 111]	120	[175, 176]	177	[121, 10]	234	[259, 201]
7	[15, 16]	64	[112, 113]	121	[177, 178]	178	[215, 142]	235	[196, 73]
8	[17, 18]	65	[114, 115]	122	[179, 154]	179	[135, 223]	236	[260, 75]
9	[19, 20]	66	[114, 36]	123	[28, 180]	180	[40, 52]	237	[261, 254]
10	[21, 22]	67	[36, 116]	124	[181, 77]	181	[224, 225]	238	[90, 262]

11	[23, 24]	68	[44, 43]	125	[22, 67]	182	[40, 145]	239	[218, 107]
12	[25, 26]	69	[117, 118]	126	[20, 182]	183	[226, 40]	240	[116, 216]
13	[27, 28]	70	[49, 110]	127	[183, 184]	184	[220, 121]	241	[263, 131]
14	[29, 30]	71	[119, 120]	128	[185, 186]	185	[222, 136]	242	[264, 113]
15	[31, 32]	72	[33, 10]	129	[187, 168]	186	[32, 58]	243	[217, 114]
16	[33, 34]	73	[121, 34]	130	[188, 189]	187	[124, 227]	244	[265, 136]
17	[35, 36]	74	[122, 123]	131	[190, 153]	188	[228, 136]	245	[130, 140]
18	[37, 38]	75	[124, 113]	132	[191, 192]	189	[229, 121]	246	[147, 131]
19	[39, 40]	76	[115, 125]	133	[193, 89]	190	[230, 108]	247	[55, 105]
20	[14, 41]	77	[8, 110]	134	[178, 194]	191	[231, 225]	248	[266, 227]
21	[37, 42]	78	[10, 126]	135	[195, 91]	192	[232, 39]	249	[214, 105]
22	[43, 44]	79	[127, 128]	136	[194, 16]	193	[233, 128]	250	[119, 108]
23	[45, 46]	80	[55, 129]	137	[154, 196]	194	[10, 134]	251	[264, 227]
24	[47, 48]	81	[130, 131]	138	[156, 156]	195	[234, 133]	252	[267, 149]
25	[49, 50]	82	[132, 133]	139	[197, 159]	196	[129, 150]	253	[146, 36]
26	[51, 52]	83	[34, 134]	140	[198, 177]	197	[224, 235]	254	[132, 149]
27	[53, 40]	84	[135, 136]	141	[199, 200]	198	[233, 215]	255	[268, 105]
28	[54, 55]	85	[122, 41]	142	[201, 172]	199	[112, 227]	256	[269, 38]
29	[56, 57]	86	[105, 137]	143	[202, 203]	200	[125, 114]	257	[138, 216]
30	[47, 58]	87	[138, 114]	144	[204, 205]	201	[8, 50]	258	[104, 129]
31	[59, 60]	88	[139, 140]	145	[74, 83]	202	[231, 235]	259	[265, 223]
32	[61, 62]	89	[50, 141]	146	[92, 65]	203	[229, 33]	260	[35, 115]
33	[63, 24]	90	[128, 142]	147	[206, 207]	204	[54, 214]	261	[269, 42]
34	[64, 65]	91	[143, 42]	148	[208, 184]	205	[111, 152]	262	[141, 238]
35	[66, 22]	92	[116, 114]	149	[209, 95]	206	[219, 118]	263	[270, 25]
36	[67, 68]	93	[128, 144]	150	[210, 211]	207	[236, 39]	264	[271, 272]
37	[69, 70]	94	[51, 145]	151	[212, 213]	208	[237, 57]	265	[273, 75]
38	[71, 72]	95	[39, 51]	152	[85, 101]	209	[230, 120]	266	[274, 28]
39	[73, 74]	96	[146, 115]	153	[214, 129]	210	[14, 123]	267	[275, 204]
40	[75, 76]	97	[147, 140]	154	[215, 144]	211	[111, 238]	268	[102, 30]
41	[77, 78]	98	[120, 12]	155	[143, 38]	212	[110, 141]	269	[276, 277]
42	[79, 80]	99	[148, 149]	156	[216, 115]	213	[108, 12]	270	[278, 107]
43	[43, 43]	100	[139, 131]	157	[217, 216]	214	[173, 210]	271	[228, 223]
44	[76, 66]	101	[105, 150]	158	[218, 57]	215	[239, 93]	272	[232, 151]
45	[81, 82]	102	[151, 40]	159	[106, 214]	216	[68, 240]	273	[221, 38]
46	[83, 19]	103	[141, 152]	160	[34, 126]	217	[170, 168]	274	[278, 57]
47	[84, 85]	104	[153, 154]	161	[32, 48]	218	[241, 21]	275	[117, 46]
48	[60, 86]	105	[155, 156]	162	[36, 125]	219	[242, 88]	276	[237, 107]
49	[87, 88]	106	[157, 155]	163	[219, 46]	220	[243, 64]	277	[236, 151]
50	[89, 90]	107	[78, 63]	164	[220, 33]	221	[244, 60]	278	[279, 64]
51	[91, 92]	108	[158, 20]	165	[127, 215]	222	[245, 246]	279	[266, 113]
52	[93, 94]	109	[159, 160]	166	[151, 51]	223	[86, 247]		
53	[95, 93]	110	[16, 161]	167	[115, 116]	224	[248, 155]		
54	[96, 97]	111	[18, 162]	168	[44, 44]	225	[160, 249]		

55	[72, 98]	112	[163, 164]	169	[216, 36]	226	[247, 186]
56	[99, 100]	113	[165, 166]	170	[125, 216]	227	[250, 212]

Output function $n \mapsto \tau(n)$:

n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$
0	0	41	0	82	0	123	1	164	0	205	1	246	0
1	0	42	1	83	0	124	1	165	1	206	0	247	1
2	0	43	0	84	1	125	0	166	1	207	0	248	1
3	0	44	1	85	1	126	0	167	1	208	0	249	0
4	0	45	0	86	0	127	1	168	1	209	1	250	0
5	0	46	0	87	1	128	0	169	1	210	0	251	0
6	0	47	1	88	1	129	1	170	0	211	1	252	1
7	0	48	0	89	0	130	0	171	1	212	1	253	1
8	0	49	1	90	0	131	1	172	1	213	1	254	0
9	0	50	0	91	0	132	0	173	0	214	0	255	1
10	1	51	0	92	1	133	0	174	0	215	1	256	0
11	0	52	0	93	0	134	1	175	0	216	1	257	1
12	1	53	0	94	0	135	1	176	0	217	0	258	0
13	0	54	1	95	0	136	1	177	0	218	1	259	1
14	0	55	1	96	1	137	1	178	1	219	0	260	0
15	0	56	0	97	0	138	1	179	1	220	0	261	0
16	1	57	0	98	0	139	1	180	1	221	1	262	0
17	0	58	1	99	0	140	0	181	1	222	0	263	1
18	1	59	0	100	1	141	0	182	1	223	0	264	0
19	0	60	0	101	0	142	0	183	1	224	1	265	1
20	0	61	0	102	1	143	0	184	0	225	0	266	1
21	1	62	1	103	0	144	1	185	0	226	1	267	1
22	0	63	1	104	0	145	1	186	0	227	0	268	1
23	0	64	0	105	0	146	1	187	1	228	0	269	0
24	1	65	0	106	0	147	0	188	0	229	1	270	1
25	1	66	0	107	1	148	0	189	1	230	1	271	0
26	0	67	0	108	1	149	1	190	1	231	0	272	1
27	0	68	1	109	0	150	0	191	0	232	1	273	1
28	1	69	1	110	1	151	1	192	1	233	0	274	1
29	0	70	1	111	1	152	1	193	0	234	1	275	1
30	1	71	0	112	0	153	0	194	1	235	1	276	0
31	0	72	1	113	1	154	1	195	1	236	0	277	0
32	0	73	0	114	0	155	0	196	1	237	0	278	1
33	1	74	1	115	1	156	1	197	1	238	0	279	1
34	0	75	1	116	1	157	0	198	0	239	1		
35	0	76	1	117	1	158	1	199	0	240	1		
36	0	77	0	118	1	159	0	200	0	241	1		
37	1	78	1	119	0	160	0	201	0	242	0		
38	0	79	1	120	0	161	0	202	0	243	0		

39	0	80	1	121	0	162	0	203	1	244	1	
40	1	81	0	122	1	163	0	204	1	245	0	

UNIVERSITÉ DE STRASBOURG, CNRS, IRMA UMR 7501, F-67000 STRASBOURG, FRANCE
Email address: `guoniu.han@unistra.fr`

SCHOOL OF MATHEMATICS AND STATISTICS, HUAZHONG UNIVERSITY OF SCIENCE AND TECHNOLOGY, WUHAN, PR CHINA
Email address: `huyining@hust.edu.cn`