

PROOF OF THE CONTINUED FRACTION CONJECTURE FOR

$$(a, b) = (z^5 + z^4 + z^3 + z, z + 1)$$

GUO-NIU HAN* AND YINING HU*

ABSTRACT. The continued fraction conjecture claims that the continued fraction defined by the (a, b) -Thue-Morse sequence is algebraic over $\mathbb{F}_2(x)$ for all pairs of distinct elements (a, b) in $\mathbb{F}_2[x] \setminus \mathbb{F}_2$. In this paper we prove the conjecture for $(a, b) = (z^5 + z^4 + z^3 + z, z + 1)$.

1. INTRODUCTION

Let $\mathbb{F}_2$ be the finite field of cardinality 2. Given a sequence of polynomials $\mathbf{a} = (a_0(z), a_1(z), a_2(z), \dots)$ of elements from $\mathbb{F}_2[z] \setminus \mathbb{F}_2$, we define the infinite continued fraction

$$(1.1) \quad \text{CF}(\mathbf{a}(z)) = \frac{1}{a_0(z) + \frac{1}{a_1(z) + \frac{1}{a_2(z) + \frac{1}{\ddots}}}}$$

as the limit of the finite continued fractions

$$(1.2) \quad \text{CF}_n(\mathbf{a}(z)) = \frac{1}{a_0(z) + \frac{1}{a_1(z) + \frac{1}{\ddots + \frac{1}{a_n(z)}}}} \in \mathbb{F}_2((1/z)).$$

Conversely, each power series in $\mathbb{F}_2[[1/z]]$ without constant term

$$(1.3) \quad c_{-1}z^{-1} + c_{-2}z^{-2} + c_{-3}z^{-3} + \dots$$

admits a continued fraction expansion of form (1.1).

Let a and b be two distinct elements from $\mathbb{F}_2[z] \setminus \mathbb{F}_2$. We consider the continued fraction (1.1) associated with the (a, b) -Thue-Morse sequence

$$\mathbf{t} = (t_0(z), t_1(z), t_2(z), \dots) = (a, b, b, a, b, a, a, b, \dots)$$

that we shall denote

$$(1.4) \quad \text{CF}(a, b) := \text{CF}(\mathbf{t}).$$

Date: 9/04/2020 15:46:58 .

2010 Mathematics Subject Classification. 11B85, 11J70, 11B50, 11Y65, 05A15.

Key words and phrases. automatic sequence, continued fraction, Thue-Morse sequence.

* This paper was automatically generated from a computer program developed by the authors, with 8.1 minutes CPU time used.

Theorem 1.1. *When $(a, b) = (z^5 + z^4 + z^3 + z, z + 1)$, the two power series $\text{CF}(a, b)$ and $\text{CF}(b, a)$ are algebraic over $\mathbb{F}_2(z)$, with minimal polynomials of the form*

$$p_4(z)y^4 + p_3(z)y^3 + p_2(z)y^2 + p_1(z)y + p_0(z) = 0$$

where, for $\text{CF}(a, b)$,

$$\begin{aligned} p_0(z) &= z^{12} + z^9 + z^8 + z^7 + z^4 + z^3 + z^2 + z + 1, \\ p_1(z) &= z^{17} + z^{16} + z^{15} + z^{14} + z^{11} + z^7 + z^4 + z^3 + z^2 + z, \\ p_2(z) &= z^{18} + z^{10} + z^8 + z^6 + z^4 + z^2, \\ p_3(z) &= z^{17} + z^{16} + z^{15} + z^{14} + z^{11} + z^7 + z^4 + z^3 + z^2 + z, \\ p_4(z) &= z^{16} + z^{14} + z^{10} + z^9 + z^7 + z^4 + z^3 + z^2 + z, \end{aligned}$$

and for $\text{CF}(b, a)$,

$$\begin{aligned} p_0(z) &= z^{14} + z^{10} + z^9 + z^7 + z^4 + z^3 + z, \\ p_1(z) &= z^{17} + z^{16} + z^{15} + z^{14} + z^{11} + z^7 + z^4 + z^3 + z^2 + z, \\ p_2(z) &= z^{18} + z^{10} + z^8 + z^6 + z^4 + z^2, \\ p_3(z) &= z^{17} + z^{16} + z^{15} + z^{14} + z^{11} + z^7 + z^4 + z^3 + z^2 + z, \\ p_4(z) &= z^{16} + z^{12} + z^9 + z^8 + z^7 + z^4 + z^3 + z + 1. \end{aligned}$$

2. PROOF

Define the sequences of polynomials $(P_n(z))_{n \geq 0}$ and $(Q_n(z))_{n \geq 0}$ by

$$\begin{pmatrix} P_n(z) & Q_n(z) \\ P_{n-1}(z) & Q_{n-1}(z) \end{pmatrix} := \begin{pmatrix} t_n(z) & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} t_{n-1}(z) & 1 \\ 1 & 0 \end{pmatrix} \cdots \begin{pmatrix} t_0(z) & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}.$$

By the basic properties of continued fractions, we have

$$(2.1) \quad \text{CF}_n(a, b) = \frac{P_n(z)}{Q_n(z)} \in \mathbb{F}_2((1/z)).$$

As formula (2.1) is not efficient for calculating $\text{CF}_n(a, b)$, a fast method is to be derived and described as follows. Define

$$\begin{aligned} M_n(x) &= x^{\deg(t_{2^n-1})} \begin{pmatrix} t_{2^n-1}(1/x) & 1 \\ 1 & 0 \end{pmatrix} \times x^{\deg(t_{2^n-2})} \begin{pmatrix} t_{2^n-2}(1/x) & 1 \\ 1 & 0 \end{pmatrix} \times \\ &\quad \cdots \times x^{\deg(t_0)} \begin{pmatrix} t_0(1/x) & 1 \\ 1 & 0 \end{pmatrix}, \\ W_n(x) &= x^{\deg(\bar{t}_{2^n-1})} \begin{pmatrix} \bar{t}_{2^n-1}(1/x) & 1 \\ 1 & 0 \end{pmatrix} \times x^{\deg(\bar{t}_{2^n-2})} \begin{pmatrix} \bar{t}_{2^n-2}(1/x) & 1 \\ 1 & 0 \end{pmatrix} \times \\ &\quad \cdots \times x^{\deg(\bar{t}_0)} \begin{pmatrix} \bar{t}_0(1/x) & 1 \\ 1 & 0 \end{pmatrix}, \end{aligned}$$

where $\bar{t}$ is the (b, a) -Thue-Morse sequence. By the properties of the Thue-Morse sequence, we have for $n \geq 0$

$$\begin{aligned} M_{n+1}(x) &= W_n(x) \cdot M_n(x), \\ W_{n+1}(x) &= M_n(x) \cdot W_n(x). \end{aligned}$$

Let $x := 1/z$. For each non-zero polynomial $P(z)$, define $\tilde{P}(x)$ to be $P(1/x)$. Then,

$$\text{CF}_n(a, b) = \frac{P_n(z)}{Q_n(z)} = \frac{\tilde{P}_n(x)}{\tilde{Q}_n(x)} \in \mathbb{F}_2((x)) = \mathbb{F}_2((1/z)).$$

Comparing the definition of $M_n(x)$ with the definition of $P_n(x)$ and $Q_n(x)$, we see that

$$\begin{aligned} M_n(x)_{0,1} &= x^{d_n} \tilde{P}_{2^n-1}(x), \\ M_n(x)_{0,0} &= x^{d_n} \tilde{Q}_{2^n-1}(x), \end{aligned}$$

for some positive integer d_n . Hence,

$$(2.2) \quad \text{CF}_{2^{2^n}-1}(a, b) = \frac{\tilde{P}_{2^{2^n}-1}(x)}{\tilde{Q}_{2^{2^n}-1}(x)} = \frac{M_{2^n}(x)_{0,1}}{M_{2^n}(x)_{0,0}}.$$

Therefore, by the convergence theorem of continued fraction, the algebraicity of $\text{CF}(a, b)$ will be established if it is shown that both $M_{2^n}(x)_{0,1}$ and $M_{2^n}(x)_{0,0}$ converge to algebraic series in $\mathbb{F}_2[[x]]$.

Actually, we will prove that for all $0 \leq i, j \leq 1$ the four sequences $(M_{2^n}(x)_{i,j})_n$, $(M_{2^{n+1}}(x)_{i,j})_n$, $(W_{2^n}(x)_{i,j})_n$, and $(W_{2^{n+1}}(x)_{i,j})_n$ converge to algebraic series in $\mathbb{F}_2[[x]]$. For this purpose, we define four 2×2 matrices M^e, M^o, W^e, W^o as follows: For each $T \in \{M^e, M^o, W^e, W^o\}$ and $0 \leq i, j \leq 1$, $T_{i,j}$ is defined to be the unique solution in $\mathbb{F}_2[[x]]$ of the polynomial $\phi(T, i, j)$ under certain initial conditions; the polynomials $\phi(T, i, j)$ and initial conditions are given in Section 3. We will prove that these four matrices, whose components are algebraic by definition, are the limits of $(M_{2^n}(x))_n$, $(M_{2^{n+1}}(x))_n$, $(W_{2^n}(x))_n$, and $(W_{2^{n+1}}(x))_n$.

Let us explain how the polynomials $\phi(T, i, j)$ and initial conditions are found, and why the solutions exist and are unique. For $0 \leq i, j \leq 1$, the coefficients of the polynomial $\phi(M^e, i, j)$ (resp. $\phi(M^o, i, j)$, $\phi(W^e, i, j)$, and $\phi(W^o, i, j)$) are the Padé-Hermite approximants of type

$$(150, 150, 150, 150, 150)$$

of the vector

$$(1, T^3, T^6, T^9, T^{12}),$$

where $T = M_{12,i,j}$ (resp. $M_{11,i,j}$, $W_{12,i,j}$, and $W_{11,i,j}$), found by the Derksen algorithm. We take the first 8 terms of $M_{12,i,j}$ (resp. $M_{11,i,j}$, $W_{12,i,j}$, and $W_{11,i,j}$) as initial conditions for $\phi(M^e, i, j)$ (resp. $\phi(M^o, i, j)$, $\phi(W^e, i, j)$, and $\phi(W^o, i, j)$). The following fact will be used to ensure that the solution exists and is unique: let $P(x, y) \in \mathbb{F}_2[x, y]$ and for each series $f(x) = \sum_{n=0}^{\infty} a_n x^n \in \mathbb{F}_2[[x]]$ denote the partial sum $\sum_{j=0}^{n-1} a_j x^j$ by $f_n(x)$ for $n \geq 0$. If for some $n \geq 0$ and $a_0, a_1, \dots, a_{n-1} \in \mathbb{F}_2$ $P(x, \sum_{j=0}^{n-1} a_j x^j) = O(x^n)$ and $Q(x, y) := P(x, \sum_{j=0}^{n-1} a_j x^j + x^n y)$ can be written as $x^m \sum_{j=0}^{\infty} q_j(x) y^j$ where $q_j(x)$ are polynomials for $j \geq 0$, $q_1(0) = 1$, and $q_j(0) = 0$ for $j > 1$, then there exists a unique solution $f(x) \in \mathbb{F}_2[[x]]$ of $P(x, f(x)) = 0$ that satisfies the initial condition $f_n(x) = \sum_{j=0}^{n-1} a_j x^j$.

We state two lemmas concerning the four matrices M^e, M^o, W^e, W^o . The first one is about relations between them; the second, about the structure of the each matrix.

Lemma 2.1. *We have*

$$(2.3) \quad M^e = W^o \cdot M^o,$$

$$(2.4) \quad M^o = W^e \cdot M^e,$$

$$(2.5) \quad W^e = M^o \cdot W^o,$$

$$(2.6) \quad W^o = M^e \cdot W^e.$$

Proof. We give the proof of the identity

$$M_{0,0}^e = W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o,$$

the proofs of the others being similar.

First, we compute the minimal polynomials of $W_{0,0}^o M_{0,0}^o$ and $W_{0,1}^o M_{1,0}^o$. We know that

$$P(x, y) = \text{Res}_z (\phi(W^o, 0, 0)(x, z), z^{12} \cdot \phi(M^o, 0, 0)(x, y/z))$$

is an annihilating polynomial of $W_{0,0}^o M_{0,0}^o$. We use Padé-Hermite approximation to find a candidate for the minimal polynomial of $W_{0,0}^o M_{0,0}^o$, that will be called $\phi_0(x, y)$. To prove that $\phi_0(x, y)$ is indeed the minimal polynomial, it suffices to prove that it is an irreducible factor of $P(x, y)$ of multiplicity m and that $Q(x, y) := P(x, y)/\phi_0(x, y)^m$ is not an annihilating polynomial of $W_{0,0}^o M_{0,0}^o$. We verify the first point directly. For the second point, we truncate $W_{0,0}^o M_{0,0}^o$ to order 540 and substitute it for y in $Q(x, y)$. We get a series of valuation less than 540, which proves that $Q(x, y)$ is not an annihilating polynomial of $W_{0,0}^o M_{0,0}^o$. We find the minimal polynomial $\phi_1(x, y)$ of $W_{0,1}^o M_{1,0}^o$ in a similar way.

Now we prove that $\phi(M^e, 0, 0)$ is the minimal polynomial of $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$. We know that

$$S(x, y) = \text{Res}_z (\phi_0(x, z), \phi_1(x, y + z))$$

is an annihilating of $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$. We verify that $\phi(M^e, 0, 0)$ is an irreducible factor of $S(x, y)$ of multiplicity μ , and that $Q(x, y) := S(x, y)/\phi(M^e, 0, 0)^\mu$ is not an annihilating polynomial of $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$. To see the last point, we truncate $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$ to order 660 and substitute it for y in $Q(x, y)$. We get a series of valuation less than 660, and therefore $Q(x, y)$ is not an annihilating polynomial of $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$.

Finally, the first 16 terms of $M_{0,0}^e$ and $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$ coincide. As these first terms determine a unique solution of $\phi(M^e, 0, 0)$, we know that the two series are one and the same. $\square$

Define

$$R^e = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \quad \text{and} \quad R^o = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}.$$

Lemma 2.2. *For all integers $k \geq 2$ and $u = 2^{2k-1}$, the following identities hold.*

$$\begin{aligned} M^e[:12u] &= M^e[:6u] + x^{2u} M^e[:4u] + x^{3u} M^e[:3u] + x^{4u} M^e[:2u] + x^{2u} M^e[:6u] \\ &\quad + x^{4u} M^e[:4u] + x^{5u} M^e[:3u] + x^{6u} M^e[:2u] + x^{3u} M^e[:6u] \\ &\quad + x^{5u} M^e[:4u] + x^{6u} M^e[:3u] + x^{7u} M^e[:2u] + x^{6u} M^e[:6u] \\ &\quad + (x^{6u} + x^{8u} + x^{9u}) R^e, \\ W^e[:12u] &= W^e[:6u] + x^{2u} W^e[:4u] + x^{3u} W^e[:3u] + x^{4u} W^e[:2u] + x^{2u} W^e[:6u] \\ &\quad + x^{4u} W^e[:4u] + x^{5u} W^e[:3u] + x^{6u} W^e[:2u] + x^{3u} W^e[:6u] \end{aligned}$$

$$\begin{aligned}
& + x^{5u}W^e[:4u] + x^{6u}W^e[:3u] + x^{7u}W^e[:2u] + x^{6u}W^e[:6u] \\
& + (x^{6u} + x^{8u} + x^{9u})R^e.
\end{aligned}$$

For all integers $k \geq 2$ and $u = 2^{2k}$, the following identities hold.

$$\begin{aligned}
M^o[:12u] &= M^o[:6u] + x^{2u}M^o[:4u] + x^{3u}M^o[:3u] + x^{4u}M^o[:2u] + x^{2u}M^o[:6u] \\
&+ x^{4u}M^o[:4u] + x^{5u}M^o[:3u] + x^{6u}M^o[:2u] + x^{3u}M^o[:6u] \\
&+ x^{5u}M^o[:4u] + x^{6u}M^o[:3u] + x^{7u}M^o[:2u] + x^{6u}M^o[:6u] \\
&+ (x^{6u} + x^{8u} + x^{9u})R^o, \\
W^o[:12u] &= W^o[:6u] + x^{2u}W^o[:4u] + x^{3u}W^o[:3u] + x^{4u}W^o[:2u] + x^{2u}W^o[:6u] \\
&+ x^{4u}W^o[:4u] + x^{5u}W^o[:3u] + x^{6u}W^o[:2u] + x^{3u}W^o[:6u] \\
&+ x^{5u}W^o[:4u] + x^{6u}W^o[:3u] + x^{7u}W^o[:2u] + x^{6u}W^o[:6u] \\
&+ (x^{6u} + x^{8u} + x^{9u})R^o.
\end{aligned}$$

Proof. To prove Lemma 2.2 we first construct an automaton for each sequence concerned, and then transform the conditions on infinitely many k 's into finitely many conditions on the states of the automaton. In the following, we will prove that for $T = M_{0,0}^o$, for all integer $k \geq 2$ and $u = 2^{2k}$,

$$\begin{aligned}
T[:12u] &= T[:6u] + x^{2u}T[:4u] + x^{3u}T[:3u] + x^{4u}T[:2u] + x^{2u}T[:6u] + x^{4u}T[:4u] \\
&+ x^{5u}T[:3u] + x^{6u}T[:2u] + x^{3u}T[:6u] + x^{5u}T[:4u] + x^{6u}T[:3u] \\
&+ x^{7u}T[:2u] + x^{6u}T[:6u].
\end{aligned}$$

The proof of the other 15 cases are similar. We break down the above identity into 6 parts:

$$\begin{aligned}
0 &= x^{6u}T[:u] + x^{4u}T[2u:3u] + x^{3u}T[3u:4u] + x^{2u}T[4u:5u] + T[6u:7u], \\
0 &= x^{7u}T[:u] + x^{6u}T[u:2u] + x^{4u}T[3u:4u] + x^{3u}T[4u:5u] \\
&+ x^{2u}T[5u:6u] + T[7u:8u], \\
0 &= x^{7u}T[u:2u] + x^{5u}T[3u:4u] + x^{3u}T[5u:6u] + T[8u:9u], \\
0 &= x^{6u}T[3u:4u] + T[9u:10u], \\
0 &= x^{6u}T[4u:5u] + T[10u:11u], \\
0 &= x^{6u}T[5u:6u] + T[11u:12u],
\end{aligned}$$

which can be rewritten as

$$(2.7) \quad 0 = T[[w]_2] + T[[10w]_2] + T[[11w]_2] + T[[100w]_2] + T[[110w]_2],$$

$$0 = T[[w]_2] + T[[1w]_2] + T[[11w]_2] + T[[100w]_2] + T[[101w]_2]$$

$$(2.8) \quad + T[[111w]_2],$$

$$(2.9) \quad 0 = T[[1w]_2] + T[[11w]_2] + T[[101w]_2] + T[[1000w]_2],$$

$$(2.10) \quad 0 = T[[11w]_2] + T[[1001w]_2],$$

$$(2.11) \quad 0 = T[[100w]_2] + T[[1010w]_2],$$

$$(2.12) \quad 0 = T[[101w]_2] + T[[1011w]_2],$$

for all binary word w of length $2k$ and $w \neq 0^{2k}$, and

$$(2.13) \quad 0 = T[[w]_2] + T[[10w]_2] + T[[11w]_2] + T[[100w]_2] + T[[110w]_2],$$

$$0 = T[[w]_2] + T[[1w]_2] + T[[11w]_2] + T[[100w]_2] + T[[101w]_2]$$

$$(2.14) \quad + T[[111w]_2],$$

$$(2.15) \quad 0 = T[[1w]_2] + T[[11w]_2] + T[[101w]_2] + T[[1000w]_2],$$

$$(2.16) \quad 0 = T[[11w]_2] + T[[1001w]_2],$$

$$(2.17) \quad 0 = T[[100w]_2] + T[[1010w]_2],$$

$$(2.18) \quad 0 = T[[101w]_2] + T[[1011w]_2],$$

for $w = 0^{2k}$.

First we calculate an 2-automaton that generates T from its minimal polynomial and its first terms. This automaton has 379 states; its transition function and output function can be found in the annex. Let $A(s, w)$ denote the state reached after reading w from right to left starting from the state s , and τ the output function. Define

$$E_{2k} = \{A(i, w) : |w| = 2k, w \neq 0^{2k}\}.$$

Identities (2.7) through (2.18) can be written as

$$(2.19) \quad \begin{aligned} 0 &= \tau(A(s, \epsilon)) + \tau(A(s, 10)) + \tau(A(s, 11)) + \tau(A(s, 100)) \\ &\quad + \tau(A(s, 110)), \end{aligned}$$

$$(2.20) \quad \begin{aligned} 0 &= \tau(A(s, \epsilon)) + \tau(A(s, 1)) + \tau(A(s, 11)) + \tau(A(s, 100)) \\ &\quad + \tau(A(s, 101)) + \tau(A(s, 111)), \end{aligned}$$

$$(2.21) \quad 0 = \tau(A(s, 1)) + \tau(A(s, 11)) + \tau(A(s, 101)) + \tau(A(s, 1000)),$$

$$(2.22) \quad 0 = \tau(A(s, 11)) + \tau(A(s, 1001)),$$

$$(2.23) \quad 0 = \tau(A(s, 100)) + \tau(A(s, 1010)),$$

$$(2.24) \quad 0 = \tau(A(s, 101)) + \tau(A(s, 1011)),$$

for all $s \in E_{2k}$, and

$$(2.25) \quad \begin{aligned} 0 &= \tau(A(s, \epsilon)) + \tau(A(s, 10)) + \tau(A(s, 11)) + \tau(A(s, 100)) \\ &\quad + \tau(A(s, 110)), \end{aligned}$$

$$(2.26) \quad \begin{aligned} 0 &= \tau(A(s, \epsilon)) + \tau(A(s, 1)) + \tau(A(s, 11)) + \tau(A(s, 100)) \\ &\quad + \tau(A(s, 101)) + \tau(A(s, 111)), \end{aligned}$$

$$(2.27) \quad 0 = \tau(A(s, 1)) + \tau(A(s, 11)) + \tau(A(s, 101)) + \tau(A(s, 1000)),$$

$$(2.28) \quad 0 = \tau(A(s, 11)) + \tau(A(s, 1001)),$$

$$(2.29) \quad 0 = \tau(A(s, 100)) + \tau(A(s, 1010)),$$

$$(2.30) \quad 0 = \tau(A(s, 101)) + \tau(A(s, 1011)),$$

for $s = A(i, 0^{2k})$. We find that $(A(i, 0^{16}), E_{16}) = (A(i, 0^{12}), E_{12})$, so that we only have to verify that identities (2.19) through (2.30) hold for $2 \leq k \leq 8$, which turns out to be true. $\square$

In the following lemma, we express M_{2k} , M_{2k+1} , W_{2k} , and W_{2k+1} in terms of M^e , M^o , W^e , and W^o .

Lemma 2.3. *For all integer $k \geq 2$, and $u = 2^{2k-1}$,*

$$\begin{aligned} M_{2k} &= M^e[:6u] + x^{2u}M^e[:4u] + x^{3u}M^e[:3u] + x^{4u}M^e[:2u] + x^{6u}R^e, \\ W_{2k} &= W^e[:6u] + x^{2u}W^e[:4u] + x^{3u}W^e[:3u] + x^{4u}W^e[:2u] + x^{6u}R^e. \end{aligned}$$

For all integer $k \geq 2$, and $u = 2^{2k}$,

$$\begin{aligned} M_{2k+1} &= M^o[:6u] + x^{2u}M^o[:4u] + x^{3u}M^o[:3u] + x^{4u}M^o[:2u] + x^{6u}R^o, \\ W_{2k+1} &= W^o[:6u] + x^{2u}W^o[:4u] + x^{3u}W^o[:3u] + x^{4u}W^o[:2u] + x^{6u}R^o. \end{aligned}$$

Proof. We call the four identities in Lemma 2.3 also by the name M_{2k} , W_{2k} , M_{2k+1} , and W_{2k+1} . For $n = 2$, the identities can be verified directly. For $n \geq 2$, we claim that

$$\begin{aligned} M_{2k} \wedge W_{2k} &\Rightarrow M_{2k+1} \wedge W_{2k+1}, \\ M_{2k+1} \wedge W_{2k+1} &\Rightarrow M_{2k+2} \wedge W_{2k+2}. \end{aligned}$$

We give the proof of

$$M_{2k} \wedge W_{2k} \Rightarrow M_{2k+1},$$

the other ones being similar. Set $u = 2^{2k}$ and $v = 2^{2k-1}$. By definition and induction hypothesis, the left side of identity M_{2k+1} is equal to

$$\begin{aligned} (2.31) \quad &W_{2k}M_{2k} = (W^e[:6v] + x^{2v}W^e[:4v] + x^{3v}W^e[:3v] + x^{4v}W^e[:2v] + x^{6v}R^e \\ &\quad) \times (M^e[:6v] + x^{2v}M^e[:4v] + x^{3v}M^e[:3v] + x^{4v}M^e[:2v] + x^{6v}R^e \\ &\quad); \end{aligned}$$

Call this expression *lhs*. Note that both sides of identity M_{2k+1} have the same term of highest degree $x^{12v}R^o$. Therefore we only need to prove that their difference is $O(x^{12v})$. Using Lemma 2.1 it can be seen that the right side of identity M_{2k+1} is congruent, modulo x^{12v} , to

$$\begin{aligned} &W^e[:12v]M^e[:12v] + x^{4v}W^e[:8v]M^e[:8v] + x^{6v}W^e[:6v]M^e[:6v] \\ &\quad + x^{8v}W^e[:4v]M^e[:4v]. \end{aligned}$$

For all $n \leq 12$, replace the occurrences of $W^e[:n \cdot v]$ and $M^e[:n \cdot v]$ in the above expression by the reduction modulo $x^{n \cdot v}$ of the right side of the corresponding identity in Lemma 2.2 and get a new expression, which we call *rhs*. Define

$$\begin{aligned} X &:= x^v, \\ a_n &:= W^e[n \cdot v : (n+1) \cdot v] / X^n, \\ b_n &:= M^e[n \cdot v : (n+1) \cdot v] / X^n, \\ c &:= R^e. \end{aligned}$$

Using the notation introduced above, we can represent the expressions *lhs* (2.31) and *rhs* as polynomials in $\mathbb{F}_2[a_1, \dots, a_{12}, b_1, \dots, b_{12}, c][X]$. Note that it is not a problem that a_j commutes with b_k while W^e does not commute with M^e , because in the expressions concerned, the products of W^e -terms and M^e -terms are always in the same order. We let the computer do the simplification and check that the difference between these two polynomials is indeed $O(X^{12})$, which completes the proof. $\square$

Proof of Theorem 1.1. We prove the theorem for $\text{CF}(a, b)$; for $\text{CF}(b, a)$, the proof is similar. By Lemma 2.3, we have For all $0 \leq j, k \leq 1$,

$$\begin{aligned}\lim_{n \rightarrow \infty} M_{2n, j, k} &= M_{j, k}^e, \\ \lim_{n \rightarrow \infty} M_{2n+1, j, k} &= M_{j, k}^o, \\ \lim_{n \rightarrow \infty} W_{2n, j, k} &= W_{j, k}^e, \\ \lim_{n \rightarrow \infty} W_{2n+1, j, k} &= W_{j, k}^o.\end{aligned}$$

Let $z = 1/x$. By the convergence theorem and identity (2.2),

$$\text{CF}(a, b) = \frac{M_{0,1}^e(x)}{M_{0,0}^e(x)}.$$

By definition, $\phi(M^e, 0, 1)$ and $\phi(M^e, 0, 0)$ are minimal polynomials of $M_{0,1}^e$ and $M_{0,0}^e$. Therefore

$$P(x, y) = \text{Res}_t (\phi(M^e, 0, 1)(x, t), y^{12} \phi(M^e, 0, 1)(x, t/y))$$

is an annihilating polynomial of $f(x) = M_{0,1}^e/M_{0,0}^e$.

Define

$$Q(x, y) = q_4(x)y^4 + q_3(x)y^3 + q_2(x)y^2 + q_1(x)y + q_0(x),$$

where

$$\begin{aligned}q_0(x) &= x^{18} + x^{17} + x^{16} + x^{15} + x^{14} + x^{11} + x^{10} + x^9 + x^6, \\ q_1(x) &= x^{17} + x^{16} + x^{15} + x^{14} + x^{11} + x^7 + x^4 + x^3 + x^2 + x, \\ q_2(x) &= x^{16} + x^{14} + x^{12} + x^{10} + x^8 + 1, \\ q_3(x) &= x^{17} + x^{16} + x^{15} + x^{14} + x^{11} + x^7 + x^4 + x^3 + x^2 + x, \\ q_4(x) &= x^{17} + x^{16} + x^{15} + x^{14} + x^{11} + x^9 + x^8 + x^4 + x^2.\end{aligned}$$

The polynomial $Q(x, y)$ is the candidate for the minimal polynomial of $f(x)$ found by Padé-Hermite approximation. To prove that it is indeed the minimal polynomial of $f(x)$, we only need to prove that it is an irreducible factor of $P(x, y)$ of multiplicity m and $R(x, y) := P(x, y)/Q(x, y)^m$ is not an annihilating polynomial of $f(x)$. We verify the first point directly. For the second point, we find that when we truncate $f(x)$ to order 186, and substitute it for y in $R(z, y)$, we get a series with valuation smaller than 186, which proves that $R(z, y)$ is not an annihilating polynomial of $f(x)$. Finally, $z^{12}Q(1/z, y)$ is the minimal polynomial of $\text{CF}(a, b) = f(1/z)$. $\square$

3. ANNEXE

All 16 polynomials are of the form

$$p_0(x) + p_3(x)y^3 + p_6(x)y^6 + p_9(x)y^9 + p_{12}(x)y^{12}.$$

The coefficients $p_j(x)$ and the 16 initial terms to determine the solutions uniquely are given below.

For $\phi(M^e, 0, 0)$:

$$\begin{aligned}p_0(x) &= x^{84} + x^{80} + x^{78} + x^{68} + x^{56} + x^{48} + x^{40} + x^{38} + x^{36} + x^{34} + x^{28} \\ &\quad + x^{24} + x^{22} + x^{18} + x^{12}, \\ p_3(x) &= x^{80} + x^{78} + x^{74} + x^{72} + x^{66} + x^{62} + x^{58} + x^{56} + x^{52} + x^{50} + x^{48}\end{aligned}$$

$$\begin{aligned}
& + x^{42} + x^{40} + x^{36} + x^{32} + x^{30} + x^{28} + x^{16} + x^{14} + x^{12} + x^{10} + x^8, \\
p_6(x) &= x^{78} + x^{76} + x^{74} + x^{70} + x^{60} + x^{58} + x^{56} + x^{42} + x^{32} + x^{30} + x^{28} \\
& + x^{24} + x^{22} + x^{18} + x^{12} + x^{10} + x^2 + 1, \\
p_9(x) &= x^{82} + x^{76} + x^{74} + x^{70} + x^{66} + x^{64} + x^{52} + x^{46} + x^{44} + x^{40} + x^{38} \\
& + x^{32} + x^{20} + x^{18} + x^{14} + x^{10} + x^8 + x^2, \\
p_{12}(x) &= x^{82} + x^{80} + x^{78} + x^{74} + x^{70} + x^{66} + x^{62} + x^{58} + x^{54} + x^{48} + x^{34} \\
& + x^{32} + x^{30} + x^{26} + x^{22} + x^{18} + x^{14} + x^{10} + x^6 + 1,
\end{aligned}$$

and the initial terms are $[1, 0, 1, 0, 0, 0, 0, 0, 1, 0, 1, 0, 1, 0, 0, 0]$.

For $\phi(M^e, 0, 1)$:

$$\begin{aligned}
p_0(x) &= x^{96} + x^{95} + x^{94} + x^{87} + x^{85} + x^{84} + x^{82} + x^{80} + x^{79} + x^{77} + x^{75} \\
& + x^{67} + x^{66} + x^{64} + x^{60} + x^{59} + x^{58} + x^{50} + x^{49} + x^{48} + x^{47} + x^{46} \\
& + x^{44} + x^{43} + x^{42} + x^{41} + x^{39} + x^{37} + x^{36} + x^{35} + x^{34} + x^{33} + x^{32} \\
& + x^{31} + x^{30} + x^{28} + x^{24}, \\
p_3(x) &= x^{94} + x^{92} + x^{91} + x^{84} + x^{82} + x^{81} + x^{78} + x^{75} + x^{74} + x^{73} + x^{62} \\
& + x^{60} + x^{59} + x^{54} + x^{51} + x^{50} + x^{49} + x^{22} + x^{20} + x^{19} + x^{14} + x^{11} \\
& + x^{10} + x^9, \\
p_6(x) &= x^{93} + x^{92} + x^{91} + x^{89} + x^{87} + x^{85} + x^{83} + x^{81} + x^{78} + x^{73} + x^{72} \\
& + x^{71} + x^{69} + x^{67} + x^{64} + x^{59} + x^{58} + x^{56} + x^{53} + x^{52} + x^{50} + x^{43} \\
& + x^{42} + x^{40} + x^{39} + x^{38} + x^{37} + x^{35} + x^{32} + x^{29} + x^{28} + x^{26} + x^{25} \\
& + x^{24} + x^{23} + x^{20} + x^{19} + x^{18} + x^{17} + x^{14} + x^{11} + x^{10} + x^9 + x^6, \\
p_9(x) &= x^{92} + x^{89} + x^{88} + x^{87} + x^{82} + x^{80} + x^{79} + x^{76} + x^{73} + x^{71} + x^{69} \\
& + x^{67} + x^{66} + x^{65} + x^{62} + x^{59} + x^{58} + x^{57} + x^{52} + x^{50} + x^{49} + x^{48} \\
& + x^{45} + x^{43} + x^{42} + x^{41} + x^{40} + x^{38} + x^{37} + x^{36} + x^{34} + x^{33} + x^{30} \\
& + x^{28} + x^{27} + x^{26} + x^{24} + x^{23} + x^{20} + x^{17} + x^{15} + x^{13} + x^{12} + x^{11} \\
& + x^{10} + x^8 + x^7 + x^6 + x^4 + x^3, \\
p_{12}(x) &= x^{91} + x^{90} + x^{88} + x^{79} + x^{78} + x^{76} + x^{75} + x^{74} + x^{72} + x^{71} + x^{70} \\
& + x^{68} + x^{67} + x^{66} + x^{64} + x^{63} + x^{62} + x^{60} + x^{55} + x^{54} + x^{52} + x^{51} \\
& + x^{50} + x^{48} + x^{43} + x^{42} + x^{40} + x^{31} + x^{30} + x^{28} + x^{27} + x^{26} + x^{24} \\
& + x^{23} + x^{22} + x^{20} + x^{19} + x^{18} + x^{16} + x^{15} + x^{14} + x^{12} + x^7 + x^6 + x^4 \\
& + x^3 + x^2 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 1, 1, 1, 0, 0, 1, 1, 0, 0, 0, 0]$.

For $\phi(M^e, 1, 0)$:

$$\begin{aligned}
p_0(x) &= x^{96} + x^{95} + x^{94} + x^{87} + x^{85} + x^{84} + x^{82} + x^{80} + x^{79} + x^{77} + x^{75} \\
& + x^{67} + x^{66} + x^{64} + x^{60} + x^{59} + x^{58} + x^{50} + x^{49} + x^{48} + x^{47} + x^{46} \\
& + x^{44} + x^{43} + x^{42} + x^{41} + x^{39} + x^{37} + x^{36} + x^{35} + x^{34} + x^{33} + x^{32}
\end{aligned}$$

$$\begin{aligned}
& + x^{31} + x^{30} + x^{28} + x^{24}, \\
p_3(x) &= x^{94} + x^{92} + x^{91} + x^{84} + x^{82} + x^{81} + x^{78} + x^{75} + x^{74} + x^{73} + x^{62} \\
& + x^{60} + x^{59} + x^{54} + x^{51} + x^{50} + x^{49} + x^{22} + x^{20} + x^{19} + x^{14} + x^{11} \\
& + x^{10} + x^9, \\
p_6(x) &= x^{93} + x^{92} + x^{91} + x^{89} + x^{87} + x^{85} + x^{83} + x^{81} + x^{78} + x^{73} + x^{72} \\
& + x^{71} + x^{69} + x^{67} + x^{64} + x^{59} + x^{58} + x^{56} + x^{53} + x^{52} + x^{50} + x^{43} \\
& + x^{42} + x^{40} + x^{39} + x^{38} + x^{37} + x^{35} + x^{32} + x^{29} + x^{28} + x^{26} + x^{25} \\
& + x^{24} + x^{23} + x^{20} + x^{19} + x^{18} + x^{17} + x^{14} + x^{11} + x^{10} + x^9 + x^6, \\
p_9(x) &= x^{92} + x^{89} + x^{88} + x^{87} + x^{82} + x^{80} + x^{79} + x^{76} + x^{73} + x^{71} + x^{69} \\
& + x^{67} + x^{66} + x^{65} + x^{62} + x^{59} + x^{58} + x^{57} + x^{52} + x^{50} + x^{49} + x^{48} \\
& + x^{45} + x^{43} + x^{42} + x^{41} + x^{40} + x^{38} + x^{37} + x^{36} + x^{34} + x^{33} + x^{30} \\
& + x^{28} + x^{27} + x^{26} + x^{24} + x^{23} + x^{20} + x^{17} + x^{15} + x^{13} + x^{12} + x^{11} \\
& + x^{10} + x^8 + x^7 + x^6 + x^4 + x^3, \\
p_{12}(x) &= x^{91} + x^{90} + x^{88} + x^{79} + x^{78} + x^{76} + x^{75} + x^{74} + x^{72} + x^{71} + x^{70} \\
& + x^{68} + x^{67} + x^{66} + x^{64} + x^{63} + x^{62} + x^{60} + x^{55} + x^{54} + x^{52} + x^{51} \\
& + x^{50} + x^{48} + x^{43} + x^{42} + x^{40} + x^{31} + x^{30} + x^{28} + x^{27} + x^{26} + x^{24} \\
& + x^{23} + x^{22} + x^{20} + x^{19} + x^{18} + x^{16} + x^{15} + x^{14} + x^{12} + x^7 + x^6 + x^4 \\
& + x^3 + x^2 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 1, 1, 1, 0, 0, 1, 1, 0, 0, 0, 0]$.

For $\phi(M^e, 1, 1)$:

$$\begin{aligned}
p_0(x) &= x^{108} + x^{106} + x^{100} + x^{96} + x^{94} + x^{88} + x^{86} + x^{84} + x^{80} + x^{78} + x^{74} \\
& + x^{68} + x^{58} + x^{56} + x^{48} + x^{46} + x^{44} + x^{42} + x^{36}, \\
p_3(x) &= x^{102} + x^{100} + x^{98} + x^{90} + x^{88} + x^{86} + x^{82} + x^{80} + x^{78} + x^{76} + x^{74} \\
& + x^{72} + x^{70} + x^{68} + x^{66} + x^{62} + x^{60} + x^{56} + x^{50} + x^{48} + x^{40} + x^{38} \\
& + x^{36} + x^{30} + x^{26} + x^{22} + x^{20} + x^{14} + x^{10} + x^6, \\
p_6(x) &= x^{102} + x^{96} + x^{90} + x^{82} + x^{80} + x^{78} + x^{76} + x^{72} + x^{70} + x^{68} + x^{66} \\
& + x^{64} + x^{62} + x^{58} + x^{56} + x^{50} + x^{48} + x^{46} + x^{38} + x^{30} + x^{28} + x^{20} \\
& + x^{16} + x^6 + x^2 + 1, \\
p_9(x) &= x^{100} + x^{98} + x^{96} + x^{94} + x^{90} + x^{88} + x^{84} + x^{82} + x^{78} + x^{74} + x^{60} \\
& + x^{58} + x^{56} + x^{54} + x^{50} + x^{48} + x^{44} + x^{42} + x^{38} + x^{36} + x^{32} + x^{30} \\
& + x^{28} + x^{22} + x^{20} + x^{16} + x^{14} + x^{12} + x^6 + x^2, \\
p_{12}(x) &= x^{100} + x^{96} + x^{76} + x^{72} + x^{60} + x^{56} + x^{28} + x^{24} + x^{12} + x^8 + x^4 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 1, 0, 0, 0, 1, 0]$.

For $\phi(W^e, 0, 0)$:

$$p_0(x) = x^{108} + x^{106} + x^{104} + x^{100} + x^{98} + x^{82} + x^{80} + x^{78} + x^{76} + x^{74} + x^{64}$$

$$\begin{aligned}
& + x^{60} + x^{58} + x^{46} + x^{44} + x^{40} + x^{38} + x^{30} + x^{26} + x^{20} + x^{12}, \\
p_3(x) &= x^{102} + x^{100} + x^{96} + x^{90} + x^{88} + x^{80} + x^{74} + x^{70} + x^{66} + x^{64} + x^{62} \\
& + x^{58} + x^{56} + x^{52} + x^{50} + x^{46} + x^{44} + x^{42} + x^{36} + x^{34} + x^{32} + x^{30} \\
& + x^{24} + x^{20} + x^{18} + x^{14} + x^{12} + x^8, \\
p_6(x) &= x^{102} + x^{98} + x^{96} + x^{90} + x^{88} + x^{86} + x^{72} + x^{70} + x^{68} + x^{66} + x^{64} \\
& + x^{62} + x^{56} + x^{50} + x^{42} + x^{40} + x^{36} + x^{34} + x^{30} + x^{28} + x^{26} + x^{24} \\
& + x^{22} + x^{20} + x^{18} + x^{16} + x^{12} + x^{10} + x^8 + x^4 + x^2 + 1, \\
p_9(x) &= x^{100} + x^{98} + x^{96} + x^{94} + x^{90} + x^{88} + x^{84} + x^{82} + x^{78} + x^{74} + x^{60} \\
& + x^{58} + x^{56} + x^{54} + x^{50} + x^{48} + x^{44} + x^{42} + x^{38} + x^{36} + x^{32} + x^{30} \\
& + x^{28} + x^{22} + x^{20} + x^{16} + x^{14} + x^{12} + x^6 + x^2, \\
p_{12}(x) &= x^{100} + x^{96} + x^{76} + x^{72} + x^{60} + x^{56} + x^{28} + x^{24} + x^{12} + x^8 + x^4 + 1,
\end{aligned}$$

and the initial terms are $[1, 0, 0, 0, 1, 0, 0, 0, 1, 0, 1, 0, 1, 0, 1, 0]$.

For $\phi(W^e, 0, 1)$:

$$\begin{aligned}
p_0(x) &= x^{96} + x^{94} + x^{93} + x^{91} + x^{90} + x^{88} + x^{86} + x^{85} + x^{83} + x^{81} + x^{79} \\
& + x^{78} + x^{74} + x^{72} + x^{70} + x^{69} + x^{68} + x^{65} + x^{64} + x^{63} + x^{62} + x^{61} \\
& + x^{59} + x^{58} + x^{57} + x^{52} + x^{51} + x^{46} + x^{44} + x^{42} + x^{40} + x^{36} + x^{33} \\
& + x^{28} + x^{25} + x^{23} + x^{21} + x^{20} + x^{18}, \\
p_3(x) &= x^{94} + x^{92} + x^{91} + x^{84} + x^{82} + x^{81} + x^{78} + x^{75} + x^{74} + x^{73} + x^{62} \\
& + x^{60} + x^{59} + x^{54} + x^{51} + x^{50} + x^{49} + x^{22} + x^{20} + x^{19} + x^{14} + x^{11} \\
& + x^{10} + x^9, \\
p_6(x) &= x^{93} + x^{92} + x^{91} + x^{89} + x^{87} + x^{85} + x^{83} + x^{81} + x^{78} + x^{73} + x^{72} \\
& + x^{71} + x^{69} + x^{67} + x^{64} + x^{59} + x^{58} + x^{56} + x^{53} + x^{52} + x^{50} + x^{43} \\
& + x^{42} + x^{40} + x^{39} + x^{38} + x^{37} + x^{35} + x^{32} + x^{29} + x^{28} + x^{26} + x^{25} \\
& + x^{24} + x^{23} + x^{20} + x^{19} + x^{18} + x^{17} + x^{14} + x^{11} + x^{10} + x^9 + x^6, \\
p_9(x) &= x^{92} + x^{89} + x^{88} + x^{87} + x^{82} + x^{80} + x^{79} + x^{76} + x^{73} + x^{71} + x^{69} \\
& + x^{67} + x^{66} + x^{65} + x^{62} + x^{59} + x^{58} + x^{57} + x^{52} + x^{50} + x^{49} + x^{48} \\
& + x^{45} + x^{43} + x^{42} + x^{41} + x^{40} + x^{38} + x^{37} + x^{36} + x^{34} + x^{33} + x^{30} \\
& + x^{28} + x^{27} + x^{26} + x^{24} + x^{23} + x^{20} + x^{17} + x^{15} + x^{13} + x^{12} + x^{11} \\
& + x^{10} + x^8 + x^7 + x^6 + x^4 + x^3, \\
p_{12}(x) &= x^{91} + x^{90} + x^{88} + x^{79} + x^{78} + x^{76} + x^{75} + x^{74} + x^{72} + x^{71} + x^{70} \\
& + x^{68} + x^{67} + x^{66} + x^{64} + x^{63} + x^{62} + x^{60} + x^{55} + x^{54} + x^{52} + x^{51} \\
& + x^{50} + x^{48} + x^{43} + x^{42} + x^{40} + x^{31} + x^{30} + x^{28} + x^{27} + x^{26} + x^{24} \\
& + x^{23} + x^{22} + x^{20} + x^{19} + x^{18} + x^{16} + x^{15} + x^{14} + x^{12} + x^7 + x^6 + x^4 \\
& + x^3 + x^2 + 1,
\end{aligned}$$

and the initial terms are $[0, 1, 1, 1, 1, 0, 0, 1, 1, 0, 1, 0, 1, 0, 1, 1]$.

For $\phi(W^e, 1, 0)$:

$$\begin{aligned}
p_0(x) &= x^{96} + x^{94} + x^{93} + x^{91} + x^{90} + x^{88} + x^{86} + x^{85} + x^{83} + x^{81} + x^{79} \\
&\quad + x^{78} + x^{74} + x^{72} + x^{70} + x^{69} + x^{68} + x^{65} + x^{64} + x^{63} + x^{62} + x^{61} \\
&\quad + x^{59} + x^{58} + x^{57} + x^{52} + x^{51} + x^{46} + x^{44} + x^{42} + x^{40} + x^{36} + x^{33} \\
&\quad + x^{28} + x^{25} + x^{23} + x^{21} + x^{20} + x^{18}, \\
p_3(x) &= x^{94} + x^{92} + x^{91} + x^{84} + x^{82} + x^{81} + x^{78} + x^{75} + x^{74} + x^{73} + x^{62} \\
&\quad + x^{60} + x^{59} + x^{54} + x^{51} + x^{50} + x^{49} + x^{22} + x^{20} + x^{19} + x^{14} + x^{11} \\
&\quad + x^{10} + x^9, \\
p_6(x) &= x^{93} + x^{92} + x^{91} + x^{89} + x^{87} + x^{85} + x^{83} + x^{81} + x^{78} + x^{73} + x^{72} \\
&\quad + x^{71} + x^{69} + x^{67} + x^{64} + x^{59} + x^{58} + x^{56} + x^{53} + x^{52} + x^{50} + x^{43} \\
&\quad + x^{42} + x^{40} + x^{39} + x^{38} + x^{37} + x^{35} + x^{32} + x^{29} + x^{28} + x^{26} + x^{25} \\
&\quad + x^{24} + x^{23} + x^{20} + x^{19} + x^{18} + x^{17} + x^{14} + x^{11} + x^{10} + x^9 + x^6, \\
p_9(x) &= x^{92} + x^{89} + x^{88} + x^{87} + x^{82} + x^{80} + x^{79} + x^{76} + x^{73} + x^{71} + x^{69} \\
&\quad + x^{67} + x^{66} + x^{65} + x^{62} + x^{59} + x^{58} + x^{57} + x^{52} + x^{50} + x^{49} + x^{48} \\
&\quad + x^{45} + x^{43} + x^{42} + x^{41} + x^{40} + x^{38} + x^{37} + x^{36} + x^{34} + x^{33} + x^{30} \\
&\quad + x^{28} + x^{27} + x^{26} + x^{24} + x^{23} + x^{20} + x^{17} + x^{15} + x^{13} + x^{12} + x^{11} \\
&\quad + x^{10} + x^8 + x^7 + x^6 + x^4 + x^3, \\
p_{12}(x) &= x^{91} + x^{90} + x^{88} + x^{79} + x^{78} + x^{76} + x^{75} + x^{74} + x^{72} + x^{71} + x^{70} \\
&\quad + x^{68} + x^{67} + x^{66} + x^{64} + x^{63} + x^{62} + x^{60} + x^{55} + x^{54} + x^{52} + x^{51} \\
&\quad + x^{50} + x^{48} + x^{43} + x^{42} + x^{40} + x^{31} + x^{30} + x^{28} + x^{27} + x^{26} + x^{24} \\
&\quad + x^{23} + x^{22} + x^{20} + x^{19} + x^{18} + x^{16} + x^{15} + x^{14} + x^{12} + x^7 + x^6 + x^4 \\
&\quad + x^3 + x^2 + 1,
\end{aligned}$$

and the initial terms are $[0, 1, 1, 1, 1, 0, 0, 1, 1, 0, 1, 0, 1, 0, 1, 1]$.

For $\phi(W^e, 1, 1)$:

$$\begin{aligned}
p_0(x) &= x^{84} + x^{82} + x^{80} + x^{78} + x^{76} + x^{74} + x^{72} + x^{64} + x^{62} + x^{58} + x^{54} \\
&\quad + x^{48} + x^{46} + x^{44} + x^{42} + x^{40} + x^{38} + x^{36} + x^{34} + x^{32} + x^{30} + x^{28} \\
&\quad + x^{24}, \\
p_3(x) &= x^{78} + x^{76} + x^{72} + x^{66} + x^{64} + x^{62} + x^{60} + x^{52} + x^{46} + x^{42} + x^{36} \\
&\quad + x^{34} + x^{24} + x^{22} + x^{16} + x^{14} + x^{10} + x^6, \\
p_6(x) &= x^{80} + x^{76} + x^{72} + x^{70} + x^{64} + x^{62} + x^{60} + x^{56} + x^{54} + x^{52} + x^{48} \\
&\quad + x^{46} + x^{44} + x^{42} + x^{40} + x^{38} + x^{36} + x^{28} + x^{26} + x^{24} + x^{20} + x^{18} \\
&\quad + x^6 + x^4 + x^2 + 1, \\
p_9(x) &= x^{82} + x^{76} + x^{74} + x^{70} + x^{66} + x^{64} + x^{52} + x^{46} + x^{44} + x^{40} + x^{38} \\
&\quad + x^{32} + x^{20} + x^{18} + x^{14} + x^{10} + x^8 + x^2, \\
p_{12}(x) &= x^{82} + x^{80} + x^{78} + x^{74} + x^{70} + x^{66} + x^{62} + x^{58} + x^{54} + x^{48} + x^{34}
\end{aligned}$$

$$+ x^{32} + x^{30} + x^{26} + x^{22} + x^{18} + x^{14} + x^{10} + x^6 + 1,$$

and the initial terms are $[0, 0, 1, 0, 1, 0, 0, 0, 0, 1, 0, 0, 0, 0, 0]$.

For $\phi(M^o, 0, 0)$:

$$\begin{aligned} p_0(x) = & x^{96} + x^{95} + x^{92} + x^{90} + x^{87} + x^{86} + x^{84} + x^{81} + x^{79} + x^{75} + x^{68} \\ & + x^{66} + x^{65} + x^{61} + x^{59} + x^{57} + x^{54} + x^{53} + x^{51} + x^{50} + x^{47} + x^{44} \\ & + x^{42} + x^{40} + x^{39} + x^{38} + x^{36} + x^{35} + x^{28} + x^{27} + x^{26} + x^{25} + x^{23} \\ & + x^{20} + x^{18} + x^{17} + x^{16} + x^{15} + x^{12}, \end{aligned}$$

$$\begin{aligned} p_3(x) = & x^{95} + x^{93} + x^{92} + x^{91} + x^{90} + x^{89} + x^{88} + x^{85} + x^{84} + x^{83} + x^{81} \\ & + x^{79} + x^{78} + x^{76} + x^{75} + x^{74} + x^{72} + x^{71} + x^{70} + x^{68} + x^{65} + x^{64} \\ & + x^{61} + x^{56} + x^{53} + x^{49} + x^{44} + x^{42} + x^{40} + x^{39} + x^{37} + x^{36} + x^{33} \\ & + x^{31} + x^{30} + x^{29} + x^{28} + x^{27} + x^{26} + x^{25} + x^{24} + x^{22} + x^{21} + x^{20} \\ & + x^{19} + x^{16} + x^{15} + x^{13} + x^{11} + x^8, \end{aligned}$$

$$\begin{aligned} p_6(x) = & x^{95} + x^{93} + x^{92} + x^{91} + x^{88} + x^{86} + x^{85} + x^{84} + x^{83} + x^{82} + x^{80} \\ & + x^{79} + x^{78} + x^{76} + x^{75} + x^{73} + x^{72} + x^{71} + x^{68} + x^{65} + x^{63} + x^{62} \\ & + x^{61} + x^{59} + x^{55} + x^{54} + x^{50} + x^{49} + x^{48} + x^{47} + x^{44} + x^{40} + x^{39} \\ & + x^{38} + x^{36} + x^{35} + x^{34} + x^{33} + x^{32} + x^{30} + x^{28} + x^{24} + x^{23} + x^{21} \\ & + x^{20} + x^{19} + x^{17} + x^{14} + x^{11} + x^9 + x^6 + x^3 + x^2 + 1, \end{aligned}$$

$$\begin{aligned} p_9(x) = & x^{95} + x^{94} + x^{92} + x^{91} + x^{90} + x^{88} + x^{83} + x^{82} + x^{80} + x^{79} + x^{78} \\ & + x^{76} + x^{75} + x^{74} + x^{72} + x^{71} + x^{70} + x^{68} + x^{67} + x^{66} + x^{64} + x^{55} \\ & + x^{54} + x^{52} + x^{47} + x^{46} + x^{44} + x^{43} + x^{42} + x^{40} + x^{35} + x^{34} + x^{32} \\ & + x^{31} + x^{30} + x^{28} + x^{27} + x^{26} + x^{24} + x^{23} + x^{22} + x^{20} + x^{19} + x^{18} \\ & + x^{16} + x^7 + x^6 + x^4, \end{aligned}$$

$$\begin{aligned} p_{12}(x) = & x^{95} + x^{94} + x^{92} + x^{83} + x^{82} + x^{80} + x^{63} + x^{62} + x^{60} + x^{51} + x^{50} \\ & + x^{48} + x^{47} + x^{46} + x^{44} + x^{35} + x^{34} + x^{32} + x^{15} + x^{14} + x^{12} + x^3 \\ & + x^2 + 1, \end{aligned}$$

and the initial terms are $[1, 0, 1, 0, 1, 0, 0, 0, 1, 1, 1, 0, 1, 0, 1, 1]$.

For $\phi(M^o, 0, 1)$:

$$\begin{aligned} p_0(x) = & x^{108} + x^{105} + x^{103} + x^{102} + x^{100} + x^{95} + x^{94} + x^{89} + x^{88} + x^{87} + x^{86} \\ & + x^{84} + x^{83} + x^{82} + x^{80} + x^{78} + x^{77} + x^{76} + x^{75} + x^{73} + x^{72} + x^{71} \\ & + x^{70} + x^{65} + x^{62} + x^{60} + x^{59} + x^{58} + x^{57} + x^{53} + x^{47} + x^{46} + x^{45} \\ & + x^{44} + x^{43} + x^{42} + x^{40} + x^{38} + x^{37} + x^{34} + x^{32} + x^{30} + x^{29} + x^{27} \\ & + x^{24}, \end{aligned}$$

$$\begin{aligned} p_3(x) = & x^{101} + x^{100} + x^{95} + x^{93} + x^{91} + x^{90} + x^{88} + x^{87} + x^{85} + x^{82} + x^{81} \\ & + x^{80} + x^{78} + x^{77} + x^{76} + x^{75} + x^{74} + x^{73} + x^{69} + x^{68} + x^{63} + x^{60} \\ & + x^{59} + x^{58} + x^{56} + x^{53} + x^{52} + x^{51} + x^{49} + x^{46} + x^{42} + x^{38} + x^{34} \end{aligned}$$

$$\begin{aligned}
& + x^{30} + x^{29} + x^{28} + x^{26} + x^{23} + x^{22} + x^{20} + x^{19} + x^{16} + x^{14} + x^{13} \\
& + x^{12} + x^{11} + x^{10} + x^9, \\
p_6(x) &= x^{102} + x^{100} + x^{90} + x^{84} + x^{78} + x^{74} + x^{72} + x^{70} + x^{66} + x^{62} + x^{58} \\
& + x^{56} + x^{52} + x^{50} + x^{46} + x^{42} + x^{38} + x^{36} + x^{34} + x^{30} + x^{24} + x^{18} \\
& + x^8 + x^6, \\
p_9(x) &= x^{103} + x^{102} + x^{101} + x^{100} + x^{97} + x^{92} + x^{90} + x^{89} + x^{88} + x^{87} + x^{86} \\
& + x^{84} + x^{83} + x^{82} + x^{78} + x^{74} + x^{71} + x^{69} + x^{68} + x^{66} + x^{65} + x^{62} \\
& + x^{60} + x^{57} + x^{56} + x^{54} + x^{53} + x^{51} + x^{49} + x^{44} + x^{42} + x^{41} + x^{40} \\
& + x^{39} + x^{38} + x^{36} + x^{35} + x^{34} + x^{30} + x^{26} + x^{23} + x^{21} + x^{20} + x^{18} \\
& + x^{17} + x^{14} + x^{12} + x^9 + x^8 + x^7 + x^4 + x^3, \\
p_{12}(x) &= x^{104} + x^{100} + x^{88} + x^{80} + x^{72} + x^{68} + x^{52} + x^{48} + x^{40} + x^{32} + x^{24} \\
& + x^{20} + x^8 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 1, 1, 1, 0, 1, 0, 1, 1, 0, 1, 0]$.

For $\phi(M^o, 1, 0)$:

$$\begin{aligned}
p_0(x) &= x^{90} + x^{89} + x^{88} + x^{86} + x^{84} + x^{81} + x^{79} + x^{78} + x^{75} + x^{74} + x^{73} \\
& + x^{72} + x^{68} + x^{65} + x^{62} + x^{61} + x^{58} + x^{56} + x^{55} + x^{54} + x^{51} + x^{47} \\
& + x^{46} + x^{44} + x^{40} + x^{38} + x^{36} + x^{32} + x^{31} + x^{28} + x^{27} + x^{26} + x^{25} \\
& + x^{24} + x^{23} + x^{20} + x^{18}, \\
p_3(x) &= x^{89} + x^{88} + x^{85} + x^{84} + x^{83} + x^{80} + x^{78} + x^{77} + x^{76} + x^{75} + x^{74} \\
& + x^{73} + x^{57} + x^{56} + x^{53} + x^{52} + x^{51} + x^{49} + x^{46} + x^{42} + x^{38} + x^{34} \\
& + x^{30} + x^{26} + x^{22} + x^{18} + x^{17} + x^{16} + x^{14} + x^{13} + x^{12} + x^{11} + x^{10} \\
& + x^9, \\
p_6(x) &= x^{90} + x^{88} + x^{86} + x^{84} + x^{82} + x^{80} + x^{78} + x^{70} + x^{68} + x^{66} + x^{64} \\
& + x^{56} + x^{50} + x^{40} + x^{36} + x^{34} + x^{32} + x^{26} + x^{22} + x^{20} + x^{16} + x^{14} \\
& + x^8 + x^6, \\
p_9(x) &= x^{91} + x^{90} + x^{89} + x^{88} + x^{87} + x^{86} + x^{84} + x^{83} + x^{82} + x^{78} + x^{74} \\
& + x^{70} + x^{66} + x^{62} + x^{59} + x^{57} + x^{56} + x^{55} + x^{52} + x^{51} + x^{43} + x^{42} \\
& + x^{41} + x^{40} + x^{39} + x^{38} + x^{36} + x^{35} + x^{34} + x^{30} + x^{26} + x^{22} + x^{18} \\
& + x^{14} + x^{11} + x^9 + x^8 + x^7 + x^4 + x^3, \\
p_{12}(x) &= x^{92} + x^{80} + x^{60} + x^{48} + x^{44} + x^{32} + x^{12} + 1,
\end{aligned}$$

and the initial terms are $[0, 1, 1, 0, 0, 1, 1, 0, 0, 0, 0, 1, 1, 1, 1, 1]$.

For $\phi(M^o, 1, 1)$:

$$\begin{aligned}
p_0(x) &= x^{96} + x^{92} + x^{91} + x^{88} + x^{87} + x^{86} + x^{83} + x^{81} + x^{80} + x^{77} + x^{73} \\
& + x^{72} + x^{71} + x^{67} + x^{66} + x^{61} + x^{58} + x^{57} + x^{56} + x^{55} + x^{53} + x^{52} \\
& + x^{51} + x^{48} + x^{47} + x^{45} + x^{39} + x^{38} + x^{35} + x^{32} + x^{30},
\end{aligned}$$

$$\begin{aligned}
p_3(x) &= x^{95} + x^{91} + x^{89} + x^{88} + x^{85} + x^{84} + x^{82} + x^{81} + x^{80} + x^{79} + x^{78} \\
&\quad + x^{77} + x^{74} + x^{71} + x^{70} + x^{68} + x^{65} + x^{64} + x^{60} + x^{58} + x^{56} + x^{52} \\
&\quad + x^{51} + x^{49} + x^{48} + x^{44} + x^{42} + x^{40} + x^{39} + x^{37} + x^{36} + x^{33} + x^{31} \\
&\quad + x^{30} + x^{29} + x^{28} + x^{27} + x^{26} + x^{25} + x^{24} + x^{22} + x^{19} + x^{18} + x^{16} \\
&\quad + x^{15} + x^{12}, \\
p_6(x) &= x^{95} + x^{91} + x^{90} + x^{89} + x^{82} + x^{81} + x^{80} + x^{78} + x^{75} + x^{71} + x^{70} \\
&\quad + x^{69} + x^{66} + x^{65} + x^{61} + x^{60} + x^{58} + x^{56} + x^{53} + x^{51} + x^{50} + x^{49} \\
&\quad + x^{46} + x^{40} + x^{37} + x^{36} + x^{35} + x^{33} + x^{32} + x^{31} + x^{29} + x^{28} + x^{27} \\
&\quad + x^{25} + x^{24} + x^{21} + x^{19} + x^{16} + x^{11} + x^8 + x^7 + x^6 + x^4 + x^3 + x^2 + 1, \\
p_9(x) &= x^{95} + x^{94} + x^{92} + x^{91} + x^{90} + x^{88} + x^{83} + x^{82} + x^{80} + x^{79} + x^{78} \\
&\quad + x^{76} + x^{75} + x^{74} + x^{72} + x^{71} + x^{70} + x^{68} + x^{67} + x^{66} + x^{64} + x^{55} \\
&\quad + x^{54} + x^{52} + x^{47} + x^{46} + x^{44} + x^{43} + x^{42} + x^{40} + x^{35} + x^{34} + x^{32} \\
&\quad + x^{31} + x^{30} + x^{28} + x^{27} + x^{26} + x^{24} + x^{23} + x^{22} + x^{20} + x^{19} + x^{18} \\
&\quad + x^{16} + x^7 + x^6 + x^4, \\
p_{12}(x) &= x^{95} + x^{94} + x^{92} + x^{83} + x^{82} + x^{80} + x^{63} + x^{62} + x^{60} + x^{51} + x^{50} \\
&\quad + x^{48} + x^{47} + x^{46} + x^{44} + x^{35} + x^{34} + x^{32} + x^{15} + x^{14} + x^{12} + x^3 \\
&\quad + x^2 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 0, 1, 0, 1, 1, 0, 0, 0, 0, 1, 1]$.

For $\phi(W^\circ, 0, 0)$:

$$\begin{aligned}
p_0(x) &= x^{96} + x^{95} + x^{92} + x^{90} + x^{87} + x^{86} + x^{84} + x^{81} + x^{79} + x^{75} + x^{68} \\
&\quad + x^{66} + x^{65} + x^{61} + x^{59} + x^{57} + x^{54} + x^{53} + x^{51} + x^{50} + x^{47} + x^{44} \\
&\quad + x^{42} + x^{40} + x^{39} + x^{38} + x^{36} + x^{35} + x^{28} + x^{27} + x^{26} + x^{25} + x^{23} \\
&\quad + x^{20} + x^{18} + x^{17} + x^{16} + x^{15} + x^{12}, \\
p_3(x) &= x^{95} + x^{93} + x^{92} + x^{91} + x^{90} + x^{89} + x^{88} + x^{85} + x^{84} + x^{83} + x^{81} \\
&\quad + x^{79} + x^{78} + x^{76} + x^{75} + x^{74} + x^{72} + x^{71} + x^{70} + x^{68} + x^{65} + x^{64} \\
&\quad + x^{61} + x^{56} + x^{53} + x^{49} + x^{44} + x^{42} + x^{40} + x^{39} + x^{37} + x^{36} + x^{33} \\
&\quad + x^{31} + x^{30} + x^{29} + x^{28} + x^{27} + x^{26} + x^{25} + x^{24} + x^{22} + x^{21} + x^{20} \\
&\quad + x^{19} + x^{16} + x^{15} + x^{13} + x^{11} + x^8, \\
p_6(x) &= x^{95} + x^{93} + x^{92} + x^{91} + x^{88} + x^{86} + x^{85} + x^{84} + x^{83} + x^{82} + x^{80} \\
&\quad + x^{79} + x^{78} + x^{76} + x^{75} + x^{73} + x^{72} + x^{71} + x^{68} + x^{65} + x^{63} + x^{62} \\
&\quad + x^{61} + x^{59} + x^{55} + x^{54} + x^{50} + x^{49} + x^{48} + x^{47} + x^{44} + x^{40} + x^{39} \\
&\quad + x^{38} + x^{36} + x^{35} + x^{34} + x^{33} + x^{32} + x^{30} + x^{28} + x^{24} + x^{23} + x^{21} \\
&\quad + x^{20} + x^{19} + x^{17} + x^{14} + x^{11} + x^9 + x^6 + x^3 + x^2 + 1, \\
p_9(x) &= x^{95} + x^{94} + x^{92} + x^{91} + x^{90} + x^{88} + x^{83} + x^{82} + x^{80} + x^{79} + x^{78} \\
&\quad + x^{76} + x^{75} + x^{74} + x^{72} + x^{71} + x^{70} + x^{68} + x^{67} + x^{66} + x^{64} + x^{55}
\end{aligned}$$

$$\begin{aligned}
& + x^{54} + x^{52} + x^{47} + x^{46} + x^{44} + x^{43} + x^{42} + x^{40} + x^{35} + x^{34} + x^{32} \\
& + x^{31} + x^{30} + x^{28} + x^{27} + x^{26} + x^{24} + x^{23} + x^{22} + x^{20} + x^{19} + x^{18} \\
& + x^{16} + x^7 + x^6 + x^4, \\
p_{12}(x) = & x^{95} + x^{94} + x^{92} + x^{83} + x^{82} + x^{80} + x^{63} + x^{62} + x^{60} + x^{51} + x^{50} \\
& + x^{48} + x^{47} + x^{46} + x^{44} + x^{35} + x^{34} + x^{32} + x^{15} + x^{14} + x^{12} + x^3 \\
& + x^2 + 1,
\end{aligned}$$

and the initial terms are $[1, 0, 1, 0, 1, 0, 0, 0, 1, 1, 1, 0, 1, 0, 1, 1]$.

For $\phi(W^o, 0, 1)$:

$$\begin{aligned}
p_0(x) = & x^{90} + x^{89} + x^{88} + x^{86} + x^{84} + x^{81} + x^{79} + x^{78} + x^{75} + x^{74} + x^{73} \\
& + x^{72} + x^{68} + x^{65} + x^{62} + x^{61} + x^{58} + x^{56} + x^{55} + x^{54} + x^{51} + x^{47} \\
& + x^{46} + x^{44} + x^{40} + x^{38} + x^{36} + x^{32} + x^{31} + x^{28} + x^{27} + x^{26} + x^{25} \\
& + x^{24} + x^{23} + x^{20} + x^{18}, \\
p_3(x) = & x^{89} + x^{88} + x^{85} + x^{84} + x^{83} + x^{80} + x^{78} + x^{77} + x^{76} + x^{75} + x^{74} \\
& + x^{73} + x^{57} + x^{56} + x^{53} + x^{52} + x^{51} + x^{49} + x^{46} + x^{42} + x^{38} + x^{34} \\
& + x^{30} + x^{26} + x^{22} + x^{18} + x^{17} + x^{16} + x^{14} + x^{13} + x^{12} + x^{11} + x^{10} \\
& + x^9, \\
p_6(x) = & x^{90} + x^{88} + x^{86} + x^{84} + x^{82} + x^{80} + x^{78} + x^{70} + x^{68} + x^{66} + x^{64} \\
& + x^{56} + x^{50} + x^{40} + x^{36} + x^{34} + x^{32} + x^{26} + x^{22} + x^{20} + x^{16} + x^{14} \\
& + x^8 + x^6, \\
p_9(x) = & x^{91} + x^{90} + x^{89} + x^{88} + x^{87} + x^{86} + x^{84} + x^{83} + x^{82} + x^{78} + x^{74} \\
& + x^{70} + x^{66} + x^{62} + x^{59} + x^{57} + x^{56} + x^{55} + x^{52} + x^{51} + x^{43} + x^{42} \\
& + x^{41} + x^{40} + x^{39} + x^{38} + x^{36} + x^{35} + x^{34} + x^{30} + x^{26} + x^{22} + x^{18} \\
& + x^{14} + x^{11} + x^9 + x^8 + x^7 + x^4 + x^3, \\
p_{12}(x) = & x^{92} + x^{80} + x^{60} + x^{48} + x^{44} + x^{32} + x^{12} + 1,
\end{aligned}$$

and the initial terms are $[0, 1, 1, 0, 0, 1, 1, 0, 0, 0, 0, 1, 1, 1, 1, 1]$.

For $\phi(W^o, 1, 0)$:

$$\begin{aligned}
p_0(x) = & x^{108} + x^{105} + x^{103} + x^{102} + x^{100} + x^{95} + x^{94} + x^{89} + x^{88} + x^{87} + x^{86} \\
& + x^{84} + x^{83} + x^{82} + x^{80} + x^{78} + x^{77} + x^{76} + x^{75} + x^{73} + x^{72} + x^{71} \\
& + x^{70} + x^{65} + x^{62} + x^{60} + x^{59} + x^{58} + x^{57} + x^{53} + x^{47} + x^{46} + x^{45} \\
& + x^{44} + x^{43} + x^{42} + x^{40} + x^{38} + x^{37} + x^{34} + x^{32} + x^{30} + x^{29} + x^{27} \\
& + x^{24}, \\
p_3(x) = & x^{101} + x^{100} + x^{95} + x^{93} + x^{91} + x^{90} + x^{88} + x^{87} + x^{85} + x^{82} + x^{81} \\
& + x^{80} + x^{78} + x^{77} + x^{76} + x^{75} + x^{74} + x^{73} + x^{69} + x^{68} + x^{63} + x^{60} \\
& + x^{59} + x^{58} + x^{56} + x^{53} + x^{52} + x^{51} + x^{49} + x^{46} + x^{42} + x^{38} + x^{34} \\
& + x^{30} + x^{29} + x^{28} + x^{26} + x^{23} + x^{22} + x^{20} + x^{19} + x^{16} + x^{14} + x^{13}
\end{aligned}$$

$$\begin{aligned}
& + x^{12} + x^{11} + x^{10} + x^9, \\
p_6(x) &= x^{102} + x^{100} + x^{90} + x^{84} + x^{78} + x^{74} + x^{72} + x^{70} + x^{66} + x^{62} + x^{58} \\
& + x^{56} + x^{52} + x^{50} + x^{46} + x^{42} + x^{38} + x^{36} + x^{34} + x^{30} + x^{24} + x^{18} \\
& + x^8 + x^6, \\
p_9(x) &= x^{103} + x^{102} + x^{101} + x^{100} + x^{97} + x^{92} + x^{90} + x^{89} + x^{88} + x^{87} + x^{86} \\
& + x^{84} + x^{83} + x^{82} + x^{78} + x^{74} + x^{71} + x^{69} + x^{68} + x^{66} + x^{65} + x^{62} \\
& + x^{60} + x^{57} + x^{56} + x^{54} + x^{53} + x^{51} + x^{49} + x^{44} + x^{42} + x^{41} + x^{40} \\
& + x^{39} + x^{38} + x^{36} + x^{35} + x^{34} + x^{30} + x^{26} + x^{23} + x^{21} + x^{20} + x^{18} \\
& + x^{17} + x^{14} + x^{12} + x^9 + x^8 + x^7 + x^4 + x^3, \\
p_{12}(x) &= x^{104} + x^{100} + x^{88} + x^{80} + x^{72} + x^{68} + x^{52} + x^{48} + x^{40} + x^{32} + x^{24} \\
& + x^{20} + x^8 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 1, 1, 1, 0, 1, 0, 1, 1, 0, 1, 0]$.

For $\phi(W^o, 1, 1)$:

$$\begin{aligned}
p_0(x) &= x^{96} + x^{92} + x^{91} + x^{88} + x^{87} + x^{86} + x^{83} + x^{81} + x^{80} + x^{77} + x^{73} \\
& + x^{72} + x^{71} + x^{67} + x^{66} + x^{61} + x^{58} + x^{57} + x^{56} + x^{55} + x^{53} + x^{52} \\
& + x^{51} + x^{48} + x^{47} + x^{45} + x^{39} + x^{38} + x^{35} + x^{32} + x^{30}, \\
p_3(x) &= x^{95} + x^{91} + x^{89} + x^{88} + x^{85} + x^{84} + x^{82} + x^{81} + x^{80} + x^{79} + x^{78} \\
& + x^{77} + x^{74} + x^{71} + x^{70} + x^{68} + x^{65} + x^{64} + x^{60} + x^{58} + x^{56} + x^{52} \\
& + x^{51} + x^{49} + x^{48} + x^{44} + x^{42} + x^{40} + x^{39} + x^{37} + x^{36} + x^{33} + x^{31} \\
& + x^{30} + x^{29} + x^{28} + x^{27} + x^{26} + x^{25} + x^{24} + x^{22} + x^{19} + x^{18} + x^{16} \\
& + x^{15} + x^{12}, \\
p_6(x) &= x^{95} + x^{91} + x^{90} + x^{89} + x^{82} + x^{81} + x^{80} + x^{78} + x^{75} + x^{71} + x^{70} \\
& + x^{69} + x^{66} + x^{65} + x^{61} + x^{60} + x^{58} + x^{56} + x^{53} + x^{51} + x^{50} + x^{49} \\
& + x^{46} + x^{40} + x^{37} + x^{36} + x^{35} + x^{33} + x^{32} + x^{31} + x^{29} + x^{28} + x^{27} \\
& + x^{25} + x^{24} + x^{21} + x^{19} + x^{16} + x^{11} + x^8 + x^7 + x^6 + x^4 + x^3 + x^2 + 1, \\
p_9(x) &= x^{95} + x^{94} + x^{92} + x^{91} + x^{90} + x^{88} + x^{83} + x^{82} + x^{80} + x^{79} + x^{78} \\
& + x^{76} + x^{75} + x^{74} + x^{72} + x^{71} + x^{70} + x^{68} + x^{67} + x^{66} + x^{64} + x^{55} \\
& + x^{54} + x^{52} + x^{47} + x^{46} + x^{44} + x^{43} + x^{42} + x^{40} + x^{35} + x^{34} + x^{32} \\
& + x^{31} + x^{30} + x^{28} + x^{27} + x^{26} + x^{24} + x^{23} + x^{22} + x^{20} + x^{19} + x^{18} \\
& + x^{16} + x^7 + x^6 + x^4, \\
p_{12}(x) &= x^{95} + x^{94} + x^{92} + x^{83} + x^{82} + x^{80} + x^{63} + x^{62} + x^{60} + x^{51} + x^{50} \\
& + x^{48} + x^{47} + x^{46} + x^{44} + x^{35} + x^{34} + x^{32} + x^{15} + x^{14} + x^{12} + x^3 \\
& + x^2 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 0, 1, 0, 1, 1, 0, 0, 0, 0, 1, 1]$.

Below is the transition function and output function of an 2-automaton that generates $T = M_{0,0}^o$:

Transition function $(n, j) \mapsto \delta(n, j)$ ($\Lambda(n) := [\delta(n, 0), \delta(n, 1)]$):

n	$\Lambda(n)$	n	$\Lambda(n)$	n	$\Lambda(n)$	n	$\Lambda(n)$	n	$\Lambda(n)$
0	[1, 2]	76	[137, 135]	152	[91, 220]	228	[302, 171]	304	[201, 234]
1	[3, 4]	77	[138, 113]	153	[225, 226]	229	[303, 137]	305	[230, 331]
2	[5, 6]	78	[139, 140]	154	[227, 176]	230	[304, 248]	306	[232, 88]
3	[7, 8]	79	[141, 106]	155	[221, 228]	231	[305, 53]	307	[328, 24]
4	[9, 10]	80	[142, 55]	156	[76, 229]	232	[306, 42]	308	[58, 228]
5	[11, 12]	81	[143, 110]	157	[230, 218]	233	[307, 307]	309	[338, 338]
6	[13, 14]	82	[39, 144]	158	[87, 231]	234	[308, 255]	310	[347, 348]
7	[15, 16]	83	[145, 146]	159	[205, 226]	235	[241, 52]	311	[104, 349]
8	[17, 18]	84	[147, 148]	160	[232, 231]	236	[309, 136]	312	[350, 254]
9	[19, 20]	85	[149, 150]	161	[193, 75]	237	[167, 137]	313	[351, 105]
10	[21, 22]	86	[151, 9]	162	[233, 234]	238	[12, 143]	314	[352, 258]
11	[23, 24]	87	[152, 153]	163	[235, 84]	239	[310, 298]	315	[353, 143]
12	[25, 26]	88	[154, 121]	164	[15, 99]	240	[159, 272]	316	[269, 284]
13	[27, 28]	89	[155, 30]	165	[207, 236]	241	[216, 193]	317	[41, 154]
14	[29, 21]	90	[156, 157]	166	[237, 209]	242	[58, 311]	318	[354, 139]
15	[30, 31]	91	[158, 159]	167	[236, 237]	243	[312, 18]	319	[355, 117]
16	[32, 33]	92	[126, 114]	168	[238, 224]	244	[313, 87]	320	[356, 281]
17	[34, 11]	93	[160, 104]	169	[239, 184]	245	[314, 99]	321	[357, 244]
18	[35, 36]	94	[108, 161]	170	[213, 57]	246	[219, 206]	322	[358, 306]
19	[37, 33]	95	[162, 163]	171	[220, 240]	247	[315, 316]	323	[2, 301]
20	[38, 39]	96	[164, 103]	172	[146, 241]	248	[317, 87]	324	[51, 153]
21	[40, 41]	97	[165, 166]	173	[31, 242]	249	[318, 199]	325	[359, 169]
22	[42, 43]	98	[166, 167]	174	[8, 45]	250	[319, 96]	326	[161, 146]
23	[44, 45]	99	[168, 169]	175	[243, 244]	251	[320, 76]	327	[360, 47]
24	[46, 47]	100	[170, 115]	176	[245, 246]	252	[94, 200]	328	[361, 132]
25	[48, 49]	101	[153, 171]	177	[247, 248]	253	[321, 322]	329	[362, 103]
26	[50, 51]	102	[47, 142]	178	[249, 246]	254	[323, 322]	330	[363, 250]
27	[52, 53]	103	[102, 172]	179	[250, 251]	255	[324, 21]	331	[364, 306]
28	[54, 55]	104	[99, 173]	180	[148, 252]	256	[68, 182]	332	[365, 9]
29	[56, 42]	105	[174, 74]	181	[253, 254]	257	[325, 20]	333	[264, 277]
30	[57, 58]	106	[98, 98]	182	[10, 255]	258	[326, 71]	334	[366, 39]
31	[59, 29]	107	[98, 97]	183	[256, 257]	259	[84, 79]	335	[115, 30]
32	[60, 61]	108	[175, 176]	184	[258, 10]	260	[95, 327]	336	[367, 274]
33	[62, 63]	109	[177, 178]	185	[106, 157]	261	[229, 75]	337	[368, 134]
34	[64, 65]	110	[177, 26]	186	[259, 142]	262	[328, 329]	338	[369, 135]
35	[66, 28]	111	[179, 180]	187	[260, 261]	263	[320, 216]	339	[370, 275]
36	[67, 58]	112	[181, 182]	188	[262, 34]	264	[330, 186]	340	[119, 148]
37	[68, 69]	113	[183, 184]	189	[263, 40]	265	[208, 237]	341	[284, 299]
38	[70, 71]	114	[185, 186]	190	[36, 264]	266	[217, 331]	342	[242, 371]
39	[27, 72]	115	[187, 22]	191	[134, 123]	267	[211, 61]	343	[372, 291]
40	[73, 74]	116	[188, 89]	192	[7, 119]	268	[332, 333]	344	[373, 268]

41	[75, 76]	117	[189, 93]	193	[136, 265]	269	[221, 311]	345	[261, 40]
42	[77, 78]	118	[190, 77]	194	[266, 251]	270	[174, 192]	346	[251, 250]
43	[79, 80]	119	[191, 94]	195	[267, 268]	271	[77, 334]	347	[110, 253]
44	[81, 82]	120	[73, 192]	196	[150, 269]	272	[231, 101]	348	[374, 371]
45	[83, 84]	121	[76, 193]	197	[270, 161]	273	[223, 335]	349	[172, 80]
46	[85, 86]	122	[93, 194]	198	[271, 272]	274	[336, 63]	350	[23, 329]
47	[87, 88]	123	[195, 196]	199	[273, 274]	275	[206, 240]	351	[216, 229]
48	[81, 89]	124	[25, 178]	200	[123, 275]	276	[215, 337]	352	[186, 197]
49	[90, 91]	125	[197, 198]	201	[276, 253]	277	[66, 72]	353	[64, 337]
50	[92, 86]	126	[186, 179]	202	[277, 126]	278	[67, 221]	354	[340, 197]
51	[93, 94]	127	[97, 97]	203	[121, 108]	279	[60, 212]	355	[312, 342]
52	[95, 96]	128	[191, 194]	204	[278, 38]	280	[208, 210]	356	[237, 207]
53	[97, 98]	129	[199, 78]	205	[279, 257]	281	[338, 288]	357	[188, 82]
54	[99, 100]	130	[194, 200]	206	[157, 261]	282	[288, 338]	358	[318, 77]
55	[88, 101]	131	[201, 202]	207	[280, 167]	283	[187, 339]	359	[347, 375]
56	[102, 79]	132	[203, 93]	208	[265, 281]	284	[340, 179]	360	[204, 225]
57	[103, 104]	133	[204, 205]	209	[281, 166]	285	[341, 225]	361	[315, 6]
58	[105, 106]	134	[91, 206]	210	[282, 265]	286	[5, 316]	362	[190, 199]
59	[107, 108]	135	[207, 208]	211	[254, 34]	287	[209, 236]	363	[229, 320]
60	[109, 110]	136	[209, 208]	212	[283, 284]	288	[288, 288]	364	[314, 16]
61	[111, 14]	137	[210, 209]	213	[53, 105]	289	[17, 342]	365	[376, 375]
62	[112, 113]	138	[211, 212]	214	[285, 51]	290	[341, 205]	366	[59, 324]
63	[114, 115]	139	[213, 67]	215	[286, 163]	291	[84, 172]	367	[376, 348]
64	[116, 117]	140	[197, 180]	216	[287, 288]	292	[227, 214]	368	[85, 195]
65	[118, 119]	141	[175, 214]	217	[289, 49]	293	[319, 327]	369	[236, 210]
66	[120, 121]	142	[86, 196]	218	[290, 291]	294	[193, 320]	370	[16, 100]
67	[122, 123]	143	[215, 65]	219	[246, 129]	295	[321, 343]	371	[377, 27]
68	[124, 11]	144	[71, 27]	220	[292, 41]	296	[179, 198]	372	[92, 195]
69	[125, 126]	145	[75, 216]	221	[293, 241]	297	[238, 335]	373	[181, 69]
70	[127, 52]	146	[217, 218]	222	[294, 154]	298	[344, 333]	374	[21, 339]
71	[128, 129]	147	[219, 220]	223	[295, 12]	299	[345, 29]	375	[140, 36]
72	[129, 130]	148	[16, 173]	224	[296, 144]	300	[233, 202]	376	[378, 307]
73	[131, 132]	149	[57, 221]	225	[297, 298]	301	[346, 91]	377	[291, 159]
74	[133, 134]	150	[222, 67]	226	[299, 111]	302	[199, 334]	378	[323, 343]
75	[135, 136]	151	[223, 224]	227	[300, 301]	303	[210, 207]		

Output function $n \mapsto \tau(n)$:

n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$
0	0	55	0	110	0	165	0	220	0	275	1	330	0
1	0	56	0	111	1	166	1	221	1	276	1	331	1
2	1	57	0	112	0	167	1	222	1	277	1	332	1
3	0	58	0	113	1	168	1	223	0	278	1	333	0
4	1	59	1	114	1	169	0	224	1	279	0	334	1
5	1	60	0	115	0	170	0	225	1	280	0	335	0

6	0	61	1	116	0	171	0	226	0	281	1	336	1
7	0	62	0	117	1	172	0	227	0	282	0	337	0
8	0	63	1	118	1	173	1	228	0	283	0	338	1
9	1	64	0	119	0	174	0	229	0	284	0	339	0
10	1	65	1	120	1	175	1	230	1	285	0	340	0
11	1	66	1	121	0	176	1	231	1	286	1	341	0
12	1	67	1	122	1	177	0	232	1	287	1	342	0
13	0	68	1	123	1	178	0	233	0	288	0	343	1
14	0	69	0	124	1	179	1	234	0	289	0	344	0
15	0	70	0	125	0	180	0	235	1	290	0	345	0
16	0	71	0	126	1	181	0	236	1	291	1	346	1
17	0	72	0	127	0	182	1	237	1	292	0	347	0
18	1	73	1	128	0	183	1	238	1	293	1	348	1
19	1	74	1	129	0	184	1	239	0	294	1	349	0
20	0	75	0	130	0	185	1	240	0	295	0	350	1
21	1	76	0	131	1	186	1	241	1	296	1	351	1
22	1	77	1	132	0	187	0	242	0	297	1	352	1
23	1	78	0	133	1	188	0	243	1	298	0	353	0
24	0	79	1	134	0	189	1	244	1	299	0	354	0
25	1	80	0	135	0	190	1	245	1	300	0	355	1
26	1	81	1	136	1	191	0	246	1	301	1	356	1
27	0	82	0	137	0	192	0	247	0	302	0	357	0
28	1	83	0	138	1	193	1	248	0	303	0	358	0
29	0	84	1	139	0	194	0	249	0	304	1	359	0
30	0	85	0	140	0	195	1	250	1	305	1	360	1
31	1	86	0	141	1	196	1	251	1	306	1	361	0
32	0	87	0	142	0	197	0	252	1	307	0	362	1
33	0	88	0	143	1	198	1	253	0	308	0	363	0
34	0	89	1	144	0	199	0	254	1	309	1	364	1
35	1	90	0	145	0	200	1	255	1	310	0	365	1
36	1	91	0	146	0	201	1	256	1	311	1	366	1
37	1	92	1	147	1	202	1	257	0	312	1	367	1
38	0	93	1	148	0	203	0	258	1	313	1	368	0
39	0	94	1	149	0	204	1	259	1	314	1	369	1
40	1	95	0	150	1	205	0	260	0	315	0	370	0
41	0	96	0	151	0	206	1	261	0	316	1	371	1
42	1	97	0	152	0	207	0	262	0	317	0	372	1
43	1	98	1	153	1	208	0	263	1	318	0	373	0
44	1	99	1	154	0	209	1	264	0	319	1	374	1
45	0	100	0	155	1	210	0	265	0	320	1	375	0
46	0	101	1	156	0	211	1	266	0	321	0	376	1
47	0	102	0	157	1	212	0	267	1	322	0	377	1
48	1	103	0	158	0	213	0	268	1	323	1	378	1
49	0	104	1	159	0	214	0	269	1	324	1		

50	1	105	0	160	1	215	1	270	0	325	0	
51	1	106	1	161	1	216	1	271	1	326	1	
52	0	107	1	162	0	217	0	272	1	327	1	
53	0	108	1	163	1	218	0	273	0	328	0	
54	1	109	0	164	0	219	1	274	1	329	1	

UNIVERSITÉ DE STRASBOURG, CNRS, IRMA UMR 7501, F-67000 STRASBOURG, FRANCE

Email address: `guoniu.han@unistra.fr`

SCHOOL OF MATHEMATICS AND STATISTICS, HUAZHONG UNIVERSITY OF SCIENCE AND TECHNOLOGY, WUHAN, PR CHINA

Email address: `huyining@hust.edu.cn`