

PROOF OF THE CONTINUED FRACTION CONJECTURE FOR

$$(a, b) = (z^3 + z^2 + z, z^3 + z^2 + 1)$$

GUO-NIU HAN* AND YINING HU*

ABSTRACT. The continued fraction conjecture claims that the continued fraction defined by the (a, b) -Thue-Morse sequence is algebraic over $\mathbb{F}_2(x)$ for all pairs of distinct elements (a, b) in $\mathbb{F}_2[x] \setminus \mathbb{F}_2$. In this paper we prove the conjecture for $(a, b) = (z^3 + z^2 + z, z^3 + z^2 + 1)$.

1. INTRODUCTION

Let $\mathbb{F}_2$ be the finite field of cardinality 2. Given a sequence of polynomials $\mathbf{a} = (a_0(z), a_1(z), a_2(z), \dots)$ of elements from $\mathbb{F}_2[z] \setminus \mathbb{F}_2$, we define the infinite continued fraction

$$(1.1) \quad \text{CF}(\mathbf{a}(z)) = \frac{1}{a_0(z) + \frac{1}{a_1(z) + \frac{1}{a_2(z) + \frac{1}{\ddots}}}}$$

as the limit of the finite continued fractions

$$(1.2) \quad \text{CF}_n(\mathbf{a}(z)) = \frac{1}{a_0(z) + \frac{1}{a_1(z) + \frac{1}{\ddots + \frac{1}{a_n(z)}}}} \in \mathbb{F}_2((1/z)).$$

Conversely, each power series in $\mathbb{F}_2[[1/z]]$ without constant term

$$(1.3) \quad c_{-1}z^{-1} + c_{-2}z^{-2} + c_{-3}z^{-3} + \dots$$

admits a continued fraction expansion of form (1.1).

Let a and b be two distinct elements from $\mathbb{F}_2[z] \setminus \mathbb{F}_2$. We consider the continued fraction (1.1) associated with the (a, b) -Thue-Morse sequence

$$\mathbf{t} = (t_0(z), t_1(z), t_2(z), \dots) = (a, b, b, a, b, a, a, b, \dots)$$

that we shall denote

$$(1.4) \quad \text{CF}(a, b) := \text{CF}(\mathbf{t}).$$

Date: 9/04/2020 18:54:56 .

2010 Mathematics Subject Classification. 11B85, 11J70, 11B50, 11Y65, 05A15.

Key words and phrases. automatic sequence, continued fraction, Thue-Morse sequence.

* This paper was automatically generated from a computer program developed by the authors, with 7.0 minutes CPU time used.

Theorem 1.1. *When $(a, b) = (z^3 + z^2 + z, z^3 + z^2 + 1)$, the two power series $\text{CF}(a, b)$ and $\text{CF}(b, a)$ are algebraic over $\mathbb{F}_2(z)$, with minimal polynomials of the form*

$$p_4(z)y^4 + p_3(z)y^3 + p_2(z)y^2 + p_1(z)y + p_0(z) = 0$$

where, for $\text{CF}(a, b)$,

$$\begin{aligned} p_0(z) &= z^{18} + z^{14} + z^{13} + z^{10} + z^8 + z^5 + z^4 + z + 1, \\ p_1(z) &= z^{19} + z^{18} + z^{15} + z^{13} + z^{11} + z^{10} + z^6 + z^5 + z^3 + z, \\ p_2(z) &= z^{24} + z^{14} + z^{12} + z^8 + z^6 + z^4 + z^2, \\ p_3(z) &= z^{19} + z^{18} + z^{15} + z^{13} + z^{11} + z^{10} + z^6 + z^5 + z^3 + z, \\ p_4(z) &= z^{18} + z^{13} + z^{12} + z^{10} + z^8 + z^6 + z^5 + z^2 + z, \end{aligned}$$

and for $\text{CF}(b, a)$,

$$\begin{aligned} p_0(z) &= z^{18} + z^{14} + z^{13} + z^{10} + z^8 + z^5 + z, \\ p_1(z) &= z^{19} + z^{18} + z^{15} + z^{13} + z^{11} + z^{10} + z^6 + z^5 + z^3 + z, \\ p_2(z) &= z^{24} + z^{14} + z^{12} + z^8 + z^6 + z^4 + z^2, \\ p_3(z) &= z^{19} + z^{18} + z^{15} + z^{13} + z^{11} + z^{10} + z^6 + z^5 + z^3 + z, \\ p_4(z) &= z^{18} + z^{13} + z^{12} + z^{10} + z^8 + z^6 + z^5 + z^4 + z^2 + z + 1. \end{aligned}$$

2. PROOF

Define the sequences of polynomials $(P_n(z))_{n \geq 0}$ and $(Q_n(z))_{n \geq 0}$ by

$$\begin{pmatrix} P_n(z) & Q_n(z) \\ P_{n-1}(z) & Q_{n-1}(z) \end{pmatrix} := \begin{pmatrix} t_n(z) & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} t_{n-1}(z) & 1 \\ 1 & 0 \end{pmatrix} \cdots \begin{pmatrix} t_0(z) & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}.$$

By the basic properties of continued fractions, we have

$$(2.1) \quad \text{CF}_n(a, b) = \frac{P_n(z)}{Q_n(z)} \in \mathbb{F}_2((1/z)).$$

As formula (2.1) is not efficient for calculating $\text{CF}_n(a, b)$, a fast method is to be derived and described as follows. Define

$$\begin{aligned} M_n(x) &= x^{\deg(t_{2^n-1})} \begin{pmatrix} t_{2^n-1}(1/x) & 1 \\ 1 & 0 \end{pmatrix} \times x^{\deg(t_{2^n-2})} \begin{pmatrix} t_{2^n-2}(1/x) & 1 \\ 1 & 0 \end{pmatrix} \times \\ &\quad \cdots \times x^{\deg(t_0)} \begin{pmatrix} t_0(1/x) & 1 \\ 1 & 0 \end{pmatrix}, \\ W_n(x) &= x^{\deg(\bar{t}_{2^n-1})} \begin{pmatrix} \bar{t}_{2^n-1}(1/x) & 1 \\ 1 & 0 \end{pmatrix} \times x^{\deg(\bar{t}_{2^n-2})} \begin{pmatrix} \bar{t}_{2^n-2}(1/x) & 1 \\ 1 & 0 \end{pmatrix} \times \\ &\quad \cdots \times x^{\deg(\bar{t}_0)} \begin{pmatrix} \bar{t}_0(1/x) & 1 \\ 1 & 0 \end{pmatrix}, \end{aligned}$$

where $\bar{t}$ is the (b, a) -Thue-Morse sequence. By the properties of the Thue-Morse sequence, we have for $n \geq 0$

$$\begin{aligned} M_{n+1}(x) &= W_n(x) \cdot M_n(x), \\ W_{n+1}(x) &= M_n(x) \cdot W_n(x). \end{aligned}$$

Let $x := 1/z$. For each non-zero polynomial $P(z)$, define $\tilde{P}(x)$ to be $P(1/x)$. Then,

$$\text{CF}_n(a, b) = \frac{P_n(z)}{Q_n(z)} = \frac{\tilde{P}_n(x)}{\tilde{Q}_n(x)} \in \mathbb{F}_2((x)) = \mathbb{F}_2((1/z)).$$

Comparing the definition of $M_n(x)$ with the definition of $P_n(x)$ and $Q_n(x)$, we see that

$$\begin{aligned} M_n(x)_{0,1} &= x^{d_n} \tilde{P}_{2^n-1}(x), \\ M_n(x)_{0,0} &= x^{d_n} \tilde{Q}_{2^n-1}(x), \end{aligned}$$

for some positive integer d_n . Hence,

$$(2.2) \quad \text{CF}_{2^{2^n}-1}(a, b) = \frac{\tilde{P}_{2^{2^n}-1}(x)}{\tilde{Q}_{2^{2^n}-1}(x)} = \frac{M_{2^n}(x)_{0,1}}{M_{2^n}(x)_{0,0}}.$$

Therefore, by the convergence theorem of continued fraction, the algebraicity of $\text{CF}(a, b)$ will be established if it is shown that both $M_{2^n}(x)_{0,1}$ and $M_{2^n}(x)_{0,0}$ converge to algebraic series in $\mathbb{F}_2[[x]]$.

Actually, we will prove that for all $0 \leq i, j \leq 1$ the four sequences $(M_{2^n}(x)_{i,j})_n$, $(M_{2^{n+1}}(x)_{i,j})_n$, $(W_{2^n}(x)_{i,j})_n$, and $(W_{2^{n+1}}(x)_{i,j})_n$ converge to algebraic series in $\mathbb{F}_2[[x]]$. For this purpose, we define four 2×2 matrices M^e, M^o, W^e, W^o as follows: For each $T \in \{M^e, M^o, W^e, W^o\}$ and $0 \leq i, j \leq 1$, $T_{i,j}$ is defined to be the unique solution in $\mathbb{F}_2[[x]]$ of the polynomial $\phi(T, i, j)$ under certain initial conditions; the polynomials $\phi(T, i, j)$ and initial conditions are given in Section 3. We will prove that these four matrices, whose components are algebraic by definition, are the limits of $(M_{2^n}(x))_n$, $(M_{2^{n+1}}(x))_n$, $(W_{2^n}(x))_n$, and $(W_{2^{n+1}}(x))_n$.

Let us explain how the polynomials $\phi(T, i, j)$ and initial conditions are found, and why the solutions exist and are unique. For $0 \leq i, j \leq 1$, the coefficients of the polynomial $\phi(M^e, i, j)$ (resp. $\phi(M^o, i, j)$, $\phi(W^e, i, j)$, and $\phi(W^o, i, j)$) are the Padé-Hermite approximants of type

$$(150, 150, 150, 150, 150)$$

of the vector

$$(1, T^3, T^6, T^9, T^{12}),$$

where $T = M_{12,i,j}$ (resp. $M_{11,i,j}$, $W_{12,i,j}$, and $W_{11,i,j}$), found by the Derksen algorithm. We take the first 8 terms of $M_{12,i,j}$ (resp. $M_{11,i,j}$, $W_{12,i,j}$, and $W_{11,i,j}$) as initial conditions for $\phi(M^e, i, j)$ (resp. $\phi(M^o, i, j)$, $\phi(W^e, i, j)$, and $\phi(W^o, i, j)$). The following fact will be used to ensure that the solution exists and is unique: let $P(x, y) \in \mathbb{F}_2[x, y]$ and for each series $f(x) = \sum_{n=0}^{\infty} a_n x^n \in \mathbb{F}_2[[x]]$ denote the partial sum $\sum_{j=0}^{n-1} a_j x^j$ by $f_n(x)$ for $n \geq 0$. If for some $n \geq 0$ and $a_0, a_1, \dots, a_{n-1} \in \mathbb{F}_2$ $P(x, \sum_{j=0}^{n-1} a_j x^j) = O(x^n)$ and $Q(x, y) := P(x, \sum_{j=0}^{n-1} a_j x^j + x^n y)$ can be written as $x^m \sum_{j=0}^{\infty} q_j(x) y^j$ where $q_j(x)$ are polynomials for $j \geq 0$, $q_1(0) = 1$, and $q_j(0) = 0$ for $j > 1$, then there exists a unique solution $f(x) \in \mathbb{F}_2[[x]]$ of $P(x, f(x)) = 0$ that satisfies the initial condition $f_n(x) = \sum_{j=0}^{n-1} a_j x^j$.

We state two lemmas concerning the four matrices M^e, M^o, W^e, W^o . The first one is about relations between them; the second, about the structure of the each matrix.

Lemma 2.1. *We have*

$$(2.3) \quad M^e = W^o \cdot M^o,$$

$$(2.4) \quad M^o = W^e \cdot M^e,$$

$$(2.5) \quad W^e = M^o \cdot W^o,$$

$$(2.6) \quad W^o = M^e \cdot W^e.$$

Proof. We give the proof of the identity

$$M_{0,0}^e = W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o,$$

the proofs of the others being similar.

First, we compute the minimal polynomials of $W_{0,0}^o M_{0,0}^o$ and $W_{0,1}^o M_{1,0}^o$. We know that

$$P(x, y) = \text{Res}_z (\phi(W^o, 0, 0)(x, z), z^{12} \cdot \phi(M^o, 0, 0)(x, y/z))$$

is an annihilating polynomial of $W_{0,0}^o M_{0,0}^o$. We use Padé-Hermite approximation to find a candidate for the minimal polynomial of $W_{0,0}^o M_{0,0}^o$, that will be called $\phi_0(x, y)$. To prove that $\phi_0(x, y)$ is indeed the minimal polynomial, it suffices to prove that it is an irreducible factor of $P(x, y)$ of multiplicity m and that $Q(x, y) := P(x, y)/\phi_0(x, y)^m$ is not an annihilating polynomial of $W_{0,0}^o M_{0,0}^o$. We verify the first point directly. For the second point, we truncate $W_{0,0}^o M_{0,0}^o$ to order 540 and substitute it for y in $Q(x, y)$. We get a series of valuation less than 540, which proves that $Q(x, y)$ is not an annihilating polynomial of $W_{0,0}^o M_{0,0}^o$. We find the minimal polynomial $\phi_1(x, y)$ of $W_{0,1}^o M_{1,0}^o$ in a similar way.

Now we prove that $\phi(M^e, 0, 0)$ is the minimal polynomial of $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$. We know that

$$S(x, y) = \text{Res}_z (\phi_0(x, z), \phi_1(x, y + z))$$

is an annihilating of $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$. We verify that $\phi(M^e, 0, 0)$ is an irreducible factor of $S(x, y)$ of multiplicity μ , and that $Q(x, y) := S(x, y)/\phi(M^e, 0, 0)^\mu$ is not an annihilating polynomial of $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$. To see the last point, we truncate $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$ to order 660 and substitute it for y in $Q(x, y)$. We get a series of valuation less than 660, and therefore $Q(x, y)$ is not an annihilating polynomial of $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$.

Finally, the first 24 terms of $M_{0,0}^e$ and $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$ coincide. As these first terms determine a unique solution of $\phi(M^e, 0, 0)$, we know that the two series are one and the same. $\square$

Define

$$R^e = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \quad \text{and} \quad R^o = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}.$$

Lemma 2.2. *For all integers $k \geq 2$ and $u = 2^{2k-1}$, the following identities hold.*

$$\begin{aligned} M^e[:12u] &= M^e[:6u] + x^{4u} M^e[:2u] + x^{5u} M^e[:u] + x^{4u} M^e[:6u] + x^{8u} M^e[:2u] \\ &\quad + x^{9u} M^e[:u] + x^{5u} M^e[:6u] + x^{9u} M^e[:2u] + x^{10u} M^e[:u] \\ &\quad + x^{8u} M^e[:4u] + (x^{6u} + x^{10u} + x^{11u}) R^e, \\ W^e[:12u] &= W^e[:6u] + x^{4u} W^e[:2u] + x^{5u} W^e[:u] + x^{4u} W^e[:6u] + x^{8u} W^e[:2u] \\ &\quad + x^{9u} W^e[:u] + x^{5u} W^e[:6u] + x^{9u} W^e[:2u] + x^{10u} W^e[:u] \\ &\quad + x^{8u} W^e[:4u] + (x^{6u} + x^{10u} + x^{11u}) R^e. \end{aligned}$$

For all integers $k \geq 2$ and $u = 2^{2k}$, the following identities hold.

$$\begin{aligned}
M^o[:12u] &= M^o[:6u] + x^{4u}M^o[:2u] + x^{5u}M^o[:u] + x^{4u}M^o[:6u] + x^{8u}M^o[:2u] \\
&\quad + x^{9u}M^o[:u] + x^{5u}M^o[:6u] + x^{9u}M^o[:2u] + x^{10u}M^o[:u] \\
&\quad + x^{8u}M^o[:4u] + (x^{6u} + x^{10u} + x^{11u})R^o, \\
W^o[:12u] &= W^o[:6u] + x^{4u}W^o[:2u] + x^{5u}W^o[:u] + x^{4u}W^o[:6u] + x^{8u}W^o[:2u] \\
&\quad + x^{9u}W^o[:u] + x^{5u}W^o[:6u] + x^{9u}W^o[:2u] + x^{10u}W^o[:u] \\
&\quad + x^{8u}W^o[:4u] + (x^{6u} + x^{10u} + x^{11u})R^o.
\end{aligned}$$

Proof. To prove Lemma 2.2 we first construct an automaton for each sequence concerned, and then transform the conditions on infinitely many k 's into finitely many conditions on the states of the automaton. In the following, we will prove that for $T = M_{0,0}^o$, for all integer $k \geq 2$ and $u = 2^{2k}$,

$$\begin{aligned}
T[:12u] &= T[:6u] + x^{4u}T[:2u] + x^{5u}T[:u] + x^{4u}T[:6u] + x^{8u}T[:2u] + x^{9u}T[:u] \\
&\quad + x^{5u}T[:6u] + x^{9u}T[:2u] + x^{10u}T[:u] + x^{8u}T[:4u].
\end{aligned}$$

The proof of the other 15 cases are similar. We break down the above identity into 6 parts:

$$\begin{aligned}
0 &= x^{5u}T[u:2u] + x^{4u}T[2u:3u] + T[6u:7u], \\
0 &= x^{5u}T[2u:3u] + x^{4u}T[3u:4u] + T[7u:8u], \\
0 &= x^{5u}T[3u:4u] + x^{4u}T[4u:5u] + T[8u:9u], \\
0 &= x^{5u}T[4u:5u] + x^{4u}T[5u:6u] + T[9u:10u], \\
0 &= x^{10u}T[:u] + x^{9u}T[u:2u] + x^{8u}T[2u:3u] + x^{5u}T[5u:6u] \\
&\quad + T[10u:11u], \\
0 &= x^{8u}T[3u:4u] + T[11u:12u],
\end{aligned}$$

which can be rewritten as

$$(2.7) \quad 0 = T[[1w]_2] + T[[10w]_2] + T[[110w]_2],$$

$$(2.8) \quad 0 = T[[10w]_2] + T[[11w]_2] + T[[111w]_2],$$

$$(2.9) \quad 0 = T[[11w]_2] + T[[100w]_2] + T[[1000w]_2],$$

$$(2.10) \quad 0 = T[[100w]_2] + T[[101w]_2] + T[[1001w]_2],$$

$$(2.11) \quad 0 = T[[w]_2] + T[[1w]_2] + T[[10w]_2] + T[[101w]_2] + T[[1010w]_2],$$

$$(2.12) \quad 0 = T[[11w]_2] + T[[1011w]_2],$$

for all binary word w of length $2k$ and $w \neq 0^{2k}$, and

$$(2.13) \quad 0 = T[[1w]_2] + T[[10w]_2] + T[[110w]_2],$$

$$(2.14) \quad 0 = T[[10w]_2] + T[[11w]_2] + T[[111w]_2],$$

$$(2.15) \quad 0 = T[[11w]_2] + T[[100w]_2] + T[[1000w]_2],$$

$$(2.16) \quad 0 = T[[100w]_2] + T[[101w]_2] + T[[1001w]_2],$$

$$(2.17) \quad 0 = T[[w]_2] + T[[1w]_2] + T[[10w]_2] + T[[101w]_2] + T[[1010w]_2],$$

$$(2.18) \quad 0 = T[[11w]_2] + T[[1011w]_2],$$

for $w = 0^{2k}$.

First we calculate an 2-automaton that generates T from its minimal polynomial and its first terms. This automaton has 246 states; its transition function and output function can be found in the annex. Let $A(s, w)$ denote the state reached after reading w from right to left starting from the state s , and τ the output function. Define

$$E_{2k} = \{A(i, w) : |w| = 2k, w \neq 0^{2k}\}.$$

Identities (2.7) through (2.18) can be written as

$$(2.19) \quad 0 = \tau(A(s, 1)) + \tau(A(s, 10)) + \tau(A(s, 110)),$$

$$(2.20) \quad 0 = \tau(A(s, 10)) + \tau(A(s, 11)) + \tau(A(s, 111)),$$

$$(2.21) \quad 0 = \tau(A(s, 11)) + \tau(A(s, 100)) + \tau(A(s, 1000)),$$

$$(2.22) \quad 0 = \tau(A(s, 100)) + \tau(A(s, 101)) + \tau(A(s, 1001)),$$

$$(2.23) \quad \begin{aligned} 0 &= \tau(A(s, \epsilon)) + \tau(A(s, 1)) + \tau(A(s, 10)) + \tau(A(s, 101)) \\ &\quad + \tau(A(s, 1010)), \end{aligned}$$

$$(2.24) \quad 0 = \tau(A(s, 11)) + \tau(A(s, 1011)),$$

for all $s \in E_{2k}$, and

$$(2.25) \quad 0 = \tau(A(s, 1)) + \tau(A(s, 10)) + \tau(A(s, 110)),$$

$$(2.26) \quad 0 = \tau(A(s, 10)) + \tau(A(s, 11)) + \tau(A(s, 111)),$$

$$(2.27) \quad 0 = \tau(A(s, 11)) + \tau(A(s, 100)) + \tau(A(s, 1000)),$$

$$(2.28) \quad 0 = \tau(A(s, 100)) + \tau(A(s, 101)) + \tau(A(s, 1001)),$$

$$(2.29) \quad \begin{aligned} 0 &= \tau(A(s, \epsilon)) + \tau(A(s, 1)) + \tau(A(s, 10)) + \tau(A(s, 101)) \\ &\quad + \tau(A(s, 1010)), \end{aligned}$$

$$(2.30) \quad 0 = \tau(A(s, 11)) + \tau(A(s, 1011)),$$

for $s = A(i, 0^{2k})$. We find that $(A(i, 0^{20}), E_{20}) = (A(i, 0^{12}), E_{12})$, so that we only have to verify that identities (2.19) through (2.30) hold for $2 \leq k \leq 10$, which turns out to be true. $\square$

In the following lemma, we express M_{2k} , M_{2k+1} , W_{2k} , and W_{2k+1} in terms of M^e , M^o , W^e , and W^o .

Lemma 2.3. *For all integer $k \geq 2$, and $u = 2^{2k-1}$,*

$$M_{2k} = M^e[:6u] + x^{4u} M^e[:2u] + x^{5u} M^e[:u] + x^{6u} R^e,$$

$$W_{2k} = W^e[:6u] + x^{4u} W^e[:2u] + x^{5u} W^e[:u] + x^{6u} R^e.$$

For all integer $k \geq 2$, and $u = 2^{2k}$,

$$M_{2k+1} = M^o[:6u] + x^{4u} M^o[:2u] + x^{5u} M^o[:u] + x^{6u} R^o,$$

$$W_{2k+1} = W^o[:6u] + x^{4u} W^o[:2u] + x^{5u} W^o[:u] + x^{6u} R^o.$$

Proof. We call the four identities in Lemma 2.3 also by the name M_{2k} , W_{2k} , M_{2k+1} , and W_{2k+1} . For $n = 2$, the identities can be verified directly. For $n \geq 2$, we claim that

$$\begin{aligned} M_{2k} \wedge W_{2k} &\Rightarrow M_{2k+1} \wedge W_{2k+1}, \\ M_{2k+1} \wedge W_{2k+1} &\Rightarrow M_{2k+2} \wedge W_{2k+2}. \end{aligned}$$

We give the proof of

$$M_{2k} \wedge W_{2k} \Rightarrow M_{2k+1},$$

the other ones being similar. Set $u = 2^{2k}$ and $v = 2^{2k-1}$. By definition and induction hypothesis, the left side of identity M_{2k+1} is equal to

$$(2.31) \quad \begin{aligned} W_{2k} M_{2k} = & (W^e[:6v] + x^{4v} W^e[:2v] + x^{5v} W^e[:v] + x^{6u} R^e) \times (M^e[:6v] \\ & + x^{4v} M^e[:2v] + x^{5v} M^e[:v] + x^{6u} R^e); \end{aligned}$$

Call this expression *lhs*. Note that both sides of identity M_{2k+1} have the same term of highest degree $x^{12v} R^o$. Therefore we only need to prove that their difference is $O(x^{12v})$. Using Lemma 2.1 it can be seen that the right side of identity M_{2k+1} is congruent, modulo x^{12v} , to

$$W^e[:12v] M^e[:12v] + x^{8v} W^e[:4v] M^e[:4v] + x^{10v} W^e[:2v] M^e[:2v].$$

For all $n \leq 12$, replace the occurrences of $W^e[:n \cdot v]$ and $M^e[:n \cdot v]$ in the above expression by the reduction modulo $x^{n \cdot v}$ of the right side of the corresponding identity in Lemma 2.2 and get a new expression, which we call *rhs*. Define

$$\begin{aligned} X &:= x^v, \\ a_n &:= W^e[n \cdot v : (n+1) \cdot v] / X^n, \\ b_n &:= M^e[n \cdot v : (n+1) \cdot v] / X^n, \\ c &:= R^e. \end{aligned}$$

Using the notation introduced above, we can represent the expressions *lhs* (2.31) and *rhs* as polynomials in $\mathbb{F}_2[a_1, \dots, a_{12}, b_1, \dots, b_{12}, c][X]$. Note that it is not a problem that a_j commutes with b_k while W^e does not commute with M^e , because in the expressions concerned, the products of W^e -terms and M^e -terms are always in the same order. We let the computer do the simplification and check that the difference between these two polynomials is indeed $O(X^{12})$, which completes the proof. $\square$

Proof of Theorem 1.1. We prove the theorem for $\text{CF}(a, b)$; for $\text{CF}(b, a)$, the proof is similar. By Lemma 2.3, we have For all $0 \leq j, k \leq 1$,

$$\begin{aligned} \lim_{n \rightarrow \infty} M_{2n,j,k} &= M_{j,k}^e, \\ \lim_{n \rightarrow \infty} M_{2n+1,j,k} &= M_{j,k}^o, \\ \lim_{n \rightarrow \infty} W_{2n,j,k} &= W_{j,k}^e, \\ \lim_{n \rightarrow \infty} W_{2n+1,j,k} &= W_{j,k}^o. \end{aligned}$$

Let $z = 1/x$. By the convergence theorem and identity (2.2),

$$\text{CF}(a, b) = \frac{M_{0,1}^e(x)}{M_{0,0}^e(x)}.$$

By definition, $\phi(M^e, 0, 1)$ and $\phi(M^e, 0, 0)$ are minimal polynomials of $M_{0,1}^e$ and $M_{0,0}^e$. Therefore

$$P(x, y) = \text{Res}_t (\phi(M^e, 0, 1)(x, t), y^{12} \phi(M^e, 0, 1)(x, t/y))$$

is an annihilating polynomial of $f(x) = M_{0,1}^e / M_{0,0}^e$.

Define

$$Q(x, y) = q_4(x)y^4 + q_3(x)y^3 + q_2(x)y^2 + q_1(x)y + q_0(x),$$

where

$$\begin{aligned}
q_0(x) &= x^{24} + x^{23} + x^{20} + x^{19} + x^{16} + x^{14} + x^{11} + x^{10} + x^6, \\
q_1(x) &= x^{23} + x^{21} + x^{19} + x^{18} + x^{14} + x^{13} + x^{11} + x^9 + x^6 + x^5, \\
q_2(x) &= x^{22} + x^{20} + x^{18} + x^{16} + x^{12} + x^{10} + 1, \\
q_3(x) &= x^{23} + x^{21} + x^{19} + x^{18} + x^{14} + x^{13} + x^{11} + x^9 + x^6 + x^5, \\
q_4(x) &= x^{23} + x^{22} + x^{19} + x^{18} + x^{16} + x^{14} + x^{12} + x^{11} + x^6.
\end{aligned}$$

The polynomial $Q(x, y)$ is the candidate for the minimal polynomial of $f(x)$ found by Padé-Hermite approximation. To prove that it is indeed the minimal polynomial of $f(x)$, we only need to prove that it is an irreducible factor of $P(x, y)$ of multiplicity m and $R(x, y) := P(x, y)/Q(x, y)^m$ is not an annihilating polynomial of $f(x)$. We verify the first point directly. For the second point, we find that when we truncate $f(x)$ to order 186, and substitute it for y in $R(z, y)$, we get a series with valuation smaller than 186, which proves that $R(z, y)$ is not an annihilating polynomial of $f(x)$. Finally, $z^{12}Q(1/z, y)$ is the minimal polynomial of $\text{CF}(a, b) = f(1/z)$. $\square$

3. ANNEXE

All 16 polynomials are of the form

$$p_0(x) + p_3(x)y^3 + p_6(x)y^6 + p_9(x)y^9 + p_{12}(x)y^{12}.$$

The coefficients $p_j(x)$ and the 24 initial terms to determine the solutions uniquely are given below.

For $\phi(M^e, 0, 0)$:

$$\begin{aligned}
p_0(x) &= x^{126} + x^{122} + x^{120} + x^{116} + x^{112} + x^{110} + x^{108} + x^{104} + x^{102} + x^{100} \\
&\quad + x^{98} + x^{88} + x^{86} + x^{84} + x^{82} + x^{80} + x^{76} + x^{72} + x^{70} + x^{68} + x^{64} \\
&\quad + x^{62} + x^{60} + x^{56} + x^{48} + x^{44} + x^{40} + x^{38} + x^{36}, \\
p_3(x) &= x^{114} + x^{104} + x^{98} + x^{96} + x^{94} + x^{86} + x^{84} + x^{80} + x^{76} + x^{72} + x^{68} \\
&\quad + x^{66} + x^{62} + x^{60} + x^{56} + x^{54} + x^{48} + x^{46} + x^{40} + x^{38} + x^{36} + x^{30} \\
&\quad + x^{28} + x^{22}, \\
p_6(x) &= x^{122} + x^{118} + x^{116} + x^{112} + x^{110} + x^{108} + x^{106} + x^{104} + x^{94} + x^{90} \\
&\quad + x^{88} + x^{70} + x^{50} + x^{46} + x^{44} + x^{42} + x^{40} + x^{38} + x^{34} + x^{28} + x^{22} \\
&\quad + x^{14} + x^8 + x^4 + x^2 + 1, \\
p_9(x) &= x^{124} + x^{120} + x^{118} + x^{116} + x^{114} + x^{110} + x^{104} + x^{98} + x^{76} + x^{72} \\
&\quad + x^{70} + x^{66} + x^{64} + x^{58} + x^{52} + x^{48} + x^{46} + x^{44} + x^{42} + x^{38} + x^{36} \\
&\quad + x^{30} + x^{28} + x^{22} + x^{16} + x^{10}, \\
p_{12}(x) &= x^{124} + x^{122} + x^{120} + x^{108} + x^{106} + x^{104} + x^{100} + x^{98} + x^{96} + x^{92} \\
&\quad + x^{90} + x^{88} + x^{76} + x^{74} + x^{72} + x^{68} + x^{66} + x^{64} + x^{28} + x^{26} + x^{24} \\
&\quad + x^{12} + x^{10} + x^8 + x^4 + x^2 + 1,
\end{aligned}$$

and the initial terms are $[1, 0, 0, 0, 0, 0, 1, 0, 1, 0, 1, 0, 1, 0, 0, 0, 0, 0, 1, 0, 0, 0]$.

For $\phi(M^e, 0, 1)$:

$$\begin{aligned}
p_0(x) &= x^{126} + x^{125} + x^{124} + x^{123} + x^{121} + x^{117} + x^{116} + x^{113} + x^{112} + x^{110} \\
&\quad + x^{109} + x^{108} + x^{107} + x^{104} + x^{103} + x^{102} + x^{97} + x^{92} + x^{91} + x^{89} \\
&\quad + x^{86} + x^{83} + x^{82} + x^{80} + x^{79} + x^{77} + x^{76} + x^{75} + x^{74} + x^{73} + x^{70} \\
&\quad + x^{67} + x^{66} + x^{64} + x^{62} + x^{61} + x^{60} + x^{59} + x^{55} + x^{54} + x^{53} + x^{52} \\
&\quad + x^{51} + x^{50} + x^{48} + x^{46} + x^{42} + x^{41} + x^{36}, \\
p_3(x) &= x^{124} + x^{123} + x^{122} + x^{121} + x^{120} + x^{119} + x^{118} + x^{117} + x^{116} + x^{115} \\
&\quad + x^{110} + x^{109} + x^{108} + x^{107} + x^{100} + x^{99} + x^{90} + x^{89} + x^{88} + x^{87} \\
&\quad + x^{86} + x^{85} + x^{78} + x^{77} + x^{76} + x^{75} + x^{74} + x^{73} + x^{72} + x^{71} + x^{70} \\
&\quad + x^{69} + x^{68} + x^{67} + x^{62} + x^{61} + x^{58} + x^{57} + x^{56} + x^{55} + x^{54} + x^{53} \\
&\quad + x^{46} + x^{45}, \\
p_6(x) &= x^{123} + x^{122} + x^{118} + x^{113} + x^{112} + x^{111} + x^{110} + x^{109} + x^{107} + x^{105} \\
&\quad + x^{103} + x^{101} + x^{98} + x^{97} + x^{95} + x^{94} + x^{93} + x^{91} + x^{88} + x^{86} + x^{83} \\
&\quad + x^{82} + x^{78} + x^{73} + x^{72} + x^{71} + x^{70} + x^{69} + x^{67} + x^{65} + x^{63} + x^{61} \\
&\quad + x^{58} + x^{57} + x^{55} + x^{54} + x^{53} + x^{50} + x^{48} + x^{43} + x^{42} + x^{41} + x^{40} \\
&\quad + x^{39} + x^{37} + x^{35} + x^{32} + x^{30}, \\
p_9(x) &= x^{122} + x^{121} + x^{120} + x^{119} + x^{110} + x^{109} + x^{108} + x^{107} + x^{106} + x^{105} \\
&\quad + x^{104} + x^{103} + x^{98} + x^{97} + x^{96} + x^{95} + x^{90} + x^{89} + x^{88} + x^{87} + x^{86} \\
&\quad + x^{85} + x^{84} + x^{83} + x^{82} + x^{81} + x^{80} + x^{79} + x^{74} + x^{73} + x^{72} + x^{71} \\
&\quad + x^{70} + x^{69} + x^{68} + x^{67} + x^{66} + x^{65} + x^{64} + x^{63} + x^{58} + x^{57} + x^{56} \\
&\quad + x^{55} + x^{46} + x^{45} + x^{44} + x^{43} + x^{42} + x^{41} + x^{40} + x^{39} + x^{38} + x^{37} \\
&\quad + x^{36} + x^{35} + x^{30} + x^{29} + x^{28} + x^{27} + x^{26} + x^{25} + x^{24} + x^{23} + x^{18} \\
&\quad + x^{17} + x^{16} + x^{15}, \\
p_{12}(x) &= x^{121} + x^{120} + x^{117} + x^{112} + x^{101} + x^{100} + x^{96} + x^{89} + x^{88} + x^{85} \\
&\quad + x^{80} + x^{69} + x^{68} + x^{64} + x^{25} + x^{24} + x^{21} + x^{16} + x^5 + x^4 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 1, 1, 0, 1, 1, 0, 1, 1, 1, 0, 1, 0, 1, 1, 0, 1, 0, 1, 0, 0, 1]$.

For $\phi(M^e, 1, 0)$:

$$\begin{aligned}
p_0(x) &= x^{126} + x^{125} + x^{124} + x^{123} + x^{121} + x^{117} + x^{116} + x^{113} + x^{112} + x^{110} \\
&\quad + x^{109} + x^{108} + x^{107} + x^{104} + x^{103} + x^{102} + x^{97} + x^{92} + x^{91} + x^{89} \\
&\quad + x^{86} + x^{83} + x^{82} + x^{80} + x^{79} + x^{77} + x^{76} + x^{75} + x^{74} + x^{73} + x^{70} \\
&\quad + x^{67} + x^{66} + x^{64} + x^{62} + x^{61} + x^{60} + x^{59} + x^{55} + x^{54} + x^{53} + x^{52} \\
&\quad + x^{51} + x^{50} + x^{48} + x^{46} + x^{42} + x^{41} + x^{36}, \\
p_3(x) &= x^{124} + x^{123} + x^{122} + x^{121} + x^{120} + x^{119} + x^{118} + x^{117} + x^{116} + x^{115} \\
&\quad + x^{110} + x^{109} + x^{108} + x^{107} + x^{100} + x^{99} + x^{90} + x^{89} + x^{88} + x^{87}
\end{aligned}$$

$$\begin{aligned}
& + x^{86} + x^{85} + x^{78} + x^{77} + x^{76} + x^{75} + x^{74} + x^{73} + x^{72} + x^{71} + x^{70} \\
& + x^{69} + x^{68} + x^{67} + x^{62} + x^{61} + x^{58} + x^{57} + x^{56} + x^{55} + x^{54} + x^{53} \\
& + x^{46} + x^{45}, \\
p_6(x) &= x^{123} + x^{122} + x^{118} + x^{113} + x^{112} + x^{111} + x^{110} + x^{109} + x^{107} + x^{105} \\
& + x^{103} + x^{101} + x^{98} + x^{97} + x^{95} + x^{94} + x^{93} + x^{91} + x^{88} + x^{86} + x^{83} \\
& + x^{82} + x^{78} + x^{73} + x^{72} + x^{71} + x^{70} + x^{69} + x^{67} + x^{65} + x^{63} + x^{61} \\
& + x^{58} + x^{57} + x^{55} + x^{54} + x^{53} + x^{50} + x^{48} + x^{43} + x^{42} + x^{41} + x^{40} \\
& + x^{39} + x^{37} + x^{35} + x^{32} + x^{30}, \\
p_9(x) &= x^{122} + x^{121} + x^{120} + x^{119} + x^{110} + x^{109} + x^{108} + x^{107} + x^{106} + x^{105} \\
& + x^{104} + x^{103} + x^{98} + x^{97} + x^{96} + x^{95} + x^{90} + x^{89} + x^{88} + x^{87} + x^{86} \\
& + x^{85} + x^{84} + x^{83} + x^{82} + x^{81} + x^{80} + x^{79} + x^{74} + x^{73} + x^{72} + x^{71} \\
& + x^{70} + x^{69} + x^{68} + x^{67} + x^{66} + x^{65} + x^{64} + x^{63} + x^{58} + x^{57} + x^{56} \\
& + x^{55} + x^{46} + x^{45} + x^{44} + x^{43} + x^{42} + x^{41} + x^{40} + x^{39} + x^{38} + x^{37} \\
& + x^{36} + x^{35} + x^{30} + x^{29} + x^{28} + x^{27} + x^{26} + x^{25} + x^{24} + x^{23} + x^{18} \\
& + x^{17} + x^{16} + x^{15}, \\
p_{12}(x) &= x^{121} + x^{120} + x^{117} + x^{112} + x^{101} + x^{100} + x^{96} + x^{89} + x^{88} + x^{85} \\
& + x^{80} + x^{69} + x^{68} + x^{64} + x^{25} + x^{24} + x^{21} + x^{16} + x^5 + x^4 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 1, 1, 0, 1, 1, 0, 1, 1, 1, 0, 1, 0, 1, 1, 0, 1, 0, 1, 0, 0, 1]$.

For $\phi(M^e, 1, 1)$:

$$\begin{aligned}
p_0(x) &= x^{126} + x^{124} + x^{120} + x^{118} + x^{116} + x^{114} + x^{112} + x^{110} + x^{104} + x^{94} \\
& + x^{90} + x^{86} + x^{82} + x^{80} + x^{76} + x^{72} + x^{70} + x^{66} + x^{64} + x^{62} + x^{60} \\
& + x^{56} + x^{54} + x^{52} + x^{48} + x^{46} + x^{42} + x^{38} + x^{36}, \\
p_3(x) &= x^{120} + x^{114} + x^{112} + x^{96} + x^{94} + x^{90} + x^{86} + x^{76} + x^{64} + x^{62} + x^{60} \\
& + x^{58} + x^{56} + x^{54} + x^{50} + x^{46} + x^{42} + x^{40} + x^{38} + x^{34} + x^{30} + x^{28} \\
& + x^{26} + x^{22}, \\
p_6(x) &= x^{120} + x^{118} + x^{116} + x^{108} + x^{104} + x^{102} + x^{98} + x^{94} + x^{88} + x^{86} \\
& + x^{82} + x^{80} + x^{74} + x^{70} + x^{68} + x^{66} + x^{64} + x^{60} + x^{58} + x^{56} + x^{50} \\
& + x^{48} + x^{36} + x^{34} + x^{30} + x^{28} + x^{26} + x^{22} + x^{20} + x^{18} + x^{14} + x^{12} \\
& + x^8 + x^6 + x^2 + 1, \\
p_9(x) &= x^{118} + x^{116} + x^{112} + x^{104} + x^{100} + x^{98} + x^{96} + x^{88} + x^{84} + x^{82} + x^{80} \\
& + x^{78} + x^{76} + x^{70} + x^{66} + x^{64} + x^{60} + x^{58} + x^{56} + x^{48} + x^{46} + x^{42} \\
& + x^{38} + x^{36} + x^{32} + x^{30} + x^{28} + x^{22} + x^{18} + x^{16} + x^{14} + x^{10}, \\
p_{12}(x) &= x^{118} + x^{114} + x^{112} + x^{110} + x^{106} + x^{104} + x^{102} + x^{98} + x^{96} + x^{86} \\
& + x^{82} + x^{80} + x^{78} + x^{74} + x^{72} + x^{70} + x^{66} + x^{64} + x^{22} + x^{18} + x^{16}
\end{aligned}$$

$$+ x^{14} + x^{10} + x^8 + x^6 + x^2 + 1,$$

and the initial terms are $[0, 0, 0, 0, 0, 0, 1, 0, 1, 0, 0, 0, 0, 0, 1, 0, 0, 0, 1, 0, 0, 0, 1, 0]$.

For $\phi(W^e, 0, 0)$:

$$p_0(x) = x^{126} + x^{124} + x^{122} + x^{120} + x^{116} + x^{114} + x^{108} + x^{106} + x^{100} + x^{94} \\ + x^{78} + x^{58} + x^{52} + x^{44} + x^{42} + x^{38} + x^{36},$$

$$p_3(x) = x^{120} + x^{116} + x^{114} + x^{110} + x^{106} + x^{104} + x^{98} + x^{96} + x^{94} + x^{90} \\ + x^{88} + x^{86} + x^{82} + x^{80} + x^{76} + x^{74} + x^{62} + x^{60} + x^{54} + x^{52} + x^{48} \\ + x^{40} + x^{38} + x^{36} + x^{34} + x^{28} + x^{26} + x^{22},$$

$$p_6(x) = x^{120} + x^{118} + x^{114} + x^{112} + x^{106} + x^{102} + x^{98} + x^{94} + x^{92} + x^{90} \\ + x^{88} + x^{80} + x^{78} + x^{76} + x^{74} + x^{70} + x^{64} + x^{62} + x^{60} + x^{56} + x^{54} \\ + x^{40} + x^{38} + x^{34} + x^{32} + x^{26} + x^{24} + x^{18} + x^{14} + x^{12} + x^8 + x^6 + x^2 \\ + 1,$$

$$p_9(x) = x^{118} + x^{116} + x^{112} + x^{104} + x^{100} + x^{98} + x^{96} + x^{88} + x^{84} + x^{82} + x^{80} \\ + x^{78} + x^{76} + x^{70} + x^{66} + x^{64} + x^{60} + x^{58} + x^{56} + x^{48} + x^{46} + x^{42} \\ + x^{38} + x^{36} + x^{32} + x^{30} + x^{28} + x^{22} + x^{18} + x^{16} + x^{14} + x^{10},$$

$$p_{12}(x) = x^{118} + x^{114} + x^{112} + x^{110} + x^{106} + x^{104} + x^{102} + x^{98} + x^{96} + x^{86} \\ + x^{82} + x^{80} + x^{78} + x^{74} + x^{72} + x^{70} + x^{66} + x^{64} + x^{22} + x^{18} + x^{16} \\ + x^{14} + x^{10} + x^8 + x^6 + x^2 + 1,$$

and the initial terms are $[1, 0, 0, 0, 0, 0, 1, 0, 1, 0, 0, 0, 0, 0, 1, 0, 0, 0, 1, 0, 1, 0, 1, 0]$.

For $\phi(W^e, 0, 1)$:

$$p_0(x) = x^{126} + x^{124} + x^{123} + x^{119} + x^{118} + x^{117} + x^{116} + x^{112} + x^{111} + x^{110} \\ + x^{109} + x^{107} + x^{106} + x^{104} + x^{103} + x^{102} + x^{99} + x^{97} + x^{96} + x^{86} \\ + x^{84} + x^{83} + x^{81} + x^{80} + x^{79} + x^{78} + x^{77} + x^{76} + x^{75} + x^{74} + x^{73} \\ + x^{70} + x^{69} + x^{68} + x^{67} + x^{66} + x^{65} + x^{64} + x^{63} + x^{60} + x^{56} + x^{55} \\ + x^{53} + x^{52} + x^{51} + x^{50} + x^{48} + x^{46} + x^{42} + x^{41} + x^{36},$$

$$p_3(x) = x^{124} + x^{123} + x^{122} + x^{121} + x^{120} + x^{119} + x^{118} + x^{117} + x^{116} + x^{115} \\ + x^{110} + x^{109} + x^{108} + x^{107} + x^{100} + x^{99} + x^{90} + x^{89} + x^{88} + x^{87} \\ + x^{86} + x^{85} + x^{78} + x^{77} + x^{76} + x^{75} + x^{74} + x^{73} + x^{72} + x^{71} + x^{70} \\ + x^{69} + x^{68} + x^{67} + x^{62} + x^{61} + x^{58} + x^{57} + x^{56} + x^{55} + x^{54} + x^{53} \\ + x^{46} + x^{45},$$

$$p_6(x) = x^{123} + x^{122} + x^{118} + x^{113} + x^{112} + x^{111} + x^{110} + x^{109} + x^{107} + x^{105} \\ + x^{103} + x^{101} + x^{98} + x^{97} + x^{95} + x^{94} + x^{93} + x^{91} + x^{88} + x^{86} + x^{83} \\ + x^{82} + x^{78} + x^{73} + x^{72} + x^{71} + x^{70} + x^{69} + x^{67} + x^{65} + x^{63} + x^{61} \\ + x^{58} + x^{57} + x^{55} + x^{54} + x^{53} + x^{50} + x^{48} + x^{43} + x^{42} + x^{41} + x^{40}$$

$$\begin{aligned}
& + x^{39} + x^{37} + x^{35} + x^{32} + x^{30}, \\
p_9(x) = & x^{122} + x^{121} + x^{120} + x^{119} + x^{110} + x^{109} + x^{108} + x^{107} + x^{106} + x^{105} \\
& + x^{104} + x^{103} + x^{98} + x^{97} + x^{96} + x^{95} + x^{90} + x^{89} + x^{88} + x^{87} + x^{86} \\
& + x^{85} + x^{84} + x^{83} + x^{82} + x^{81} + x^{80} + x^{79} + x^{74} + x^{73} + x^{72} + x^{71} \\
& + x^{70} + x^{69} + x^{68} + x^{67} + x^{66} + x^{65} + x^{64} + x^{63} + x^{58} + x^{57} + x^{56} \\
& + x^{55} + x^{46} + x^{45} + x^{44} + x^{43} + x^{42} + x^{41} + x^{40} + x^{39} + x^{38} + x^{37} \\
& + x^{36} + x^{35} + x^{30} + x^{29} + x^{28} + x^{27} + x^{26} + x^{25} + x^{24} + x^{23} + x^{18} \\
& + x^{17} + x^{16} + x^{15}, \\
p_{12}(x) = & x^{121} + x^{120} + x^{117} + x^{112} + x^{101} + x^{100} + x^{96} + x^{89} + x^{88} + x^{85} \\
& + x^{80} + x^{69} + x^{68} + x^{64} + x^{25} + x^{24} + x^{21} + x^{16} + x^5 + x^4 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 1, 1, 1, 0, 1, 0, 0, 1, 0, 0, 0, 1, 0, 0, 1, 1, 0, 1, 0, 1, 1]$.

For $\phi(W^e, 1, 0)$:

$$\begin{aligned}
p_0(x) = & x^{126} + x^{124} + x^{123} + x^{119} + x^{118} + x^{117} + x^{116} + x^{112} + x^{111} + x^{110} \\
& + x^{109} + x^{107} + x^{106} + x^{104} + x^{103} + x^{102} + x^{99} + x^{97} + x^{96} + x^{86} \\
& + x^{84} + x^{83} + x^{81} + x^{80} + x^{79} + x^{78} + x^{77} + x^{76} + x^{75} + x^{74} + x^{73} \\
& + x^{70} + x^{69} + x^{68} + x^{67} + x^{66} + x^{65} + x^{64} + x^{63} + x^{60} + x^{56} + x^{55} \\
& + x^{53} + x^{52} + x^{51} + x^{50} + x^{48} + x^{46} + x^{42} + x^{41} + x^{36}, \\
p_3(x) = & x^{124} + x^{123} + x^{122} + x^{121} + x^{120} + x^{119} + x^{118} + x^{117} + x^{116} + x^{115} \\
& + x^{110} + x^{109} + x^{108} + x^{107} + x^{100} + x^{99} + x^{90} + x^{89} + x^{88} + x^{87} \\
& + x^{86} + x^{85} + x^{78} + x^{77} + x^{76} + x^{75} + x^{74} + x^{73} + x^{72} + x^{71} + x^{70} \\
& + x^{69} + x^{68} + x^{67} + x^{62} + x^{61} + x^{58} + x^{57} + x^{56} + x^{55} + x^{54} + x^{53} \\
& + x^{46} + x^{45}, \\
p_6(x) = & x^{123} + x^{122} + x^{118} + x^{113} + x^{112} + x^{111} + x^{110} + x^{109} + x^{107} + x^{105} \\
& + x^{103} + x^{101} + x^{98} + x^{97} + x^{95} + x^{94} + x^{93} + x^{91} + x^{88} + x^{86} + x^{83} \\
& + x^{82} + x^{78} + x^{73} + x^{72} + x^{71} + x^{70} + x^{69} + x^{67} + x^{65} + x^{63} + x^{61} \\
& + x^{58} + x^{57} + x^{55} + x^{54} + x^{53} + x^{50} + x^{48} + x^{43} + x^{42} + x^{41} + x^{40} \\
& + x^{39} + x^{37} + x^{35} + x^{32} + x^{30}, \\
p_9(x) = & x^{122} + x^{121} + x^{120} + x^{119} + x^{110} + x^{109} + x^{108} + x^{107} + x^{106} + x^{105} \\
& + x^{104} + x^{103} + x^{98} + x^{97} + x^{96} + x^{95} + x^{90} + x^{89} + x^{88} + x^{87} + x^{86} \\
& + x^{85} + x^{84} + x^{83} + x^{82} + x^{81} + x^{80} + x^{79} + x^{74} + x^{73} + x^{72} + x^{71} \\
& + x^{70} + x^{69} + x^{68} + x^{67} + x^{66} + x^{65} + x^{64} + x^{63} + x^{58} + x^{57} + x^{56} \\
& + x^{55} + x^{46} + x^{45} + x^{44} + x^{43} + x^{42} + x^{41} + x^{40} + x^{39} + x^{38} + x^{37} \\
& + x^{36} + x^{35} + x^{30} + x^{29} + x^{28} + x^{27} + x^{26} + x^{25} + x^{24} + x^{23} + x^{18} \\
& + x^{17} + x^{16} + x^{15},
\end{aligned}$$

$$p_{12}(x) = x^{121} + x^{120} + x^{117} + x^{112} + x^{101} + x^{100} + x^{96} + x^{89} + x^{88} + x^{85} \\ + x^{80} + x^{69} + x^{68} + x^{64} + x^{25} + x^{24} + x^{21} + x^{16} + x^5 + x^4 + 1,$$

and the initial terms are $[0, 0, 0, 1, 1, 1, 0, 1, 0, 0, 1, 0, 0, 0, 1, 0, 0, 1, 1, 0, 1, 0, 1, 1]$.

For $\phi(W^e, 1, 1)$:

$$p_0(x) = x^{126} + x^{124} + x^{122} + x^{112} + x^{110} + x^{106} + x^{102} + x^{100} + x^{98} + x^{82} \\ + x^{78} + x^{68} + x^{64} + x^{58} + x^{52} + x^{50} + x^{48} + x^{46} + x^{40} + x^{38} + x^{36}, \\ p_3(x) = x^{122} + x^{120} + x^{118} + x^{110} + x^{108} + x^{106} + x^{104} + x^{102} + x^{88} + x^{86} \\ + x^{82} + x^{78} + x^{74} + x^{70} + x^{66} + x^{62} + x^{54} + x^{46} + x^{44} + x^{40} + x^{38} \\ + x^{34} + x^{28} + x^{22}, \\ p_6(x) = x^{118} + x^{114} + x^{110} + x^{108} + x^{104} + x^{102} + x^{100} + x^{90} + x^{84} + x^{82} \\ + x^{80} + x^{78} + x^{72} + x^{70} + x^{68} + x^{66} + x^{64} + x^{58} + x^{52} + x^{48} + x^{44} \\ + x^{42} + x^{36} + x^{34} + x^{28} + x^{26} + x^{20} + x^{14} + x^8 + x^4 + x^2 + 1, \\ p_9(x) = x^{124} + x^{120} + x^{118} + x^{116} + x^{114} + x^{110} + x^{104} + x^{98} + x^{76} + x^{72} \\ + x^{70} + x^{66} + x^{64} + x^{58} + x^{52} + x^{48} + x^{46} + x^{44} + x^{42} + x^{38} + x^{36} \\ + x^{30} + x^{28} + x^{22} + x^{16} + x^{10}, \\ p_{12}(x) = x^{124} + x^{122} + x^{120} + x^{108} + x^{106} + x^{104} + x^{100} + x^{98} + x^{96} + x^{92} \\ + x^{90} + x^{88} + x^{76} + x^{74} + x^{72} + x^{68} + x^{66} + x^{64} + x^{28} + x^{26} + x^{24} \\ + x^{12} + x^{10} + x^8 + x^4 + x^2 + 1,$$

and the initial terms are $[0, 0, 0, 0, 0, 0, 1, 0, 1, 0, 1, 0, 1, 0, 1, 0, 0, 0, 0, 0, 0, 0, 0, 0]$.

For $\phi(M^o, 0, 0)$:

$$p_0(x) = x^{126} + x^{125} + x^{123} + x^{120} + x^{119} + x^{118} + x^{117} + x^{116} + x^{115} + x^{113} \\ + x^{112} + x^{109} + x^{107} + x^{106} + x^{105} + x^{104} + x^{103} + x^{101} + x^{99} + x^{98} \\ + x^{96} + x^{95} + x^{89} + x^{86} + x^{85} + x^{83} + x^{79} + x^{78} + x^{77} + x^{76} + x^{75} \\ + x^{72} + x^{71} + x^{70} + x^{69} + x^{67} + x^{66} + x^{63} + x^{62} + x^{60} + x^{59} + x^{53} \\ + x^{52} + x^{51} + x^{48} + x^{46} + x^{41} + x^{40} + x^{36}, \\ p_3(x) = x^{125} + x^{123} + x^{117} + x^{115} + x^{114} + x^{113} + x^{111} + x^{110} + x^{107} + x^{105} \\ + x^{104} + x^{103} + x^{101} + x^{100} + x^{98} + x^{96} + x^{93} + x^{92} + x^{90} + x^{87} \\ + x^{85} + x^{84} + x^{83} + x^{81} + x^{79} + x^{78} + x^{77} + x^{76} + x^{75} + x^{74} + x^{73} \\ + x^{72} + x^{67} + x^{66} + x^{65} + x^{64} + x^{63} + x^{62} + x^{61} + x^{57} + x^{56} + x^{55} \\ + x^{52} + x^{51} + x^{49} + x^{48} + x^{47} + x^{46} + x^{44} + x^{42} + x^{41} + x^{40} + x^{37} \\ + x^{32}, \\ p_6(x) = x^{125} + x^{123} + x^{120} + x^{119} + x^{115} + x^{114} + x^{112} + x^{109} + x^{106} + x^{105} \\ + x^{104} + x^{100} + x^{98} + x^{96} + x^{95} + x^{94} + x^{93} + x^{89} + x^{86} + x^{85} + x^{84} \\ + x^{80} + x^{79} + x^{78} + x^{77} + x^{76} + x^{75} + x^{74} + x^{73} + x^{72} + x^{66} + x^{65}$$

$$\begin{aligned}
& + x^{61} + x^{58} + x^{55} + x^{54} + x^{51} + x^{49} + x^{48} + x^{47} + x^{46} + x^{45} + x^{44} \\
& + x^{43} + x^{42} + x^{39} + x^{38} + x^{36} + x^{35} + x^{29} + x^{28} + x^{27} + x^{25} + x^{22} \\
& + x^{21} + x^{17} + x^{12} + x^5 + x^4 + 1, \\
p_9(x) &= x^{125} + x^{124} + x^{121} + x^{116} + x^{93} + x^{92} + x^{89} + x^{84} + x^{29} + x^{28} + x^{25} \\
& + x^{20}, \\
p_{12}(x) &= x^{125} + x^{124} + x^{120} + x^{117} + x^{116} + x^{112} + x^{101} + x^{100} + x^{96} + x^{93} \\
& + x^{92} + x^{88} + x^{85} + x^{84} + x^{80} + x^{69} + x^{68} + x^{64} + x^{29} + x^{28} + x^{24} \\
& + x^{21} + x^{20} + x^{16} + x^5 + x^4 + 1,
\end{aligned}$$

and the initial terms are $[1, 0, 0, 0, 0, 0, 1, 0, 0, 0, 0, 1, 0, 0, 1, 0, 0, 0, 1, 1, 1, 0, 0, 1]$.

For $\phi(M^o, 0, 1)$:

$$\begin{aligned}
p_0(x) &= x^{132} + x^{129} + x^{126} + x^{123} + x^{122} + x^{120} + x^{119} + x^{117} + x^{115} + x^{114} \\
& + x^{109} + x^{108} + x^{106} + x^{105} + x^{104} + x^{103} + x^{101} + x^{100} + x^{97} + x^{94} \\
& + x^{93} + x^{92} + x^{86} + x^{82} + x^{81} + x^{80} + x^{79} + x^{75} + x^{74} + x^{72} + x^{70} \\
& + x^{67} + x^{66} + x^{65} + x^{63} + x^{60} + x^{58} + x^{55} + x^{52} + x^{51} + x^{49} + x^{47} \\
& + x^{46} + x^{45} + x^{44} + x^{42} + x^{36}, \\
p_3(x) &= x^{125} + x^{120} + x^{119} + x^{118} + x^{116} + x^{114} + x^{113} + x^{111} + x^{110} + x^{109} \\
& + x^{108} + x^{107} + x^{105} + x^{103} + x^{100} + x^{99} + x^{97} + x^{95} + x^{93} + x^{92} \\
& + x^{91} + x^{89} + x^{88} + x^{86} + x^{83} + x^{82} + x^{78} + x^{72} + x^{71} + x^{70} + x^{68} \\
& + x^{66} + x^{65} + x^{63} + x^{62} + x^{60} + x^{59} + x^{57} + x^{56} + x^{54} + x^{51} + x^{50} \\
& + x^{46} + x^{45}, \\
p_6(x) &= x^{126} + x^{122} + x^{118} + x^{116} + x^{114} + x^{108} + x^{106} + x^{104} + x^{100} + x^{96} \\
& + x^{88} + x^{82} + x^{78} + x^{76} + x^{74} + x^{68} + x^{66} + x^{64} + x^{60} + x^{56} + x^{54} \\
& + x^{50} + x^{48} + x^{46} + x^{44} + x^{38} + x^{32} + x^{30}, \\
p_9(x) &= x^{127} + x^{122} + x^{121} + x^{120} + x^{119} + x^{118} + x^{117} + x^{116} + x^{114} + x^{113} \\
& + x^{112} + x^{111} + x^{95} + x^{90} + x^{89} + x^{88} + x^{87} + x^{86} + x^{85} + x^{84} + x^{82} \\
& + x^{81} + x^{80} + x^{79} + x^{31} + x^{26} + x^{25} + x^{24} + x^{23} + x^{22} + x^{21} + x^{20} \\
& + x^{18} + x^{17} + x^{16} + x^{15}, \\
p_{12}(x) &= x^{128} + x^{124} + x^{116} + x^{112} + x^{104} + x^{100} + x^{92} + x^{84} + x^{80} + x^{72} \\
& + x^{68} + x^{64} + x^{32} + x^{28} + x^{20} + x^{16} + x^8 + x^4 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 1, 1, 0, 1, 1, 0, 1, 1, 0, 1, 0, 1, 0, 1, 1, 1, 0, 0, 0]$.

For $\phi(M^o, 1, 0)$:

$$\begin{aligned}
p_0(x) &= x^{130} + x^{129} + x^{128} + x^{127} + x^{126} + x^{125} + x^{124} + x^{123} + x^{121} + x^{120} \\
& + x^{118} + x^{117} + x^{113} + x^{111} + x^{109} + x^{108} + x^{107} + x^{106} + x^{105} \\
& + x^{104} + x^{101} + x^{100} + x^{98} + x^{97} + x^{92} + x^{91} + x^{90} + x^{89} + x^{88} + x^{82}
\end{aligned}$$

$$\begin{aligned}
& + x^{74} + x^{71} + x^{70} + x^{69} + x^{67} + x^{65} + x^{64} + x^{63} + x^{62} + x^{61} + x^{60} \\
& + x^{59} + x^{57} + x^{53} + x^{52} + x^{51} + x^{50} + x^{49} + x^{47} + x^{46} + x^{45} + x^{42} \\
& + x^{36}, \\
p_3(x) &= x^{129} + x^{125} + x^{124} + x^{123} + x^{122} + x^{121} + x^{119} + x^{114} + x^{112} + x^{111} \\
& + x^{110} + x^{107} + x^{104} + x^{103} + x^{101} + x^{99} + x^{97} + x^{96} + x^{95} + x^{92} \\
& + x^{90} + x^{89} + x^{88} + x^{85} + x^{83} + x^{82} + x^{81} + x^{78} + x^{76} + x^{75} + x^{74} \\
& + x^{73} + x^{71} + x^{66} + x^{65} + x^{64} + x^{63} + x^{62} + x^{61} + x^{60} + x^{58} + x^{57} \\
& + x^{56} + x^{53} + x^{51} + x^{50} + x^{46} + x^{45}, \\
p_6(x) &= x^{130} + x^{122} + x^{120} + x^{116} + x^{114} + x^{108} + x^{104} + x^{102} + x^{100} + x^{94} \\
& + x^{90} + x^{88} + x^{86} + x^{82} + x^{80} + x^{76} + x^{74} + x^{68} + x^{64} + x^{62} + x^{60} \\
& + x^{58} + x^{54} + x^{46} + x^{44} + x^{40} + x^{32} + x^{30}, \\
p_9(x) &= x^{131} + x^{127} + x^{126} + x^{125} + x^{124} + x^{118} + x^{117} + x^{116} + x^{114} + x^{113} \\
& + x^{112} + x^{111} + x^{99} + x^{95} + x^{94} + x^{93} + x^{92} + x^{86} + x^{85} + x^{84} + x^{82} \\
& + x^{81} + x^{80} + x^{79} + x^{35} + x^{31} + x^{30} + x^{29} + x^{28} + x^{22} + x^{21} + x^{20} \\
& + x^{18} + x^{17} + x^{16} + x^{15}, \\
p_{12}(x) &= x^{132} + x^{120} + x^{116} + x^{112} + x^{108} + x^{96} + x^{88} + x^{84} + x^{80} + x^{76} + x^{68} \\
& + x^{64} + x^{36} + x^{24} + x^{20} + x^{16} + x^{12} + x^4 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 1, 1, 1, 0, 1, 0, 0, 1, 1, 1, 1, 0, 0, 1, 0, 0, 0, 1, 0, 1, 1]$.

For $\phi(M^o, 1, 1)$:

$$\begin{aligned}
p_0(x) &= x^{126} + x^{121} + x^{120} + x^{118} + x^{117} + x^{116} + x^{112} + x^{111} + x^{110} + x^{109} \\
& + x^{108} + x^{107} + x^{106} + x^{103} + x^{102} + x^{99} + x^{98} + x^{97} + x^{91} + x^{90} \\
& + x^{89} + x^{88} + x^{84} + x^{83} + x^{82} + x^{77} + x^{74} + x^{73} + x^{71} + x^{70} + x^{67} \\
& + x^{65} + x^{61} + x^{60} + x^{57} + x^{54} + x^{52} + x^{51} + x^{49} + x^{48} + x^{46} + x^{44} \\
& + x^{41} + x^{40} + x^{36}, \\
p_3(x) &= x^{125} + x^{122} + x^{121} + x^{120} + x^{119} + x^{114} + x^{113} + x^{112} + x^{111} + x^{109} \\
& + x^{108} + x^{106} + x^{105} + x^{103} + x^{102} + x^{101} + x^{100} + x^{99} + x^{96} + x^{94} \\
& + x^{93} + x^{92} + x^{90} + x^{89} + x^{88} + x^{86} + x^{84} + x^{83} + x^{82} + x^{81} + x^{80} \\
& + x^{79} + x^{78} + x^{72} + x^{71} + x^{69} + x^{66} + x^{65} + x^{63} + x^{60} + x^{56} + x^{54} \\
& + x^{53} + x^{52} + x^{51} + x^{50} + x^{49} + x^{47} + x^{46} + x^{45} + x^{42} + x^{41} + x^{37} \\
& + x^{32}, \\
p_6(x) &= x^{125} + x^{122} + x^{121} + x^{119} + x^{118} + x^{116} + x^{115} + x^{114} + x^{113} + x^{111} \\
& + x^{110} + x^{107} + x^{106} + x^{104} + x^{103} + x^{97} + x^{93} + x^{92} + x^{91} + x^{88} \\
& + x^{84} + x^{83} + x^{82} + x^{79} + x^{77} + x^{76} + x^{75} + x^{74} + x^{71} + x^{70} + x^{69} \\
& + x^{67} + x^{66} + x^{63} + x^{61} + x^{60} + x^{57} + x^{56} + x^{52} + x^{51} + x^{50} + x^{49}
\end{aligned}$$

$$\begin{aligned}
& + x^{48} + x^{47} + x^{46} + x^{41} + x^{38} + x^{37} + x^{36} + x^{32} + x^{30} + x^{27} + x^{22} \\
& + x^{21} + x^{20} + x^{17} + x^{12} + x^5 + x^4 + 1, \\
p_9(x) &= x^{125} + x^{124} + x^{121} + x^{116} + x^{93} + x^{92} + x^{89} + x^{84} + x^{29} + x^{28} + x^{25} \\
& + x^{20}, \\
p_{12}(x) &= x^{125} + x^{124} + x^{120} + x^{117} + x^{116} + x^{112} + x^{101} + x^{100} + x^{96} + x^{93} \\
& + x^{92} + x^{88} + x^{85} + x^{84} + x^{80} + x^{69} + x^{68} + x^{64} + x^{29} + x^{28} + x^{24} \\
& + x^{21} + x^{20} + x^{16} + x^5 + x^4 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 0, 1, 0, 0, 0, 1, 1, 1, 0, 1, 0, 0, 0, 0, 1, 1, 0, 1, 1]$.

For $\phi(W^o, 0, 0)$:

$$\begin{aligned}
p_0(x) &= x^{126} + x^{125} + x^{123} + x^{120} + x^{119} + x^{118} + x^{117} + x^{116} + x^{115} + x^{113} \\
& + x^{112} + x^{109} + x^{107} + x^{106} + x^{105} + x^{104} + x^{103} + x^{101} + x^{99} + x^{98} \\
& + x^{96} + x^{95} + x^{89} + x^{86} + x^{85} + x^{83} + x^{79} + x^{78} + x^{77} + x^{76} + x^{75} \\
& + x^{72} + x^{71} + x^{70} + x^{69} + x^{67} + x^{66} + x^{63} + x^{62} + x^{60} + x^{59} + x^{53} \\
& + x^{52} + x^{51} + x^{48} + x^{46} + x^{41} + x^{40} + x^{36}, \\
p_3(x) &= x^{125} + x^{123} + x^{117} + x^{115} + x^{114} + x^{113} + x^{111} + x^{110} + x^{107} + x^{105} \\
& + x^{104} + x^{103} + x^{101} + x^{100} + x^{98} + x^{96} + x^{93} + x^{92} + x^{90} + x^{87} \\
& + x^{85} + x^{84} + x^{83} + x^{81} + x^{79} + x^{78} + x^{77} + x^{76} + x^{75} + x^{74} + x^{73} \\
& + x^{72} + x^{67} + x^{66} + x^{65} + x^{64} + x^{63} + x^{62} + x^{61} + x^{57} + x^{56} + x^{55} \\
& + x^{52} + x^{51} + x^{49} + x^{48} + x^{47} + x^{46} + x^{44} + x^{42} + x^{41} + x^{40} + x^{37} \\
& + x^{32}, \\
p_6(x) &= x^{125} + x^{123} + x^{120} + x^{119} + x^{115} + x^{114} + x^{112} + x^{109} + x^{106} + x^{105} \\
& + x^{104} + x^{100} + x^{98} + x^{96} + x^{95} + x^{94} + x^{93} + x^{89} + x^{86} + x^{85} + x^{84} \\
& + x^{80} + x^{79} + x^{78} + x^{77} + x^{76} + x^{75} + x^{74} + x^{73} + x^{72} + x^{66} + x^{65} \\
& + x^{61} + x^{58} + x^{55} + x^{54} + x^{51} + x^{49} + x^{48} + x^{47} + x^{46} + x^{45} + x^{44} \\
& + x^{43} + x^{42} + x^{39} + x^{38} + x^{36} + x^{35} + x^{29} + x^{28} + x^{27} + x^{25} + x^{22} \\
& + x^{21} + x^{17} + x^{12} + x^5 + x^4 + 1, \\
p_9(x) &= x^{125} + x^{124} + x^{121} + x^{116} + x^{93} + x^{92} + x^{89} + x^{84} + x^{29} + x^{28} + x^{25} \\
& + x^{20}, \\
p_{12}(x) &= x^{125} + x^{124} + x^{120} + x^{117} + x^{116} + x^{112} + x^{101} + x^{100} + x^{96} + x^{93} \\
& + x^{92} + x^{88} + x^{85} + x^{84} + x^{80} + x^{69} + x^{68} + x^{64} + x^{29} + x^{28} + x^{24} \\
& + x^{21} + x^{20} + x^{16} + x^5 + x^4 + 1,
\end{aligned}$$

and the initial terms are $[1, 0, 0, 0, 0, 0, 1, 0, 0, 0, 0, 1, 0, 0, 1, 0, 0, 0, 1, 1, 1, 0, 0, 1]$.

For $\phi(W^o, 0, 1)$:

$$p_0(x) = x^{130} + x^{129} + x^{128} + x^{127} + x^{126} + x^{125} + x^{124} + x^{123} + x^{121} + x^{120}$$

$$\begin{aligned}
& + x^{118} + x^{117} + x^{113} + x^{111} + x^{109} + x^{108} + x^{107} + x^{106} + x^{105} \\
& + x^{104} + x^{101} + x^{100} + x^{98} + x^{97} + x^{92} + x^{91} + x^{90} + x^{89} + x^{88} + x^{82} \\
& + x^{74} + x^{71} + x^{70} + x^{69} + x^{67} + x^{65} + x^{64} + x^{63} + x^{62} + x^{61} + x^{60} \\
& + x^{59} + x^{57} + x^{53} + x^{52} + x^{51} + x^{50} + x^{49} + x^{47} + x^{46} + x^{45} + x^{42} \\
& + x^{36}, \\
p_3(x) &= x^{129} + x^{125} + x^{124} + x^{123} + x^{122} + x^{121} + x^{119} + x^{114} + x^{112} + x^{111} \\
& + x^{110} + x^{107} + x^{104} + x^{103} + x^{101} + x^{99} + x^{97} + x^{96} + x^{95} + x^{92} \\
& + x^{90} + x^{89} + x^{88} + x^{85} + x^{83} + x^{82} + x^{81} + x^{78} + x^{76} + x^{75} + x^{74} \\
& + x^{73} + x^{71} + x^{66} + x^{65} + x^{64} + x^{63} + x^{62} + x^{61} + x^{60} + x^{58} + x^{57} \\
& + x^{56} + x^{53} + x^{51} + x^{50} + x^{46} + x^{45}, \\
p_6(x) &= x^{130} + x^{122} + x^{120} + x^{116} + x^{114} + x^{108} + x^{104} + x^{102} + x^{100} + x^{94} \\
& + x^{90} + x^{88} + x^{86} + x^{82} + x^{80} + x^{76} + x^{74} + x^{68} + x^{64} + x^{62} + x^{60} \\
& + x^{58} + x^{54} + x^{46} + x^{44} + x^{40} + x^{32} + x^{30}, \\
p_9(x) &= x^{131} + x^{127} + x^{126} + x^{125} + x^{124} + x^{118} + x^{117} + x^{116} + x^{114} + x^{113} \\
& + x^{112} + x^{111} + x^{99} + x^{95} + x^{94} + x^{93} + x^{92} + x^{86} + x^{85} + x^{84} + x^{82} \\
& + x^{81} + x^{80} + x^{79} + x^{35} + x^{31} + x^{30} + x^{29} + x^{28} + x^{22} + x^{21} + x^{20} \\
& + x^{18} + x^{17} + x^{16} + x^{15}, \\
p_{12}(x) &= x^{132} + x^{120} + x^{116} + x^{112} + x^{108} + x^{96} + x^{88} + x^{84} + x^{80} + x^{76} + x^{68} \\
& + x^{64} + x^{36} + x^{24} + x^{20} + x^{16} + x^{12} + x^4 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 1, 1, 1, 0, 1, 0, 0, 1, 1, 1, 1, 0, 0, 1, 0, 0, 0, 1, 0, 1, 0, 1, 1]$.

For $\phi(W^o, 1, 0)$:

$$\begin{aligned}
p_0(x) &= x^{132} + x^{129} + x^{126} + x^{123} + x^{122} + x^{120} + x^{119} + x^{117} + x^{115} + x^{114} \\
& + x^{109} + x^{108} + x^{106} + x^{105} + x^{104} + x^{103} + x^{101} + x^{100} + x^{97} + x^{94} \\
& + x^{93} + x^{92} + x^{86} + x^{82} + x^{81} + x^{80} + x^{79} + x^{75} + x^{74} + x^{72} + x^{70} \\
& + x^{67} + x^{66} + x^{65} + x^{63} + x^{60} + x^{58} + x^{55} + x^{52} + x^{51} + x^{49} + x^{47} \\
& + x^{46} + x^{45} + x^{44} + x^{42} + x^{36}, \\
p_3(x) &= x^{125} + x^{120} + x^{119} + x^{118} + x^{116} + x^{114} + x^{113} + x^{111} + x^{110} + x^{109} \\
& + x^{108} + x^{107} + x^{105} + x^{103} + x^{100} + x^{99} + x^{97} + x^{95} + x^{93} + x^{92} \\
& + x^{91} + x^{89} + x^{88} + x^{86} + x^{83} + x^{82} + x^{78} + x^{72} + x^{71} + x^{70} + x^{68} \\
& + x^{66} + x^{65} + x^{63} + x^{62} + x^{60} + x^{59} + x^{57} + x^{56} + x^{54} + x^{51} + x^{50} \\
& + x^{46} + x^{45}, \\
p_6(x) &= x^{126} + x^{122} + x^{118} + x^{116} + x^{114} + x^{108} + x^{106} + x^{104} + x^{100} + x^{96} \\
& + x^{88} + x^{82} + x^{78} + x^{76} + x^{74} + x^{68} + x^{66} + x^{64} + x^{60} + x^{56} + x^{54} \\
& + x^{50} + x^{48} + x^{46} + x^{44} + x^{38} + x^{32} + x^{30},
\end{aligned}$$

$$\begin{aligned}
p_9(x) &= x^{127} + x^{122} + x^{121} + x^{120} + x^{119} + x^{118} + x^{117} + x^{116} + x^{114} + x^{113} \\
&\quad + x^{112} + x^{111} + x^{95} + x^{90} + x^{89} + x^{88} + x^{87} + x^{86} + x^{85} + x^{84} + x^{82} \\
&\quad + x^{81} + x^{80} + x^{79} + x^{31} + x^{26} + x^{25} + x^{24} + x^{23} + x^{22} + x^{21} + x^{20} \\
&\quad + x^{18} + x^{17} + x^{16} + x^{15}, \\
p_{12}(x) &= x^{128} + x^{124} + x^{116} + x^{112} + x^{104} + x^{100} + x^{92} + x^{84} + x^{80} + x^{72} \\
&\quad + x^{68} + x^{64} + x^{32} + x^{28} + x^{20} + x^{16} + x^8 + x^4 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 1, 1, 0, 1, 1, 0, 1, 1, 0, 1, 0, 1, 0, 1, 1, 1, 0, 0, 0]$.

For $\phi(W^o, 1, 1)$:

$$\begin{aligned}
p_0(x) &= x^{126} + x^{121} + x^{120} + x^{118} + x^{117} + x^{116} + x^{112} + x^{111} + x^{110} + x^{109} \\
&\quad + x^{108} + x^{107} + x^{106} + x^{103} + x^{102} + x^{99} + x^{98} + x^{97} + x^{91} + x^{90} \\
&\quad + x^{89} + x^{88} + x^{84} + x^{83} + x^{82} + x^{77} + x^{74} + x^{73} + x^{71} + x^{70} + x^{67} \\
&\quad + x^{65} + x^{61} + x^{60} + x^{57} + x^{54} + x^{52} + x^{51} + x^{49} + x^{48} + x^{46} + x^{44} \\
&\quad + x^{41} + x^{40} + x^{36}, \\
p_3(x) &= x^{125} + x^{122} + x^{121} + x^{120} + x^{119} + x^{114} + x^{113} + x^{112} + x^{111} + x^{109} \\
&\quad + x^{108} + x^{106} + x^{105} + x^{103} + x^{102} + x^{101} + x^{100} + x^{99} + x^{96} + x^{94} \\
&\quad + x^{93} + x^{92} + x^{90} + x^{89} + x^{88} + x^{86} + x^{84} + x^{83} + x^{82} + x^{81} + x^{80} \\
&\quad + x^{79} + x^{78} + x^{72} + x^{71} + x^{69} + x^{66} + x^{65} + x^{63} + x^{60} + x^{56} + x^{54} \\
&\quad + x^{53} + x^{52} + x^{51} + x^{50} + x^{49} + x^{47} + x^{46} + x^{45} + x^{42} + x^{41} + x^{37} \\
&\quad + x^{32}, \\
p_6(x) &= x^{125} + x^{122} + x^{121} + x^{119} + x^{118} + x^{116} + x^{115} + x^{114} + x^{113} + x^{111} \\
&\quad + x^{110} + x^{107} + x^{106} + x^{104} + x^{103} + x^{97} + x^{93} + x^{92} + x^{91} + x^{88} \\
&\quad + x^{84} + x^{83} + x^{82} + x^{79} + x^{77} + x^{76} + x^{75} + x^{74} + x^{71} + x^{70} + x^{69} \\
&\quad + x^{67} + x^{66} + x^{63} + x^{61} + x^{60} + x^{57} + x^{56} + x^{52} + x^{51} + x^{50} + x^{49} \\
&\quad + x^{48} + x^{47} + x^{46} + x^{41} + x^{38} + x^{37} + x^{36} + x^{32} + x^{30} + x^{27} + x^{22} \\
&\quad + x^{21} + x^{20} + x^{17} + x^{12} + x^5 + x^4 + 1, \\
p_9(x) &= x^{125} + x^{124} + x^{121} + x^{116} + x^{93} + x^{92} + x^{89} + x^{84} + x^{29} + x^{28} + x^{25} \\
&\quad + x^{20}, \\
p_{12}(x) &= x^{125} + x^{124} + x^{120} + x^{117} + x^{116} + x^{112} + x^{101} + x^{100} + x^{96} + x^{93} \\
&\quad + x^{92} + x^{88} + x^{85} + x^{84} + x^{80} + x^{69} + x^{68} + x^{64} + x^{29} + x^{28} + x^{24} \\
&\quad + x^{21} + x^{20} + x^{16} + x^5 + x^4 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 0, 1, 0, 0, 0, 1, 1, 1, 0, 1, 0, 0, 0, 0, 1, 1, 0, 1, 1]$.

Below is the transition function and output function of an 2-automaton that generates $T = M_{0,0}^o$:

Transition function $(n, j) \mapsto \delta(n, j)$ ($\Lambda(n) := [\delta(n, 0), \delta(n, 1)]$):

n	$\Lambda(n)$	n	$\Lambda(n)$	n	$\Lambda(n)$	n	$\Lambda(n)$	n	$\Lambda(n)$
0	[1, 2]	50	[81, 82]	100	[138, 72]	150	[193, 116]	200	[217, 218]
1	[3, 4]	51	[83, 31]	101	[139, 140]	151	[194, 195]	201	[219, 220]
2	[5, 6]	52	[84, 85]	102	[26, 141]	152	[34, 196]	202	[221, 101]
3	[7, 8]	53	[8, 86]	103	[142, 143]	153	[30, 197]	203	[167, 222]
4	[9, 10]	54	[87, 8]	104	[15, 126]	154	[66, 198]	204	[178, 217]
5	[11, 12]	55	[88, 89]	105	[144, 145]	155	[150, 149]	205	[223, 190]
6	[13, 14]	56	[90, 91]	106	[146, 140]	156	[199, 143]	206	[224, 186]
7	[5, 15]	57	[92, 93]	107	[142, 147]	157	[61, 58]	207	[225, 226]
8	[16, 17]	58	[94, 95]	108	[26, 135]	158	[200, 201]	208	[227, 228]
9	[18, 16]	59	[96, 97]	109	[148, 149]	159	[202, 76]	209	[229, 230]
10	[19, 20]	60	[14, 42]	110	[22, 137]	160	[70, 203]	210	[231, 168]
11	[21, 19]	61	[98, 99]	111	[51, 18]	161	[130, 54]	211	[118, 46]
12	[22, 23]	62	[100, 98]	112	[6, 126]	162	[139, 151]	212	[232, 233]
13	[24, 25]	63	[101, 102]	113	[150, 124]	163	[15, 26]	213	[86, 234]
14	[6, 26]	64	[103, 104]	114	[136, 23]	164	[51, 16]	214	[155, 110]
15	[27, 28]	65	[105, 106]	115	[130, 53]	165	[48, 131]	215	[199, 147]
16	[29, 30]	66	[107, 108]	116	[146, 151]	166	[204, 63]	216	[145, 61]
17	[31, 32]	67	[93, 109]	117	[145, 50]	167	[127, 75]	217	[205, 59]
18	[18, 18]	68	[110, 111]	118	[65, 76]	168	[74, 128]	218	[25, 135]
19	[33, 34]	69	[80, 112]	119	[123, 133]	169	[17, 54]	219	[212, 153]
20	[35, 36]	70	[82, 113]	120	[53, 18]	170	[205, 56]	220	[24, 26]
21	[37, 35]	71	[114, 115]	121	[152, 153]	171	[24, 126]	221	[235, 69]
22	[38, 39]	72	[116, 88]	122	[57, 131]	172	[126, 58]	222	[53, 17]
23	[40, 41]	73	[106, 117]	123	[154, 155]	173	[202, 66]	223	[50, 236]
24	[42, 43]	74	[45, 118]	124	[156, 157]	174	[206, 71]	224	[149, 237]
25	[44, 45]	75	[119, 120]	125	[43, 94]	175	[51, 17]	225	[60, 208]
26	[46, 47]	76	[121, 122]	126	[12, 10]	176	[5, 24]	226	[208, 135]
27	[15, 48]	77	[123, 68]	127	[158, 159]	177	[126, 207]	227	[206, 203]
28	[49, 50]	78	[124, 20]	128	[160, 161]	178	[132, 49]	228	[19, 129]
29	[17, 51]	79	[60, 57]	129	[162, 163]	179	[51, 130]	229	[15, 25]
30	[16, 18]	80	[125, 126]	130	[164, 87]	180	[149, 73]	230	[126, 131]
31	[52, 53]	81	[127, 128]	131	[28, 165]	181	[69, 61]	231	[238, 200]
32	[52, 54]	82	[124, 129]	132	[166, 167]	182	[208, 141]	232	[208, 239]
33	[55, 56]	83	[130, 52]	133	[168, 169]	183	[48, 209]	233	[144, 6]
34	[57, 58]	84	[53, 130]	134	[170, 171]	184	[210, 145]	234	[200, 129]
35	[55, 59]	85	[18, 17]	135	[112, 172]	185	[63, 73]	235	[173, 114]
36	[60, 61]	86	[54, 53]	136	[10, 173]	186	[148, 124]	236	[240, 241]
37	[62, 60]	87	[54, 54]	137	[174, 175]	187	[145, 208]	237	[242, 243]
38	[63, 64]	88	[61, 131]	138	[176, 170]	188	[211, 49]	238	[196, 33]
39	[65, 66]	89	[132, 15]	139	[177, 178]	189	[66, 20]	239	[244, 245]
40	[67, 68]	90	[53, 16]	140	[179, 180]	190	[211, 15]	240	[24, 48]
41	[17, 53]	91	[66, 129]	141	[181, 182]	191	[212, 213]	241	[50, 141]
42	[49, 61]	92	[67, 133]	142	[183, 184]	192	[25, 141]	242	[152, 213]

43	[69, 50]	93	[76, 134]	143	[32, 185]	193	[214, 125]	243	[60, 50]
44	[70, 71]	94	[125, 26]	144	[186, 81]	194	[17, 52]	244	[145, 57]
45	[72, 73]	95	[50, 135]	145	[187, 181]	195	[200, 20]	245	[48, 58]
46	[74, 75]	96	[130, 51]	146	[108, 188]	196	[210, 6]		
47	[76, 73]	97	[63, 134]	147	[85, 189]	197	[149, 134]		
48	[77, 78]	98	[136, 137]	148	[190, 162]	198	[215, 216]		
49	[79, 80]	99	[72, 134]	149	[191, 192]	199	[192, 105]		

Output function $n \mapsto \tau(n)$:

n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$
0	0	36	1	72	0	108	0	144	0	180	0	216	0
1	0	37	0	73	0	109	0	145	0	181	0	217	1
2	0	38	1	74	0	110	1	146	0	182	0	218	1
3	0	39	0	75	0	111	0	147	0	183	0	219	0
4	0	40	1	76	1	112	1	148	0	184	1	220	1
5	0	41	1	77	0	113	1	149	0	185	1	221	1
6	1	42	1	78	1	114	0	150	1	186	0	222	1
7	0	43	0	79	1	115	0	151	1	187	0	223	1
8	1	44	1	80	0	116	0	152	1	188	0	224	0
9	0	45	0	81	1	117	0	153	1	189	0	225	1
10	0	46	0	82	1	118	0	154	0	190	0	226	0
11	0	47	1	83	0	119	0	155	1	191	0	227	0
12	1	48	0	84	1	120	1	156	1	192	1	228	0
13	1	49	1	85	0	121	1	157	0	193	1	229	0
14	1	50	1	86	0	122	1	158	1	194	1	230	1
15	0	51	0	87	0	123	0	159	1	195	1	231	1
16	1	52	1	88	0	124	1	160	1	196	1	232	0
17	1	53	1	89	1	125	0	161	0	197	0	233	0
18	0	54	0	90	1	126	1	162	1	198	1	234	1
19	0	55	0	91	0	127	1	163	0	199	1	235	1
20	0	56	1	92	1	128	1	164	0	200	1	236	1
21	0	57	1	93	1	129	1	165	0	201	0	237	1
22	1	58	0	94	0	130	0	166	1	202	1	238	1
23	1	59	0	95	1	131	1	167	1	203	1	239	0
24	1	60	1	96	0	132	1	168	0	204	1	240	1
25	1	61	0	97	1	133	0	169	1	205	1	241	1
26	0	62	0	98	0	134	1	170	1	206	0	242	1
27	0	63	1	99	0	135	1	171	1	207	1	243	1
28	1	64	0	100	0	136	0	172	1	208	0	244	0
29	1	65	0	101	1	137	0	173	1	209	0	245	0
30	1	66	0	102	0	138	0	174	0	210	1		
31	1	67	1	103	0	139	1	175	0	211	0		
32	1	68	1	104	0	140	0	176	0	212	0		
33	0	69	0	105	0	141	0	177	1	213	0		

34	1	70	1	106	0	142	0	178	1	214	1	
35	0	71	0	107	0	143	1	179	0	215	1	

UNIVERSITÉ DE STRASBOURG, CNRS, IRMA UMR 7501, F-67000 STRASBOURG, FRANCE
Email address: `guoniu.han@unistra.fr`

SCHOOL OF MATHEMATICS AND STATISTICS, HUAZHONG UNIVERSITY OF SCIENCE AND TECHNOLOGY, WUHAN, PR CHINA
Email address: `huyining@hust.edu.cn`