

PROOF OF THE CONTINUED FRACTION CONJECTURE FOR

$$(a, b) = (z^5 + z^3 + z^2 + 1, z + 1)$$

GUO-NIU HAN* AND YINING HU*

ABSTRACT. The continued fraction conjecture claims that the continued fraction defined by the (a, b) -Thue-Morse sequence is algebraic over $\mathbb{F}_2(x)$ for all pairs of distinct elements (a, b) in $\mathbb{F}_2[x] \setminus \mathbb{F}_2$. In this paper we prove the conjecture for $(a, b) = (z^5 + z^3 + z^2 + 1, z + 1)$.

1. INTRODUCTION

Let $\mathbb{F}_2$ be the finite field of cardinality 2. Given a sequence of polynomials $\mathbf{a} = (a_0(z), a_1(z), a_2(z), \dots)$ of elements from $\mathbb{F}_2[z] \setminus \mathbb{F}_2$, we define the infinite continued fraction

$$(1.1) \quad \text{CF}(\mathbf{a}(z)) = \frac{1}{a_0(z) + \frac{1}{a_1(z) + \frac{1}{a_2(z) + \frac{1}{\ddots}}}}$$

as the limit of the finite continued fractions

$$(1.2) \quad \text{CF}_n(\mathbf{a}(z)) = \frac{1}{a_0(z) + \frac{1}{a_1(z) + \frac{1}{\ddots + \frac{1}{a_n(z)}}}} \in \mathbb{F}_2((1/z)).$$

Conversely, each power series in $\mathbb{F}_2[[1/z]]$ without constant term

$$(1.3) \quad c_{-1}z^{-1} + c_{-2}z^{-2} + c_{-3}z^{-3} + \dots$$

admits a continued fraction expansion of form (1.1).

Let a and b be two distinct elements from $\mathbb{F}_2[z] \setminus \mathbb{F}_2$. We consider the continued fraction (1.1) associated with the (a, b) -Thue-Morse sequence

$$\mathbf{t} = (t_0(z), t_1(z), t_2(z), \dots) = (a, b, b, a, b, a, a, b, \dots)$$

that we shall denote

$$(1.4) \quad \text{CF}(a, b) := \text{CF}(\mathbf{t}).$$

Date: 9/04/2020 15:49:34 .

2010 Mathematics Subject Classification. 11B85, 11J70, 11B50, 11Y65, 05A15.

Key words and phrases. automatic sequence, continued fraction, Thue-Morse sequence.

* This paper was automatically generated from a computer program developed by the authors, with 9.2 minutes CPU time used.

Theorem 1.1. *When $(a, b) = (z^5 + z^3 + z^2 + 1, z + 1)$, the two power series $\text{CF}(a, b)$ and $\text{CF}(b, a)$ are algebraic over $\mathbb{F}_2(z)$, with minimal polynomials of the form*

$$p_4(z)y^4 + p_3(z)y^3 + p_2(z)y^2 + p_1(z)y + p_0(z) = 0$$

where, for $\text{CF}(a, b)$,

$$\begin{aligned} p_0(z) &= z^{14} + z^{13} + z^{12} + z^{10} + z^9 + z^8 + z^7 + z + 1, \\ p_1(z) &= z^{19} + z^{18} + z^{14} + z^{12} + z^{11} + z^6 + z^3 + z, \\ p_2(z) &= z^{20} + z^{18} + z^{16} + z^{14} + z^{10} + z^8 + z^2 + 1, \\ p_3(z) &= z^{19} + z^{18} + z^{14} + z^{12} + z^{11} + z^6 + z^3 + z, \\ p_4(z) &= z^{18} + z^{13} + z^{12} + z^9 + z^8 + z^7 + z^4 + z^2 + z + 1, \end{aligned}$$

and for $\text{CF}(b, a)$,

$$\begin{aligned} p_0(z) &= z^{16} + z^{14} + z^{13} + z^{10} + z^9 + z^8 + z^7 + z^4 + z + 1, \\ p_1(z) &= z^{19} + z^{18} + z^{14} + z^{12} + z^{11} + z^6 + z^3 + z, \\ p_2(z) &= z^{20} + z^{18} + z^{16} + z^{14} + z^{10} + z^8 + z^2 + 1, \\ p_3(z) &= z^{19} + z^{18} + z^{14} + z^{12} + z^{11} + z^6 + z^3 + z, \\ p_4(z) &= z^{18} + z^{16} + z^{13} + z^9 + z^8 + z^7 + z^2 + z + 1. \end{aligned}$$

2. PROOF

Define the sequences of polynomials $(P_n(z))_{n \geq 0}$ and $(Q_n(z))_{n \geq 0}$ by

$$\begin{pmatrix} P_n(z) & Q_n(z) \\ P_{n-1}(z) & Q_{n-1}(z) \end{pmatrix} := \begin{pmatrix} t_n(z) & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} t_{n-1}(z) & 1 \\ 1 & 0 \end{pmatrix} \cdots \begin{pmatrix} t_0(z) & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}.$$

By the basic properties of continued fractions, we have

$$(2.1) \quad \text{CF}_n(a, b) = \frac{P_n(z)}{Q_n(z)} \in \mathbb{F}_2((1/z)).$$

As formula (2.1) is not efficient for calculating $\text{CF}_n(a, b)$, a fast method is to be derived and described as follows. Define

$$\begin{aligned} M_n(x) &= x^{\deg(t_{2^n-1})} \begin{pmatrix} t_{2^n-1}(1/x) & 1 \\ 1 & 0 \end{pmatrix} \times x^{\deg(t_{2^n-2})} \begin{pmatrix} t_{2^n-2}(1/x) & 1 \\ 1 & 0 \end{pmatrix} \times \\ &\quad \cdots \times x^{\deg(t_0)} \begin{pmatrix} t_0(1/x) & 1 \\ 1 & 0 \end{pmatrix}, \\ W_n(x) &= x^{\deg(\bar{t}_{2^n-1})} \begin{pmatrix} \bar{t}_{2^n-1}(1/x) & 1 \\ 1 & 0 \end{pmatrix} \times x^{\deg(\bar{t}_{2^n-2})} \begin{pmatrix} \bar{t}_{2^n-2}(1/x) & 1 \\ 1 & 0 \end{pmatrix} \times \\ &\quad \cdots \times x^{\deg(\bar{t}_0)} \begin{pmatrix} \bar{t}_0(1/x) & 1 \\ 1 & 0 \end{pmatrix}, \end{aligned}$$

where $\bar{t}$ is the (b, a) -Thue-Morse sequence. By the properties of the Thue-Morse sequence, we have for $n \geq 0$

$$\begin{aligned} M_{n+1}(x) &= W_n(x) \cdot M_n(x), \\ W_{n+1}(x) &= M_n(x) \cdot W_n(x). \end{aligned}$$

Let $x := 1/z$. For each non-zero polynomial $P(z)$, define $\tilde{P}(x)$ to be $P(1/x)$. Then,

$$\text{CF}_n(a, b) = \frac{P_n(z)}{Q_n(z)} = \frac{\tilde{P}_n(x)}{\tilde{Q}_n(x)} \in \mathbb{F}_2((x)) = \mathbb{F}_2((1/z)).$$

Comparing the definition of $M_n(x)$ with the definition of $P_n(x)$ and $Q_n(x)$, we see that

$$\begin{aligned} M_n(x)_{0,1} &= x^{d_n} \tilde{P}_{2^n-1}(x), \\ M_n(x)_{0,0} &= x^{d_n} \tilde{Q}_{2^n-1}(x), \end{aligned}$$

for some positive integer d_n . Hence,

$$(2.2) \quad \text{CF}_{2^{2^n}-1}(a, b) = \frac{\tilde{P}_{2^{2^n}-1}(x)}{\tilde{Q}_{2^{2^n}-1}(x)} = \frac{M_{2^n}(x)_{0,1}}{M_{2^n}(x)_{0,0}}.$$

Therefore, by the convergence theorem of continued fraction, the algebraicity of $\text{CF}(a, b)$ will be established if it is shown that both $M_{2^n}(x)_{0,1}$ and $M_{2^n}(x)_{0,0}$ converge to algebraic series in $\mathbb{F}_2[[x]]$.

Actually, we will prove that for all $0 \leq i, j \leq 1$ the four sequences $(M_{2^n}(x)_{i,j})_n$, $(M_{2^{n+1}}(x)_{i,j})_n$, $(W_{2^n}(x)_{i,j})_n$, and $(W_{2^{n+1}}(x)_{i,j})_n$ converge to algebraic series in $\mathbb{F}_2[[x]]$. For this purpose, we define four 2×2 matrices M^e, M^o, W^e, W^o as follows: For each $T \in \{M^e, M^o, W^e, W^o\}$ and $0 \leq i, j \leq 1$, $T_{i,j}$ is defined to be the unique solution in $\mathbb{F}_2[[x]]$ of the polynomial $\phi(T, i, j)$ under certain initial conditions; the polynomials $\phi(T, i, j)$ and initial conditions are given in Section 3. We will prove that these four matrices, whose components are algebraic by definition, are the limits of $(M_{2^n}(x))_n$, $(M_{2^{n+1}}(x))_n$, $(W_{2^n}(x))_n$, and $(W_{2^{n+1}}(x))_n$.

Let us explain how the polynomials $\phi(T, i, j)$ and initial conditions are found, and why the solutions exist and are unique. For $0 \leq i, j \leq 1$, the coefficients of the polynomial $\phi(M^e, i, j)$ (resp. $\phi(M^o, i, j)$, $\phi(W^e, i, j)$, and $\phi(W^o, i, j)$) are the Padé-Hermite approximants of type

$$(150, 150, 150, 150, 150)$$

of the vector

$$(1, T^3, T^6, T^9, T^{12}),$$

where $T = M_{12,i,j}$ (resp. $M_{11,i,j}$, $W_{12,i,j}$, and $W_{11,i,j}$), found by the Derksen algorithm. We take the first 8 terms of $M_{12,i,j}$ (resp. $M_{11,i,j}$, $W_{12,i,j}$, and $W_{11,i,j}$) as initial conditions for $\phi(M^e, i, j)$ (resp. $\phi(M^o, i, j)$, $\phi(W^e, i, j)$, and $\phi(W^o, i, j)$). The following fact will be used to ensure that the solution exists and is unique: let $P(x, y) \in \mathbb{F}_2[x, y]$ and for each series $f(x) = \sum_{n=0}^{\infty} a_n x^n \in \mathbb{F}_2[[x]]$ denote the partial sum $\sum_{j=0}^{n-1} a_j x^j$ by $f_n(x)$ for $n \geq 0$. If for some $n \geq 0$ and $a_0, a_1, \dots, a_{n-1} \in \mathbb{F}_2$ $P(x, \sum_{j=0}^{n-1} a_j x^j) = O(x^n)$ and $Q(x, y) := P(x, \sum_{j=0}^{n-1} a_j x^j + x^n y)$ can be written as $x^m \sum_{j=0}^{\infty} q_j(x) y^j$ where $q_j(x)$ are polynomials for $j \geq 0$, $q_1(0) = 1$, and $q_j(0) = 0$ for $j > 1$, then there exists a unique solution $f(x) \in \mathbb{F}_2[[x]]$ of $P(x, f(x)) = 0$ that satisfies the initial condition $f_n(x) = \sum_{j=0}^{n-1} a_j x^j$.

We state two lemmas concerning the four matrices M^e, M^o, W^e, W^o . The first one is about relations between them; the second, about the structure of the each matrix.

Lemma 2.1. *We have*

$$(2.3) \quad M^e = W^o \cdot M^o,$$

$$(2.4) \quad M^o = W^e \cdot M^e,$$

$$(2.5) \quad W^e = M^o \cdot W^o,$$

$$(2.6) \quad W^o = M^e \cdot W^e.$$

Proof. We give the proof of the identity

$$M_{0,0}^e = W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o,$$

the proofs of the others being similar.

First, we compute the minimal polynomials of $W_{0,0}^o M_{0,0}^o$ and $W_{0,1}^o M_{1,0}^o$. We know that

$$P(x, y) = \text{Res}_z (\phi(W^o, 0, 0)(x, z), z^{12} \cdot \phi(M^o, 0, 0)(x, y/z))$$

is an annihilating polynomial of $W_{0,0}^o M_{0,0}^o$. We use Padé-Hermite approximation to find a candidate for the minimal polynomial of $W_{0,0}^o M_{0,0}^o$, that will be called $\phi_0(x, y)$. To prove that $\phi_0(x, y)$ is indeed the minimal polynomial, it suffices to prove that it is an irreducible factor of $P(x, y)$ of multiplicity m and that $Q(x, y) := P(x, y)/\phi_0(x, y)^m$ is not an annihilating polynomial of $W_{0,0}^o M_{0,0}^o$. We verify the first point directly. For the second point, we truncate $W_{0,0}^o M_{0,0}^o$ to order 540 and substitute it for y in $Q(x, y)$. We get a series of valuation less than 540, which proves that $Q(x, y)$ is not an annihilating polynomial of $W_{0,0}^o M_{0,0}^o$. We find the minimal polynomial $\phi_1(x, y)$ of $W_{0,1}^o M_{1,0}^o$ in a similar way.

Now we prove that $\phi(M^e, 0, 0)$ is the minimal polynomial of $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$. We know that

$$S(x, y) = \text{Res}_z (\phi_0(x, z), \phi_1(x, y + z))$$

is an annihilating of $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$. We verify that $\phi(M^e, 0, 0)$ is an irreducible factor of $S(x, y)$ of multiplicity μ , and that $Q(x, y) := S(x, y)/\phi(M^e, 0, 0)^\mu$ is not an annihilating polynomial of $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$. To see the last point, we truncate $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$ to order 660 and substitute it for y in $Q(x, y)$. We get a series of valuation less than 660, and therefore $Q(x, y)$ is not an annihilating polynomial of $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$.

Finally, the first 16 terms of $M_{0,0}^e$ and $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$ coincide. As these first terms determine a unique solution of $\phi(M^e, 0, 0)$, we know that the two series are one and the same. $\square$

Define

$$R^e = \begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix} \quad \text{and} \quad R^o = \begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix}.$$

Lemma 2.2. *For all integers $k \geq 2$ and $u = 2^{2k-1}$, the following identities hold.*

$$\begin{aligned} M^e[:12u] &= M^e[:6u] + x^u M^e[:5u] + x^{3u} M^e[:3u] + x^{5u} M^e[:u] + x^u M^e[:6u] \\ &\quad + x^{2u} M^e[:5u] + x^{4u} M^e[:3u] + x^{6u} M^e[:u] + x^{2u} M^e[:6u] \\ &\quad + x^{3u} M^e[:5u] + x^{5u} M^e[:3u] + x^{7u} M^e[:u] + x^{4u} M^e[:6u] \\ &\quad + x^{5u} M^e[:5u] + x^{7u} M^e[:3u] + x^{9u} M^e[:u] + x^{5u} M^e[:6u] \\ &\quad + x^{6u} M^e[:5u] + x^{8u} M^e[:3u] + x^{10u} M^e[:u] + x^{7u} M^e[:5u] \\ &\quad + x^{10u} M^e[:2u] + x^{9u} M^e[:3u] + x^{11u} M^e[:u] \end{aligned}$$

$$\begin{aligned}
& + (x^{6u} + x^{7u} + x^{8u} + x^{10u} + x^{11u})R^e, \\
W^e[:12u] &= W^e[:6u] + x^u W^e[:5u] + x^{3u} W^e[:3u] + x^{5u} W^e[:u] + x^u W^e[:6u] \\
& + x^{2u} W^e[:5u] + x^{4u} W^e[:3u] + x^{6u} W^e[:u] + x^{2u} W^e[:6u] \\
& + x^{3u} W^e[:5u] + x^{5u} W^e[:3u] + x^{7u} W^e[:u] + x^{4u} W^e[:6u] \\
& + x^{5u} W^e[:5u] + x^{7u} W^e[:3u] + x^{9u} W^e[:u] + x^{5u} W^e[:6u] \\
& + x^{6u} W^e[:5u] + x^{8u} W^e[:3u] + x^{10u} W^e[:u] + x^{7u} W^e[:5u] \\
& + x^{10u} W^e[:2u] + x^{9u} W^e[:3u] + x^{11u} W^e[:u] \\
& + (x^{6u} + x^{7u} + x^{8u} + x^{10u} + x^{11u})R^e.
\end{aligned}$$

For all integers $k \geq 2$ and $u = 2^{2k}$, the following identities hold.

$$\begin{aligned}
M^o[:12u] &= M^o[:6u] + x^u M^o[:5u] + x^{3u} M^o[:3u] + x^{5u} M^o[:u] + x^u M^o[:6u] \\
& + x^{2u} M^o[:5u] + x^{4u} M^o[:3u] + x^{6u} M^o[:u] + x^{2u} M^o[:6u] \\
& + x^{3u} M^o[:5u] + x^{5u} M^o[:3u] + x^{7u} M^o[:u] + x^{4u} M^o[:6u] \\
& + x^{5u} M^o[:5u] + x^{7u} M^o[:3u] + x^{9u} M^o[:u] + x^{5u} M^o[:6u] \\
& + x^{6u} M^o[:5u] + x^{8u} M^o[:3u] + x^{10u} M^o[:u] + x^{7u} M^o[:5u] \\
& + x^{10u} M^o[:2u] + x^{9u} M^o[:3u] + x^{11u} M^o[:u] \\
& + (x^{6u} + x^{7u} + x^{8u} + x^{10u} + x^{11u})R^o, \\
W^o[:12u] &= W^o[:6u] + x^u W^o[:5u] + x^{3u} W^o[:3u] + x^{5u} W^o[:u] + x^u W^o[:6u] \\
& + x^{2u} W^o[:5u] + x^{4u} W^o[:3u] + x^{6u} W^o[:u] + x^{2u} W^o[:6u] \\
& + x^{3u} W^o[:5u] + x^{5u} W^o[:3u] + x^{7u} W^o[:u] + x^{4u} W^o[:6u] \\
& + x^{5u} W^o[:5u] + x^{7u} W^o[:3u] + x^{9u} W^o[:u] + x^{5u} W^o[:6u] \\
& + x^{6u} W^o[:5u] + x^{8u} W^o[:3u] + x^{10u} W^o[:u] + x^{7u} W^o[:5u] \\
& + x^{10u} W^o[:2u] + x^{9u} W^o[:3u] + x^{11u} W^o[:u] \\
& + (x^{6u} + x^{7u} + x^{8u} + x^{10u} + x^{11u})R^o.
\end{aligned}$$

Proof. To prove Lemma 2.2 we first construct an automaton for each sequence concerned, and then transform the conditions on infinitely many k 's into finitely many conditions on the states of the automaton. In the following, we will prove that for $T = M_{0,0}^o$, for all integer $k \geq 2$ and $u = 2^{2k}$,

$$\begin{aligned}
T[:12u] &= T[:6u] + x^u T[:5u] + x^{3u} T[:3u] + x^{5u} T[:u] + x^u T[:6u] + x^{2u} T[:5u] \\
& + x^{4u} T[:3u] + x^{6u} T[:u] + x^{2u} T[:6u] + x^{3u} T[:5u] + x^{5u} T[:3u] \\
& + x^{7u} T[:u] + x^{4u} T[:6u] + x^{5u} T[:5u] + x^{7u} T[:3u] + x^{9u} T[:u] \\
& + x^{5u} T[:6u] + x^{6u} T[:5u] + x^{8u} T[:3u] + x^{10u} T[:u] + x^{7u} T[:5u] \\
& + x^{10u} T[:2u] + x^{9u} T[:3u] + x^{11u} T[:u] \\
& + (x^{6u} + x^{7u} + x^{8u} + x^{10u} + x^{11u}).
\end{aligned}$$

The proof of the other 15 cases are similar. We break down the above identity into 6 parts:

$$x^6 u = x^{5u} T[u:2u] + x^{3u} T[3u:4u] + x^u T[5u:6u] + T[6u:7u],$$

$$\begin{aligned}
x^7u &= x^{7u}T[:u] + x^{6u}T[u:2u] + x^{5u}T[2u:3u] + x^{4u}T[3u:4u] \\
&\quad + x^{3u}T[4u:5u] + x^{2u}T[5u:6u] + T[7u:8u], \\
x^8u &= x^{8u}T[:u] + x^{6u}T[2u:3u] + x^{4u}T[4u:5u] + T[8u:9u], \\
0 &= x^{8u}T[u:2u] + x^{6u}T[3u:4u] + x^{4u}T[5u:6u] + T[9u:10u], \\
x^10u &= x^{9u}T[u:2u] + x^{8u}T[2u:3u] + x^{7u}T[3u:4u] + x^{6u}T[4u:5u] \\
&\quad + x^{5u}T[5u:6u] + T[10u:11u], \\
x^11u &= x^{11u}T[:u] + x^{10u}T[u:2u] + x^{9u}T[2u:3u] + x^{7u}T[4u:5u] \\
&\quad + T[11u:12u],
\end{aligned}$$

which can be rewritten as

$$\begin{aligned}
(2.7) \quad 0 &= T[[1w]_2] + T[[11w]_2] + T[[101w]_2] + T[[110w]_2], \\
0 &= T[[w]_2] + T[[1w]_2] + T[[10w]_2] + T[[11w]_2] + T[[100w]_2] \\
(2.8) \quad &+ T[[101w]_2] + T[[111w]_2], \\
(2.9) \quad 0 &= T[[w]_2] + T[[10w]_2] + T[[100w]_2] + T[[1000w]_2], \\
(2.10) \quad 0 &= T[[1w]_2] + T[[11w]_2] + T[[101w]_2] + T[[1001w]_2], \\
0 &= T[[1w]_2] + T[[10w]_2] + T[[11w]_2] + T[[100w]_2] + T[[101w]_2] \\
(2.11) \quad &+ T[[1010w]_2], \\
(2.12) \quad 0 &= T[[w]_2] + T[[1w]_2] + T[[10w]_2] + T[[100w]_2] + T[[1011w]_2],
\end{aligned}$$

for all binary word w of length $2k$ and $w \neq 0^{2k}$, and

$$\begin{aligned}
(2.13) \quad 1 &= T[[1w]_2] + T[[11w]_2] + T[[101w]_2] + T[[110w]_2], \\
1 &= T[[w]_2] + T[[1w]_2] + T[[10w]_2] + T[[11w]_2] + T[[100w]_2] \\
(2.14) \quad &+ T[[101w]_2] + T[[111w]_2], \\
(2.15) \quad 1 &= T[[w]_2] + T[[10w]_2] + T[[100w]_2] + T[[1000w]_2], \\
(2.16) \quad 0 &= T[[1w]_2] + T[[11w]_2] + T[[101w]_2] + T[[1001w]_2], \\
1 &= T[[1w]_2] + T[[10w]_2] + T[[11w]_2] + T[[100w]_2] + T[[101w]_2] \\
(2.17) \quad &+ T[[1010w]_2], \\
(2.18) \quad 1 &= T[[w]_2] + T[[1w]_2] + T[[10w]_2] + T[[100w]_2] + T[[1011w]_2],
\end{aligned}$$

for $w = 0^{2k}$.

First we calculate an 2-automaton that generates T from its minimal polynomial and its first terms. This automaton has 435 states; its transition function and output function can be found in the annex. Let $A(s, w)$ denote the state reached after reading w from right to left starting from the state s , and τ the output function. Define

$$E_{2k} = \{A(i, w) : |w| = 2k, w \neq 0^{2k}\}.$$

Identities (2.7) through (2.18) can be written as

$$\begin{aligned}
(2.19) \quad 0 &= \tau(A(s, 1)) + \tau(A(s, 11)) + \tau(A(s, 101)) + \tau(A(s, 110)), \\
0 &= \tau(A(s, \epsilon)) + \tau(A(s, 1)) + \tau(A(s, 10)) + \tau(A(s, 11)) + \tau(A(s, 100)) \\
(2.20) \quad &+ \tau(A(s, 101)) + \tau(A(s, 111)), \\
(2.21) \quad 0 &= \tau(A(s, \epsilon)) + \tau(A(s, 10)) + \tau(A(s, 100)) + \tau(A(s, 1000)),
\end{aligned}$$

$$(2.22) \quad 0 = \tau(A(s, 1)) + \tau(A(s, 11)) + \tau(A(s, 101)) + \tau(A(s, 1001)),$$

$$0 = \tau(A(s, 1)) + \tau(A(s, 10)) + \tau(A(s, 11)) + \tau(A(s, 100))$$

$$(2.23) \quad + \tau(A(s, 101)) + \tau(A(s, 1010)),$$

$$0 = \tau(A(s, \epsilon)) + \tau(A(s, 1)) + \tau(A(s, 10)) + \tau(A(s, 100))$$

$$(2.24) \quad + \tau(A(s, 1011)),$$

for all $s \in E_{2k}$, and

$$(2.25) \quad 1 = \tau(A(s, 1)) + \tau(A(s, 11)) + \tau(A(s, 101)) + \tau(A(s, 110)),$$

$$1 = \tau(A(s, \epsilon)) + \tau(A(s, 1)) + \tau(A(s, 10)) + \tau(A(s, 11)) + \tau(A(s, 100))$$

$$(2.26) \quad + \tau(A(s, 101)) + \tau(A(s, 111)),$$

$$(2.27) \quad 1 = \tau(A(s, \epsilon)) + \tau(A(s, 10)) + \tau(A(s, 100)) + \tau(A(s, 1000)),$$

$$(2.28) \quad 0 = \tau(A(s, 1)) + \tau(A(s, 11)) + \tau(A(s, 101)) + \tau(A(s, 1001)),$$

$$1 = \tau(A(s, 1)) + \tau(A(s, 10)) + \tau(A(s, 11)) + \tau(A(s, 100))$$

$$(2.29) \quad + \tau(A(s, 101)) + \tau(A(s, 1010)),$$

$$1 = \tau(A(s, \epsilon)) + \tau(A(s, 1)) + \tau(A(s, 10)) + \tau(A(s, 100))$$

$$(2.30) \quad + \tau(A(s, 1011)),$$

for $s = A(i, 0^{2k})$. We find that $(A(i, 0^{20}), E_{20}) = (A(i, 0^{16}), E_{16})$, so that we only have to verify that identities (2.19) through (2.30) hold for $2 \leq k \leq 10$, which turns out to be true. $\square$

In the following lemma, we express M_{2k} , M_{2k+1} , W_{2k} , and W_{2k+1} in terms of M^e , M^o , W^e , and W^o .

Lemma 2.3. *For all integer $k \geq 2$, and $u = 2^{2k-1}$,*

$$M_{2k} = M^e[:6u] + x^u M^e[:5u] + x^{3u} M^e[:3u] + x^{5u} M^e[:u] + x^{6u} R^e,$$

$$W_{2k} = W^e[:6u] + x^u W^e[:5u] + x^{3u} W^e[:3u] + x^{5u} W^e[:u] + x^{6u} R^e.$$

For all integer $k \geq 2$, and $u = 2^{2k}$,

$$M_{2k+1} = M^o[:6u] + x^u M^o[:5u] + x^{3u} M^o[:3u] + x^{5u} M^o[:u] + x^{6u} R^o,$$

$$W_{2k+1} = W^o[:6u] + x^u W^o[:5u] + x^{3u} W^o[:3u] + x^{5u} W^o[:u] + x^{6u} R^o.$$

Proof. We call the four identities in Lemma 2.3 also by the name M_{2k} , W_{2k} , M_{2k+1} , and W_{2k+1} . For $n = 2$, the identities can be verified directly. For $n \geq 2$, we claim that

$$\begin{aligned} M_{2k} \wedge W_{2k} &\Rightarrow M_{2k+1} \wedge W_{2k+1}, \\ M_{2k+1} \wedge W_{2k+1} &\Rightarrow M_{2k+2} \wedge W_{2k+2}. \end{aligned}$$

We give the proof of

$$M_{2k} \wedge W_{2k} \Rightarrow M_{2k+1},$$

the other ones being similar. Set $u = 2^{2k}$ and $v = 2^{2k-1}$. By definition and induction hypothesis, the left side of identity M_{2k+1} is equal to

$$(2.31) \quad \begin{aligned} W_{2k} M_{2k} &= (W^e[:6v] + x^v W^e[:5v] + x^{3v} W^e[:3v] + x^{5v} W^e[:v] + x^{6v} R^e) \times (\\ &M^e[:6v] + x^v M^e[:5v] + x^{3v} M^e[:3v] + x^{5v} M^e[:v] + x^{6v} R^e); \end{aligned}$$

Call this expression *lhs*. Note that both sides of identity M_{2k+1} have the same term of highest degree $x^{12v}R^o$. Therefore we only need to prove that their difference is $O(x^{12v})$. Using Lemma 2.1 it can be seen that the right side of identity M_{2k+1} is congruent, modulo x^{12v} , to

$$W^e[:12v]M^e[:12v] + x^{2v}W^e[:10v]M^e[:10v] + x^{6v}W^e[:6v]M^e[:6v] + x^{10v}W^e[:2v]M^e[:2v].$$

For all $n \leq 12$, replace the occurrences of $W^e[:n \cdot v]$ and $M^e[:n \cdot v]$ in the above expression by the reduction modulo $x^{n \cdot v}$ of the right side of the corresponding identity in Lemma 2.2 and get a new expression, which we call *rhs*. Define

$$\begin{aligned} X &:= x^v, \\ a_n &:= W^e[n \cdot v : (n+1) \cdot v] / X^n, \\ b_n &:= M^e[n \cdot v : (n+1) \cdot v] / X^n, \\ c &:= R^e. \end{aligned}$$

Using the notation introduced above, we can represent the expressions *lhs* (2.31) and *rhs* as polynomials in $\mathbb{F}_2[a_1, \dots, a_{12}, b_1, \dots, b_{12}, c][X]$. Note that it is not a problem that a_j commutes with b_k while W^e does not commute with M^e , because in the expressions concerned, the products of W^e -terms and M^e -terms are always in the same order. We let the computer do the simplification and check that the difference between these two polynomials is indeed $O(X^{12})$, which completes the proof. $\square$

Proof of Theorem 1.1. We prove the theorem for $\text{CF}(a, b)$; for $\text{CF}(b, a)$, the proof is similar. By Lemma 2.3, we have For all $0 \leq j, k \leq 1$,

$$\begin{aligned} \lim_{n \rightarrow \infty} M_{2n,j,k} &= M_{j,k}^e, \\ \lim_{n \rightarrow \infty} M_{2n+1,j,k} &= M_{j,k}^o, \\ \lim_{n \rightarrow \infty} W_{2n,j,k} &= W_{j,k}^e, \\ \lim_{n \rightarrow \infty} W_{2n+1,j,k} &= W_{j,k}^o. \end{aligned}$$

Let $z = 1/x$. By the convergence theorem and identity (2.2),

$$\text{CF}(a, b) = \frac{M_{0,1}^e(x)}{M_{0,0}^e(x)}.$$

By definition, $\phi(M^e, 0, 1)$ and $\phi(M^e, 0, 0)$ are minimal polynomials of $M_{0,1}^e$ and $M_{0,0}^e$. Therefore

$$P(x, y) = \text{Res}_t (\phi(M^e, 0, 1)(x, t), y^{12}\phi(M^e, 0, 1)(x, t/y))$$

is an annihilating polynomial of $f(x) = M_{0,1}^e/M_{0,0}^e$.

Define

$$Q(x, y) = q_4(x)y^4 + q_3(x)y^3 + q_2(x)y^2 + q_1(x)y + q_0(x),$$

where

$$\begin{aligned} q_0(x) &= x^{20} + x^{19} + x^{13} + x^{12} + x^{11} + x^{10} + x^8 + x^7 + x^6, \\ q_1(x) &= x^{19} + x^{17} + x^{14} + x^9 + x^8 + x^6 + x^2 + x, \\ q_2(x) &= x^{20} + x^{18} + x^{12} + x^{10} + x^6 + x^4 + x^2 + 1, \end{aligned}$$

$$\begin{aligned}
q_3(x) &= x^{19} + x^{17} + x^{14} + x^9 + x^8 + x^6 + x^2 + x, \\
q_4(x) &= x^{20} + x^{19} + x^{18} + x^{16} + x^{13} + x^{12} + x^{11} + x^8 + x^7 + x^2.
\end{aligned}$$

The polynomial $Q(x, y)$ is the candidate for the minimal polynomial of $f(x)$ found by Padé-Hermite approximation. To prove that it is indeed the minimal polynomial of $f(x)$, we only need to prove that it is an irreducible factor of $P(x, y)$ of multiplicity m and $R(x, y) := P(x, y)/Q(x, y)^m$ is not an annihilating polynomial of $f(x)$. We verify the first point directly. For the second point, we find that when we truncate $f(x)$ to order 186, and substitute it for y in $R(z, y)$, we get a series with valuation smaller than 186, which proves that $R(z, y)$ is not an annihilating polynomial of $f(x)$. Finally, $z^{12}Q(1/z, y)$ is the minimal polynomial of $\text{CF}(a, b) = f(1/z)$. $\square$

3. ANNEXE

All 16 polynomials are of the form

$$p_0(x) + p_3(x)y^3 + p_6(x)y^6 + p_9(x)y^9 + p_{12}(x)y^{12}.$$

The coefficients $p_j(x)$ and the 16 initial terms to determine the solutions uniquely are given below.

For $\phi(M^e, 0, 0)$:

$$\begin{aligned}
p_0(x) &= x^{96} + x^{92} + x^{88} + x^{86} + x^{84} + x^{82} + x^{80} + x^{76} + x^{74} + x^{72} + x^{66} \\
&\quad + x^{64} + x^{56} + x^{48} + x^{36} + x^{34} + x^{30} + x^{28} + x^{18} + x^{14} + x^{12}, \\
p_3(x) &= x^{94} + x^{90} + x^{84} + x^{82} + x^{80} + x^{78} + x^{76} + x^{72} + x^{66} + x^{62} + x^{60} \\
&\quad + x^{54} + x^{50} + x^{46} + x^{44} + x^{42} + x^{40} + x^{38} + x^{36} + x^{34} + x^{30} + x^{28} \\
&\quad + x^{22} + x^8, \\
p_6(x) &= x^{94} + x^{92} + x^{88} + x^{84} + x^{76} + x^{70} + x^{68} + x^{64} + x^{62} + x^{52} + x^{48} \\
&\quad + x^{46} + x^{40} + x^{36} + x^{32} + x^{20} + x^{16} + x^8 + x^4 + 1, \\
p_9(x) &= x^{94} + x^{88} + x^{86} + x^{84} + x^{82} + x^{80} + x^{78} + x^{70} + x^{60} + x^{56} + x^{48} \\
&\quad + x^{38} + x^{36} + x^{32} + x^{30} + x^{28} + x^{24} + x^{20} + x^{18} + x^{16} + x^{10} + x^8 \\
&\quad + x^4 + x^2, \\
p_{12}(x) &= x^{96} + x^{94} + x^{90} + x^{86} + x^{82} + x^{78} + x^{74} + x^{70} + x^{66} + x^{62} + x^{58} \\
&\quad + x^{54} + x^{50} + x^{48} + x^{32} + x^{30} + x^{26} + x^{22} + x^{18} + x^{14} + x^{10} + x^6 \\
&\quad + x^2 + 1,
\end{aligned}$$

and the initial terms are $[1, 0, 1, 0, 0, 0, 0, 0, 1, 0, 1, 0, 1, 0, 1, 0]$.

For $\phi(M^e, 0, 1)$:

$$\begin{aligned}
p_0(x) &= x^{108} + x^{107} + x^{103} + x^{102} + x^{101} + x^{100} + x^{97} + x^{96} + x^{95} + x^{94} \\
&\quad + x^{92} + x^{79} + x^{77} + x^{76} + x^{75} + x^{74} + x^{73} + x^{72} + x^{68} + x^{66} + x^{65} \\
&\quad + x^{63} + x^{61} + x^{52} + x^{51} + x^{49} + x^{48} + x^{45} + x^{44} + x^{41} + x^{40} + x^{39} \\
&\quad + x^{37} + x^{34} + x^{33} + x^{29} + x^{28} + x^{26} + x^{24}, \\
p_3(x) &= x^{99} + x^{98} + x^{97} + x^{94} + x^{93} + x^{91} + x^{85} + x^{84} + x^{82} + x^{78} + x^{77} \\
&\quad + x^{75} + x^{69} + x^{68} + x^{66} + x^{62} + x^{61} + x^{58} + x^{57} + x^{54} + x^{52} + x^{51}
\end{aligned}$$

$$\begin{aligned}
& + x^{50} + x^{46} + x^{44} + x^{43} + x^{42} + x^{38} + x^{36} + x^{35} + x^{34} + x^{30} + x^{28} \\
& + x^{27} + x^{26} + x^{22} + x^{20} + x^{17} + x^{13} + x^{12} + x^{11} + x^9, \\
p_6(x) &= x^{102} + x^{101} + x^{99} + x^{98} + x^{94} + x^{93} + x^{92} + x^{86} + x^{85} + x^{84} + x^{82} \\
& + x^{81} + x^{79} + x^{78} + x^{76} + x^{75} + x^{73} + x^{72} + x^{70} + x^{69} + x^{67} + x^{65} \\
& + x^{64} + x^{60} + x^{59} + x^{57} + x^{56} + x^{54} + x^{53} + x^{51} + x^{50} + x^{48} + x^{47} \\
& + x^{45} + x^{44} + x^{36} + x^{35} + x^{34} + x^{30} + x^{29} + x^{27} + x^{25} + x^{23} + x^{21} \\
& + x^{20} + x^{18} + x^{17} + x^{16} + x^{12} + x^{11} + x^{10} + x^8 + x^7 + x^6, \\
p_9(x) &= x^{105} + x^{104} + x^{103} + x^{101} + x^{97} + x^{96} + x^{94} + x^{91} + x^{87} + x^{86} + x^{84} \\
& + x^{83} + x^{82} + x^{81} + x^{80} + x^{77} + x^{75} + x^{74} + x^{73} + x^{70} + x^{69} + x^{66} \\
& + x^{64} + x^{60} + x^{58} + x^{55} + x^{53} + x^{52} + x^{51} + x^{48} + x^{46} + x^{43} + x^{39} \\
& + x^{38} + x^{36} + x^{32} + x^{30} + x^{26} + x^{24} + x^{23} + x^{22} + x^{18} + x^{17} + x^{15} \\
& + x^9 + x^8 + x^6 + x^3, \\
p_{12}(x) &= x^{108} + x^{107} + x^{105} + x^{103} + x^{101} + x^{99} + x^{97} + x^{96} + x^{60} + x^{59} + x^{57} \\
& + x^{55} + x^{53} + x^{51} + x^{49} + x^{48} + x^{44} + x^{43} + x^{41} + x^{39} + x^{37} + x^{35} \\
& + x^{33} + x^{32} + x^{12} + x^{11} + x^9 + x^7 + x^5 + x^3 + x + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 1, 0, 0, 1, 0, 0, 0, 1, 0, 0, 1]$.

For $\phi(M^e, 1, 0)$:

$$\begin{aligned}
p_0(x) &= x^{108} + x^{107} + x^{103} + x^{102} + x^{101} + x^{100} + x^{97} + x^{96} + x^{95} + x^{94} \\
& + x^{92} + x^{79} + x^{77} + x^{76} + x^{75} + x^{74} + x^{73} + x^{72} + x^{68} + x^{66} + x^{65} \\
& + x^{63} + x^{61} + x^{52} + x^{51} + x^{49} + x^{48} + x^{45} + x^{44} + x^{41} + x^{40} + x^{39} \\
& + x^{37} + x^{34} + x^{33} + x^{29} + x^{28} + x^{26} + x^{24}, \\
p_3(x) &= x^{99} + x^{98} + x^{97} + x^{94} + x^{93} + x^{91} + x^{85} + x^{84} + x^{82} + x^{78} + x^{77} \\
& + x^{75} + x^{69} + x^{68} + x^{66} + x^{62} + x^{61} + x^{58} + x^{57} + x^{54} + x^{52} + x^{51} \\
& + x^{50} + x^{46} + x^{44} + x^{43} + x^{42} + x^{38} + x^{36} + x^{35} + x^{34} + x^{30} + x^{28} \\
& + x^{27} + x^{26} + x^{22} + x^{20} + x^{17} + x^{13} + x^{12} + x^{11} + x^9, \\
p_6(x) &= x^{102} + x^{101} + x^{99} + x^{98} + x^{94} + x^{93} + x^{92} + x^{86} + x^{85} + x^{84} + x^{82} \\
& + x^{81} + x^{79} + x^{78} + x^{76} + x^{75} + x^{73} + x^{72} + x^{70} + x^{69} + x^{67} + x^{65} \\
& + x^{64} + x^{60} + x^{59} + x^{57} + x^{56} + x^{54} + x^{53} + x^{51} + x^{50} + x^{48} + x^{47} \\
& + x^{45} + x^{44} + x^{36} + x^{35} + x^{34} + x^{30} + x^{29} + x^{27} + x^{25} + x^{23} + x^{21} \\
& + x^{20} + x^{18} + x^{17} + x^{16} + x^{12} + x^{11} + x^{10} + x^8 + x^7 + x^6, \\
p_9(x) &= x^{105} + x^{104} + x^{103} + x^{101} + x^{97} + x^{96} + x^{94} + x^{91} + x^{87} + x^{86} + x^{84} \\
& + x^{83} + x^{82} + x^{81} + x^{80} + x^{77} + x^{75} + x^{74} + x^{73} + x^{70} + x^{69} + x^{66} \\
& + x^{64} + x^{60} + x^{58} + x^{55} + x^{53} + x^{52} + x^{51} + x^{48} + x^{46} + x^{43} + x^{39} \\
& + x^{38} + x^{36} + x^{32} + x^{30} + x^{26} + x^{24} + x^{23} + x^{22} + x^{18} + x^{17} + x^{15} \\
& + x^9 + x^8 + x^6 + x^3,
\end{aligned}$$

$$\begin{aligned}
p_{12}(x) = & x^{108} + x^{107} + x^{105} + x^{103} + x^{101} + x^{99} + x^{97} + x^{96} + x^{60} + x^{59} + x^{57} \\
& + x^{55} + x^{53} + x^{51} + x^{49} + x^{48} + x^{44} + x^{43} + x^{41} + x^{39} + x^{37} + x^{35} \\
& + x^{33} + x^{32} + x^{12} + x^{11} + x^9 + x^7 + x^5 + x^3 + x + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 1, 0, 0, 1, 0, 0, 0, 1, 0, 0, 1]$.

For $\phi(M^e, 1, 1)$:

$$\begin{aligned}
p_0(x) = & x^{120} + x^{118} + x^{116} + x^{114} + x^{106} + x^{104} + x^{98} + x^{94} + x^{92} + x^{86} \\
& + x^{84} + x^{82} + x^{80} + x^{76} + x^{72} + x^{70} + x^{64} + x^{62} + x^{60} + x^{56} + x^{54} \\
& + x^{52} + x^{48} + x^{44} + x^{42} + x^{38} + x^{36}, \\
p_3(x) = & x^{118} + x^{116} + x^{114} + x^{112} + x^{110} + x^{102} + x^{100} + x^{92} + x^{82} + x^{78} \\
& + x^{74} + x^{72} + x^{70} + x^{66} + x^{60} + x^{58} + x^{54} + x^{52} + x^{50} + x^{44} + x^{42} \\
& + x^{36} + x^{34} + x^{32} + x^{28} + x^{26} + x^{24} + x^{22} + x^{18} + x^6, \\
p_6(x) = & x^{118} + x^{116} + x^{114} + x^{112} + x^{110} + x^{108} + x^{106} + x^{104} + x^{96} + x^{92} \\
& + x^{90} + x^{88} + x^{84} + x^{74} + x^{72} + x^{68} + x^{66} + x^{64} + x^{60} + x^{56} + x^{54} \\
& + x^{52} + x^{50} + x^{48} + x^{44} + x^{34} + x^{32} + x^{30} + x^{28} + x^{20} + x^{18} + x^{12} \\
& + x^{10} + x^4 + x^2 + 1, \\
p_9(x) = & x^{118} + x^{116} + x^{114} + x^{112} + x^{110} + x^{106} + x^{104} + x^{98} + x^{96} + x^{92} \\
& + x^{90} + x^{88} + x^{86} + x^{84} + x^{80} + x^{78} + x^{72} + x^{66} + x^{64} + x^{62} + x^{54} \\
& + x^{48} + x^{46} + x^{44} + x^{42} + x^{38} + x^{36} + x^{24} + x^{20} + x^{16} + x^{10} + x^2, \\
p_{12}(x) = & x^{120} + x^{96} + x^{72} + x^{56} + x^{48} + x^{32} + x^{24} + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 0, 0, 0, 0, 1, 0, 1, 0, 1, 0]$.

For $\phi(W^e, 0, 0)$:

$$\begin{aligned}
p_0(x) = & x^{120} + x^{118} + x^{114} + x^{110} + x^{108} + x^{106} + x^{104} + x^{100} + x^{96} + x^{92} \\
& + x^{88} + x^{82} + x^{80} + x^{78} + x^{72} + x^{66} + x^{64} + x^{62} + x^{60} + x^{58} + x^{56} \\
& + x^{50} + x^{46} + x^{44} + x^{40} + x^{38} + x^{36} + x^{34} + x^{24} + x^{22} + x^{20} + x^{16} \\
& + x^{12}, \\
p_3(x) = & x^{118} + x^{116} + x^{112} + x^{110} + x^{106} + x^{104} + x^{102} + x^{86} + x^{80} + x^{70} \\
& + x^{68} + x^{66} + x^{52} + x^{46} + x^{30} + x^{26} + x^{20} + x^{18} + x^{14} + x^8, \\
p_6(x) = & x^{118} + x^{112} + x^{110} + x^{90} + x^{88} + x^{86} + x^{82} + x^{78} + x^{68} + x^{66} + x^{60} \\
& + x^{58} + x^{56} + x^{46} + x^{44} + x^{42} + x^{40} + x^{28} + x^{24} + x^{22} + x^{20} + x^{18} \\
& + x^{14} + x^{12} + x^{10} + x^6 + x^2 + 1, \\
p_9(x) = & x^{118} + x^{116} + x^{114} + x^{112} + x^{110} + x^{106} + x^{104} + x^{98} + x^{96} + x^{92} \\
& + x^{90} + x^{88} + x^{86} + x^{84} + x^{80} + x^{78} + x^{72} + x^{66} + x^{64} + x^{62} + x^{54} \\
& + x^{48} + x^{46} + x^{44} + x^{42} + x^{38} + x^{36} + x^{24} + x^{20} + x^{16} + x^{10} + x^2, \\
p_{12}(x) = & x^{120} + x^{96} + x^{72} + x^{56} + x^{48} + x^{32} + x^{24} + 1,
\end{aligned}$$

and the initial terms are $[1, 0, 0, 0, 1, 0, 0, 0, 1, 0, 1, 0, 0, 0, 1, 0]$.

For $\phi(W^e, 0, 1)$:

$$\begin{aligned} p_0(x) = & x^{108} + x^{107} + x^{103} + x^{100} + x^{99} + x^{96} + x^{93} + x^{92} + x^{89} + x^{88} + x^{87} \\ & + x^{84} + x^{83} + x^{80} + x^{78} + x^{77} + x^{75} + x^{73} + x^{71} + x^{70} + x^{68} + x^{63} \\ & + x^{61} + x^{60} + x^{59} + x^{58} + x^{57} + x^{54} + x^{53} + x^{52} + x^{50} + x^{49} + x^{48} \\ & + x^{47} + x^{44} + x^{43} + x^{42} + x^{41} + x^{40} + x^{39} + x^{36} + x^{35} + x^{33} + x^{26} \\ & + x^{24} + x^{19} + x^{18}, \end{aligned}$$

$$\begin{aligned} p_3(x) = & x^{99} + x^{98} + x^{97} + x^{94} + x^{93} + x^{91} + x^{85} + x^{84} + x^{82} + x^{78} + x^{77} \\ & + x^{75} + x^{69} + x^{68} + x^{66} + x^{62} + x^{61} + x^{58} + x^{57} + x^{54} + x^{52} + x^{51} \\ & + x^{50} + x^{46} + x^{44} + x^{43} + x^{42} + x^{38} + x^{36} + x^{35} + x^{34} + x^{30} + x^{28} \\ & + x^{27} + x^{26} + x^{22} + x^{20} + x^{17} + x^{13} + x^{12} + x^{11} + x^9, \end{aligned}$$

$$\begin{aligned} p_6(x) = & x^{102} + x^{101} + x^{99} + x^{98} + x^{94} + x^{93} + x^{92} + x^{86} + x^{85} + x^{84} + x^{82} \\ & + x^{81} + x^{79} + x^{78} + x^{76} + x^{75} + x^{73} + x^{72} + x^{70} + x^{69} + x^{67} + x^{65} \\ & + x^{64} + x^{60} + x^{59} + x^{57} + x^{56} + x^{54} + x^{53} + x^{51} + x^{50} + x^{48} + x^{47} \\ & + x^{45} + x^{44} + x^{36} + x^{35} + x^{34} + x^{30} + x^{29} + x^{27} + x^{25} + x^{23} + x^{21} \\ & + x^{20} + x^{18} + x^{17} + x^{16} + x^{12} + x^{11} + x^{10} + x^8 + x^7 + x^6, \end{aligned}$$

$$\begin{aligned} p_9(x) = & x^{105} + x^{104} + x^{103} + x^{101} + x^{97} + x^{96} + x^{94} + x^{91} + x^{87} + x^{86} + x^{84} \\ & + x^{83} + x^{82} + x^{81} + x^{80} + x^{77} + x^{75} + x^{74} + x^{73} + x^{70} + x^{69} + x^{66} \\ & + x^{64} + x^{60} + x^{58} + x^{55} + x^{53} + x^{52} + x^{51} + x^{48} + x^{46} + x^{43} + x^{39} \\ & + x^{38} + x^{36} + x^{32} + x^{30} + x^{26} + x^{24} + x^{23} + x^{22} + x^{18} + x^{17} + x^{15} \\ & + x^9 + x^8 + x^6 + x^3, \end{aligned}$$

$$\begin{aligned} p_{12}(x) = & x^{108} + x^{107} + x^{105} + x^{103} + x^{101} + x^{99} + x^{97} + x^{96} + x^{60} + x^{59} + x^{57} \\ & + x^{55} + x^{53} + x^{51} + x^{49} + x^{48} + x^{44} + x^{43} + x^{41} + x^{39} + x^{37} + x^{35} \\ & + x^{33} + x^{32} + x^{12} + x^{11} + x^9 + x^7 + x^5 + x^3 + x + 1, \end{aligned}$$

and the initial terms are $[0, 1, 1, 1, 1, 0, 0, 1, 0, 1, 0, 0, 0, 0, 1, 0]$.

For $\phi(W^e, 1, 0)$:

$$\begin{aligned} p_0(x) = & x^{108} + x^{107} + x^{103} + x^{100} + x^{99} + x^{96} + x^{93} + x^{92} + x^{89} + x^{88} + x^{87} \\ & + x^{84} + x^{83} + x^{80} + x^{78} + x^{77} + x^{75} + x^{73} + x^{71} + x^{70} + x^{68} + x^{63} \\ & + x^{61} + x^{60} + x^{59} + x^{58} + x^{57} + x^{54} + x^{53} + x^{52} + x^{50} + x^{49} + x^{48} \\ & + x^{47} + x^{44} + x^{43} + x^{42} + x^{41} + x^{40} + x^{39} + x^{36} + x^{35} + x^{33} + x^{26} \\ & + x^{24} + x^{19} + x^{18}, \end{aligned}$$

$$\begin{aligned} p_3(x) = & x^{99} + x^{98} + x^{97} + x^{94} + x^{93} + x^{91} + x^{85} + x^{84} + x^{82} + x^{78} + x^{77} \\ & + x^{75} + x^{69} + x^{68} + x^{66} + x^{62} + x^{61} + x^{58} + x^{57} + x^{54} + x^{52} + x^{51} \\ & + x^{50} + x^{46} + x^{44} + x^{43} + x^{42} + x^{38} + x^{36} + x^{35} + x^{34} + x^{30} + x^{28} \\ & + x^{27} + x^{26} + x^{22} + x^{20} + x^{17} + x^{13} + x^{12} + x^{11} + x^9, \end{aligned}$$

$$p_6(x) = x^{102} + x^{101} + x^{99} + x^{98} + x^{94} + x^{93} + x^{92} + x^{86} + x^{85} + x^{84} + x^{82}$$

$$\begin{aligned}
& + x^{81} + x^{79} + x^{78} + x^{76} + x^{75} + x^{73} + x^{72} + x^{70} + x^{69} + x^{67} + x^{65} \\
& + x^{64} + x^{60} + x^{59} + x^{57} + x^{56} + x^{54} + x^{53} + x^{51} + x^{50} + x^{48} + x^{47} \\
& + x^{45} + x^{44} + x^{36} + x^{35} + x^{34} + x^{30} + x^{29} + x^{27} + x^{25} + x^{23} + x^{21} \\
& + x^{20} + x^{18} + x^{17} + x^{16} + x^{12} + x^{11} + x^{10} + x^8 + x^7 + x^6, \\
p_9(x) = & x^{105} + x^{104} + x^{103} + x^{101} + x^{97} + x^{96} + x^{94} + x^{91} + x^{87} + x^{86} + x^{84} \\
& + x^{83} + x^{82} + x^{81} + x^{80} + x^{77} + x^{75} + x^{74} + x^{73} + x^{70} + x^{69} + x^{66} \\
& + x^{64} + x^{60} + x^{58} + x^{55} + x^{53} + x^{52} + x^{51} + x^{48} + x^{46} + x^{43} + x^{39} \\
& + x^{38} + x^{36} + x^{32} + x^{30} + x^{26} + x^{24} + x^{23} + x^{22} + x^{18} + x^{17} + x^{15} \\
& + x^9 + x^8 + x^6 + x^3, \\
p_{12}(x) = & x^{108} + x^{107} + x^{105} + x^{103} + x^{101} + x^{99} + x^{97} + x^{96} + x^{60} + x^{59} + x^{57} \\
& + x^{55} + x^{53} + x^{51} + x^{49} + x^{48} + x^{44} + x^{43} + x^{41} + x^{39} + x^{37} + x^{35} \\
& + x^{33} + x^{32} + x^{12} + x^{11} + x^9 + x^7 + x^5 + x^3 + x + 1,
\end{aligned}$$

and the initial terms are $[0, 1, 1, 1, 1, 0, 0, 1, 0, 1, 0, 0, 0, 1, 0]$.

For $\phi(W^e, 1, 1)$:

$$\begin{aligned}
p_0(x) = & x^{96} + x^{90} + x^{88} + x^{86} + x^{84} + x^{82} + x^{80} + x^{78} + x^{74} + x^{70} + x^{62} \\
& + x^{56} + x^{48} + x^{40} + x^{34} + x^{30} + x^{28} + x^{26} + x^{24}, \\
p_3(x) = & x^{94} + x^{88} + x^{72} + x^{68} + x^{66} + x^{64} + x^{62} + x^{60} + x^{58} + x^{54} + x^{48} \\
& + x^{44} + x^{42} + x^{38} + x^{32} + x^{30} + x^{28} + x^{26} + x^{20} + x^{18} + x^{12} + x^{10} \\
& + x^8 + x^6, \\
p_6(x) = & x^{94} + x^{86} + x^{84} + x^{82} + x^{80} + x^{78} + x^{76} + x^{74} + x^{72} + x^{70} + x^{68} \\
& + x^{66} + x^{56} + x^{48} + x^{40} + x^{38} + x^{36} + x^{34} + x^{32} + x^{30} + x^{20} + x^{18} \\
& + x^{16} + x^{12} + x^{10} + 1, \\
p_9(x) = & x^{94} + x^{88} + x^{86} + x^{84} + x^{82} + x^{80} + x^{78} + x^{70} + x^{60} + x^{56} + x^{48} \\
& + x^{38} + x^{36} + x^{32} + x^{30} + x^{28} + x^{24} + x^{20} + x^{18} + x^{16} + x^{10} + x^8 \\
& + x^4 + x^2, \\
p_{12}(x) = & x^{96} + x^{94} + x^{90} + x^{86} + x^{82} + x^{78} + x^{74} + x^{70} + x^{66} + x^{62} + x^{58} \\
& + x^{54} + x^{50} + x^{48} + x^{32} + x^{30} + x^{26} + x^{22} + x^{18} + x^{14} + x^{10} + x^6 \\
& + x^2 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 1, 0, 1, 0, 0, 0, 0, 0, 1, 0, 0, 0, 1, 0]$.

For $\phi(M^o, 0, 0)$:

$$\begin{aligned}
p_0(x) = & x^{108} + x^{107} + x^{106} + x^{104} + x^{102} + x^{101} + x^{100} + x^{94} + x^{92} + x^{86} \\
& + x^{85} + x^{83} + x^{80} + x^{79} + x^{78} + x^{77} + x^{76} + x^{75} + x^{73} + x^{72} + x^{71} \\
& + x^{69} + x^{67} + x^{65} + x^{64} + x^{63} + x^{62} + x^{60} + x^{57} + x^{54} + x^{53} + x^{52} \\
& + x^{51} + x^{49} + x^{48} + x^{47} + x^{46} + x^{44} + x^{42} + x^{40} + x^{39} + x^{38} + x^{37} \\
& + x^{36} + x^{34} + x^{32} + x^{31} + x^{30} + x^{29} + x^{25} + x^{23} + x^{22} + x^{17} + x^{16}
\end{aligned}$$

$$\begin{aligned}
& + x^{14} + x^{13} + x^{12}, \\
p_3(x) &= x^{104} + x^{103} + x^{102} + x^{99} + x^{97} + x^{96} + x^{95} + x^{93} + x^{92} + x^{90} + x^{89} \\
& + x^{87} + x^{86} + x^{85} + x^{84} + x^{83} + x^{81} + x^{78} + x^{77} + x^{76} + x^{72} + x^{71} \\
& + x^{70} + x^{69} + x^{68} + x^{67} + x^{65} + x^{62} + x^{61} + x^{56} + x^{53} + x^{48} + x^{45} \\
& + x^{39} + x^{38} + x^{37} + x^{36} + x^{35} + x^{33} + x^{32} + x^{31} + x^{28} + x^{26} + x^{25} \\
& + x^{21} + x^{19} + x^{18} + x^{16} + x^{15} + x^{10} + x^9 + x^8, \\
p_6(x) &= x^{106} + x^{105} + x^{104} + x^{103} + x^{102} + x^{100} + x^{99} + x^{97} + x^{96} + x^{95} \\
& + x^{94} + x^{92} + x^{91} + x^{87} + x^{84} + x^{82} + x^{81} + x^{80} + x^{75} + x^{74} + x^{73} \\
& + x^{70} + x^{69} + x^{61} + x^{58} + x^{57} + x^{56} + x^{55} + x^{54} + x^{53} + x^{51} + x^{47} \\
& + x^{46} + x^{41} + x^{40} + x^{37} + x^{36} + x^{32} + x^{30} + x^{29} + x^{24} + x^{20} + x^{17} \\
& + x^{15} + x^{14} + x^9 + x^7 + x^6 + x^4 + x^3 + x + 1, \\
p_9(x) &= x^{104} + x^{103} + x^{101} + x^{99} + x^{97} + x^{95} + x^{93} + x^{92} + x^{84} + x^{83} + x^{81} \\
& + x^{79} + x^{77} + x^{76} + x^{68} + x^{67} + x^{65} + x^{63} + x^{61} + x^{60} + x^{56} + x^{55} \\
& + x^{53} + x^{52} + x^{40} + x^{39} + x^{37} + x^{35} + x^{33} + x^{31} + x^{29} + x^{28} + x^{20} \\
& + x^{19} + x^{17} + x^{15} + x^{13} + x^{12} + x^8 + x^7 + x^5 + x^4, \\
p_{12}(x) &= x^{108} + x^{107} + x^{105} + x^{104} + x^{96} + x^{95} + x^{93} + x^{92} + x^{84} + x^{83} + x^{81} \\
& + x^{79} + x^{77} + x^{76} + x^{68} + x^{67} + x^{65} + x^{63} + x^{61} + x^{59} + x^{57} + x^{56} \\
& + x^{52} + x^{51} + x^{49} + x^{48} + x^{44} + x^{43} + x^{41} + x^{40} + x^{32} + x^{31} + x^{29} \\
& + x^{28} + x^{20} + x^{19} + x^{17} + x^{15} + x^{13} + x^{11} + x^9 + x^8 + x^4 + x^3 + x + 1,
\end{aligned}$$

and the initial terms are $[1, 0, 1, 0, 1, 0, 0, 1, 1, 0, 0, 1, 1, 1, 0]$.

For $\phi(M^o, 0, 1)$:

$$\begin{aligned}
p_0(x) &= x^{120} + x^{117} + x^{114} + x^{107} + x^{106} + x^{105} + x^{104} + x^{99} + x^{97} + x^{94} \\
& + x^{93} + x^{91} + x^{90} + x^{86} + x^{85} + x^{83} + x^{82} + x^{80} + x^{78} + x^{75} + x^{71} \\
& + x^{69} + x^{66} + x^{65} + x^{62} + x^{61} + x^{58} + x^{57} + x^{55} + x^{51} + x^{45} + x^{44} \\
& + x^{43} + x^{42} + x^{37} + x^{35} + x^{33} + x^{29} + x^{26} + x^{25} + x^{24}, \\
p_3(x) &= x^{111} + x^{106} + x^{104} + x^{102} + x^{101} + x^{99} + x^{98} + x^{96} + x^{93} + x^{92} + x^{89} \\
& + x^{88} + x^{85} + x^{84} + x^{81} + x^{80} + x^{77} + x^{76} + x^{73} + x^{72} + x^{71} + x^{69} \\
& + x^{68} + x^{66} + x^{65} + x^{62} + x^{60} + x^{59} + x^{58} + x^{57} + x^{31} + x^{26} + x^{24} \\
& + x^{23} + x^{22} + x^{21} + x^{19} + x^{14} + x^{12} + x^{11} + x^{10} + x^9, \\
p_6(x) &= x^{114} + x^{110} + x^{106} + x^{104} + x^{102} + x^{96} + x^{92} + x^{90} + x^{88} + x^{86} + x^{78} \\
& + x^{70} + x^{68} + x^{66} + x^{64} + x^{62} + x^{54} + x^{50} + x^{46} + x^{44} + x^{42} + x^{36} \\
& + x^{34} + x^{32} + x^{28} + x^{24} + x^{22} + x^{16} + x^{12} + x^{10} + x^8 + x^6, \\
p_9(x) &= x^{117} + x^{112} + x^{110} + x^{109} + x^{108} + x^{105} + x^{103} + x^{102} + x^{101} + x^{98} \\
& + x^{97} + x^{96} + x^{95} + x^{94} + x^{93} + x^{92} + x^{91} + x^{90} + x^{89} + x^{88} + x^{87} \\
& + x^{86} + x^{85} + x^{84} + x^{83} + x^{82} + x^{81} + x^{80} + x^{79} + x^{78} + x^{77} + x^{76}
\end{aligned}$$

$$\begin{aligned}
& + x^{75} + x^{74} + x^{73} + x^{72} + x^{71} + x^{70} + x^{68} + x^{67} + x^{66} + x^{65} + x^{63} \\
& + x^{59} + x^{58} + x^{56} + x^{53} + x^{52} + x^{51} + x^{48} + x^{46} + x^{45} + x^{44} + x^{41} \\
& + x^{39} + x^{38} + x^{37} + x^{34} + x^{33} + x^{32} + x^{31} + x^{30} + x^{29} + x^{28} + x^{27} \\
& + x^{26} + x^{25} + x^{24} + x^{23} + x^{22} + x^{20} + x^{19} + x^{18} + x^{17} + x^{15} + x^{11} \\
& + x^{10} + x^8 + x^4 + x^3, \\
p_{12}(x) = & x^{120} + x^{116} + x^{92} + x^{88} + x^{84} + x^{80} + x^{76} + x^{64} + x^{60} + x^{48} + x^{28} \\
& + x^{16} + x^{12} + x^8 + x^4 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 1, 0, 0, 1, 1, 0, 1, 1, 1, 0, 0]$.

For $\phi(M^o, 1, 0)$:

$$\begin{aligned}
p_0(x) = & x^{100} + x^{97} + x^{89} + x^{88} + x^{87} + x^{85} + x^{84} + x^{82} + x^{79} + x^{75} + x^{74} \\
& + x^{72} + x^{71} + x^{68} + x^{67} + x^{65} + x^{61} + x^{57} + x^{56} + x^{55} + x^{54} + x^{53} \\
& + x^{51} + x^{50} + x^{48} + x^{47} + x^{46} + x^{45} + x^{43} + x^{40} + x^{39} + x^{37} + x^{36} \\
& + x^{35} + x^{32} + x^{31} + x^{30} + x^{29} + x^{28} + x^{27} + x^{26} + x^{25} + x^{23} + x^{22} \\
& + x^{21} + x^{20} + x^{18}, \\
p_3(x) = & x^{91} + x^{86} + x^{84} + x^{83} + x^{82} + x^{81} + x^{75} + x^{70} + x^{68} + x^{67} + x^{66} \\
& + x^{65} + x^{59} + x^{54} + x^{52} + x^{50} + x^{49} + x^{46} + x^{44} + x^{42} + x^{41} + x^{38} \\
& + x^{36} + x^{34} + x^{33} + x^{30} + x^{28} + x^{26} + x^{25} + x^{22} + x^{20} + x^{18} + x^{17} \\
& + x^{14} + x^{12} + x^{11} + x^{10} + x^9, \\
p_6(x) = & x^{94} + x^{90} + x^{84} + x^{82} + x^{78} + x^{68} + x^{64} + x^{62} + x^{60} + x^{54} + x^{48} \\
& + x^{42} + x^{36} + x^{34} + x^{22} + x^{14} + x^{12} + x^{10} + x^8 + x^6, \\
p_9(x) = & x^{97} + x^{92} + x^{90} + x^{88} + x^{84} + x^{83} + x^{81} + x^{76} + x^{74} + x^{72} + x^{68} \\
& + x^{67} + x^{65} + x^{60} + x^{58} + x^{56} + x^{52} + x^{51} + x^{33} + x^{28} + x^{26} + x^{24} \\
& + x^{20} + x^{19} + x^{17} + x^{12} + x^{10} + x^8 + x^4 + x^3, \\
p_{12}(x) = & x^{100} + x^{96} + x^{92} + x^{76} + x^{60} + x^{52} + x^{48} + x^{36} + x^{32} + x^{28} + x^{12} \\
& + x^4 + 1,
\end{aligned}$$

and the initial terms are $[0, 1, 1, 0, 0, 1, 1, 0, 0, 0, 0, 1, 1, 0, 0, 0]$.

For $\phi(M^o, 1, 1)$:

$$\begin{aligned}
p_0(x) = & x^{108} + x^{107} + x^{106} + x^{103} + x^{101} + x^{100} + x^{95} + x^{92} + x^{84} + x^{83} \\
& + x^{82} + x^{79} + x^{77} + x^{75} + x^{72} + x^{71} + x^{68} + x^{67} + x^{65} + x^{61} + x^{60} \\
& + x^{58} + x^{52} + x^{50} + x^{49} + x^{48} + x^{47} + x^{46} + x^{45} + x^{44} + x^{43} + x^{42} \\
& + x^{41} + x^{40} + x^{39} + x^{35} + x^{34} + x^{33} + x^{30}, \\
p_3(x) = & x^{104} + x^{103} + x^{102} + x^{100} + x^{98} + x^{97} + x^{96} + x^{95} + x^{93} + x^{91} + x^{89} \\
& + x^{87} + x^{84} + x^{80} + x^{78} + x^{77} + x^{75} + x^{74} + x^{72} + x^{71} + x^{68} + x^{64} \\
& + x^{62} + x^{61} + x^{56} + x^{54} + x^{52} + x^{50} + x^{49} + x^{46} + x^{44} + x^{42} + x^{41} \\
& + x^{40} + x^{39} + x^{35} + x^{34} + x^{31} + x^{30} + x^{29} + x^{24} + x^{22} + x^{18} + x^{17}
\end{aligned}$$

$$\begin{aligned}
& + x^{15} + x^{14} + x^{13} + x^{12}, \\
p_6(x) &= x^{106} + x^{105} + x^{98} + x^{97} + x^{89} + x^{86} + x^{84} + x^{80} + x^{79} + x^{78} + x^{76} \\
& + x^{74} + x^{72} + x^{68} + x^{64} + x^{63} + x^{59} + x^{58} + x^{55} + x^{50} + x^{48} + x^{46} \\
& + x^{45} + x^{43} + x^{37} + x^{36} + x^{34} + x^{33} + x^{32} + x^{31} + x^{29} + x^{24} + x^{20} \\
& + x^{18} + x^{14} + x^{13} + x^{12} + x^{10} + x^8 + x^7 + x^5 + x^3 + x + 1, \\
p_9(x) &= x^{104} + x^{103} + x^{101} + x^{99} + x^{97} + x^{95} + x^{93} + x^{92} + x^{84} + x^{83} + x^{81} \\
& + x^{79} + x^{77} + x^{76} + x^{68} + x^{67} + x^{65} + x^{63} + x^{61} + x^{60} + x^{56} + x^{55} \\
& + x^{53} + x^{52} + x^{40} + x^{39} + x^{37} + x^{35} + x^{33} + x^{31} + x^{29} + x^{28} + x^{20} \\
& + x^{19} + x^{17} + x^{15} + x^{13} + x^{12} + x^8 + x^7 + x^5 + x^4, \\
p_{12}(x) &= x^{108} + x^{107} + x^{105} + x^{104} + x^{96} + x^{95} + x^{93} + x^{92} + x^{84} + x^{83} + x^{81} \\
& + x^{79} + x^{77} + x^{76} + x^{68} + x^{67} + x^{65} + x^{63} + x^{61} + x^{59} + x^{57} + x^{56} \\
& + x^{52} + x^{51} + x^{49} + x^{48} + x^{44} + x^{43} + x^{41} + x^{40} + x^{32} + x^{31} + x^{29} \\
& + x^{28} + x^{20} + x^{19} + x^{17} + x^{15} + x^{13} + x^{11} + x^9 + x^8 + x^4 + x^3 + x + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 0, 1, 1, 1, 0, 1, 1, 1, 1, 0, 0]$.

For $\phi(W^o, 0, 0)$:

$$\begin{aligned}
p_0(x) &= x^{108} + x^{107} + x^{106} + x^{104} + x^{102} + x^{101} + x^{100} + x^{94} + x^{92} + x^{86} \\
& + x^{85} + x^{83} + x^{80} + x^{79} + x^{78} + x^{77} + x^{76} + x^{75} + x^{73} + x^{72} + x^{71} \\
& + x^{69} + x^{67} + x^{65} + x^{64} + x^{63} + x^{62} + x^{60} + x^{57} + x^{54} + x^{53} + x^{52} \\
& + x^{51} + x^{49} + x^{48} + x^{47} + x^{46} + x^{44} + x^{42} + x^{40} + x^{39} + x^{38} + x^{37} \\
& + x^{36} + x^{34} + x^{32} + x^{31} + x^{30} + x^{29} + x^{25} + x^{23} + x^{22} + x^{17} + x^{16} \\
& + x^{14} + x^{13} + x^{12}, \\
p_3(x) &= x^{104} + x^{103} + x^{102} + x^{99} + x^{97} + x^{96} + x^{95} + x^{93} + x^{92} + x^{90} + x^{89} \\
& + x^{87} + x^{86} + x^{85} + x^{84} + x^{83} + x^{81} + x^{78} + x^{77} + x^{76} + x^{72} + x^{71} \\
& + x^{70} + x^{69} + x^{68} + x^{67} + x^{65} + x^{62} + x^{61} + x^{56} + x^{53} + x^{48} + x^{45} \\
& + x^{39} + x^{38} + x^{37} + x^{36} + x^{35} + x^{33} + x^{32} + x^{31} + x^{28} + x^{26} + x^{25} \\
& + x^{21} + x^{19} + x^{18} + x^{16} + x^{15} + x^{10} + x^9 + x^8, \\
p_6(x) &= x^{106} + x^{105} + x^{104} + x^{103} + x^{102} + x^{100} + x^{99} + x^{97} + x^{96} + x^{95} \\
& + x^{94} + x^{92} + x^{91} + x^{87} + x^{84} + x^{82} + x^{81} + x^{80} + x^{75} + x^{74} + x^{73} \\
& + x^{70} + x^{69} + x^{61} + x^{58} + x^{57} + x^{56} + x^{55} + x^{54} + x^{53} + x^{51} + x^{47} \\
& + x^{46} + x^{41} + x^{40} + x^{37} + x^{36} + x^{32} + x^{30} + x^{29} + x^{24} + x^{20} + x^{17} \\
& + x^{15} + x^{14} + x^9 + x^7 + x^6 + x^4 + x^3 + x + 1, \\
p_9(x) &= x^{104} + x^{103} + x^{101} + x^{99} + x^{97} + x^{95} + x^{93} + x^{92} + x^{84} + x^{83} + x^{81} \\
& + x^{79} + x^{77} + x^{76} + x^{68} + x^{67} + x^{65} + x^{63} + x^{61} + x^{60} + x^{56} + x^{55} \\
& + x^{53} + x^{52} + x^{40} + x^{39} + x^{37} + x^{35} + x^{33} + x^{31} + x^{29} + x^{28} + x^{20} \\
& + x^{19} + x^{17} + x^{15} + x^{13} + x^{12} + x^8 + x^7 + x^5 + x^4,
\end{aligned}$$

$$\begin{aligned}
p_{12}(x) = & x^{108} + x^{107} + x^{105} + x^{104} + x^{96} + x^{95} + x^{93} + x^{92} + x^{84} + x^{83} + x^{81} \\
& + x^{79} + x^{77} + x^{76} + x^{68} + x^{67} + x^{65} + x^{63} + x^{61} + x^{59} + x^{57} + x^{56} \\
& + x^{52} + x^{51} + x^{49} + x^{48} + x^{44} + x^{43} + x^{41} + x^{40} + x^{32} + x^{31} + x^{29} \\
& + x^{28} + x^{20} + x^{19} + x^{17} + x^{15} + x^{13} + x^{11} + x^9 + x^8 + x^4 + x^3 + x + 1,
\end{aligned}$$

and the initial terms are $[1, 0, 1, 0, 1, 0, 0, 1, 1, 0, 0, 1, 1, 1, 0]$.

For $\phi(W^o, 0, 1)$:

$$\begin{aligned}
p_0(x) = & x^{100} + x^{97} + x^{89} + x^{88} + x^{87} + x^{85} + x^{84} + x^{82} + x^{79} + x^{75} + x^{74} \\
& + x^{72} + x^{71} + x^{68} + x^{67} + x^{65} + x^{61} + x^{57} + x^{56} + x^{55} + x^{54} + x^{53} \\
& + x^{51} + x^{50} + x^{48} + x^{47} + x^{46} + x^{45} + x^{43} + x^{40} + x^{39} + x^{37} + x^{36} \\
& + x^{35} + x^{32} + x^{31} + x^{30} + x^{29} + x^{28} + x^{27} + x^{26} + x^{25} + x^{23} + x^{22} \\
& + x^{21} + x^{20} + x^{18}, \\
p_3(x) = & x^{91} + x^{86} + x^{84} + x^{83} + x^{82} + x^{81} + x^{75} + x^{70} + x^{68} + x^{67} + x^{66} \\
& + x^{65} + x^{59} + x^{54} + x^{52} + x^{50} + x^{49} + x^{46} + x^{44} + x^{42} + x^{41} + x^{38} \\
& + x^{36} + x^{34} + x^{33} + x^{30} + x^{28} + x^{26} + x^{25} + x^{22} + x^{20} + x^{18} + x^{17} \\
& + x^{14} + x^{12} + x^{11} + x^{10} + x^9, \\
p_6(x) = & x^{94} + x^{90} + x^{84} + x^{82} + x^{78} + x^{68} + x^{64} + x^{62} + x^{60} + x^{54} + x^{48} \\
& + x^{42} + x^{36} + x^{34} + x^{22} + x^{14} + x^{12} + x^{10} + x^8 + x^6, \\
p_9(x) = & x^{97} + x^{92} + x^{90} + x^{88} + x^{84} + x^{83} + x^{81} + x^{76} + x^{74} + x^{72} + x^{68} \\
& + x^{67} + x^{65} + x^{60} + x^{58} + x^{56} + x^{52} + x^{51} + x^{33} + x^{28} + x^{26} + x^{24} \\
& + x^{20} + x^{19} + x^{17} + x^{12} + x^{10} + x^8 + x^4 + x^3, \\
p_{12}(x) = & x^{100} + x^{96} + x^{92} + x^{76} + x^{60} + x^{52} + x^{48} + x^{36} + x^{32} + x^{28} + x^{12} \\
& + x^4 + 1,
\end{aligned}$$

and the initial terms are $[0, 1, 1, 0, 0, 1, 1, 0, 0, 0, 0, 1, 1, 0, 0, 0]$.

For $\phi(W^o, 1, 0)$:

$$\begin{aligned}
p_0(x) = & x^{120} + x^{117} + x^{114} + x^{107} + x^{106} + x^{105} + x^{104} + x^{99} + x^{97} + x^{94} \\
& + x^{93} + x^{91} + x^{90} + x^{86} + x^{85} + x^{83} + x^{82} + x^{80} + x^{78} + x^{75} + x^{71} \\
& + x^{69} + x^{66} + x^{65} + x^{62} + x^{61} + x^{58} + x^{57} + x^{55} + x^{51} + x^{45} + x^{44} \\
& + x^{43} + x^{42} + x^{37} + x^{35} + x^{33} + x^{29} + x^{26} + x^{25} + x^{24}, \\
p_3(x) = & x^{111} + x^{106} + x^{104} + x^{102} + x^{101} + x^{99} + x^{98} + x^{96} + x^{93} + x^{92} + x^{89} \\
& + x^{88} + x^{85} + x^{84} + x^{81} + x^{80} + x^{77} + x^{76} + x^{73} + x^{72} + x^{71} + x^{69} \\
& + x^{68} + x^{66} + x^{65} + x^{62} + x^{60} + x^{59} + x^{58} + x^{57} + x^{31} + x^{26} + x^{24} \\
& + x^{23} + x^{22} + x^{21} + x^{19} + x^{14} + x^{12} + x^{11} + x^{10} + x^9, \\
p_6(x) = & x^{114} + x^{110} + x^{106} + x^{104} + x^{102} + x^{96} + x^{92} + x^{90} + x^{88} + x^{86} + x^{78} \\
& + x^{70} + x^{68} + x^{66} + x^{64} + x^{62} + x^{54} + x^{50} + x^{46} + x^{44} + x^{42} + x^{36} \\
& + x^{34} + x^{32} + x^{28} + x^{24} + x^{22} + x^{16} + x^{12} + x^{10} + x^8 + x^6,
\end{aligned}$$

$$\begin{aligned}
p_9(x) &= x^{117} + x^{112} + x^{110} + x^{109} + x^{108} + x^{105} + x^{103} + x^{102} + x^{101} + x^{98} \\
&\quad + x^{97} + x^{96} + x^{95} + x^{94} + x^{93} + x^{92} + x^{91} + x^{90} + x^{89} + x^{88} + x^{87} \\
&\quad + x^{86} + x^{85} + x^{84} + x^{83} + x^{82} + x^{81} + x^{80} + x^{79} + x^{78} + x^{77} + x^{76} \\
&\quad + x^{75} + x^{74} + x^{73} + x^{72} + x^{71} + x^{70} + x^{68} + x^{67} + x^{66} + x^{65} + x^{63} \\
&\quad + x^{59} + x^{58} + x^{56} + x^{53} + x^{52} + x^{51} + x^{48} + x^{46} + x^{45} + x^{44} + x^{41} \\
&\quad + x^{39} + x^{38} + x^{37} + x^{34} + x^{33} + x^{32} + x^{31} + x^{30} + x^{29} + x^{28} + x^{27} \\
&\quad + x^{26} + x^{25} + x^{24} + x^{23} + x^{22} + x^{20} + x^{19} + x^{18} + x^{17} + x^{15} + x^{11} \\
&\quad + x^{10} + x^8 + x^4 + x^3, \\
p_{12}(x) &= x^{120} + x^{116} + x^{92} + x^{88} + x^{84} + x^{80} + x^{76} + x^{64} + x^{60} + x^{48} + x^{28} \\
&\quad + x^{16} + x^{12} + x^8 + x^4 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 1, 0, 0, 1, 1, 0, 1, 1, 1, 0, 0]$.

For $\phi(W^o, 1, 1)$:

$$\begin{aligned}
p_0(x) &= x^{108} + x^{107} + x^{106} + x^{103} + x^{101} + x^{100} + x^{95} + x^{92} + x^{84} + x^{83} \\
&\quad + x^{82} + x^{79} + x^{77} + x^{75} + x^{72} + x^{71} + x^{68} + x^{67} + x^{65} + x^{61} + x^{60} \\
&\quad + x^{58} + x^{52} + x^{50} + x^{49} + x^{48} + x^{47} + x^{46} + x^{45} + x^{44} + x^{43} + x^{42} \\
&\quad + x^{41} + x^{40} + x^{39} + x^{35} + x^{34} + x^{33} + x^{30}, \\
p_3(x) &= x^{104} + x^{103} + x^{102} + x^{100} + x^{98} + x^{97} + x^{96} + x^{95} + x^{93} + x^{91} + x^{89} \\
&\quad + x^{87} + x^{84} + x^{80} + x^{78} + x^{77} + x^{75} + x^{74} + x^{72} + x^{71} + x^{68} + x^{64} \\
&\quad + x^{62} + x^{61} + x^{56} + x^{54} + x^{52} + x^{50} + x^{49} + x^{46} + x^{44} + x^{42} + x^{41} \\
&\quad + x^{40} + x^{39} + x^{35} + x^{34} + x^{31} + x^{30} + x^{29} + x^{24} + x^{22} + x^{18} + x^{17} \\
&\quad + x^{15} + x^{14} + x^{13} + x^{12}, \\
p_6(x) &= x^{106} + x^{105} + x^{98} + x^{97} + x^{89} + x^{86} + x^{84} + x^{80} + x^{79} + x^{78} + x^{76} \\
&\quad + x^{74} + x^{72} + x^{68} + x^{64} + x^{63} + x^{59} + x^{58} + x^{55} + x^{50} + x^{48} + x^{46} \\
&\quad + x^{45} + x^{43} + x^{37} + x^{36} + x^{34} + x^{33} + x^{32} + x^{31} + x^{29} + x^{24} + x^{20} \\
&\quad + x^{18} + x^{14} + x^{13} + x^{12} + x^{10} + x^8 + x^7 + x^5 + x^3 + x + 1, \\
p_9(x) &= x^{104} + x^{103} + x^{101} + x^{99} + x^{97} + x^{95} + x^{93} + x^{92} + x^{84} + x^{83} + x^{81} \\
&\quad + x^{79} + x^{77} + x^{76} + x^{68} + x^{67} + x^{65} + x^{63} + x^{61} + x^{60} + x^{56} + x^{55} \\
&\quad + x^{53} + x^{52} + x^{40} + x^{39} + x^{37} + x^{35} + x^{33} + x^{31} + x^{29} + x^{28} + x^{20} \\
&\quad + x^{19} + x^{17} + x^{15} + x^{13} + x^{12} + x^8 + x^7 + x^5 + x^4, \\
p_{12}(x) &= x^{108} + x^{107} + x^{105} + x^{104} + x^{96} + x^{95} + x^{93} + x^{92} + x^{84} + x^{83} + x^{81} \\
&\quad + x^{79} + x^{77} + x^{76} + x^{68} + x^{67} + x^{65} + x^{63} + x^{61} + x^{59} + x^{57} + x^{56} \\
&\quad + x^{52} + x^{51} + x^{49} + x^{48} + x^{44} + x^{43} + x^{41} + x^{40} + x^{32} + x^{31} + x^{29} \\
&\quad + x^{28} + x^{20} + x^{19} + x^{17} + x^{15} + x^{13} + x^{11} + x^9 + x^8 + x^4 + x^3 + x + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 0, 1, 1, 1, 0, 1, 1, 1, 1, 0, 0]$.

Below is the transition function and output function of an 2-automaton that generates $T = M_{0,0}^o$:

Transition function $(n, j) \mapsto \delta(n, j)$ ($\Lambda(n) := [\delta(n, 0), \delta(n, 1)]$):

n	$\Lambda(n)$	n	$\Lambda(n)$	n	$\Lambda(n)$	n	$\Lambda(n)$	n	$\Lambda(n)$
0	[1, 2]	88	[164, 10]	176	[16, 103]	264	[353, 318]	352	[409, 288]
1	[3, 4]	89	[13, 165]	177	[280, 281]	265	[168, 125]	353	[78, 249]
2	[5, 6]	90	[166, 140]	178	[282, 283]	266	[151, 57]	354	[205, 382]
3	[7, 8]	91	[141, 167]	179	[284, 285]	267	[347, 34]	355	[232, 269]
4	[9, 10]	92	[168, 169]	180	[286, 247]	268	[307, 192]	356	[386, 77]
5	[11, 12]	93	[144, 170]	181	[287, 97]	269	[171, 37]	357	[410, 243]
6	[13, 14]	94	[171, 49]	182	[288, 289]	270	[354, 346]	358	[96, 409]
7	[15, 16]	95	[135, 150]	183	[290, 291]	271	[355, 36]	359	[283, 411]
8	[17, 18]	96	[147, 172]	184	[292, 293]	272	[41, 128]	360	[412, 286]
9	[19, 20]	97	[173, 163]	185	[28, 263]	273	[42, 129]	361	[230, 413]
10	[21, 22]	98	[174, 141]	186	[294, 79]	274	[325, 356]	362	[414, 415]
11	[23, 24]	99	[175, 126]	187	[295, 290]	275	[313, 54]	363	[390, 388]
12	[25, 26]	100	[8, 176]	188	[260, 294]	276	[324, 321]	364	[400, 416]
13	[27, 28]	101	[177, 178]	189	[248, 239]	277	[164, 357]	365	[417, 67]
14	[29, 30]	102	[130, 179]	190	[296, 4]	278	[328, 183]	366	[380, 25]
15	[31, 32]	103	[32, 120]	191	[297, 298]	279	[355, 358]	367	[119, 122]
16	[33, 34]	104	[123, 32]	192	[299, 300]	280	[349, 349]	368	[304, 305]
17	[35, 36]	105	[180, 48]	193	[301, 222]	281	[160, 158]	369	[149, 136]
18	[37, 38]	106	[181, 42]	194	[302, 296]	282	[359, 165]	370	[325, 323]
19	[39, 40]	107	[182, 178]	195	[303, 304]	283	[57, 360]	371	[314, 143]
20	[41, 42]	108	[183, 184]	196	[121, 53]	284	[155, 350]	372	[11, 418]
21	[43, 44]	109	[185, 144]	197	[120, 123]	285	[361, 362]	373	[419, 43]
22	[45, 46]	110	[179, 44]	198	[120, 304]	286	[45, 184]	374	[350, 351]
23	[47, 48]	111	[39, 186]	199	[122, 304]	287	[363, 168]	375	[420, 45]
24	[49, 50]	112	[187, 188]	200	[123, 305]	288	[156, 351]	376	[187, 192]
25	[51, 52]	113	[152, 189]	201	[304, 32]	289	[361, 134]	377	[421, 332]
26	[53, 53]	114	[126, 176]	202	[180, 306]	290	[316, 364]	378	[351, 155]
27	[54, 55]	115	[177, 190]	203	[307, 188]	291	[339, 141]	379	[305, 120]
28	[56, 57]	116	[191, 177]	204	[188, 193]	292	[175, 8]	380	[58, 310]
29	[58, 59]	117	[192, 193]	205	[132, 308]	293	[314, 55]	381	[35, 358]
30	[33, 60]	118	[194, 177]	206	[47, 306]	294	[133, 362]	382	[146, 342]
31	[61, 62]	119	[195, 196]	207	[309, 310]	295	[363, 124]	383	[421, 333]
32	[63, 64]	120	[62, 197]	208	[53, 121]	296	[153, 364]	384	[334, 40]
33	[65, 66]	121	[104, 198]	209	[159, 311]	297	[365, 54]	385	[422, 144]
34	[67, 68]	122	[199, 200]	210	[312, 164]	298	[173, 366]	386	[354, 330]
35	[69, 70]	123	[196, 201]	211	[313, 314]	299	[156, 159]	387	[319, 418]
36	[71, 72]	124	[202, 203]	212	[315, 11]	300	[157, 161]	388	[179, 336]
37	[73, 74]	125	[204, 205]	213	[316, 154]	301	[162, 366]	389	[194, 182]
38	[75, 76]	126	[206, 93]	214	[139, 175]	302	[312, 9]	390	[423, 49]
39	[77, 78]	127	[207, 208]	215	[178, 317]	303	[367, 63]	391	[8, 127]
40	[79, 80]	128	[209, 71]	216	[318, 319]	304	[64, 199]	392	[192, 424]

41	[81, 82]	129	[82, 113]	217	[320, 321]	305	[200, 368]	393	[419, 179]
42	[83, 84]	130	[210, 87]	218	[37, 50]	306	[86, 369]	394	[128, 343]
43	[85, 86]	131	[211, 212]	219	[129, 322]	307	[266, 301]	395	[191, 182]
44	[87, 88]	132	[213, 214]	220	[321, 323]	308	[370, 265]	396	[423, 37]
45	[89, 90]	133	[108, 215]	221	[324, 325]	309	[371, 372]	397	[166, 175]
46	[91, 92]	134	[216, 217]	222	[57, 189]	310	[110, 373]	398	[425, 138]
47	[93, 94]	135	[218, 219]	223	[326, 327]	311	[374, 280]	399	[152, 360]
48	[95, 96]	136	[220, 221]	224	[328, 45]	312	[375, 21]	400	[326, 52]
49	[97, 98]	137	[222, 81]	225	[329, 330]	313	[70, 231]	401	[309, 59]
50	[99, 65]	138	[223, 16]	226	[331, 332]	314	[279, 376]	402	[174, 340]
51	[4, 100]	139	[224, 225]	227	[305, 122]	315	[201, 377]	403	[359, 14]
52	[101, 102]	140	[197, 226]	228	[331, 333]	316	[92, 297]	404	[422, 335]
53	[103, 104]	141	[227, 228]	229	[334, 186]	317	[289, 378]	405	[426, 148]
54	[105, 106]	142	[229, 230]	230	[335, 170]	318	[379, 380]	406	[9, 357]
55	[107, 108]	143	[231, 232]	231	[43, 336]	319	[381, 261]	407	[350, 159]
56	[109, 110]	144	[233, 234]	232	[132, 337]	320	[382, 271]	408	[311, 156]
57	[111, 112]	145	[235, 236]	233	[137, 338]	321	[208, 383]	409	[426, 172]
58	[113, 114]	146	[237, 238]	234	[339, 340]	322	[369, 255]	410	[182, 190]
59	[115, 116]	147	[239, 240]	235	[315, 319]	323	[384, 218]	411	[365, 314]
60	[117, 118]	148	[241, 242]	236	[124, 169]	324	[385, 386]	412	[188, 424]
61	[119, 120]	149	[243, 244]	237	[341, 164]	325	[198, 29]	413	[343, 322]
62	[121, 121]	150	[245, 246]	238	[340, 167]	326	[236, 387]	414	[321, 356]
63	[32, 122]	151	[247, 248]	239	[342, 337]	327	[388, 389]	415	[353, 315]
64	[122, 123]	152	[249, 250]	240	[343, 344]	328	[251, 245]	416	[51, 327]
65	[124, 125]	153	[88, 251]	241	[140, 8]	329	[24, 390]	417	[185, 335]
66	[126, 127]	154	[226, 252]	242	[345, 315]	330	[72, 95]	418	[377, 63]
67	[128, 129]	155	[253, 254]	243	[342, 308]	331	[293, 391]	419	[411, 213]
68	[130, 43]	156	[255, 253]	244	[190, 317]	332	[392, 393]	420	[427, 91]
69	[49, 38]	157	[256, 257]	245	[318, 11]	333	[394, 395]	421	[399, 428]
70	[131, 132]	158	[258, 259]	246	[345, 318]	334	[22, 396]	422	[429, 235]
71	[133, 134]	159	[254, 260]	247	[131, 342]	335	[298, 397]	423	[430, 216]
72	[135, 136]	160	[261, 262]	248	[42, 343]	336	[398, 399]	424	[431, 406]
73	[137, 138]	161	[263, 264]	249	[329, 346]	337	[212, 371]	425	[76, 275]
74	[139, 140]	162	[265, 266]	250	[181, 128]	338	[228, 400]	426	[18, 432]
75	[141, 142]	163	[267, 267]	251	[56, 152]	339	[94, 270]	427	[433, 168]
76	[54, 143]	164	[225, 268]	252	[347, 60]	340	[368, 401]	428	[319, 12]
77	[144, 145]	165	[252, 223]	253	[348, 349]	341	[396, 394]	429	[433, 124]
78	[146, 132]	166	[269, 17]	254	[311, 350]	342	[6, 402]	430	[341, 9]
79	[147, 148]	167	[270, 69]	255	[351, 311]	343	[403, 277]	431	[425, 338]
80	[149, 150]	168	[271, 272]	256	[183, 46]	344	[84, 299]	432	[178, 352]
81	[151, 152]	169	[273, 256]	257	[129, 344]	345	[404, 381]	433	[434, 273]
82	[153, 154]	170	[274, 27]	258	[140, 126]	346	[281, 405]	434	[420, 183]
83	[155, 156]	171	[275, 75]	259	[320, 325]	347	[406, 5]		
84	[157, 158]	172	[238, 276]	260	[349, 348]	348	[348, 348]		

85	[159, 155]	173	[277, 211]	261	[335, 145]	349	[407, 83]
86	[160, 161]	174	[278, 279]	262	[190, 352]	350	[408, 284]
87	[162, 163]	175	[26, 207]	263	[340, 142]	351	[378, 408]

Output function $n \mapsto \tau(n)$:

n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$
0	0	63	0	126	1	189	0	252	1	315	0
1	0	64	0	127	1	190	0	253	0	316	1
2	1	65	0	128	1	191	1	254	1	317	0
3	0	66	1	129	0	192	0	255	0	318	1
4	1	67	1	130	0	193	1	256	1	319	0
5	1	68	0	131	0	194	0	257	0	320	1
6	0	69	0	132	1	195	0	258	1	321	0
7	0	70	0	133	1	196	1	259	1	322	0
8	0	71	1	134	1	197	1	260	1	323	0
9	1	72	1	135	1	198	1	261	0	324	0
10	1	73	1	136	0	199	0	262	0	325	1
11	1	74	1	137	1	200	1	263	0	326	0
12	1	75	1	138	0	201	0	264	1	327	0
13	0	76	0	139	1	202	0	265	1	328	1
14	0	77	1	140	1	203	0	266	0	329	0
15	0	78	1	141	1	204	1	267	1	330	1
16	0	79	0	142	0	205	1	268	0	331	1
17	0	80	0	143	1	206	1	269	0	332	0
18	1	81	0	144	1	207	1	270	1	333	1
19	1	82	0	145	0	208	0	271	1	334	0
20	0	83	0	146	1	209	1	272	0	335	0
21	1	84	1	147	0	210	0	273	0	336	0
22	0	85	1	148	1	211	0	274	1	337	0
23	1	86	0	149	0	212	0	275	0	338	1
24	0	87	1	150	1	213	1	276	0	339	0
25	1	88	0	151	0	214	1	277	0	340	0
26	0	89	0	152	0	215	1	278	1	341	1
27	0	90	0	153	0	216	1	279	1	342	0
28	1	91	1	154	1	217	1	280	1	343	1
29	0	92	1	155	0	218	1	281	0	344	1
30	0	93	1	156	0	219	0	282	1	345	0
31	0	94	0	157	1	220	0	283	1	346	0
32	0	95	1	158	1	221	0	284	0	347	1
33	0	96	0	159	1	222	1	285	0	348	0
34	1	97	0	160	0	223	0	286	0	349	1
35	0	98	1	161	0	224	1	287	1	350	1
36	1	99	0	162	1	225	0	288	0	351	0
37	1	100	0	163	1	226	1	289	0	352	1

38	1	101	1	164	0	227	1	290	1	353	1	416	1
39	1	102	0	165	1	228	1	291	0	354	1	417	1
40	0	103	0	166	0	229	0	292	0	355	1	418	0
41	0	104	1	167	1	230	0	293	1	356	1	419	1
42	0	105	0	168	1	231	1	294	1	357	0	420	0
43	1	106	1	169	0	232	1	295	1	358	0	421	0
44	1	107	0	170	1	233	1	296	0	359	1	422	0
45	0	108	1	171	0	234	0	297	1	360	1	423	1
46	1	109	1	172	0	235	0	298	0	361	0	424	0
47	1	110	0	173	0	236	0	299	0	362	0	425	0
48	1	111	1	174	1	237	1	300	1	363	1	426	1
49	0	112	1	175	0	238	0	301	1	364	0	427	0
50	0	113	0	176	0	239	0	302	0	365	1	428	0
51	1	114	1	177	1	240	1	303	0	366	0	429	0
52	1	115	1	178	1	241	1	304	0	367	0	430	1
53	0	116	1	179	0	242	0	305	1	368	0	431	0
54	0	117	0	180	0	243	0	306	0	369	0	432	1
55	0	118	0	181	1	244	0	307	0	370	1	433	0
56	1	119	0	182	0	245	1	308	1	371	1	434	0
57	1	120	1	183	1	246	0	309	1	372	1		
58	0	121	1	184	0	247	0	310	0	373	1		
59	1	122	0	185	1	248	0	311	1	374	1		
60	0	123	1	186	1	249	0	312	0	375	0		
61	0	124	0	187	1	250	1	313	0	376	1		
62	1	125	1	188	1	251	1	314	1	377	0		

UNIVERSITÉ DE STRASBOURG, CNRS, IRMA UMR 7501, F-67000 STRASBOURG, FRANCE

Email address: `guoniu.han@unistra.fr`

SCHOOL OF MATHEMATICS AND STATISTICS, HUAZHONG UNIVERSITY OF SCIENCE AND TECHNOLOGY, WUHAN, PR CHINA

Email address: `huyining@hust.edu.cn`