

PROOF OF THE CONTINUED FRACTION CONJECTURE FOR

$$(a, b) = (z^4, z^3 + z)$$

GUO-NIU HAN* AND YINING HU*

ABSTRACT. The continued fraction conjecture claims that the continued fraction defined by the (a, b) -Thue-Morse sequence is algebraic over $\mathbb{F}_2(x)$ for all pairs of distinct elements (a, b) in $\mathbb{F}_2[x] \setminus \mathbb{F}_2$. In this paper we prove the conjecture for $(a, b) = (z^4, z^3 + z)$.

1. INTRODUCTION

Let $\mathbb{F}_2$ be the finite field of cardinality 2. Given a sequence of polynomials $\mathbf{a} = (a_0(z), a_1(z), a_2(z), \dots)$ of elements from $\mathbb{F}_2[z] \setminus \mathbb{F}_2$, we define the infinite continued fraction

$$(1.1) \quad \text{CF}(\mathbf{a}(z)) = \frac{1}{a_0(z) + \frac{1}{a_1(z) + \frac{1}{a_2(z) + \frac{1}{\ddots}}}}$$

as the limit of the finite continued fractions

$$(1.2) \quad \text{CF}_n(\mathbf{a}(z)) = \frac{1}{a_0(z) + \frac{1}{a_1(z) + \frac{1}{\ddots + \frac{1}{a_n(z)}}}} \in \mathbb{F}_2((1/z)).$$

Conversely, each power series in $\mathbb{F}_2[[1/z]]$ without constant term

$$(1.3) \quad c_{-1}z^{-1} + c_{-2}z^{-2} + c_{-3}z^{-3} + \dots$$

admits a continued fraction expansion of form (1.1).

Let a and b be two distinct elements from $\mathbb{F}_2[z] \setminus \mathbb{F}_2$. We consider the continued fraction (1.1) associated with the (a, b) -Thue-Morse sequence

$$\mathbf{t} = (t_0(z), t_1(z), t_2(z), \dots) = (a, b, b, a, b, a, a, b, \dots)$$

that we shall denote

$$(1.4) \quad \text{CF}(a, b) := \text{CF}(\mathbf{t}).$$

Date: 9/04/2020 17:26:06 .

2010 Mathematics Subject Classification. 11B85, 11J70, 11B50, 11Y65, 05A15.

Key words and phrases. automatic sequence, continued fraction, Thue-Morse sequence.

* This paper was automatically generated from a computer program developed by the authors, with 7.9 minutes CPU time used.

Theorem 1.1. *When $(a, b) = (z^4, z^3 + z)$, the two power series $\text{CF}(a, b)$ and $\text{CF}(b, a)$ are algebraic over $\mathbb{F}_2(z)$, with minimal polynomials of the form*

$$p_4(z)y^4 + p_3(z)y^3 + p_2(z)y^2 + p_1(z)y + p_0(z) = 0$$

where, for $\text{CF}(a, b)$,

$$\begin{aligned} p_0(z) &= z^{17} + z^{15} + z^{13} + z^{10} + z^8 + z^7 + z^6 + z^5 + z^3 + 1, \\ p_1(z) &= z^{21} + z^{20} + z^{19} + z^{17} + z^{14} + z^{11} + z^{10} + z^9 + z^7 + z^4, \\ p_2(z) &= z^{24} + z^{18} + z^{14} + z^8, \\ p_3(z) &= z^{21} + z^{20} + z^{19} + z^{17} + z^{14} + z^{11} + z^{10} + z^9 + z^7 + z^4, \\ p_4(z) &= z^{18} + z^{17} + z^{16} + z^{15} + z^{14} + z^{13} + z^{12} + z^{10} + z^8 + z^7 + z^6 + z^5 + z^3, \end{aligned}$$

and for $\text{CF}(b, a)$,

$$\begin{aligned} p_0(z) &= z^{17} + z^{15} + z^{13} + z^{12} + z^{10} + z^7 + z^6 + z^5 + z^3, \\ p_1(z) &= z^{21} + z^{20} + z^{19} + z^{17} + z^{14} + z^{11} + z^{10} + z^9 + z^7 + z^4, \\ p_2(z) &= z^{24} + z^{18} + z^{14} + z^8, \\ p_3(z) &= z^{21} + z^{20} + z^{19} + z^{17} + z^{14} + z^{11} + z^{10} + z^9 + z^7 + z^4, \\ p_4(z) &= z^{18} + z^{17} + z^{16} + z^{15} + z^{14} + z^{13} + z^{10} + z^7 + z^6 + z^5 + z^3 + 1. \end{aligned}$$

2. PROOF

Define the sequences of polynomials $(P_n(z))_{n \geq 0}$ and $(Q_n(z))_{n \geq 0}$ by

$$\begin{pmatrix} P_n(z) & Q_n(z) \\ P_{n-1}(z) & Q_{n-1}(z) \end{pmatrix} := \begin{pmatrix} t_n(z) & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} t_{n-1}(z) & 1 \\ 1 & 0 \end{pmatrix} \cdots \begin{pmatrix} t_0(z) & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}.$$

By the basic properties of continued fractions, we have

$$(2.1) \quad \text{CF}_n(a, b) = \frac{P_n(z)}{Q_n(z)} \in \mathbb{F}_2((1/z)).$$

As formula (2.1) is not efficient for calculating $\text{CF}_n(a, b)$, a fast method is to be derived and described as follows. Define

$$\begin{aligned} M_n(x) &= x^{\deg(t_{2^n-1})} \begin{pmatrix} t_{2^n-1}(1/x) & 1 \\ 1 & 0 \end{pmatrix} \times x^{\deg(t_{2^n-2})} \begin{pmatrix} t_{2^n-2}(1/x) & 1 \\ 1 & 0 \end{pmatrix} \times \\ &\quad \cdots \times x^{\deg(t_0)} \begin{pmatrix} t_0(1/x) & 1 \\ 1 & 0 \end{pmatrix}, \\ W_n(x) &= x^{\deg(\bar{t}_{2^n-1})} \begin{pmatrix} \bar{t}_{2^n-1}(1/x) & 1 \\ 1 & 0 \end{pmatrix} \times x^{\deg(\bar{t}_{2^n-2})} \begin{pmatrix} \bar{t}_{2^n-2}(1/x) & 1 \\ 1 & 0 \end{pmatrix} \times \\ &\quad \cdots \times x^{\deg(\bar{t}_0)} \begin{pmatrix} \bar{t}_0(1/x) & 1 \\ 1 & 0 \end{pmatrix}, \end{aligned}$$

where $\bar{t}$ is the (b, a) -Thue-Morse sequence. By the properties of the Thue-Morse sequence, we have for $n \geq 0$

$$\begin{aligned} M_{n+1}(x) &= W_n(x) \cdot M_n(x), \\ W_{n+1}(x) &= M_n(x) \cdot W_n(x). \end{aligned}$$

Let $x := 1/z$. For each non-zero polynomial $P(z)$, define $\tilde{P}(x)$ to be $P(1/x)$. Then,

$$\text{CF}_n(a, b) = \frac{P_n(z)}{Q_n(z)} = \frac{\tilde{P}_n(x)}{\tilde{Q}_n(x)} \in \mathbb{F}_2((x)) = \mathbb{F}_2((1/z)).$$

Comparing the definition of $M_n(x)$ with the definition of $P_n(x)$ and $Q_n(x)$, we see that

$$\begin{aligned} M_n(x)_{0,1} &= x^{d_n} \tilde{P}_{2^n-1}(x), \\ M_n(x)_{0,0} &= x^{d_n} \tilde{Q}_{2^n-1}(x), \end{aligned}$$

for some positive integer d_n . Hence,

$$(2.2) \quad \text{CF}_{2^{2^n}-1}(a, b) = \frac{\tilde{P}_{2^{2^n}-1}(x)}{\tilde{Q}_{2^{2^n}-1}(x)} = \frac{M_{2^n}(x)_{0,1}}{M_{2^n}(x)_{0,0}}.$$

Therefore, by the convergence theorem of continued fraction, the algebraicity of $\text{CF}(a, b)$ will be established if it is shown that both $M_{2^n}(x)_{0,1}$ and $M_{2^n}(x)_{0,0}$ converge to algebraic series in $\mathbb{F}_2[[x]]$.

Actually, we will prove that for all $0 \leq i, j \leq 1$ the four sequences $(M_{2^n}(x)_{i,j})_n$, $(M_{2^{n+1}}(x)_{i,j})_n$, $(W_{2^n}(x)_{i,j})_n$, and $(W_{2^{n+1}}(x)_{i,j})_n$ converge to algebraic series in $\mathbb{F}_2[[x]]$. For this purpose, we define four 2×2 matrices M^e, M^o, W^e, W^o as follows: For each $T \in \{M^e, M^o, W^e, W^o\}$ and $0 \leq i, j \leq 1$, $T_{i,j}$ is defined to be the unique solution in $\mathbb{F}_2[[x]]$ of the polynomial $\phi(T, i, j)$ under certain initial conditions; the polynomials $\phi(T, i, j)$ and initial conditions are given in Section 3. We will prove that these four matrices, whose components are algebraic by definition, are the limits of $(M_{2^n}(x))_n$, $(M_{2^{n+1}}(x))_n$, $(W_{2^n}(x))_n$, and $(W_{2^{n+1}}(x))_n$.

Let us explain how the polynomials $\phi(T, i, j)$ and initial conditions are found, and why the solutions exist and are unique. For $0 \leq i, j \leq 1$, the coefficients of the polynomial $\phi(M^e, i, j)$ (resp. $\phi(M^o, i, j)$, $\phi(W^e, i, j)$, and $\phi(W^o, i, j)$) are the Padé-Hermite approximants of type

$$(175, 175, 175, 175, 175)$$

of the vector

$$(1, T^3, T^6, T^9, T^{12}),$$

where $T = M_{12,i,j}$ (resp. $M_{11,i,j}$, $W_{12,i,j}$, and $W_{11,i,j}$), found by the Derksen algorithm. We take the first 8 terms of $M_{12,i,j}$ (resp. $M_{11,i,j}$, $W_{12,i,j}$, and $W_{11,i,j}$) as initial conditions for $\phi(M^e, i, j)$ (resp. $\phi(M^o, i, j)$, $\phi(W^e, i, j)$, and $\phi(W^o, i, j)$). The following fact will be used to ensure that the solution exists and is unique: let $P(x, y) \in \mathbb{F}_2[x, y]$ and for each series $f(x) = \sum_{n=0}^{\infty} a_n x^n \in \mathbb{F}_2[[x]]$ denote the partial sum $\sum_{j=0}^{n-1} a_j x^j$ by $f_n(x)$ for $n \geq 0$. If for some $n \geq 0$ and $a_0, a_1, \dots, a_{n-1} \in \mathbb{F}_2$ $P(x, \sum_{j=0}^{n-1} a_j x^j) = O(x^n)$ and $Q(x, y) := P(x, \sum_{j=0}^{n-1} a_j x^j + x^n y)$ can be written as $x^m \sum_{j=0}^{\infty} q_j(x) y^j$ where $q_j(x)$ are polynomials for $j \geq 0$, $q_1(0) = 1$, and $q_j(0) = 0$ for $j > 1$, then there exists a unique solution $f(x) \in \mathbb{F}_2[[x]]$ of $P(x, f(x)) = 0$ that satisfies the initial condition $f_n(x) = \sum_{j=0}^{n-1} a_j x^j$.

We state two lemmas concerning the four matrices M^e, M^o, W^e, W^o . The first one is about relations between them; the second, about the structure of the each matrix.

Lemma 2.1. *We have*

$$(2.3) \quad M^e = W^o \cdot M^o,$$

$$(2.4) \quad M^o = W^e \cdot M^e,$$

$$(2.5) \quad W^e = M^o \cdot W^o,$$

$$(2.6) \quad W^o = M^e \cdot W^e.$$

Proof. We give the proof of the identity

$$M_{0,0}^e = W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o,$$

the proofs of the others being similar.

First, we compute the minimal polynomials of $W_{0,0}^o M_{0,0}^o$ and $W_{0,1}^o M_{1,0}^o$. We know that

$$P(x, y) = \text{Res}_z (\phi(W^o, 0, 0)(x, z), z^{12} \cdot \phi(M^o, 0, 0)(x, y/z))$$

is an annihilating polynomial of $W_{0,0}^o M_{0,0}^o$. We use Padé-Hermite approximation to find a candidate for the minimal polynomial of $W_{0,0}^o M_{0,0}^o$, that will be called $\phi_0(x, y)$. To prove that $\phi_0(x, y)$ is indeed the minimal polynomial, it suffices to prove that it is an irreducible factor of $P(x, y)$ of multiplicity m and that $Q(x, y) := P(x, y)/\phi_0(x, y)^m$ is not an annihilating polynomial of $W_{0,0}^o M_{0,0}^o$. We verify the first point directly. For the second point, we truncate $W_{0,0}^o M_{0,0}^o$ to order 630 and substitute it for y in $Q(x, y)$. We get a series of valuation less than 630, which proves that $Q(x, y)$ is not an annihilating polynomial of $W_{0,0}^o M_{0,0}^o$. We find the minimal polynomial $\phi_1(x, y)$ of $W_{0,1}^o M_{1,0}^o$ in a similar way.

Now we prove that $\phi(M^e, 0, 0)$ is the minimal polynomial of $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$. We know that

$$S(x, y) = \text{Res}_z (\phi_0(x, z), \phi_1(x, y + z))$$

is an annihilating of $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$. We verify that $\phi(M^e, 0, 0)$ is an irreducible factor of $S(x, y)$ of multiplicity μ , and that $Q(x, y) := S(x, y)/\phi(M^e, 0, 0)^\mu$ is not an annihilating polynomial of $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$. To see the last point, we truncate $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$ to order 770 and substitute it for y in $Q(x, y)$. We get a series of valuation less than 770, and therefore $Q(x, y)$ is not an annihilating polynomial of $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$.

Finally, the first 16 terms of $M_{0,0}^e$ and $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$ coincide. As these first terms determine a unique solution of $\phi(M^e, 0, 0)$, we know that the two series are one and the same. $\square$

Define

$$R^e = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \quad \text{and} \quad R^o = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}.$$

Lemma 2.2. *For all integers $k \geq 2$ and $u = 2^{2k-1}$, the following identities hold.*

$$\begin{aligned} M^e[:14u] &= M^e[:7u] + x^{2u} M^e[:5u] + x^{6u} M^e[:u] + x^{2u} M^e[:7u] + x^{4u} M^e[:5u] \\ &\quad + x^{8u} M^e[:u] + x^{4u} M^e[:7u] + x^{6u} M^e[:5u] + x^{10u} M^e[:u] \\ &\quad + x^{8u} M^e[:6u] + x^{10u} M^e[:4u] + (x^{7u} + x^{9u} + x^{11u}) R^e, \\ W^e[:14u] &= W^e[:7u] + x^{2u} W^e[:5u] + x^{6u} W^e[:u] + x^{2u} W^e[:7u] + x^{4u} W^e[:5u] \\ &\quad + x^{8u} W^e[:u] + x^{4u} W^e[:7u] + x^{6u} W^e[:5u] + x^{10u} W^e[:u] \\ &\quad + x^{8u} W^e[:6u] + x^{10u} W^e[:4u] + (x^{7u} + x^{9u} + x^{11u}) R^e. \end{aligned}$$

For all integers $k \geq 2$ and $u = 2^{2k}$, the following identities hold.

$$\begin{aligned}
M^o[:14u] &= M^o[:7u] + x^{2u}M^o[:5u] + x^{6u}M^o[:u] + x^{2u}M^o[:7u] + x^{4u}M^o[:5u] \\
&\quad + x^{8u}M^o[:u] + x^{4u}M^o[:7u] + x^{6u}M^o[:5u] + x^{10u}M^o[:u] \\
&\quad + x^{8u}M^o[:6u] + x^{10u}M^o[:4u] + (x^{7u} + x^{9u} + x^{11u})R^o, \\
W^o[:14u] &= W^o[:7u] + x^{2u}W^o[:5u] + x^{6u}W^o[:u] + x^{2u}W^o[:7u] + x^{4u}W^o[:5u] \\
&\quad + x^{8u}W^o[:u] + x^{4u}W^o[:7u] + x^{6u}W^o[:5u] + x^{10u}W^o[:u] \\
&\quad + x^{8u}W^o[:6u] + x^{10u}W^o[:4u] + (x^{7u} + x^{9u} + x^{11u})R^o.
\end{aligned}$$

Proof. To prove Lemma 2.2 we first construct an automaton for each sequence concerned, and then transform the conditions on infinitely many k 's into finitely many conditions on the states of the automaton. In the following, we will prove that for $T = M_{0,0}^o$, for all integer $k \geq 2$ and $u = 2^{2k}$,

$$\begin{aligned}
T[:14u] &= T[:7u] + x^{2u}T[:5u] + x^{6u}T[:u] + x^{2u}T[:7u] + x^{4u}T[:5u] + x^{8u}T[:u] \\
&\quad + x^{4u}T[:7u] + x^{6u}T[:5u] + x^{10u}T[:u] + x^{8u}T[:6u] + x^{10u}T[:4u].
\end{aligned}$$

The proof of the other 15 cases are similar. We break down the above identity into 7 parts:

$$\begin{aligned}
0 &= x^{6u}T[u:2u] + x^{2u}T[5u:6u] + T[7u:8u], \\
0 &= x^{6u}T[2u:3u] + x^{2u}T[6u:7u] + T[8u:9u], \\
0 &= x^{8u}T[u:2u] + x^{6u}T[3u:4u] + x^{4u}T[5u:6u] + T[9u:10u], \\
0 &= x^{8u}T[2u:3u] + x^{6u}T[4u:5u] + x^{4u}T[6u:7u] + T[10u:11u], \\
0 &= x^{10u}T[u:2u] + x^{8u}T[3u:4u] + T[11u:12u], \\
0 &= x^{10u}T[2u:3u] + x^{8u}T[4u:5u] + T[12u:13u], \\
0 &= x^{10u}T[3u:4u] + x^{8u}T[5u:6u] + T[13u:14u],
\end{aligned}$$

which can be rewritten as

$$(2.7) \quad 0 = T[[1w]_2] + T[[101w]_2] + T[[111w]_2],$$

$$(2.8) \quad 0 = T[[10w]_2] + T[[110w]_2] + T[[1000w]_2],$$

$$(2.9) \quad 0 = T[[1w]_2] + T[[11w]_2] + T[[101w]_2] + T[[1001w]_2],$$

$$(2.10) \quad 0 = T[[10w]_2] + T[[100w]_2] + T[[110w]_2] + T[[1010w]_2],$$

$$(2.11) \quad 0 = T[[1w]_2] + T[[11w]_2] + T[[1011w]_2],$$

$$(2.12) \quad 0 = T[[10w]_2] + T[[100w]_2] + T[[1100w]_2],$$

$$(2.13) \quad 0 = T[[11w]_2] + T[[101w]_2] + T[[1101w]_2],$$

for all binary word w of length $2k$ and $w \neq 0^{2k}$, and

$$(2.14) \quad 0 = T[[1w]_2] + T[[101w]_2] + T[[111w]_2],$$

$$(2.15) \quad 0 = T[[10w]_2] + T[[110w]_2] + T[[1000w]_2],$$

$$(2.16) \quad 0 = T[[1w]_2] + T[[11w]_2] + T[[101w]_2] + T[[1001w]_2],$$

$$(2.17) \quad 0 = T[[10w]_2] + T[[100w]_2] + T[[110w]_2] + T[[1010w]_2],$$

$$(2.18) \quad 0 = T[[1w]_2] + T[[11w]_2] + T[[1011w]_2],$$

$$(2.19) \quad 0 = T[[10w]_2] + T[[100w]_2] + T[[1100w]_2],$$

$$(2.20) \quad 0 = T[[11w]_2] + T[[101w]_2] + T[[1101w]_2],$$

for $w = 0^{2k}$.

First we calculate an 2-automaton that generates T from its minimal polynomial and its first terms. This automaton has 254 states; its transition function and output function can be found in the annex. Let $A(s, w)$ denote the state reached after reading w from right to left starting from the state s , and τ the output function. Define

$$E_{2k} = \{A(i, w) : |w| = 2k, w \neq 0^{2k}\}.$$

Identities (2.7) through (2.20) can be written as

$$(2.21) \quad 0 = \tau(A(s, 1)) + \tau(A(s, 101)) + \tau(A(s, 111)),$$

$$(2.22) \quad 0 = \tau(A(s, 10)) + \tau(A(s, 110)) + \tau(A(s, 1000)),$$

$$(2.23) \quad 0 = \tau(A(s, 1)) + \tau(A(s, 11)) + \tau(A(s, 101)) + \tau(A(s, 1001)),$$

$$(2.24) \quad 0 = \tau(A(s, 10)) + \tau(A(s, 100)) + \tau(A(s, 110)) + \tau(A(s, 1010)),$$

$$(2.25) \quad 0 = \tau(A(s, 1)) + \tau(A(s, 11)) + \tau(A(s, 1011)),$$

$$(2.26) \quad 0 = \tau(A(s, 10)) + \tau(A(s, 100)) + \tau(A(s, 1100)),$$

$$(2.27) \quad 0 = \tau(A(s, 11)) + \tau(A(s, 101)) + \tau(A(s, 1101)),$$

for all $s \in E_{2k}$, and

$$(2.28) \quad 0 = \tau(A(s, 1)) + \tau(A(s, 101)) + \tau(A(s, 111)),$$

$$(2.29) \quad 0 = \tau(A(s, 10)) + \tau(A(s, 110)) + \tau(A(s, 1000)),$$

$$(2.30) \quad 0 = \tau(A(s, 1)) + \tau(A(s, 11)) + \tau(A(s, 101)) + \tau(A(s, 1001)),$$

$$(2.31) \quad 0 = \tau(A(s, 10)) + \tau(A(s, 100)) + \tau(A(s, 110)) + \tau(A(s, 1010)),$$

$$(2.32) \quad 0 = \tau(A(s, 1)) + \tau(A(s, 11)) + \tau(A(s, 1011)),$$

$$(2.33) \quad 0 = \tau(A(s, 10)) + \tau(A(s, 100)) + \tau(A(s, 1100)),$$

$$(2.34) \quad 0 = \tau(A(s, 11)) + \tau(A(s, 101)) + \tau(A(s, 1101)),$$

for $s = A(i, 0^{2k})$. We find that $(A(i, 0^{24}), E_{24}) = (A(i, 0^{12}), E_{12})$, so that we only have to verify that identities (2.21) through (2.34) hold for $2 \leq k \leq 12$, which turns out to be true. $\square$

In the following lemma, we express M_{2k} , M_{2k+1} , W_{2k} , and W_{2k+1} in terms of M^e , M^o , W^e , and W^o .

Lemma 2.3. *For all integer $k \geq 2$, and $u = 2^{2k-1}$,*

$$M_{2k} = M^e[:7u] + x^{2u}M^e[:5u] + x^{6u}M^e[:u] + x^{7u}R^e,$$

$$W_{2k} = W^e[:7u] + x^{2u}W^e[:5u] + x^{6u}W^e[:u] + x^{7u}R^e.$$

For all integer $k \geq 2$, and $u = 2^{2k}$,

$$M_{2k+1} = M^o[:7u] + x^{2u}M^o[:5u] + x^{6u}M^o[:u] + x^{7u}R^o,$$

$$W_{2k+1} = W^o[:7u] + x^{2u}W^o[:5u] + x^{6u}W^o[:u] + x^{7u}R^o.$$

Proof. We call the four identities in Lemma 2.3 also by the name M_{2k} , W_{2k} , M_{2k+1} , and W_{2k+1} . For $n = 2$, the identities can be verified directly. For $n \geq 2$, we claim that

$$M_{2k} \wedge W_{2k} \Rightarrow M_{2k+1} \wedge W_{2k+1},$$

$$M_{2k+1} \wedge W_{2k+1} \Rightarrow M_{2k+2} \wedge W_{2k+2}.$$

We give the proof of

$$M_{2k} \wedge W_{2k} \Rightarrow M_{2k+1},$$

the other ones being similar. Set $u = 2^{2k}$ and $v = 2^{2k-1}$. By definition and induction hypothesis, the left side of identity M_{2k+1} is equal to

$$(2.35) \quad \begin{aligned} W_{2k} M_{2k} = & (W^e[:7v] + x^{2v} W^e[:5v] + x^{6v} W^e[:v] + x^{7u} R^e) \times (M^e[:7v] \\ & + x^{2v} M^e[:5v] + x^{6v} M^e[:v] + x^{7u} R^e); \end{aligned}$$

Call this expression *lhs*. Note that both sides of identity M_{2k+1} have the same term of highest degree $x^{14v} R^o$. Therefore we only need to prove that their difference is $O(x^{14v})$. Using Lemma 2.1 it can be seen that the right side of identity M_{2k+1} is congruent, modulo x^{14v} , to

$$W^e[:14v] M^e[:14v] + x^{4v} W^e[:10v] M^e[:10v] + x^{12v} W^e[:2v] M^e[:2v].$$

For all $n \leq 14$, replace the occurrences of $W^e[:n \cdot v]$ and $M^e[:n \cdot v]$ in the above expression by the reduction modulo $x^{n \cdot v}$ of the right side of the corresponding identity in Lemma 2.2 and get a new expression, which we call *rhs*. Define

$$\begin{aligned} X &:= x^v, \\ a_n &:= W^e[n \cdot v : (n+1) \cdot v] / X^n, \\ b_n &:= M^e[n \cdot v : (n+1) \cdot v] / X^n, \\ c &:= R^e. \end{aligned}$$

Using the notation introduced above, we can represent the expressions *lhs* (2.35) and *rhs* as polynomials in $\mathbb{F}_2[a_1, \dots, a_{14}, b_1, \dots, b_{14}, c][X]$. Note that it is not a problem that a_j commutes with b_k while W^e does not commute with M^e , because in the expressions concerned, the products of W^e -terms and M^e -terms are always in the same order. We let the computer do the simplification and check that the difference between these two polynomials is indeed $O(X^{14})$, which completes the proof. $\square$

Proof of Theorem 1.1. We prove the theorem for $\text{CF}(a, b)$; for $\text{CF}(b, a)$, the proof is similar. By Lemma 2.3, we have For all $0 \leq j, k \leq 1$,

$$\begin{aligned} \lim_{n \rightarrow \infty} M_{2n,j,k} &= M_{j,k}^e, \\ \lim_{n \rightarrow \infty} M_{2n+1,j,k} &= M_{j,k}^o, \\ \lim_{n \rightarrow \infty} W_{2n,j,k} &= W_{j,k}^e, \\ \lim_{n \rightarrow \infty} W_{2n+1,j,k} &= W_{j,k}^o. \end{aligned}$$

Let $z = 1/x$. By the convergence theorem and identity (2.2),

$$\text{CF}(a, b) = \frac{M_{0,1}^e(x)}{M_{0,0}^e(x)}.$$

By definition, $\phi(M^e, 0, 1)$ and $\phi(M^e, 0, 0)$ are minimal polynomials of $M_{0,1}^e$ and $M_{0,0}^e$. Therefore

$$P(x, y) = \text{Res}_t (\phi(M^e, 0, 1)(x, t), y^{12} \phi(M^e, 0, 1)(x, t/y))$$

is an annihilating polynomial of $f(x) = M_{0,1}^e / M_{0,0}^e$.

Define

$$Q(x, y) = q_4(x)y^4 + q_3(x)y^3 + q_2(x)y^2 + q_1(x)y + q_0(x),$$

where

$$\begin{aligned} q_0(x) &= x^{24} + x^{21} + x^{19} + x^{18} + x^{17} + x^{16} + x^{14} + x^{11} + x^9 + x^7, \\ q_1(x) &= x^{20} + x^{17} + x^{15} + x^{14} + x^{13} + x^{10} + x^7 + x^5 + x^4 + x^3, \\ q_2(x) &= x^{16} + x^{10} + x^6 + 1, \\ q_3(x) &= x^{20} + x^{17} + x^{15} + x^{14} + x^{13} + x^{10} + x^7 + x^5 + x^4 + x^3, \\ q_4(x) &= x^{21} + x^{19} + x^{18} + x^{17} + x^{16} + x^{14} + x^{12} + x^{11} + x^{10} + x^9 + x^8 + x^7 \\ &\quad + x^6. \end{aligned}$$

The polynomial $Q(x, y)$ is the candidate for the minimal polynomial of $f(x)$ found by Padé-Hermite approximation. To prove that it is indeed the minimal polynomial of $f(x)$, we only need to prove that it is an irreducible factor of $P(x, y)$ of multiplicity m and $R(x, y) := P(x, y)/Q(x, y)^m$ is not an annihilating polynomial of $f(x)$. We verify the first point directly. For the second point, we find that when we truncate $f(x)$ to order 216, and substitute it for y in $R(z, y)$, we get a series with valuation smaller than 216, which proves that $R(z, y)$ is not an annihilating polynomial of $f(x)$. Finally, $z^{12}Q(1/z, y)$ is the minimal polynomial of $\text{CF}(a, b) = f(1/z)$. $\square$

3. ANNEXE

All 16 polynomials are of the form

$$p_0(x) + p_3(x)y^3 + p_6(x)y^6 + p_9(x)y^9 + p_{12}(x)y^{12}.$$

The coefficients $p_j(x)$ and the 16 initial terms to determine the solutions uniquely are given below.

For $\phi(M^e, 0, 0)$:

$$\begin{aligned} p_0(x) &= x^{126} + x^{122} + x^{120} + x^{116} + x^{110} + x^{108} + x^{102} + x^{100} + x^{96} + x^{94} \\ &\quad + x^{92} + x^{90} + x^{88} + x^{86} + x^{84} + x^{82} + x^{80} + x^{78} + x^{76} + x^{72} + x^{70} \\ &\quad + x^{66} + x^{62} + x^{60} + x^{54} + x^{52} + x^{50} + x^{46} + x^{44} + x^{38} + x^{36}, \\ p_3(x) &= x^{114} + x^{112} + x^{110} + x^{106} + x^{102} + x^{98} + x^{96} + x^{94} + x^{92} + x^{90} + x^{86} \\ &\quad + x^{84} + x^{74} + x^{70} + x^{68} + x^{64} + x^{60} + x^{58} + x^{56} + x^{54} + x^{52} + x^{50} \\ &\quad + x^{46} + x^{44} + x^{32} + x^{28} + x^{26} + x^{24} + x^{22} + x^{20}, \\ p_6(x) &= x^{114} + x^{112} + x^{110} + x^{100} + x^{94} + x^{90} + x^{82} + x^{78} + x^{74} + x^{66} + x^{60} \\ &\quad + x^{54} + x^{52} + x^{48} + x^{46} + x^{42} + x^{40} + x^{38} + x^{32} + x^{30} + x^{26} + x^{12} \\ &\quad + x^{10} + x^8 + x^6 + 1, \\ p_9(x) &= x^{112} + x^{104} + x^{98} + x^{94} + x^{92} + x^{90} + x^{78} + x^{76} + x^{68} + x^{66} + x^{58} \\ &\quad + x^{56} + x^{48} + x^{46} + x^{38} + x^{36} + x^{32} + x^{28} + x^{26} + x^{24} + x^{16} + x^{14} \\ &\quad + x^{12} + x^{10} + x^8 + x^6, \\ p_{12}(x) &= x^{112} + x^{96} + x^{80} + x^{32} + x^{16} + 1, \end{aligned}$$

and the initial terms are $[1, 0, 0, 0, 0, 0, 1, 0, 0, 0, 1, 0, 1, 0, 0, 0]$.

For $\phi(M^e, 0, 1)$:

$$\begin{aligned}
p_0(x) &= x^{129} + x^{125} + x^{124} + x^{123} + x^{122} + x^{120} + x^{119} + x^{118} + x^{114} + x^{112} \\
&\quad + x^{111} + x^{110} + x^{108} + x^{106} + x^{104} + x^{103} + x^{102} + x^{97} + x^{95} + x^{94} \\
&\quad + x^{93} + x^{91} + x^{87} + x^{85} + x^{82} + x^{81} + x^{80} + x^{79} + x^{78} + x^{77} + x^{76} \\
&\quad + x^{75} + x^{74} + x^{70} + x^{68} + x^{67} + x^{66} + x^{65} + x^{64} + x^{63} + x^{61} + x^{58} \\
&\quad + x^{56} + x^{55} + x^{53} + x^{52} + x^{51} + x^{49} + x^{47} + x^{46} + x^{45} + x^{40} + x^{39}, \\
p_3(x) &= x^{118} + x^{116} + x^{115} + x^{112} + x^{110} + x^{109} + x^{108} + x^{106} + x^{105} + x^{102} \\
&\quad + x^{100} + x^{99} + x^{70} + x^{68} + x^{67} + x^{64} + x^{61} + x^{59} + x^{58} + x^{57} + x^{56} \\
&\quad + x^{54} + x^{53} + x^{52} + x^{50} + x^{49} + x^{48} + x^{46} + x^{45} + x^{44} + x^{42} + x^{41} \\
&\quad + x^{40} + x^{37} + x^{35} + x^{34} + x^{33} + x^{30} + x^{28} + x^{27}, \\
p_6(x) &= x^{114} + x^{112} + x^{108} + x^{102} + x^{98} + x^{96} + x^{92} + x^{84} + x^{80} + x^{78} + x^{76} \\
&\quad + x^{70} + x^{62} + x^{60} + x^{58} + x^{54} + x^{50} + x^{48} + x^{46} + x^{42} + x^{26} + x^{24} \\
&\quad + x^{22} + x^{18}, \\
p_9(x) &= x^{110} + x^{108} + x^{107} + x^{96} + x^{94} + x^{93} + x^{92} + x^{89} + x^{88} + x^{87} + x^{84} \\
&\quad + x^{81} + x^{80} + x^{79} + x^{66} + x^{63} + x^{62} + x^{61} + x^{58} + x^{56} + x^{55} + x^{54} \\
&\quad + x^{52} + x^{51} + x^{50} + x^{48} + x^{47} + x^{44} + x^{41} + x^{40} + x^{39} + x^{30} + x^{28} \\
&\quad + x^{27} + x^{26} + x^{23} + x^{22} + x^{21} + x^{18} + x^{15} + x^{13} + x^{11} + x^{10} + x^9, \\
p_{12}(x) &= x^{106} + x^{104} + x^{98} + x^{96} + x^{74} + x^{72} + x^{66} + x^{64} + x^{58} + x^{56} + x^{50} \\
&\quad + x^{48} + x^{42} + x^{40} + x^{34} + x^{32} + x^{10} + x^8 + x^2 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 1, 0, 0, 0, 0, 0, 1, 1, 0, 1, 1, 1]$.

For $\phi(M^e, 1, 0)$:

$$\begin{aligned}
p_0(x) &= x^{129} + x^{125} + x^{124} + x^{123} + x^{122} + x^{120} + x^{119} + x^{118} + x^{114} + x^{112} \\
&\quad + x^{111} + x^{110} + x^{108} + x^{106} + x^{104} + x^{103} + x^{102} + x^{97} + x^{95} + x^{94} \\
&\quad + x^{93} + x^{91} + x^{87} + x^{85} + x^{82} + x^{81} + x^{80} + x^{79} + x^{78} + x^{77} + x^{76} \\
&\quad + x^{75} + x^{74} + x^{70} + x^{68} + x^{67} + x^{66} + x^{65} + x^{64} + x^{63} + x^{61} + x^{58} \\
&\quad + x^{56} + x^{55} + x^{53} + x^{52} + x^{51} + x^{49} + x^{47} + x^{46} + x^{45} + x^{40} + x^{39}, \\
p_3(x) &= x^{118} + x^{116} + x^{115} + x^{112} + x^{110} + x^{109} + x^{108} + x^{106} + x^{105} + x^{102} \\
&\quad + x^{100} + x^{99} + x^{70} + x^{68} + x^{67} + x^{64} + x^{61} + x^{59} + x^{58} + x^{57} + x^{56} \\
&\quad + x^{54} + x^{53} + x^{52} + x^{50} + x^{49} + x^{48} + x^{46} + x^{45} + x^{44} + x^{42} + x^{41} \\
&\quad + x^{40} + x^{37} + x^{35} + x^{34} + x^{33} + x^{30} + x^{28} + x^{27}, \\
p_6(x) &= x^{114} + x^{112} + x^{108} + x^{102} + x^{98} + x^{96} + x^{92} + x^{84} + x^{80} + x^{78} + x^{76} \\
&\quad + x^{70} + x^{62} + x^{60} + x^{58} + x^{54} + x^{50} + x^{48} + x^{46} + x^{42} + x^{26} + x^{24} \\
&\quad + x^{22} + x^{18}, \\
p_9(x) &= x^{110} + x^{108} + x^{107} + x^{96} + x^{94} + x^{93} + x^{92} + x^{89} + x^{88} + x^{87} + x^{84} \\
&\quad + x^{81} + x^{80} + x^{79} + x^{66} + x^{63} + x^{62} + x^{61} + x^{58} + x^{56} + x^{55} + x^{54}
\end{aligned}$$

$$\begin{aligned}
& + x^{52} + x^{51} + x^{50} + x^{48} + x^{47} + x^{44} + x^{41} + x^{40} + x^{39} + x^{30} + x^{28} \\
& + x^{27} + x^{26} + x^{23} + x^{22} + x^{21} + x^{18} + x^{15} + x^{13} + x^{11} + x^{10} + x^9, \\
p_{12}(x) = & x^{106} + x^{104} + x^{98} + x^{96} + x^{74} + x^{72} + x^{66} + x^{64} + x^{58} + x^{56} + x^{50} \\
& + x^{48} + x^{42} + x^{40} + x^{34} + x^{32} + x^{10} + x^8 + x^2 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 1, 0, 0, 0, 0, 0, 1, 1, 0, 1, 1, 1]$.

For $\phi(M^e, 1, 1)$:

$$\begin{aligned}
p_0(x) = & x^{132} + x^{128} + x^{126} + x^{110} + x^{100} + x^{98} + x^{96} + x^{94} + x^{90} + x^{88} + x^{86} \\
& + x^{74} + x^{68} + x^{66} + x^{64} + x^{56} + x^{54} + x^{50} + x^{42}, \\
p_3(x) = & x^{108} + x^{104} + x^{100} + x^{96} + x^{92} + x^{88} + x^{74} + x^{68} + x^{66} + x^{64} + x^{60} \\
& + x^{58} + x^{56} + x^{52} + x^{48} + x^{34} + x^{26} + x^{18}, \\
p_6(x) = & x^{108} + x^{104} + x^{94} + x^{92} + x^{84} + x^{82} + x^{80} + x^{78} + x^{76} + x^{72} + x^{68} \\
& + x^{58} + x^{48} + x^{46} + x^{42} + x^{38} + x^{36} + x^{32} + x^{24} + x^{18} + x^{14} + x^{12} \\
& + x^8 + x^6 + x^4 + 1, \\
p_9(x) = & x^{100} + x^{96} + x^{92} + x^{88} + x^{86} + x^{84} + x^{74} + x^{72} + x^{70} + x^{68} + x^{66} \\
& + x^{64} + x^{62} + x^{60} + x^{58} + x^{48} + x^{40} + x^{36} + x^{34} + x^{30} + x^{26} + x^{22} \\
& + x^{20} + x^{18} + x^8 + x^6, \\
p_{12}(x) = & x^{100} + x^{96} + x^{68} + x^{64} + x^{52} + x^{48} + x^{36} + x^{32} + x^4 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 0, 0, 0, 1, 0, 0, 0, 0, 0, 1, 0]$.

For $\phi(W^e, 0, 0)$:

$$\begin{aligned}
p_0(x) = & x^{132} + x^{128} + x^{126} + x^{116} + x^{112} + x^{110} + x^{98} + x^{90} + x^{86} + x^{84} \\
& + x^{82} + x^{80} + x^{78} + x^{70} + x^{64} + x^{62} + x^{60} + x^{56} + x^{54} + x^{46} + x^{44} \\
& + x^{42} + x^{40} + x^{38} + x^{36}, \\
p_3(x) = & x^{108} + x^{104} + x^{100} + x^{96} + x^{86} + x^{80} + x^{76} + x^{72} + x^{68} + x^{64} + x^{62} \\
& + x^{56} + x^{46} + x^{40} + x^{36} + x^{32} + x^{22} + x^{20}, \\
p_6(x) = & x^{108} + x^{104} + x^{94} + x^{88} + x^{86} + x^{82} + x^{80} + x^{78} + x^{74} + x^{72} + x^{66} \\
& + x^{64} + x^{62} + x^{60} + x^{44} + x^{42} + x^{38} + x^{34} + x^{32} + x^{28} + x^{26} + x^{22} \\
& + x^{20} + x^{14} + x^8 + x^6 + x^4 + 1, \\
p_9(x) = & x^{100} + x^{96} + x^{92} + x^{88} + x^{86} + x^{84} + x^{74} + x^{72} + x^{70} + x^{68} + x^{66} \\
& + x^{64} + x^{62} + x^{60} + x^{58} + x^{48} + x^{40} + x^{36} + x^{34} + x^{30} + x^{26} + x^{22} \\
& + x^{20} + x^{18} + x^8 + x^6, \\
p_{12}(x) = & x^{100} + x^{96} + x^{68} + x^{64} + x^{52} + x^{48} + x^{36} + x^{32} + x^4 + 1,
\end{aligned}$$

and the initial terms are $[1, 0, 0, 0, 0, 0, 0, 0, 1, 0, 0, 0, 1, 0, 1, 0]$.

For $\phi(W^e, 0, 1)$:

$$\begin{aligned}
p_0(x) = & x^{129} + x^{125} + x^{123} + x^{118} + x^{116} + x^{113} + x^{110} + x^{109} + x^{108} + x^{98} \\
& + x^{94} + x^{92} + x^{88} + x^{86} + x^{81} + x^{80} + x^{79} + x^{77} + x^{76} + x^{75} + x^{74} \\
& + x^{72} + x^{71} + x^{70} + x^{68} + x^{66} + x^{64} + x^{61} + x^{60} + x^{59} + x^{58} + x^{57}
\end{aligned}$$

$$\begin{aligned}
& + x^{56} + x^{54} + x^{53} + x^{52} + x^{50} + x^{48} + x^{47} + x^{45} + x^{44} + x^{40} + x^{39}, \\
p_3(x) &= x^{118} + x^{116} + x^{115} + x^{112} + x^{110} + x^{109} + x^{108} + x^{106} + x^{105} + x^{102} \\
& + x^{100} + x^{99} + x^{70} + x^{68} + x^{67} + x^{64} + x^{61} + x^{59} + x^{58} + x^{57} + x^{56} \\
& + x^{54} + x^{53} + x^{52} + x^{50} + x^{49} + x^{48} + x^{46} + x^{45} + x^{44} + x^{42} + x^{41} \\
& + x^{40} + x^{37} + x^{35} + x^{34} + x^{33} + x^{30} + x^{28} + x^{27}, \\
p_6(x) &= x^{114} + x^{112} + x^{108} + x^{102} + x^{98} + x^{96} + x^{92} + x^{84} + x^{80} + x^{78} + x^{76} \\
& + x^{70} + x^{62} + x^{60} + x^{58} + x^{54} + x^{50} + x^{48} + x^{46} + x^{42} + x^{26} + x^{24} \\
& + x^{22} + x^{18}, \\
p_9(x) &= x^{110} + x^{108} + x^{107} + x^{96} + x^{94} + x^{93} + x^{92} + x^{89} + x^{88} + x^{87} + x^{84} \\
& + x^{81} + x^{80} + x^{79} + x^{66} + x^{63} + x^{62} + x^{61} + x^{58} + x^{56} + x^{55} + x^{54} \\
& + x^{52} + x^{51} + x^{50} + x^{48} + x^{47} + x^{44} + x^{41} + x^{40} + x^{39} + x^{30} + x^{28} \\
& + x^{27} + x^{26} + x^{23} + x^{22} + x^{21} + x^{18} + x^{15} + x^{13} + x^{11} + x^{10} + x^9, \\
p_{12}(x) &= x^{106} + x^{104} + x^{98} + x^{96} + x^{74} + x^{72} + x^{66} + x^{64} + x^{58} + x^{56} + x^{50} \\
& + x^{48} + x^{42} + x^{40} + x^{34} + x^{32} + x^{10} + x^8 + x^2 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 1, 0, 1, 0, 1, 0, 1, 1, 0, 0, 0, 1, 1]$.

For $\phi(W^e, 1, 0)$:

$$\begin{aligned}
p_0(x) &= x^{129} + x^{125} + x^{123} + x^{118} + x^{116} + x^{113} + x^{110} + x^{109} + x^{108} + x^{98} \\
& + x^{94} + x^{92} + x^{88} + x^{86} + x^{81} + x^{80} + x^{79} + x^{77} + x^{76} + x^{75} + x^{74} \\
& + x^{72} + x^{71} + x^{70} + x^{68} + x^{66} + x^{64} + x^{61} + x^{60} + x^{59} + x^{58} + x^{57} \\
& + x^{56} + x^{54} + x^{53} + x^{52} + x^{50} + x^{48} + x^{47} + x^{45} + x^{44} + x^{40} + x^{39}, \\
p_3(x) &= x^{118} + x^{116} + x^{115} + x^{112} + x^{110} + x^{109} + x^{108} + x^{106} + x^{105} + x^{102} \\
& + x^{100} + x^{99} + x^{70} + x^{68} + x^{67} + x^{64} + x^{61} + x^{59} + x^{58} + x^{57} + x^{56} \\
& + x^{54} + x^{53} + x^{52} + x^{50} + x^{49} + x^{48} + x^{46} + x^{45} + x^{44} + x^{42} + x^{41} \\
& + x^{40} + x^{37} + x^{35} + x^{34} + x^{33} + x^{30} + x^{28} + x^{27}, \\
p_6(x) &= x^{114} + x^{112} + x^{108} + x^{102} + x^{98} + x^{96} + x^{92} + x^{84} + x^{80} + x^{78} + x^{76} \\
& + x^{70} + x^{62} + x^{60} + x^{58} + x^{54} + x^{50} + x^{48} + x^{46} + x^{42} + x^{26} + x^{24} \\
& + x^{22} + x^{18}, \\
p_9(x) &= x^{110} + x^{108} + x^{107} + x^{96} + x^{94} + x^{93} + x^{92} + x^{89} + x^{88} + x^{87} + x^{84} \\
& + x^{81} + x^{80} + x^{79} + x^{66} + x^{63} + x^{62} + x^{61} + x^{58} + x^{56} + x^{55} + x^{54} \\
& + x^{52} + x^{51} + x^{50} + x^{48} + x^{47} + x^{44} + x^{41} + x^{40} + x^{39} + x^{30} + x^{28} \\
& + x^{27} + x^{26} + x^{23} + x^{22} + x^{21} + x^{18} + x^{15} + x^{13} + x^{11} + x^{10} + x^9, \\
p_{12}(x) &= x^{106} + x^{104} + x^{98} + x^{96} + x^{74} + x^{72} + x^{66} + x^{64} + x^{58} + x^{56} + x^{50} \\
& + x^{48} + x^{42} + x^{40} + x^{34} + x^{32} + x^{10} + x^8 + x^2 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 1, 0, 1, 0, 1, 0, 1, 1, 0, 0, 0, 1, 1]$.

For $\phi(W^e, 1, 1)$:

$$\begin{aligned}
p_0(x) &= x^{126} + x^{122} + x^{120} + x^{110} + x^{106} + x^{98} + x^{94} + x^{90} + x^{82} + x^{78} + x^{76} \\
&\quad + x^{72} + x^{68} + x^{66} + x^{62} + x^{60} + x^{56} + x^{54} + x^{52} + x^{46} + x^{42}, \\
p_3(x) &= x^{114} + x^{112} + x^{110} + x^{106} + x^{104} + x^{102} + x^{96} + x^{88} + x^{86} + x^{82} \\
&\quad + x^{80} + x^{66} + x^{64} + x^{58} + x^{56} + x^{48} + x^{46} + x^{42} + x^{40} + x^{34} + x^{32} \\
&\quad + x^{30} + x^{22} + x^{18}, \\
p_6(x) &= x^{114} + x^{112} + x^{110} + x^{104} + x^{100} + x^{98} + x^{96} + x^{92} + x^{88} + x^{78} + x^{76} \\
&\quad + x^{72} + x^{70} + x^{68} + x^{60} + x^{52} + x^{50} + x^{46} + x^{40} + x^{38} + x^{36} + x^{34} \\
&\quad + x^{28} + x^{24} + x^{18} + x^{16} + x^{10} + x^8 + x^6 + 1, \\
p_9(x) &= x^{112} + x^{104} + x^{98} + x^{94} + x^{92} + x^{90} + x^{78} + x^{76} + x^{68} + x^{66} + x^{58} \\
&\quad + x^{56} + x^{48} + x^{46} + x^{38} + x^{36} + x^{32} + x^{28} + x^{26} + x^{24} + x^{16} + x^{14} \\
&\quad + x^{12} + x^{10} + x^8 + x^6, \\
p_{12}(x) &= x^{112} + x^{96} + x^{80} + x^{32} + x^{16} + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 0, 1, 0, 0, 0, 1, 0, 0, 0, 0, 0]$.

For $\phi(M^o, 0, 0)$:

$$\begin{aligned}
p_0(x) &= x^{129} + x^{125} + x^{124} + x^{123} + x^{122} + x^{121} + x^{120} + x^{119} + x^{117} + x^{108} \\
&\quad + x^{107} + x^{105} + x^{102} + x^{100} + x^{98} + x^{97} + x^{95} + x^{91} + x^{90} + x^{88} \\
&\quad + x^{87} + x^{86} + x^{83} + x^{79} + x^{77} + x^{74} + x^{72} + x^{71} + x^{68} + x^{66} + x^{63} \\
&\quad + x^{62} + x^{61} + x^{59} + x^{54} + x^{53} + x^{52} + x^{51} + x^{50} + x^{49} + x^{48} + x^{43} \\
&\quad + x^{42} + x^{40} + x^{36}, \\
p_3(x) &= x^{122} + x^{120} + x^{119} + x^{113} + x^{110} + x^{109} + x^{106} + x^{104} + x^{103} + x^{102} \\
&\quad + x^{100} + x^{94} + x^{88} + x^{86} + x^{84} + x^{82} + x^{80} + x^{78} + x^{74} + x^{72} + x^{71} \\
&\quad + x^{68} + x^{65} + x^{64} + x^{63} + x^{62} + x^{61} + x^{60} + x^{58} + x^{57} + x^{56} + x^{54} \\
&\quad + x^{53} + x^{50} + x^{49} + x^{46} + x^{45} + x^{44} + x^{42} + x^{41} + x^{40} + x^{39} + x^{38} \\
&\quad + x^{37} + x^{31} + x^{30} + x^{28} + x^{24}, \\
p_6(x) &= x^{122} + x^{120} + x^{119} + x^{113} + x^{110} + x^{109} + x^{106} + x^{104} + x^{102} + x^{100} \\
&\quad + x^{98} + x^{97} + x^{96} + x^{94} + x^{93} + x^{92} + x^{91} + x^{90} + x^{87} + x^{86} + x^{85} \\
&\quad + x^{84} + x^{83} + x^{82} + x^{81} + x^{79} + x^{77} + x^{71} + x^{70} + x^{68} + x^{67} + x^{66} \\
&\quad + x^{63} + x^{61} + x^{59} + x^{58} + x^{54} + x^{51} + x^{50} + x^{49} + x^{47} + x^{46} + x^{44} \\
&\quad + x^{43} + x^{40} + x^{34} + x^{32} + x^{31} + x^{28} + x^{27} + x^{26} + x^{25} + x^{23} + x^{22} \\
&\quad + x^{20} + x^{19} + x^{18} + x^{12} + x^{10} + x^8 + x^2 + 1, \\
p_9(x) &= x^{122} + x^{120} + x^{114} + x^{112} + x^{110} + x^{108} + x^{90} + x^{88} + x^{82} + x^{80} \\
&\quad + x^{78} + x^{76} + x^{74} + x^{72} + x^{66} + x^{64} + x^{62} + x^{60} + x^{58} + x^{56} + x^{50} \\
&\quad + x^{48} + x^{46} + x^{44} + x^{26} + x^{24} + x^{18} + x^{16} + x^{14} + x^{12}, \\
p_{12}(x) &= x^{122} + x^{120} + x^{114} + x^{112} + x^{110} + x^{108} + x^{106} + x^{104} + x^{98} + x^{96}
\end{aligned}$$

$$\begin{aligned}
& + x^{90} + x^{88} + x^{82} + x^{80} + x^{78} + x^{76} + x^{62} + x^{60} + x^{46} + x^{44} + x^{42} \\
& + x^{40} + x^{34} + x^{32} + x^{26} + x^{24} + x^{18} + x^{16} + x^{14} + x^{12} + x^{10} + x^8 \\
& + x^2 + 1,
\end{aligned}$$

and the initial terms are $[1, 0, 0, 0, 0, 0, 1, 1, 1, 1, 1, 1, 0, 0, 0, 1]$.

For $\phi(M^o, 0, 1)$:

$$\begin{aligned}
p_0(x) &= x^{136} + x^{130} + x^{126} + x^{125} + x^{123} + x^{121} + x^{119} + x^{117} + x^{115} + x^{114} \\
&+ x^{113} + x^{110} + x^{107} + x^{106} + x^{103} + x^{101} + x^{100} + x^{99} + x^{95} + x^{94} \\
&+ x^{92} + x^{89} + x^{87} + x^{86} + x^{85} + x^{84} + x^{83} + x^{81} + x^{78} + x^{74} + x^{71} \\
&+ x^{68} + x^{67} + x^{66} + x^{64} + x^{63} + x^{62} + x^{59} + x^{58} + x^{57} + x^{56} + x^{55} \\
&+ x^{54} + x^{52} + x^{51} + x^{49} + x^{47} + x^{45} + x^{44} + x^{43} + x^{42} + x^{41} + x^{40} \\
&+ x^{39}, \\
p_3(x) &= x^{108} + x^{105} + x^{103} + x^{101} + x^{100} + x^{99} + x^{60} + x^{57} + x^{55} + x^{53} + x^{51} \\
&+ x^{49} + x^{47} + x^{45} + x^{43} + x^{41} + x^{39} + x^{37} + x^{35} + x^{33} + x^{31} + x^{29} \\
&+ x^{28} + x^{27}, \\
p_6(x) &= x^{112} + x^{106} + x^{104} + x^{102} + x^{96} + x^{90} + x^{88} + x^{86} + x^{84} + x^{78} + x^{74} \\
&+ x^{72} + x^{70} + x^{60} + x^{56} + x^{54} + x^{48} + x^{44} + x^{42} + x^{24} + x^{20} + x^{18}, \\
p_9(x) &= x^{116} + x^{113} + x^{111} + x^{110} + x^{109} + x^{108} + x^{106} + x^{105} + x^{84} + x^{81} \\
&+ x^{79} + x^{78} + x^{77} + x^{76} + x^{74} + x^{73} + x^{68} + x^{65} + x^{63} + x^{62} + x^{61} \\
&+ x^{60} + x^{58} + x^{57} + x^{52} + x^{49} + x^{47} + x^{46} + x^{45} + x^{44} + x^{42} + x^{41} \\
&+ x^{20} + x^{17} + x^{15} + x^{14} + x^{13} + x^{12} + x^{10} + x^9, \\
p_{12}(x) &= x^{120} + x^{112} + x^{108} + x^{104} + x^{96} + x^{88} + x^{80} + x^{76} + x^{60} + x^{44} + x^{40} \\
&+ x^{32} + x^{24} + x^{16} + x^{12} + x^8 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 1, 0, 0, 0, 0, 0, 1, 0, 1, 0, 1, 0]$.

For $\phi(M^o, 1, 0)$:

$$\begin{aligned}
p_0(x) &= x^{126} + x^{122} + x^{121} + x^{120} + x^{119} + x^{118} + x^{117} + x^{116} + x^{115} + x^{114} \\
&+ x^{113} + x^{110} + x^{106} + x^{103} + x^{102} + x^{100} + x^{99} + x^{96} + x^{94} + x^{93} \\
&+ x^{92} + x^{91} + x^{87} + x^{86} + x^{85} + x^{81} + x^{79} + x^{78} + x^{77} + x^{76} + x^{74} \\
&+ x^{73} + x^{72} + x^{69} + x^{68} + x^{67} + x^{65} + x^{64} + x^{62} + x^{60} + x^{58} + x^{53} \\
&+ x^{49} + x^{45} + x^{43} + x^{42} + x^{41} + x^{40} + x^{39}, \\
p_3(x) &= x^{116} + x^{113} + x^{111} + x^{109} + x^{107} + x^{105} + x^{103} + x^{101} + x^{100} + x^{99} \\
&+ x^{68} + x^{65} + x^{63} + x^{61} + x^{60} + x^{59} + x^{36} + x^{33} + x^{31} + x^{29} + x^{28} \\
&+ x^{27}, \\
p_6(x) &= x^{120} + x^{114} + x^{110} + x^{106} + x^{102} + x^{98} + x^{94} + x^{92} + x^{90} + x^{88} + x^{84} \\
&+ x^{82} + x^{80} + x^{74} + x^{72} + x^{70} + x^{68} + x^{64} + x^{62} + x^{60} + x^{54} + x^{52} \\
&+ x^{50} + x^{48} + x^{44} + x^{42} + x^{32} + x^{28} + x^{26} + x^{24} + x^{20} + x^{18},
\end{aligned}$$

$$\begin{aligned}
p_9(x) &= x^{124} + x^{121} + x^{119} + x^{118} + x^{117} + x^{114} + x^{111} + x^{110} + x^{109} + x^{108} \\
&\quad + x^{106} + x^{105} + x^{92} + x^{89} + x^{87} + x^{86} + x^{85} + x^{82} + x^{79} + x^{78} + x^{77} \\
&\quad + x^{74} + x^{71} + x^{70} + x^{69} + x^{66} + x^{63} + x^{62} + x^{61} + x^{58} + x^{55} + x^{54} \\
&\quad + x^{53} + x^{50} + x^{47} + x^{46} + x^{45} + x^{44} + x^{42} + x^{41} + x^{28} + x^{25} + x^{23} \\
&\quad + x^{22} + x^{21} + x^{18} + x^{15} + x^{14} + x^{13} + x^{12} + x^{10} + x^9, \\
p_{12}(x) &= x^{128} + x^{116} + x^{108} + x^{84} + x^{80} + x^{76} + x^{68} + x^{60} + x^{52} + x^{48} + x^{44} \\
&\quad + x^{20} + x^{12} + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 1, 0, 1, 0, 1, 0, 0, 0, 1, 0, 0, 0, 0]$.

For $\phi(M^o, 1, 1)$:

$$\begin{aligned}
p_0(x) &= x^{129} + x^{125} + x^{123} + x^{121} + x^{117} + x^{113} + x^{110} + x^{109} + x^{108} + x^{107} \\
&\quad + x^{105} + x^{98} + x^{96} + x^{95} + x^{94} + x^{93} + x^{92} + x^{90} + x^{88} + x^{87} + x^{86} \\
&\quad + x^{85} + x^{84} + x^{82} + x^{80} + x^{78} + x^{77} + x^{76} + x^{72} + x^{70} + x^{68} + x^{67} \\
&\quad + x^{65} + x^{63} + x^{61} + x^{59} + x^{58} + x^{57} + x^{56} + x^{53} + x^{51} + x^{50} + x^{48} \\
&\quad + x^{47} + x^{46} + x^{44} + x^{42}, \\
p_3(x) &= x^{122} + x^{120} + x^{119} + x^{114} + x^{113} + x^{112} + x^{110} + x^{109} + x^{108} + x^{103} \\
&\quad + x^{100} + x^{98} + x^{96} + x^{94} + x^{88} + x^{86} + x^{84} + x^{82} + x^{80} + x^{78} + x^{74} \\
&\quad + x^{72} + x^{71} + x^{68} + x^{66} + x^{65} + x^{63} + x^{62} + x^{61} + x^{58} + x^{57} + x^{56} \\
&\quad + x^{53} + x^{52} + x^{49} + x^{48} + x^{45} + x^{41} + x^{39} + x^{37} + x^{36} + x^{31} + x^{28} \\
&\quad + x^{26}, \\
p_6(x) &= x^{122} + x^{120} + x^{119} + x^{114} + x^{113} + x^{112} + x^{110} + x^{109} + x^{108} + x^{100} \\
&\quad + x^{98} + x^{97} + x^{96} + x^{94} + x^{93} + x^{91} + x^{88} + x^{87} + x^{86} + x^{85} + x^{83} \\
&\quad + x^{81} + x^{79} + x^{77} + x^{71} + x^{68} + x^{67} + x^{64} + x^{63} + x^{62} + x^{61} + x^{60} \\
&\quad + x^{59} + x^{54} + x^{52} + x^{51} + x^{50} + x^{49} + x^{47} + x^{44} + x^{43} + x^{38} + x^{36} \\
&\quad + x^{34} + x^{32} + x^{31} + x^{28} + x^{27} + x^{25} + x^{24} + x^{23} + x^{20} + x^{19} + x^{18} \\
&\quad + x^{16} + x^{14} + x^{10} + x^8 + x^2 + 1, \\
p_9(x) &= x^{122} + x^{120} + x^{114} + x^{112} + x^{110} + x^{108} + x^{90} + x^{88} + x^{82} + x^{80} \\
&\quad + x^{78} + x^{76} + x^{74} + x^{72} + x^{66} + x^{64} + x^{62} + x^{60} + x^{58} + x^{56} + x^{50} \\
&\quad + x^{48} + x^{46} + x^{44} + x^{26} + x^{24} + x^{18} + x^{16} + x^{14} + x^{12}, \\
p_{12}(x) &= x^{122} + x^{120} + x^{114} + x^{112} + x^{110} + x^{108} + x^{106} + x^{104} + x^{98} + x^{96} \\
&\quad + x^{90} + x^{88} + x^{82} + x^{80} + x^{78} + x^{76} + x^{62} + x^{60} + x^{46} + x^{44} + x^{42} \\
&\quad + x^{40} + x^{34} + x^{32} + x^{26} + x^{24} + x^{18} + x^{16} + x^{14} + x^{12} + x^{10} + x^8 \\
&\quad + x^2 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 0, 0, 1, 0, 1, 0, 1, 0, 0, 1, 1]$.

For $\phi(W^o, 0, 0)$:

$$p_0(x) = x^{129} + x^{125} + x^{124} + x^{123} + x^{122} + x^{121} + x^{120} + x^{119} + x^{117} + x^{108}$$

$$\begin{aligned}
& + x^{107} + x^{105} + x^{102} + x^{100} + x^{98} + x^{97} + x^{95} + x^{91} + x^{90} + x^{88} \\
& + x^{87} + x^{86} + x^{83} + x^{79} + x^{77} + x^{74} + x^{72} + x^{71} + x^{68} + x^{66} + x^{63} \\
& + x^{62} + x^{61} + x^{59} + x^{54} + x^{53} + x^{52} + x^{51} + x^{50} + x^{49} + x^{48} + x^{43} \\
& + x^{42} + x^{40} + x^{36}, \\
p_3(x) &= x^{122} + x^{120} + x^{119} + x^{113} + x^{110} + x^{109} + x^{106} + x^{104} + x^{103} + x^{102} \\
& + x^{100} + x^{94} + x^{88} + x^{86} + x^{84} + x^{82} + x^{80} + x^{78} + x^{74} + x^{72} + x^{71} \\
& + x^{68} + x^{65} + x^{64} + x^{63} + x^{62} + x^{61} + x^{60} + x^{58} + x^{57} + x^{56} + x^{54} \\
& + x^{53} + x^{50} + x^{49} + x^{46} + x^{45} + x^{44} + x^{42} + x^{41} + x^{40} + x^{39} + x^{38} \\
& + x^{37} + x^{31} + x^{30} + x^{28} + x^{24}, \\
p_6(x) &= x^{122} + x^{120} + x^{119} + x^{113} + x^{110} + x^{109} + x^{106} + x^{104} + x^{102} + x^{100} \\
& + x^{98} + x^{97} + x^{96} + x^{94} + x^{93} + x^{92} + x^{91} + x^{90} + x^{87} + x^{86} + x^{85} \\
& + x^{84} + x^{83} + x^{82} + x^{81} + x^{79} + x^{77} + x^{71} + x^{70} + x^{68} + x^{67} + x^{66} \\
& + x^{63} + x^{61} + x^{59} + x^{58} + x^{54} + x^{51} + x^{50} + x^{49} + x^{47} + x^{46} + x^{44} \\
& + x^{43} + x^{40} + x^{34} + x^{32} + x^{31} + x^{28} + x^{27} + x^{26} + x^{25} + x^{23} + x^{22} \\
& + x^{20} + x^{19} + x^{18} + x^{12} + x^{10} + x^8 + x^2 + 1, \\
p_9(x) &= x^{122} + x^{120} + x^{114} + x^{112} + x^{110} + x^{108} + x^{90} + x^{88} + x^{82} + x^{80} \\
& + x^{78} + x^{76} + x^{74} + x^{72} + x^{66} + x^{64} + x^{62} + x^{60} + x^{58} + x^{56} + x^{50} \\
& + x^{48} + x^{46} + x^{44} + x^{26} + x^{24} + x^{18} + x^{16} + x^{14} + x^{12}, \\
p_{12}(x) &= x^{122} + x^{120} + x^{114} + x^{112} + x^{110} + x^{108} + x^{106} + x^{104} + x^{98} + x^{96} \\
& + x^{90} + x^{88} + x^{82} + x^{80} + x^{78} + x^{76} + x^{62} + x^{60} + x^{46} + x^{44} + x^{42} \\
& + x^{40} + x^{34} + x^{32} + x^{26} + x^{24} + x^{18} + x^{16} + x^{14} + x^{12} + x^{10} + x^8 \\
& + x^2 + 1,
\end{aligned}$$

and the initial terms are $[1, 0, 0, 0, 0, 0, 1, 1, 1, 1, 1, 1, 0, 0, 0, 1]$.

For $\phi(W^o, 0, 1)$:

$$\begin{aligned}
p_0(x) &= x^{126} + x^{122} + x^{121} + x^{120} + x^{119} + x^{118} + x^{117} + x^{116} + x^{115} + x^{114} \\
& + x^{113} + x^{110} + x^{106} + x^{103} + x^{102} + x^{100} + x^{99} + x^{96} + x^{94} + x^{93} \\
& + x^{92} + x^{91} + x^{87} + x^{86} + x^{85} + x^{81} + x^{79} + x^{78} + x^{77} + x^{76} + x^{74} \\
& + x^{73} + x^{72} + x^{69} + x^{68} + x^{67} + x^{65} + x^{64} + x^{62} + x^{60} + x^{58} + x^{53} \\
& + x^{49} + x^{45} + x^{43} + x^{42} + x^{41} + x^{40} + x^{39}, \\
p_3(x) &= x^{116} + x^{113} + x^{111} + x^{109} + x^{107} + x^{105} + x^{103} + x^{101} + x^{100} + x^{99} \\
& + x^{68} + x^{65} + x^{63} + x^{61} + x^{60} + x^{59} + x^{36} + x^{33} + x^{31} + x^{29} + x^{28} \\
& + x^{27}, \\
p_6(x) &= x^{120} + x^{114} + x^{110} + x^{106} + x^{102} + x^{98} + x^{94} + x^{92} + x^{90} + x^{88} + x^{84} \\
& + x^{82} + x^{80} + x^{74} + x^{72} + x^{70} + x^{68} + x^{64} + x^{62} + x^{60} + x^{54} + x^{52} \\
& + x^{50} + x^{48} + x^{44} + x^{42} + x^{32} + x^{28} + x^{26} + x^{24} + x^{20} + x^{18},
\end{aligned}$$

$$\begin{aligned}
p_9(x) &= x^{124} + x^{121} + x^{119} + x^{118} + x^{117} + x^{114} + x^{111} + x^{110} + x^{109} + x^{108} \\
&\quad + x^{106} + x^{105} + x^{92} + x^{89} + x^{87} + x^{86} + x^{85} + x^{82} + x^{79} + x^{78} + x^{77} \\
&\quad + x^{74} + x^{71} + x^{70} + x^{69} + x^{66} + x^{63} + x^{62} + x^{61} + x^{58} + x^{55} + x^{54} \\
&\quad + x^{53} + x^{50} + x^{47} + x^{46} + x^{45} + x^{44} + x^{42} + x^{41} + x^{28} + x^{25} + x^{23} \\
&\quad + x^{22} + x^{21} + x^{18} + x^{15} + x^{14} + x^{13} + x^{12} + x^{10} + x^9, \\
p_{12}(x) &= x^{128} + x^{116} + x^{108} + x^{84} + x^{80} + x^{76} + x^{68} + x^{60} + x^{52} + x^{48} + x^{44} \\
&\quad + x^{20} + x^{12} + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 1, 0, 1, 0, 1, 0, 0, 0, 1, 0, 0, 0, 0]$.

For $\phi(W^\circ, 1, 0)$:

$$\begin{aligned}
p_0(x) &= x^{136} + x^{130} + x^{126} + x^{125} + x^{123} + x^{121} + x^{119} + x^{117} + x^{115} + x^{114} \\
&\quad + x^{113} + x^{110} + x^{107} + x^{106} + x^{103} + x^{101} + x^{100} + x^{99} + x^{95} + x^{94} \\
&\quad + x^{92} + x^{89} + x^{87} + x^{86} + x^{85} + x^{84} + x^{83} + x^{81} + x^{78} + x^{74} + x^{71} \\
&\quad + x^{68} + x^{67} + x^{66} + x^{64} + x^{63} + x^{62} + x^{59} + x^{58} + x^{57} + x^{56} + x^{55} \\
&\quad + x^{54} + x^{52} + x^{51} + x^{49} + x^{47} + x^{45} + x^{44} + x^{43} + x^{42} + x^{41} + x^{40} \\
&\quad + x^{39}, \\
p_3(x) &= x^{108} + x^{105} + x^{103} + x^{101} + x^{100} + x^{99} + x^{60} + x^{57} + x^{55} + x^{53} + x^{51} \\
&\quad + x^{49} + x^{47} + x^{45} + x^{43} + x^{41} + x^{39} + x^{37} + x^{35} + x^{33} + x^{31} + x^{29} \\
&\quad + x^{28} + x^{27}, \\
p_6(x) &= x^{112} + x^{106} + x^{104} + x^{102} + x^{96} + x^{90} + x^{88} + x^{86} + x^{84} + x^{78} + x^{74} \\
&\quad + x^{72} + x^{70} + x^{60} + x^{56} + x^{54} + x^{48} + x^{44} + x^{42} + x^{24} + x^{20} + x^{18}, \\
p_9(x) &= x^{116} + x^{113} + x^{111} + x^{110} + x^{109} + x^{108} + x^{106} + x^{105} + x^{84} + x^{81} \\
&\quad + x^{79} + x^{78} + x^{77} + x^{76} + x^{74} + x^{73} + x^{68} + x^{65} + x^{63} + x^{62} + x^{61} \\
&\quad + x^{60} + x^{58} + x^{57} + x^{52} + x^{49} + x^{47} + x^{46} + x^{45} + x^{44} + x^{42} + x^{41} \\
&\quad + x^{20} + x^{17} + x^{15} + x^{14} + x^{13} + x^{12} + x^{10} + x^9, \\
p_{12}(x) &= x^{120} + x^{112} + x^{108} + x^{104} + x^{96} + x^{88} + x^{80} + x^{76} + x^{60} + x^{44} + x^{40} \\
&\quad + x^{32} + x^{24} + x^{16} + x^{12} + x^8 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 1, 0, 0, 0, 0, 0, 1, 0, 1, 0, 1, 0]$.

For $\phi(W^\circ, 1, 1)$:

$$\begin{aligned}
p_0(x) &= x^{129} + x^{125} + x^{123} + x^{121} + x^{117} + x^{113} + x^{110} + x^{109} + x^{108} + x^{107} \\
&\quad + x^{105} + x^{98} + x^{96} + x^{95} + x^{94} + x^{93} + x^{92} + x^{90} + x^{88} + x^{87} + x^{86} \\
&\quad + x^{85} + x^{84} + x^{82} + x^{80} + x^{78} + x^{77} + x^{76} + x^{72} + x^{70} + x^{68} + x^{67} \\
&\quad + x^{65} + x^{63} + x^{61} + x^{59} + x^{58} + x^{57} + x^{56} + x^{53} + x^{51} + x^{50} + x^{48} \\
&\quad + x^{47} + x^{46} + x^{44} + x^{42}, \\
p_3(x) &= x^{122} + x^{120} + x^{119} + x^{114} + x^{113} + x^{112} + x^{110} + x^{109} + x^{108} + x^{103} \\
&\quad + x^{100} + x^{98} + x^{96} + x^{94} + x^{88} + x^{86} + x^{84} + x^{82} + x^{80} + x^{78} + x^{74}
\end{aligned}$$

$$\begin{aligned}
& + x^{72} + x^{71} + x^{68} + x^{66} + x^{65} + x^{63} + x^{62} + x^{61} + x^{58} + x^{57} + x^{56} \\
& + x^{53} + x^{52} + x^{49} + x^{48} + x^{45} + x^{41} + x^{39} + x^{37} + x^{36} + x^{31} + x^{28} \\
& + x^{26}, \\
p_6(x) = & x^{122} + x^{120} + x^{119} + x^{114} + x^{113} + x^{112} + x^{110} + x^{109} + x^{108} + x^{100} \\
& + x^{98} + x^{97} + x^{96} + x^{94} + x^{93} + x^{91} + x^{88} + x^{87} + x^{86} + x^{85} + x^{83} \\
& + x^{81} + x^{79} + x^{77} + x^{71} + x^{68} + x^{67} + x^{64} + x^{63} + x^{62} + x^{61} + x^{60} \\
& + x^{59} + x^{54} + x^{52} + x^{51} + x^{50} + x^{49} + x^{47} + x^{44} + x^{43} + x^{38} + x^{36} \\
& + x^{34} + x^{32} + x^{31} + x^{28} + x^{27} + x^{25} + x^{24} + x^{23} + x^{20} + x^{19} + x^{18} \\
& + x^{16} + x^{14} + x^{10} + x^8 + x^2 + 1, \\
p_9(x) = & x^{122} + x^{120} + x^{114} + x^{112} + x^{110} + x^{108} + x^{90} + x^{88} + x^{82} + x^{80} \\
& + x^{78} + x^{76} + x^{74} + x^{72} + x^{66} + x^{64} + x^{62} + x^{60} + x^{58} + x^{56} + x^{50} \\
& + x^{48} + x^{46} + x^{44} + x^{26} + x^{24} + x^{18} + x^{16} + x^{14} + x^{12}, \\
p_{12}(x) = & x^{122} + x^{120} + x^{114} + x^{112} + x^{110} + x^{108} + x^{106} + x^{104} + x^{98} + x^{96} \\
& + x^{90} + x^{88} + x^{82} + x^{80} + x^{78} + x^{76} + x^{62} + x^{60} + x^{46} + x^{44} + x^{42} \\
& + x^{40} + x^{34} + x^{32} + x^{26} + x^{24} + x^{18} + x^{16} + x^{14} + x^{12} + x^{10} + x^8 \\
& + x^2 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 0, 0, 1, 0, 1, 0, 1, 0, 0, 1, 1]$.

Below is the transition function and output function of an 2-automaton that generates $T = M_{0,0}^o$:

Transition function $(n, j) \mapsto \delta(n, j)$ ($\Lambda(n) := [\delta(n, 0), \delta(n, 1)]$):

n	$\Lambda(n)$	n	$\Lambda(n)$	n	$\Lambda(n)$	n	$\Lambda(n)$	n	$\Lambda(n)$
0	[1, 2]	51	[28, 40]	102	[150, 151]	153	[10, 12]	204	[177, 28]
1	[3, 4]	52	[8, 2]	103	[152, 153]	154	[186, 92]	205	[231, 8]
2	[5, 6]	53	[88, 89]	104	[154, 155]	155	[90, 97]	206	[232, 227]
3	[7, 8]	54	[29, 90]	105	[156, 157]	156	[187, 177]	207	[172, 174]
4	[9, 9]	55	[91, 92]	106	[144, 158]	157	[99, 107]	208	[29, 81]
5	[10, 11]	56	[28, 43]	107	[159, 160]	158	[177, 30]	209	[91, 230]
6	[12, 13]	57	[93, 93]	108	[161, 162]	159	[46, 81]	210	[86, 12]
7	[14, 15]	58	[94, 33]	109	[163, 164]	160	[39, 182]	211	[101, 2]
8	[16, 17]	59	[35, 36]	110	[165, 166]	161	[188, 89]	212	[39, 175]
9	[18, 19]	60	[36, 93]	111	[167, 168]	162	[46, 174]	213	[233, 101]
10	[20, 21]	61	[93, 35]	112	[169, 170]	163	[188, 189]	214	[172, 12]
11	[22, 23]	62	[95, 96]	113	[171, 22]	164	[174, 40]	215	[45, 42]
12	[24, 25]	63	[30, 97]	114	[96, 84]	165	[190, 185]	216	[174, 182]
13	[26, 27]	64	[98, 39]	115	[44, 107]	166	[98, 28]	217	[234, 41]
14	[10, 28]	65	[30, 13]	116	[172, 11]	167	[88, 189]	218	[11, 82]
15	[29, 30]	66	[12, 97]	117	[173, 109]	168	[12, 82]	219	[99, 2]
16	[31, 32]	67	[99, 100]	118	[9, 94]	169	[36, 35]	220	[11, 13]
17	[33, 34]	68	[101, 102]	119	[86, 11]	170	[34, 93]	221	[235, 83]

18	[34, 35]	69	[10, 39]	120	[174, 175]	171	[178, 41]	222	[46, 12]
19	[9, 36]	70	[103, 104]	121	[7, 99]	172	[191, 192]	223	[96, 100]
20	[37, 29]	71	[35, 94]	122	[176, 46]	173	[193, 194]	224	[90, 13]
21	[8, 38]	72	[105, 30]	123	[44, 102]	174	[195, 196]	225	[236, 237]
22	[39, 40]	73	[90, 106]	124	[177, 39]	175	[50, 197]	226	[238, 239]
23	[41, 42]	74	[41, 107]	125	[81, 175]	176	[198, 199]	227	[240, 241]
24	[11, 43]	75	[105, 81]	126	[83, 100]	177	[200, 201]	228	[242, 243]
25	[44, 42]	76	[108, 109]	127	[28, 175]	178	[202, 72]	229	[244, 245]
26	[45, 38]	77	[34, 9]	128	[178, 45]	179	[203, 204]	230	[246, 202]
27	[46, 11]	78	[110, 111]	129	[10, 174]	180	[205, 51]	231	[222, 5]
28	[47, 48]	79	[112, 112]	130	[90, 43]	181	[206, 207]	232	[247, 248]
29	[49, 50]	80	[113, 46]	131	[45, 2]	182	[21, 208]	233	[214, 249]
30	[51, 52]	81	[63, 114]	132	[41, 84]	183	[209, 210]	234	[250, 251]
31	[53, 54]	82	[115, 116]	133	[30, 106]	184	[211, 212]	235	[252, 253]
32	[55, 56]	83	[117, 118]	134	[7, 83]	185	[213, 214]	236	[235, 99]
33	[33, 33]	84	[119, 120]	135	[29, 28]	186	[215, 216]	237	[174, 43]
34	[57, 58]	85	[121, 66]	136	[112, 93]	187	[217, 218]	238	[8, 42]
35	[59, 60]	86	[122, 123]	137	[112, 35]	188	[219, 220]	239	[12, 106]
36	[58, 61]	87	[124, 125]	138	[29, 174]	189	[221, 222]	240	[95, 101]
37	[62, 63]	88	[126, 127]	139	[179, 104]	190	[223, 224]	241	[105, 28]
38	[64, 65]	89	[128, 129]	140	[36, 9]	191	[225, 29]	242	[105, 174]
39	[66, 67]	90	[130, 131]	141	[99, 102]	192	[41, 102]	243	[177, 81]
40	[68, 69]	91	[132, 133]	142	[98, 30]	193	[226, 227]	244	[233, 96]
41	[70, 71]	92	[134, 135]	143	[180, 177]	194	[105, 90]	245	[81, 43]
42	[72, 73]	93	[19, 59]	144	[96, 87]	195	[174, 82]	246	[234, 45]
43	[74, 75]	94	[136, 137]	145	[181, 32]	196	[101, 100]	247	[44, 84]
44	[76, 77]	95	[138, 119]	146	[93, 112]	197	[86, 39]	248	[81, 182]
45	[78, 79]	96	[139, 140]	147	[28, 182]	198	[228, 8]	249	[98, 81]
46	[80, 26]	97	[141, 142]	148	[183, 111]	199	[90, 82]	250	[86, 28]
47	[81, 82]	98	[143, 144]	149	[94, 34]	200	[229, 86]	251	[46, 39]
48	[83, 84]	99	[145, 146]	150	[172, 81]	201	[101, 107]	252	[177, 90]
49	[85, 86]	100	[15, 147]	151	[11, 106]	202	[98, 90]	253	[172, 39]
50	[83, 87]	101	[148, 149]	152	[184, 185]	203	[186, 230]		

Output function $n \mapsto \tau(n)$:

n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$
0	0	37	0	74	0	111	0	148	0	185	1	222	0
1	0	38	0	75	1	112	1	149	1	186	1	223	1
2	0	39	1	76	1	113	0	150	1	187	1	224	0
3	0	40	0	77	0	114	1	151	1	188	1	225	1
4	0	41	0	78	1	115	1	152	0	189	1	226	0
5	0	42	1	79	1	116	1	153	0	190	1	227	0
6	1	43	0	80	0	117	0	154	1	191	1	228	1
7	0	44	1	81	0	118	0	155	0	192	0	229	1

8	0	45	1	82	1	119	1	156	1	193	0	230	1
9	0	46	0	83	0	120	1	157	1	194	1	231	0
10	0	47	0	84	1	121	0	158	1	195	1	232	1
11	1	48	0	85	0	122	1	159	0	196	0	233	1
12	1	49	0	86	1	123	1	160	1	197	1	234	1
13	1	50	0	87	1	124	1	161	1	198	1	235	1
14	0	51	0	88	0	125	0	162	0	199	0	236	1
15	0	52	0	89	0	126	0	163	1	200	1	237	1
16	0	53	0	90	0	127	0	164	1	201	0	238	0
17	0	54	0	91	0	128	0	165	1	202	0	239	1
18	0	55	0	92	0	129	0	166	0	203	1	240	0
19	0	56	0	93	0	130	0	167	0	204	1	241	1
20	0	57	0	94	1	131	1	168	1	205	0	242	1
21	0	58	1	95	0	132	0	169	1	206	1	243	1
22	1	59	1	96	1	133	0	170	0	207	1	244	1
23	0	60	1	97	1	134	0	171	0	208	0	245	0
24	1	61	0	98	0	135	0	172	1	209	0	246	1
25	1	62	0	99	1	136	1	173	0	210	1	247	1
26	1	63	0	100	0	137	1	174	1	211	0	248	0
27	0	64	0	101	0	138	0	175	0	212	1	249	0
28	0	65	0	102	1	139	1	176	1	213	1	250	1
29	0	66	1	103	0	140	1	177	1	214	1	251	0
30	0	67	1	104	1	141	1	178	0	215	1	252	1
31	0	68	0	105	1	142	0	179	1	216	1	253	1
32	0	69	0	106	1	143	0	180	0	217	1		
33	0	70	0	107	0	144	1	181	1	218	1		
34	0	71	1	108	1	145	1	182	0	219	1		
35	1	72	1	109	1	146	0	183	0	220	1		
36	1	73	0	110	1	147	0	184	0	221	1		

UNIVERSITÉ DE STRASBOURG, CNRS, IRMA UMR 7501, F-67000 STRASBOURG, FRANCE

Email address: `guoniu.han@unistra.fr`

SCHOOL OF MATHEMATICS AND STATISTICS, HUAZHONG UNIVERSITY OF SCIENCE AND TECHNOLOGY, WUHAN, PR CHINA

Email address: `huyining@hust.edu.cn`