

PROOF OF THE CONTINUED FRACTION CONJECTURE FOR

$$(a, b) = (z^5 + z^4, z^2 + 1)$$

GUO-NIU HAN* AND YINING HU*

ABSTRACT. The continued fraction conjecture claims that the continued fraction defined by the (a, b) -Thue-Morse sequence is algebraic over $\mathbb{F}_2(x)$ for all pairs of distinct elements (a, b) in $\mathbb{F}_2[x] \setminus \mathbb{F}_2$. In this paper we prove the conjecture for $(a, b) = (z^5 + z^4, z^2 + 1)$.

1. INTRODUCTION

Let $\mathbb{F}_2$ be the finite field of cardinality 2. Given a sequence of polynomials $\mathbf{a} = (a_0(z), a_1(z), a_2(z), \dots)$ of elements from $\mathbb{F}_2[z] \setminus \mathbb{F}_2$, we define the infinite continued fraction

$$(1.1) \quad \text{CF}(\mathbf{a}(z)) = \frac{1}{a_0(z) + \frac{1}{a_1(z) + \frac{1}{a_2(z) + \frac{1}{\ddots}}}}$$

as the limit of the finite continued fractions

$$(1.2) \quad \text{CF}_n(\mathbf{a}(z)) = \frac{1}{a_0(z) + \frac{1}{a_1(z) + \frac{1}{\ddots + \frac{1}{a_n(z)}}}} \in \mathbb{F}_2((1/z)).$$

Conversely, each power series in $\mathbb{F}_2[[1/z]]$ without constant term

$$(1.3) \quad c_{-1}z^{-1} + c_{-2}z^{-2} + c_{-3}z^{-3} + \dots$$

admits a continued fraction expansion of form (1.1).

Let a and b be two distinct elements from $\mathbb{F}_2[z] \setminus \mathbb{F}_2$. We consider the continued fraction (1.1) associated with the (a, b) -Thue-Morse sequence

$$\mathbf{t} = (t_0(z), t_1(z), t_2(z), \dots) = (a, b, b, a, b, a, a, b, \dots)$$

that we shall denote

$$(1.4) \quad \text{CF}(a, b) := \text{CF}(\mathbf{t}).$$

Date: 9/04/2020 17:26:06 .

2010 Mathematics Subject Classification. 11B85, 11J70, 11B50, 11Y65, 05A15.

Key words and phrases. automatic sequence, continued fraction, Thue-Morse sequence.

* This paper was automatically generated from a computer program developed by the authors, with 8.3 minutes CPU time used.

Theorem 1.1. *When $(a, b) = (z^5 + z^4, z^2 + 1)$, the two power series $\text{CF}(a, b)$ and $\text{CF}(b, a)$ are algebraic over $\mathbb{F}_2(z)$, with minimal polynomials of the form*

$$p_4(z)y^4 + p_3(z)y^3 + p_2(z)y^2 + p_1(z)y + p_0(z) = 0$$

where, for $\text{CF}(a, b)$,

$$\begin{aligned} p_0(z) &= z^{17} + z^{16} + z^{10} + z^8 + z^7 + z^6 + z^5 + 1, \\ p_1(z) &= z^{22} + z^{20} + z^{19} + z^{18} + z^{17} + z^{16} + z^{12} + z^9 + z^5 + z^4, \\ p_2(z) &= z^{24} + z^{20} + z^{14} + z^{12} + z^{10} + z^8, \\ p_3(z) &= z^{22} + z^{20} + z^{19} + z^{18} + z^{17} + z^{16} + z^{12} + z^9 + z^5 + z^4, \\ p_4(z) &= z^{20} + z^{17} + z^{16} + z^{14} + z^{12} + z^7 + z^6 + z^5 + z^4, \end{aligned}$$

and for $\text{CF}(b, a)$,

$$\begin{aligned} p_0(z) &= z^{17} + z^{10} + z^8 + z^7 + z^6 + z^5 + z^4, \\ p_1(z) &= z^{22} + z^{20} + z^{19} + z^{18} + z^{17} + z^{16} + z^{12} + z^9 + z^5 + z^4, \\ p_2(z) &= z^{24} + z^{20} + z^{14} + z^{12} + z^{10} + z^8, \\ p_3(z) &= z^{22} + z^{20} + z^{19} + z^{18} + z^{17} + z^{16} + z^{12} + z^9 + z^5 + z^4, \\ p_4(z) &= z^{20} + z^{17} + z^{14} + z^{12} + z^7 + z^6 + z^5 + 1. \end{aligned}$$

2. PROOF

Define the sequences of polynomials $(P_n(z))_{n \geq 0}$ and $(Q_n(z))_{n \geq 0}$ by

$$\begin{pmatrix} P_n(z) & Q_n(z) \\ P_{n-1}(z) & Q_{n-1}(z) \end{pmatrix} := \begin{pmatrix} t_n(z) & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} t_{n-1}(z) & 1 \\ 1 & 0 \end{pmatrix} \cdots \begin{pmatrix} t_0(z) & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}.$$

By the basic properties of continued fractions, we have

$$(2.1) \quad \text{CF}_n(a, b) = \frac{P_n(z)}{Q_n(z)} \in \mathbb{F}_2((1/z)).$$

As formula (2.1) is not efficient for calculating $\text{CF}_n(a, b)$, a fast method is to be derived and described as follows. Define

$$\begin{aligned} M_n(x) &= x^{\deg(t_{2^n-1})} \begin{pmatrix} t_{2^n-1}(1/x) & 1 \\ 1 & 0 \end{pmatrix} \times x^{\deg(t_{2^n-2})} \begin{pmatrix} t_{2^n-2}(1/x) & 1 \\ 1 & 0 \end{pmatrix} \times \\ &\quad \cdots \times x^{\deg(t_0)} \begin{pmatrix} t_0(1/x) & 1 \\ 1 & 0 \end{pmatrix}, \\ W_n(x) &= x^{\deg(\bar{t}_{2^n-1})} \begin{pmatrix} \bar{t}_{2^n-1}(1/x) & 1 \\ 1 & 0 \end{pmatrix} \times x^{\deg(\bar{t}_{2^n-2})} \begin{pmatrix} \bar{t}_{2^n-2}(1/x) & 1 \\ 1 & 0 \end{pmatrix} \times \\ &\quad \cdots \times x^{\deg(\bar{t}_0)} \begin{pmatrix} \bar{t}_0(1/x) & 1 \\ 1 & 0 \end{pmatrix}, \end{aligned}$$

where $\bar{t}$ is the (b, a) -Thue-Morse sequence. By the properties of the Thue-Morse sequence, we have for $n \geq 0$

$$\begin{aligned} M_{n+1}(x) &= W_n(x) \cdot M_n(x), \\ W_{n+1}(x) &= M_n(x) \cdot W_n(x). \end{aligned}$$

Let $x := 1/z$. For each non-zero polynomial $P(z)$, define $\tilde{P}(x)$ to be $P(1/x)$. Then,

$$\text{CF}_n(a, b) = \frac{P_n(z)}{Q_n(z)} = \frac{\tilde{P}_n(x)}{\tilde{Q}_n(x)} \in \mathbb{F}_2((x)) = \mathbb{F}_2((1/z)).$$

Comparing the definition of $M_n(x)$ with the definition of $P_n(x)$ and $Q_n(x)$, we see that

$$\begin{aligned} M_n(x)_{0,1} &= x^{d_n} \tilde{P}_{2^n-1}(x), \\ M_n(x)_{0,0} &= x^{d_n} \tilde{Q}_{2^n-1}(x), \end{aligned}$$

for some positive integer d_n . Hence,

$$(2.2) \quad \text{CF}_{2^{2^n}-1}(a, b) = \frac{\tilde{P}_{2^{2^n}-1}(x)}{\tilde{Q}_{2^{2^n}-1}(x)} = \frac{M_{2^n}(x)_{0,1}}{M_{2^n}(x)_{0,0}}.$$

Therefore, by the convergence theorem of continued fraction, the algebraicity of $\text{CF}(a, b)$ will be established if it is shown that both $M_{2^n}(x)_{0,1}$ and $M_{2^n}(x)_{0,0}$ converge to algebraic series in $\mathbb{F}_2[[x]]$.

Actually, we will prove that for all $0 \leq i, j \leq 1$ the four sequences $(M_{2^n}(x)_{i,j})_n$, $(M_{2^{n+1}}(x)_{i,j})_n$, $(W_{2^n}(x)_{i,j})_n$, and $(W_{2^{n+1}}(x)_{i,j})_n$ converge to algebraic series in $\mathbb{F}_2[[x]]$. For this purpose, we define four 2×2 matrices M^e, M^o, W^e, W^o as follows: For each $T \in \{M^e, M^o, W^e, W^o\}$ and $0 \leq i, j \leq 1$, $T_{i,j}$ is defined to be the unique solution in $\mathbb{F}_2[[x]]$ of the polynomial $\phi(T, i, j)$ under certain initial conditions; the polynomials $\phi(T, i, j)$ and initial conditions are given in Section 3. We will prove that these four matrices, whose components are algebraic by definition, are the limits of $(M_{2^n}(x))_n$, $(M_{2^{n+1}}(x))_n$, $(W_{2^n}(x))_n$, and $(W_{2^{n+1}}(x))_n$.

Let us explain how the polynomials $\phi(T, i, j)$ and initial conditions are found, and why the solutions exist and are unique. For $0 \leq i, j \leq 1$, the coefficients of the polynomial $\phi(M^e, i, j)$ (resp. $\phi(M^o, i, j)$, $\phi(W^e, i, j)$, and $\phi(W^o, i, j)$) are the Padé-Hermite approximants of type

$$(175, 175, 175, 175, 175)$$

of the vector

$$(1, T^3, T^6, T^9, T^{12}),$$

where $T = M_{12,i,j}$ (resp. $M_{11,i,j}$, $W_{12,i,j}$, and $W_{11,i,j}$), found by the Derksen algorithm. We take the first 8 terms of $M_{12,i,j}$ (resp. $M_{11,i,j}$, $W_{12,i,j}$, and $W_{11,i,j}$) as initial conditions for $\phi(M^e, i, j)$ (resp. $\phi(M^o, i, j)$, $\phi(W^e, i, j)$, and $\phi(W^o, i, j)$). The following fact will be used to ensure that the solution exists and is unique: let $P(x, y) \in \mathbb{F}_2[x, y]$ and for each series $f(x) = \sum_{n=0}^{\infty} a_n x^n \in \mathbb{F}_2[[x]]$ denote the partial sum $\sum_{j=0}^{n-1} a_j x^j$ by $f_n(x)$ for $n \geq 0$. If for some $n \geq 0$ and $a_0, a_1, \dots, a_{n-1} \in \mathbb{F}_2$ $P(x, \sum_{j=0}^{n-1} a_j x^j) = O(x^n)$ and $Q(x, y) := P(x, \sum_{j=0}^{n-1} a_j x^j + x^n y)$ can be written as $x^m \sum_{j=0}^{\infty} q_j(x) y^j$ where $q_j(x)$ are polynomials for $j \geq 0$, $q_1(0) = 1$, and $q_j(0) = 0$ for $j > 1$, then there exists a unique solution $f(x) \in \mathbb{F}_2[[x]]$ of $P(x, f(x)) = 0$ that satisfies the initial condition $f_n(x) = \sum_{j=0}^{n-1} a_j x^j$.

We state two lemmas concerning the four matrices M^e, M^o, W^e, W^o . The first one is about relations between them; the second, about the structure of the each matrix.

Lemma 2.1. *We have*

$$(2.3) \quad M^e = W^o \cdot M^o,$$

$$(2.4) \quad M^o = W^e \cdot M^e,$$

$$(2.5) \quad W^e = M^o \cdot W^o,$$

$$(2.6) \quad W^o = M^e \cdot W^e.$$

Proof. We give the proof of the identity

$$M_{0,0}^e = W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o,$$

the proofs of the others being similar.

First, we compute the minimal polynomials of $W_{0,0}^o M_{0,0}^o$ and $W_{0,1}^o M_{1,0}^o$. We know that

$$P(x, y) = \text{Res}_z (\phi(W^o, 0, 0)(x, z), z^{12} \cdot \phi(M^o, 0, 0)(x, y/z))$$

is an annihilating polynomial of $W_{0,0}^o M_{0,0}^o$. We use Padé-Hermite approximation to find a candidate for the minimal polynomial of $W_{0,0}^o M_{0,0}^o$, that will be called $\phi_0(x, y)$. To prove that $\phi_0(x, y)$ is indeed the minimal polynomial, it suffices to prove that it is an irreducible factor of $P(x, y)$ of multiplicity m and that $Q(x, y) := P(x, y)/\phi_0(x, y)^m$ is not an annihilating polynomial of $W_{0,0}^o M_{0,0}^o$. We verify the first point directly. For the second point, we truncate $W_{0,0}^o M_{0,0}^o$ to order 630 and substitute it for y in $Q(x, y)$. We get a series of valuation less than 630, which proves that $Q(x, y)$ is not an annihilating polynomial of $W_{0,0}^o M_{0,0}^o$. We find the minimal polynomial $\phi_1(x, y)$ of $W_{0,1}^o M_{1,0}^o$ in a similar way.

Now we prove that $\phi(M^e, 0, 0)$ is the minimal polynomial of $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$. We know that

$$S(x, y) = \text{Res}_z (\phi_0(x, z), \phi_1(x, y + z))$$

is an annihilating of $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$. We verify that $\phi(M^e, 0, 0)$ is an irreducible factor of $S(x, y)$ of multiplicity μ , and that $Q(x, y) := S(x, y)/\phi(M^e, 0, 0)^\mu$ is not an annihilating polynomial of $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$. To see the last point, we truncate $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$ to order 770 and substitute it for y in $Q(x, y)$. We get a series of valuation less than 770, and therefore $Q(x, y)$ is not an annihilating polynomial of $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$.

Finally, the first 16 terms of $M_{0,0}^e$ and $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$ coincide. As these first terms determine a unique solution of $\phi(M^e, 0, 0)$, we know that the two series are one and the same. $\square$

Define

$$R^e = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \quad \text{and} \quad R^o = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}.$$

Lemma 2.2. *For all integers $k \geq 2$ and $u = 2^{2k-1}$, the following identities hold.*

$$\begin{aligned} M^e[:14u] &= M^e[:7u] + x^u M^e[:6u] + x^{2u} M^e[:5u] + x^{3u} M^e[:4u] + x^{4u} M^e[:3u] \\ &\quad + x^{5u} M^e[:2u] + x^{6u} M^e[:u] + x^u M^e[:7u] + x^{2u} M^e[:6u] \\ &\quad + x^{3u} M^e[:5u] + x^{4u} M^e[:4u] + x^{5u} M^e[:3u] + x^{6u} M^e[:2u] \\ &\quad + x^{7u} M^e[:u] + x^{8u} M^e[:6u] + x^{10u} M^e[:4u] + x^{12u} M^e[:2u] \\ &\quad + (x^{7u} + x^{8u})R^e, \\ W^e[:14u] &= W^e[:7u] + x^u W^e[:6u] + x^{2u} W^e[:5u] + x^{3u} W^e[:4u] + x^{4u} W^e[:3u] \end{aligned}$$

$$\begin{aligned}
& + x^{5u}W^e[:2u] + x^{6u}W^e[:u] + x^uW^e[:7u] + x^{2u}W^e[:6u] \\
& + x^{3u}W^e[:5u] + x^{4u}W^e[:4u] + x^{5u}W^e[:3u] + x^{6u}W^e[:2u] \\
& + x^{7u}W^e[:u] + x^{8u}W^e[:6u] + x^{10u}W^e[:4u] + x^{12u}W^e[:2u] \\
& + (x^{7u} + x^{8u})R^e.
\end{aligned}$$

For all integers $k \geq 2$ and $u = 2^{2k}$, the following identities hold.

$$\begin{aligned}
M^o[:14u] &= M^o[:7u] + x^uM^o[:6u] + x^{2u}M^o[:5u] + x^{3u}M^o[:4u] + x^{4u}M^o[:3u] \\
&+ x^{5u}M^o[:2u] + x^{6u}M^o[:u] + x^uM^o[:7u] + x^{2u}M^o[:6u] \\
&+ x^{3u}M^o[:5u] + x^{4u}M^o[:4u] + x^{5u}M^o[:3u] + x^{6u}M^o[:2u] \\
&+ x^{7u}M^o[:u] + x^{8u}M^o[:6u] + x^{10u}M^o[:4u] + x^{12u}M^o[:2u] \\
&+ (x^{7u} + x^{8u})R^o, \\
W^o[:14u] &= W^o[:7u] + x^uW^o[:6u] + x^{2u}W^o[:5u] + x^{3u}W^o[:4u] + x^{4u}W^o[:3u] \\
&+ x^{5u}W^o[:2u] + x^{6u}W^o[:u] + x^uW^o[:7u] + x^{2u}W^o[:6u] \\
&+ x^{3u}W^o[:5u] + x^{4u}W^o[:4u] + x^{5u}W^o[:3u] + x^{6u}W^o[:2u] \\
&+ x^{7u}W^o[:u] + x^{8u}W^o[:6u] + x^{10u}W^o[:4u] + x^{12u}W^o[:2u] \\
&+ (x^{7u} + x^{8u})R^o.
\end{aligned}$$

Proof. To prove Lemma 2.2 we first construct an automaton for each sequence concerned, and then transform the conditions on infinitely many k 's into finitely many conditions on the states of the automaton. In the following, we will prove that for $T = M_{0,0}^o$, for all integer $k \geq 2$ and $u = 2^{2k}$,

$$\begin{aligned}
T[:14u] &= T[:7u] + x^uT[:6u] + x^{2u}T[:5u] + x^{3u}T[:4u] + x^{4u}T[:3u] + x^{5u}T[:2u] \\
&+ x^{6u}T[:u] + x^uT[:7u] + x^{2u}T[:6u] + x^{3u}T[:5u] + x^{4u}T[:4u] \\
&+ x^{5u}T[:3u] + x^{6u}T[:2u] + x^{7u}T[:u] + x^{8u}T[:6u] + x^{10u}T[:4u] \\
&+ x^{12u}T[:2u].
\end{aligned}$$

The proof of the other 15 cases are similar. We break down the above identity into 7 parts:

$$\begin{aligned}
0 &= x^{7u}T[:u] + x^{6u}T[u:2u] + x^{5u}T[2u:3u] + x^{4u}T[3u:4u] \\
&+ x^{3u}T[4u:5u] + x^{2u}T[5u:6u] + x^uT[6u:7u] + T[7u:8u], \\
0 &= x^{8u}T[:u] + T[8u:9u], \\
0 &= x^{8u}T[u:2u] + T[9u:10u], \\
0 &= x^{10u}T[:u] + x^{8u}T[2u:3u] + T[10u:11u], \\
0 &= x^{10u}T[u:2u] + x^{8u}T[3u:4u] + T[11u:12u], \\
0 &= x^{12u}T[:u] + x^{10u}T[2u:3u] + x^{8u}T[4u:5u] + T[12u:13u], \\
0 &= x^{12u}T[u:2u] + x^{10u}T[3u:4u] + x^{8u}T[5u:6u] + T[13u:14u],
\end{aligned}$$

which can be rewritten as

$$\begin{aligned}
0 &= T[[w]_2] + T[[1w]_2] + T[[10w]_2] + T[[11w]_2] + T[[100w]_2] \\
(2.7) \quad &+ T[[101w]_2] + T[[110w]_2] + T[[111w]_2],
\end{aligned}$$

$$\begin{aligned}
(2.8) \quad & 0 = T[[w]_2] + T[[1000w]_2], \\
(2.9) \quad & 0 = T[[1w]_2] + T[[1001w]_2], \\
(2.10) \quad & 0 = T[[w]_2] + T[[10w]_2] + T[[1010w]_2], \\
(2.11) \quad & 0 = T[[1w]_2] + T[[11w]_2] + T[[1011w]_2], \\
(2.12) \quad & 0 = T[[w]_2] + T[[10w]_2] + T[[100w]_2] + T[[1100w]_2], \\
(2.13) \quad & 0 = T[[1w]_2] + T[[11w]_2] + T[[101w]_2] + T[[1101w]_2],
\end{aligned}$$

for all binary word w of length $2k$ and $w \neq 0^{2k}$, and

$$\begin{aligned}
(2.14) \quad & 0 = T[[w]_2] + T[[1w]_2] + T[[10w]_2] + T[[11w]_2] + T[[100w]_2] \\
& \quad + T[[101w]_2] + T[[110w]_2] + T[[111w]_2], \\
(2.15) \quad & 0 = T[[w]_2] + T[[1000w]_2], \\
(2.16) \quad & 0 = T[[1w]_2] + T[[1001w]_2], \\
(2.17) \quad & 0 = T[[w]_2] + T[[10w]_2] + T[[1010w]_2], \\
(2.18) \quad & 0 = T[[1w]_2] + T[[11w]_2] + T[[1011w]_2], \\
(2.19) \quad & 0 = T[[w]_2] + T[[10w]_2] + T[[100w]_2] + T[[1100w]_2], \\
(2.20) \quad & 0 = T[[1w]_2] + T[[11w]_2] + T[[101w]_2] + T[[1101w]_2],
\end{aligned}$$

for $w = 0^{2k}$.

First we calculate an 2-automaton that generates T from its minimal polynomial and its first terms. This automaton has 289 states; its transition function and output function can be found in the annex. Let $A(s, w)$ denote the state reached after reading w from right to left starting from the state s , and τ the output function. Define

$$E_{2k} = \{A(i, w) : |w| = 2k, w \neq 0^{2k}\}.$$

Identities (2.7) through (2.20) can be written as

$$\begin{aligned}
(2.21) \quad & 0 = \tau(A(s, \epsilon)) + \tau(A(s, 1)) + \tau(A(s, 10)) + \tau(A(s, 11)) + \tau(A(s, 100)) \\
& \quad + \tau(A(s, 101)) + \tau(A(s, 110)) + \tau(A(s, 111)), \\
(2.22) \quad & 0 = \tau(A(s, \epsilon)) + \tau(A(s, 1000)), \\
(2.23) \quad & 0 = \tau(A(s, 1)) + \tau(A(s, 1001)), \\
(2.24) \quad & 0 = \tau(A(s, \epsilon)) + \tau(A(s, 10)) + \tau(A(s, 1010)), \\
(2.25) \quad & 0 = \tau(A(s, 1)) + \tau(A(s, 11)) + \tau(A(s, 1011)), \\
(2.26) \quad & 0 = \tau(A(s, \epsilon)) + \tau(A(s, 10)) + \tau(A(s, 100)) + \tau(A(s, 1100)), \\
(2.27) \quad & 0 = \tau(A(s, 1)) + \tau(A(s, 11)) + \tau(A(s, 101)) + \tau(A(s, 1101)),
\end{aligned}$$

for all $s \in E_{2k}$, and

$$\begin{aligned}
(2.28) \quad & 0 = \tau(A(s, \epsilon)) + \tau(A(s, 1)) + \tau(A(s, 10)) + \tau(A(s, 11)) + \tau(A(s, 100)) \\
& \quad + \tau(A(s, 101)) + \tau(A(s, 110)) + \tau(A(s, 111)), \\
(2.29) \quad & 0 = \tau(A(s, \epsilon)) + \tau(A(s, 1000)), \\
(2.30) \quad & 0 = \tau(A(s, 1)) + \tau(A(s, 1001)), \\
(2.31) \quad & 0 = \tau(A(s, \epsilon)) + \tau(A(s, 10)) + \tau(A(s, 1010)), \\
(2.32) \quad & 0 = \tau(A(s, 1)) + \tau(A(s, 11)) + \tau(A(s, 101)) + \tau(A(s, 1101)),
\end{aligned}$$

$$(2.33) \quad 0 = \tau(A(s, \epsilon)) + \tau(A(s, 10)) + \tau(A(s, 100)) + \tau(A(s, 1100)),$$

$$(2.34) \quad 0 = \tau(A(s, 1)) + \tau(A(s, 11)) + \tau(A(s, 101)) + \tau(A(s, 1101)),$$

for $s = A(i, 0^{2k})$. We find that $(A(i, 0^{16}), E_{16}) = (A(i, 0^{14}), E_{14})$, so that we only have to verify that identities (2.21) through (2.34) hold for $2 \leq k \leq 8$, which turns out to be true. $\square$

In the following lemma, we express M_{2k} , M_{2k+1} , W_{2k} , and W_{2k+1} in terms of M^e , M^o , W^e , and W^o .

Lemma 2.3. *For all integer $k \geq 2$, and $u = 2^{2k-1}$,*

$$\begin{aligned} M_{2k} &= M^e[:7u] + x^u M^e[:6u] + x^{2u} M^e[:5u] + x^{3u} M^e[:4u] + x^{4u} M^e[:3u] \\ &\quad + x^{5u} M^e[:2u] + x^{6u} M^e[:u] + x^{7u} R^e, \\ W_{2k} &= W^e[:7u] + x^u W^e[:6u] + x^{2u} W^e[:5u] + x^{3u} W^e[:4u] + x^{4u} W^e[:3u] \\ &\quad + x^{5u} W^e[:2u] + x^{6u} W^e[:u] + x^{7u} R^e. \end{aligned}$$

For all integer $k \geq 2$, and $u = 2^{2k}$,

$$\begin{aligned} M_{2k+1} &= M^o[:7u] + x^u M^o[:6u] + x^{2u} M^o[:5u] + x^{3u} M^o[:4u] + x^{4u} M^o[:3u] \\ &\quad + x^{5u} M^o[:2u] + x^{6u} M^o[:u] + x^{7u} R^o, \\ W_{2k+1} &= W^o[:7u] + x^u W^o[:6u] + x^{2u} W^o[:5u] + x^{3u} W^o[:4u] + x^{4u} W^o[:3u] \\ &\quad + x^{5u} W^o[:2u] + x^{6u} W^o[:u] + x^{7u} R^o. \end{aligned}$$

Proof. We call the four identities in Lemma 2.3 also by the name M_{2k} , W_{2k} , M_{2k+1} , and W_{2k+1} . For $n = 2$, the identities can be verified directly. For $n \geq 2$, we claim that

$$\begin{aligned} M_{2k} \wedge W_{2k} &\Rightarrow M_{2k+1} \wedge W_{2k+1}, \\ M_{2k+1} \wedge W_{2k+1} &\Rightarrow M_{2k+2} \wedge W_{2k+2}. \end{aligned}$$

We give the proof of

$$M_{2k} \wedge W_{2k} \Rightarrow M_{2k+1},$$

the other ones being similar. Set $u = 2^{2k}$ and $v = 2^{2k-1}$. By definition and induction hypothesis, the left side of identity M_{2k+1} is equal to

$$\begin{aligned} W_{2k} M_{2k} &= (W^e[:7v] + x^v W^e[:6v] + x^{2v} W^e[:5v] + x^{3v} W^e[:4v] \\ &\quad + x^{4v} W^e[:3v] + x^{5v} W^e[:2v] + x^{6v} W^e[:v] + x^{7v} R^e) \times (M^e[:7v] \\ &\quad + x^v M^e[:6v] + x^{2v} M^e[:5v] + x^{3v} M^e[:4v] + x^{4v} M^e[:3v] \\ &\quad + x^{5v} M^e[:2v] + x^{6v} M^e[:v] + x^{7v} R^e); \end{aligned} \quad (2.35)$$

Call this expression *lhs*. Note that both sides of identity M_{2k+1} have the same term of highest degree $x^{14v} R^o$. Therefore we only need to prove that their difference is $O(x^{14v})$. Using Lemma 2.1 it can be seen that the right side of identity M_{2k+1} is congruent, modulo x^{14v} , to

$$\begin{aligned} &W^e[:14v] M^e[:14v] + x^{2v} W^e[:12v] M^e[:12v] + x^{4v} W^e[:10v] M^e[:10v] \\ &\quad + x^{6v} W^e[:8v] M^e[:8v] + x^{8v} W^e[:6v] M^e[:6v] + x^{10v} W^e[:4v] M^e[:4v] \\ &\quad + x^{12v} W^e[:2v] M^e[:2v]. \end{aligned}$$

For all $n \leq 14$, replace the occurrences of $W^e[:n \cdot v]$ and $M^e[:n \cdot v]$ in the above expression by the reduction modulo $x^{n \cdot v}$ of the right side of the corresponding identity in Lemma 2.2 and get a new expression, which we call *rhs*. Define

$$\begin{aligned} X &:= x^v, \\ a_n &:= W^e[n \cdot v : (n+1) \cdot v] / X^n, \\ b_n &:= M^e[n \cdot v : (n+1) \cdot v] / X^n, \\ c &:= R^e. \end{aligned}$$

Using the notation introduced above, we can represent the expressions *lhs* (2.35) and *rhs* as polynomials in $\mathbb{F}_2[a_1, \dots, a_{14}, b_1, \dots, b_{14}, c][X]$. Note that it is not a problem that a_j commutes with b_k while W^e does not commute with M^e , because in the expressions concerned, the products of W^e -terms and M^e -terms are always in the same order. We let the computer do the simplification and check that the difference between these two polynomials is indeed $O(X^{14})$, which completes the proof. $\square$

Proof of Theorem 1.1. We prove the theorem for $\text{CF}(a, b)$; for $\text{CF}(b, a)$, the proof is similar. By Lemma 2.3, we have For all $0 \leq j, k \leq 1$,

$$\begin{aligned} \lim_{n \rightarrow \infty} M_{2n,j,k} &= M_{j,k}^e, \\ \lim_{n \rightarrow \infty} M_{2n+1,j,k} &= M_{j,k}^o, \\ \lim_{n \rightarrow \infty} W_{2n,j,k} &= W_{j,k}^e, \\ \lim_{n \rightarrow \infty} W_{2n+1,j,k} &= W_{j,k}^o. \end{aligned}$$

Let $z = 1/x$. By the convergence theorem and identity (2.2),

$$\text{CF}(a, b) = \frac{M_{0,1}^e(x)}{M_{0,0}^e(x)}.$$

By definition, $\phi(M^e, 0, 1)$ and $\phi(M^e, 0, 0)$ are minimal polynomials of $M_{0,1}^e$ and $M_{0,0}^e$. Therefore

$$P(x, y) = \text{Res}_t (\phi(M^e, 0, 1)(x, t), y^{12} \phi(M^e, 0, 1)(x, t/y))$$

is an annihilating polynomial of $f(x) = M_{0,1}^e / M_{0,0}^e$.

Define

$$Q(x, y) = q_4(x)y^4 + q_3(x)y^3 + q_2(x)y^2 + q_1(x)y + q_0(x),$$

where

$$\begin{aligned} q_0(x) &= x^{24} + x^{19} + x^{18} + x^{17} + x^{16} + x^{14} + x^8 + x^7, \\ q_1(x) &= x^{20} + x^{19} + x^{15} + x^{12} + x^8 + x^7 + x^6 + x^5 + x^4 + x^2, \\ q_2(x) &= x^{16} + x^{14} + x^{12} + x^{10} + x^4 + 1, \\ q_3(x) &= x^{20} + x^{19} + x^{15} + x^{12} + x^8 + x^7 + x^6 + x^5 + x^4 + x^2, \\ q_4(x) &= x^{20} + x^{19} + x^{18} + x^{17} + x^{12} + x^{10} + x^8 + x^7 + x^4. \end{aligned}$$

The polynomial $Q(x, y)$ is the candidate for the minimal polynomial of $f(x)$ found by Padé-Hermite approximation. To prove that it is indeed the minimal polynomial of $f(x)$, we only need to prove that it is an irreducible factor of $P(x, y)$ of multiplicity m and $R(x, y) := P(x, y)/Q(x, y)^m$ is not an annihilating polynomial of $f(x)$. We verify the first point directly. For the second point, we find that when we truncate

$f(x)$ to order 216, and substitute it for y in $R(z, y)$, we get a series with valuation smaller than 216, which proves that $R(z, y)$ is not an annihilating polynomial of $f(x)$. Finally, $z^{12}Q(1/z, y)$ is the minimal polynomial of $\text{CF}(a, b) = f(1/z)$. $\square$

3. ANNEXE

All 16 polynomials are of the form

$$p_0(x) + p_3(x)y^3 + p_6(x)y^6 + p_9(x)y^9 + p_{12}(x)y^{12}.$$

The coefficients $p_j(x)$ and the 16 initial terms to determine the solutions uniquely are given below.

For $\phi(M^e, 0, 0)$:

$$\begin{aligned} p_0(x) &= x^{120} + x^{118} + x^{102} + x^{96} + x^{94} + x^{92} + x^{90} + x^{84} + x^{78} + x^{74} + x^{70} \\ &\quad + x^{68} + x^{64} + x^{60} + x^{52} + x^{50} + x^{48} + x^{46} + x^{44} + x^{42} + x^{32} + x^{30} \\ &\quad + x^{24}, \\ p_3(x) &= x^{108} + x^{104} + x^{100} + x^{92} + x^{86} + x^{82} + x^{78} + x^{74} + x^{70} + x^{68} + x^{66} \\ &\quad + x^{62} + x^{60} + x^{58} + x^{48} + x^{44} + x^{40} + x^{38} + x^{36} + x^{34} + x^{32} + x^{28} \\ &\quad + x^{24} + x^{22} + x^{18} + x^{16}, \\ p_6(x) &= x^{108} + x^{102} + x^{98} + x^{96} + x^{94} + x^{90} + x^{88} + x^{86} + x^{84} + x^{82} + x^{80} \\ &\quad + x^{78} + x^{76} + x^{66} + x^{64} + x^{58} + x^{54} + x^{40} + x^{38} + x^{34} + x^{32} + x^{28} \\ &\quad + x^{26} + x^{24} + x^{20} + x^{10} + x^4 + 1, \\ p_9(x) &= x^{112} + x^{104} + x^{102} + x^{92} + x^{78} + x^{76} + x^{74} + x^{66} + x^{62} + x^{60} + x^{58} \\ &\quad + x^{56} + x^{50} + x^{46} + x^{38} + x^{36} + x^{26} + x^{16} + x^{14} + x^{12} + x^{10} + x^4, \\ p_{12}(x) &= x^{112} + x^{80} + x^{16} + 1, \end{aligned}$$

and the initial terms are $[1, 0, 0, 0, 1, 0, 0, 0, 0, 0, 0, 0, 0, 1, 0]$.

For $\phi(M^e, 0, 1)$:

$$\begin{aligned} p_0(x) &= x^{129} + x^{127} + x^{125} + x^{124} + x^{121} + x^{120} + x^{117} + x^{116} + x^{115} + x^{114} \\ &\quad + x^{111} + x^{109} + x^{108} + x^{107} + x^{106} + x^{105} + x^{100} + x^{99} + x^{98} + x^{97} \\ &\quad + x^{96} + x^{95} + x^{94} + x^{92} + x^{89} + x^{84} + x^{81} + x^{79} + x^{78} + x^{77} + x^{72} \\ &\quad + x^{71} + x^{68} + x^{67} + x^{65} + x^{64} + x^{59} + x^{58} + x^{57} + x^{54} + x^{53} + x^{50} \\ &\quad + x^{49} + x^{46} + x^{45} + x^{44} + x^{43} + x^{42} + x^{41} + x^{40} + x^{37} + x^{36} + x^{33}, \\ p_3(x) &= x^{121} + x^{119} + x^{117} + x^{116} + x^{113} + x^{111} + x^{108} + x^{104} + x^{102} + x^{101} \\ &\quad + x^{99} + x^{98} + x^{97} + x^{95} + x^{92} + x^{88} + x^{86} + x^{85} + x^{83} + x^{82} + x^{77} \\ &\quad + x^{73} + x^{72} + x^{71} + x^{70} + x^{69} + x^{68} + x^{67} + x^{66} + x^{65} + x^{64} + x^{63} \\ &\quad + x^{62} + x^{61} + x^{60} + x^{59} + x^{58} + x^{56} + x^{54} + x^{51} + x^{50} + x^{49} + x^{48} \\ &\quad + x^{47} + x^{46} + x^{44} + x^{43} + x^{42} + x^{41} + x^{39} + x^{37} + x^{36} + x^{33} + x^{31} \\ &\quad + x^{28} + x^{25} + x^{24} + x^{23} + x^{22} + x^{20} + x^{19} + x^{18}, \\ p_6(x) &= x^{117} + x^{116} + x^{115} + x^{114} + x^{107} + x^{106} + x^{101} + x^{100} + x^{97} + x^{96} \\ &\quad + x^{95} + x^{94} + x^{87} + x^{86} + x^{85} + x^{84} + x^{83} + x^{82} + x^{79} + x^{78} + x^{77} \end{aligned}$$

$$\begin{aligned}
& + x^{76} + x^{71} + x^{70} + x^{69} + x^{68} + x^{67} + x^{66} + x^{65} + x^{64} + x^{59} + x^{58} \\
& + x^{53} + x^{52} + x^{49} + x^{48} + x^{47} + x^{46} + x^{41} + x^{40} + x^{37} + x^{36} + x^{35} \\
& + x^{34} + x^{33} + x^{32} + x^{31} + x^{30} + x^{27} + x^{26} + x^{25} + x^{24} + x^{19} + x^{18} \\
& + x^{13} + x^{12}, \\
p_9(x) &= x^{113} + x^{108} + x^{107} + x^{106} + x^{103} + x^{98} + x^{97} + x^{96} + x^{95} + x^{93} + x^{90} \\
& + x^{89} + x^{86} + x^{85} + x^{82} + x^{81} + x^{79} + x^{78} + x^{75} + x^{70} + x^{69} + x^{68} \\
& + x^{63} + x^{61} + x^{59} + x^{58} + x^{51} + x^{47} + x^{46} + x^{45} + x^{44} + x^{43} + x^{42} \\
& + x^{41} + x^{40} + x^{38} + x^{36} + x^{35} + x^{32} + x^{31} + x^{28} + x^{24} + x^{23} + x^{22} \\
& + x^{21} + x^{17} + x^{16} + x^{14} + x^{12} + x^8 + x^7 + x^6, \\
p_{12}(x) &= x^{109} + x^{108} + x^{105} + x^{104} + x^{101} + x^{100} + x^{97} + x^{96} + x^{93} + x^{92} \\
& + x^{89} + x^{88} + x^{85} + x^{84} + x^{81} + x^{80} + x^{13} + x^{12} + x^9 + x^8 + x^5 + x^4 \\
& + x + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 1, 1, 1, 1, 0, 0, 0, 1, 0, 0, 0]$.

For $\phi(M^e, 1, 0)$:

$$\begin{aligned}
p_0(x) &= x^{129} + x^{127} + x^{125} + x^{124} + x^{121} + x^{120} + x^{117} + x^{116} + x^{115} + x^{114} \\
& + x^{111} + x^{109} + x^{108} + x^{107} + x^{106} + x^{105} + x^{100} + x^{99} + x^{98} + x^{97} \\
& + x^{96} + x^{95} + x^{94} + x^{92} + x^{89} + x^{84} + x^{81} + x^{79} + x^{78} + x^{77} + x^{72} \\
& + x^{71} + x^{68} + x^{67} + x^{65} + x^{64} + x^{59} + x^{58} + x^{57} + x^{54} + x^{53} + x^{50} \\
& + x^{49} + x^{46} + x^{45} + x^{44} + x^{43} + x^{42} + x^{41} + x^{40} + x^{37} + x^{36} + x^{33}, \\
p_3(x) &= x^{121} + x^{119} + x^{117} + x^{116} + x^{113} + x^{111} + x^{108} + x^{104} + x^{102} + x^{101} \\
& + x^{99} + x^{98} + x^{97} + x^{95} + x^{92} + x^{88} + x^{86} + x^{85} + x^{83} + x^{82} + x^{77} \\
& + x^{73} + x^{72} + x^{71} + x^{70} + x^{69} + x^{68} + x^{67} + x^{66} + x^{65} + x^{64} + x^{63} \\
& + x^{62} + x^{61} + x^{60} + x^{59} + x^{58} + x^{56} + x^{54} + x^{51} + x^{50} + x^{49} + x^{48} \\
& + x^{47} + x^{46} + x^{44} + x^{43} + x^{42} + x^{41} + x^{39} + x^{37} + x^{36} + x^{33} + x^{31} \\
& + x^{28} + x^{25} + x^{24} + x^{23} + x^{22} + x^{20} + x^{19} + x^{18}, \\
p_6(x) &= x^{117} + x^{116} + x^{115} + x^{114} + x^{107} + x^{106} + x^{101} + x^{100} + x^{97} + x^{96} \\
& + x^{95} + x^{94} + x^{87} + x^{86} + x^{85} + x^{84} + x^{83} + x^{82} + x^{79} + x^{78} + x^{77} \\
& + x^{76} + x^{71} + x^{70} + x^{69} + x^{68} + x^{67} + x^{66} + x^{65} + x^{64} + x^{59} + x^{58} \\
& + x^{53} + x^{52} + x^{49} + x^{48} + x^{47} + x^{46} + x^{41} + x^{40} + x^{37} + x^{36} + x^{35} \\
& + x^{34} + x^{33} + x^{32} + x^{31} + x^{30} + x^{27} + x^{26} + x^{25} + x^{24} + x^{19} + x^{18} \\
& + x^{13} + x^{12}, \\
p_9(x) &= x^{113} + x^{108} + x^{107} + x^{106} + x^{103} + x^{98} + x^{97} + x^{96} + x^{95} + x^{93} + x^{90} \\
& + x^{89} + x^{86} + x^{85} + x^{82} + x^{81} + x^{79} + x^{78} + x^{75} + x^{70} + x^{69} + x^{68} \\
& + x^{63} + x^{61} + x^{59} + x^{58} + x^{51} + x^{47} + x^{46} + x^{45} + x^{44} + x^{43} + x^{42} \\
& + x^{41} + x^{40} + x^{38} + x^{36} + x^{35} + x^{32} + x^{31} + x^{28} + x^{24} + x^{23} + x^{22}
\end{aligned}$$

$$\begin{aligned}
& + x^{21} + x^{17} + x^{16} + x^{14} + x^{12} + x^8 + x^7 + x^6, \\
p_{12}(x) = & x^{109} + x^{108} + x^{105} + x^{104} + x^{101} + x^{100} + x^{97} + x^{96} + x^{93} + x^{92} \\
& + x^{89} + x^{88} + x^{85} + x^{84} + x^{81} + x^{80} + x^{13} + x^{12} + x^9 + x^8 + x^5 + x^4 \\
& + x + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 1, 1, 1, 1, 0, 0, 0, 1, 0, 0, 0]$.

For $\phi(M^e, 1, 1)$:

$$\begin{aligned}
p_0(x) = & x^{138} + x^{136} + x^{130} + x^{122} + x^{120} + x^{110} + x^{94} + x^{90} + x^{88} + x^{86} \\
& + x^{84} + x^{82} + x^{72} + x^{68} + x^{66} + x^{62} + x^{58} + x^{56} + x^{42}, \\
p_3(x) = & x^{114} + x^{104} + x^{98} + x^{96} + x^{94} + x^{90} + x^{88} + x^{86} + x^{82} + x^{80} + x^{78} \\
& + x^{74} + x^{68} + x^{58} + x^{56} + x^{54} + x^{52} + x^{46} + x^{42} + x^{40} + x^{30} + x^{26} \\
& + x^{22} + x^{20} + x^{14} + x^{12}, \\
p_6(x) = & x^{114} + x^{106} + x^{98} + x^{94} + x^{88} + x^{84} + x^{78} + x^{72} + x^{70} + x^{68} + x^{66} \\
& + x^{64} + x^{62} + x^{60} + x^{56} + x^{54} + x^{50} + x^{48} + x^{44} + x^{38} + x^{36} + x^{34} \\
& + x^{32} + x^{28} + x^{26} + x^{22} + x^{18} + x^{16} + x^{12} + x^{10} + x^6 + x^4 + x^2 + 1, \\
p_9(x) = & x^{106} + x^{104} + x^{96} + x^{94} + x^{88} + x^{84} + x^{80} + x^{76} + x^{68} + x^{66} + x^{56} \\
& + x^{44} + x^{42} + x^{40} + x^{38} + x^{36} + x^{34} + x^{30} + x^{26} + x^{22} + x^{20} + x^{14} \\
& + x^{12} + x^{10} + x^6 + x^4, \\
p_{12}(x) = & x^{106} + x^{104} + x^{98} + x^{96} + x^{90} + x^{88} + x^{82} + x^{80} + x^{10} + x^8 + x^2 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 1, 0, 1, 0, 0, 0]$.

For $\phi(W^e, 0, 0)$:

$$\begin{aligned}
p_0(x) = & x^{138} + x^{136} + x^{130} + x^{118} + x^{116} + x^{110} + x^{102} + x^{100} + x^{96} + x^{94} \\
& + x^{90} + x^{86} + x^{84} + x^{78} + x^{76} + x^{74} + x^{70} + x^{68} + x^{66} + x^{64} + x^{60} \\
& + x^{58} + x^{54} + x^{48} + x^{46} + x^{44} + x^{42} + x^{40} + x^{38} + x^{34} + x^{30} + x^{26} \\
& + x^{24}, \\
p_3(x) = & x^{114} + x^{104} + x^{96} + x^{90} + x^{74} + x^{64} + x^{58} + x^{50} + x^{26} + x^{16}, \\
p_6(x) = & x^{114} + x^{106} + x^{90} + x^{88} + x^{78} + x^{76} + x^{72} + x^{68} + x^{66} + x^{58} + x^{56} \\
& + x^{52} + x^{50} + x^{46} + x^{44} + x^{34} + x^{32} + x^{30} + x^{28} + x^{26} + x^{22} + x^{18} \\
& + x^{14} + x^8 + x^6 + x^4 + x^2 + 1, \\
p_9(x) = & x^{106} + x^{104} + x^{96} + x^{94} + x^{88} + x^{84} + x^{80} + x^{76} + x^{68} + x^{66} + x^{56} \\
& + x^{44} + x^{42} + x^{40} + x^{38} + x^{36} + x^{34} + x^{30} + x^{26} + x^{22} + x^{20} + x^{14} \\
& + x^{12} + x^{10} + x^6 + x^4, \\
p_{12}(x) = & x^{106} + x^{104} + x^{98} + x^{96} + x^{90} + x^{88} + x^{82} + x^{80} + x^{10} + x^8 + x^2 + 1,
\end{aligned}$$

and the initial terms are $[1, 0, 0, 0, 0, 0, 0, 0, 1, 0, 1, 0, 1, 0, 0, 0]$.

For $\phi(W^e, 0, 1)$:

$$\begin{aligned}
p_0(x) = & x^{129} + x^{127} + x^{121} + x^{117} + x^{116} + x^{115} + x^{114} + x^{108} + x^{107} + x^{104} \\
& + x^{101} + x^{100} + x^{99} + x^{98} + x^{92} + x^{91} + x^{87} + x^{80} + x^{79} + x^{78} + x^{76}
\end{aligned}$$

$$\begin{aligned}
& + x^{74} + x^{73} + x^{72} + x^{66} + x^{64} + x^{63} + x^{62} + x^{60} + x^{59} + x^{57} + x^{56} \\
& + x^{55} + x^{53} + x^{52} + x^{51} + x^{48} + x^{47} + x^{46} + x^{45} + x^{44} + x^{38} + x^{36} \\
& + x^{34} + x^{33}, \\
p_3(x) &= x^{121} + x^{119} + x^{117} + x^{116} + x^{113} + x^{111} + x^{108} + x^{104} + x^{102} + x^{101} \\
& + x^{99} + x^{98} + x^{97} + x^{95} + x^{92} + x^{88} + x^{86} + x^{85} + x^{83} + x^{82} + x^{77} \\
& + x^{73} + x^{72} + x^{71} + x^{70} + x^{69} + x^{68} + x^{67} + x^{66} + x^{65} + x^{64} + x^{63} \\
& + x^{62} + x^{61} + x^{60} + x^{59} + x^{58} + x^{56} + x^{54} + x^{51} + x^{50} + x^{49} + x^{48} \\
& + x^{47} + x^{46} + x^{44} + x^{43} + x^{42} + x^{41} + x^{39} + x^{37} + x^{36} + x^{33} + x^{31} \\
& + x^{28} + x^{25} + x^{24} + x^{23} + x^{22} + x^{20} + x^{19} + x^{18}, \\
p_6(x) &= x^{117} + x^{116} + x^{115} + x^{114} + x^{107} + x^{106} + x^{101} + x^{100} + x^{97} + x^{96} \\
& + x^{95} + x^{94} + x^{87} + x^{86} + x^{85} + x^{84} + x^{83} + x^{82} + x^{79} + x^{78} + x^{77} \\
& + x^{76} + x^{71} + x^{70} + x^{69} + x^{68} + x^{67} + x^{66} + x^{65} + x^{64} + x^{59} + x^{58} \\
& + x^{53} + x^{52} + x^{49} + x^{48} + x^{47} + x^{46} + x^{41} + x^{40} + x^{37} + x^{36} + x^{35} \\
& + x^{34} + x^{33} + x^{32} + x^{31} + x^{30} + x^{27} + x^{26} + x^{25} + x^{24} + x^{19} + x^{18} \\
& + x^{13} + x^{12}, \\
p_9(x) &= x^{113} + x^{108} + x^{107} + x^{106} + x^{103} + x^{98} + x^{97} + x^{96} + x^{95} + x^{93} + x^{90} \\
& + x^{89} + x^{86} + x^{85} + x^{82} + x^{81} + x^{79} + x^{78} + x^{75} + x^{70} + x^{69} + x^{68} \\
& + x^{63} + x^{61} + x^{59} + x^{58} + x^{51} + x^{47} + x^{46} + x^{45} + x^{44} + x^{43} + x^{42} \\
& + x^{41} + x^{40} + x^{38} + x^{36} + x^{35} + x^{32} + x^{31} + x^{28} + x^{24} + x^{23} + x^{22} \\
& + x^{21} + x^{17} + x^{16} + x^{14} + x^{12} + x^8 + x^7 + x^6, \\
p_{12}(x) &= x^{109} + x^{108} + x^{105} + x^{104} + x^{101} + x^{100} + x^{97} + x^{96} + x^{93} + x^{92} \\
& + x^{89} + x^{88} + x^{85} + x^{84} + x^{81} + x^{80} + x^{13} + x^{12} + x^9 + x^8 + x^5 + x^4 \\
& + x + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 1, 0, 1, 0, 1, 0, 1, 1, 1, 1, 0, 0, 0, 0]$.

For $\phi(W^e, 1, 0)$:

$$\begin{aligned}
p_0(x) &= x^{129} + x^{127} + x^{121} + x^{117} + x^{116} + x^{115} + x^{114} + x^{108} + x^{107} + x^{104} \\
& + x^{101} + x^{100} + x^{99} + x^{98} + x^{92} + x^{91} + x^{87} + x^{80} + x^{79} + x^{78} + x^{76} \\
& + x^{74} + x^{73} + x^{72} + x^{66} + x^{64} + x^{63} + x^{62} + x^{60} + x^{59} + x^{57} + x^{56} \\
& + x^{55} + x^{53} + x^{52} + x^{51} + x^{48} + x^{47} + x^{46} + x^{45} + x^{44} + x^{38} + x^{36} \\
& + x^{34} + x^{33}, \\
p_3(x) &= x^{121} + x^{119} + x^{117} + x^{116} + x^{113} + x^{111} + x^{108} + x^{104} + x^{102} + x^{101} \\
& + x^{99} + x^{98} + x^{97} + x^{95} + x^{92} + x^{88} + x^{86} + x^{85} + x^{83} + x^{82} + x^{77} \\
& + x^{73} + x^{72} + x^{71} + x^{70} + x^{69} + x^{68} + x^{67} + x^{66} + x^{65} + x^{64} + x^{63} \\
& + x^{62} + x^{61} + x^{60} + x^{59} + x^{58} + x^{56} + x^{54} + x^{51} + x^{50} + x^{49} + x^{48} \\
& + x^{47} + x^{46} + x^{44} + x^{43} + x^{42} + x^{41} + x^{39} + x^{37} + x^{36} + x^{33} + x^{31}
\end{aligned}$$

$$\begin{aligned}
& + x^{28} + x^{25} + x^{24} + x^{23} + x^{22} + x^{20} + x^{19} + x^{18}, \\
p_6(x) = & x^{117} + x^{116} + x^{115} + x^{114} + x^{107} + x^{106} + x^{101} + x^{100} + x^{97} + x^{96} \\
& + x^{95} + x^{94} + x^{87} + x^{86} + x^{85} + x^{84} + x^{83} + x^{82} + x^{79} + x^{78} + x^{77} \\
& + x^{76} + x^{71} + x^{70} + x^{69} + x^{68} + x^{67} + x^{66} + x^{65} + x^{64} + x^{59} + x^{58} \\
& + x^{53} + x^{52} + x^{49} + x^{48} + x^{47} + x^{46} + x^{41} + x^{40} + x^{37} + x^{36} + x^{35} \\
& + x^{34} + x^{33} + x^{32} + x^{31} + x^{30} + x^{27} + x^{26} + x^{25} + x^{24} + x^{19} + x^{18} \\
& + x^{13} + x^{12}, \\
p_9(x) = & x^{113} + x^{108} + x^{107} + x^{106} + x^{103} + x^{98} + x^{97} + x^{96} + x^{95} + x^{93} + x^{90} \\
& + x^{89} + x^{86} + x^{85} + x^{82} + x^{81} + x^{79} + x^{78} + x^{75} + x^{70} + x^{69} + x^{68} \\
& + x^{63} + x^{61} + x^{59} + x^{58} + x^{51} + x^{47} + x^{46} + x^{45} + x^{44} + x^{43} + x^{42} \\
& + x^{41} + x^{40} + x^{38} + x^{36} + x^{35} + x^{32} + x^{31} + x^{28} + x^{24} + x^{23} + x^{22} \\
& + x^{21} + x^{17} + x^{16} + x^{14} + x^{12} + x^8 + x^7 + x^6, \\
p_{12}(x) = & x^{109} + x^{108} + x^{105} + x^{104} + x^{101} + x^{100} + x^{97} + x^{96} + x^{93} + x^{92} \\
& + x^{89} + x^{88} + x^{85} + x^{84} + x^{81} + x^{80} + x^{13} + x^{12} + x^9 + x^8 + x^5 + x^4 \\
& + x + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 1, 0, 1, 0, 1, 0, 1, 1, 1, 1, 0, 0, 0, 0]$.

For $\phi(W^e, 1, 1)$:

$$\begin{aligned}
p_0(x) = & x^{120} + x^{118} + x^{112} + x^{100} + x^{98} + x^{96} + x^{88} + x^{86} + x^{84} + x^{82} + x^{78} \\
& + x^{70} + x^{68} + x^{66} + x^{64} + x^{62} + x^{60} + x^{56} + x^{42}, \\
p_3(x) = & x^{108} + x^{102} + x^{100} + x^{96} + x^{92} + x^{88} + x^{84} + x^{72} + x^{70} + x^{68} + x^{64} \\
& + x^{60} + x^{54} + x^{52} + x^{48} + x^{46} + x^{44} + x^{38} + x^{30} + x^{24} + x^{20} + x^{12}, \\
p_6(x) = & x^{108} + x^{104} + x^{98} + x^{96} + x^{94} + x^{92} + x^{90} + x^{76} + x^{74} + x^{72} + x^{64} \\
& + x^{62} + x^{60} + x^{52} + x^{46} + x^{44} + x^{40} + x^{38} + x^{30} + x^{28} + x^{26} + x^{24} \\
& + x^{22} + x^{20} + x^{18} + x^{12} + x^{10} + x^8 + x^4 + 1, \\
p_9(x) = & x^{112} + x^{104} + x^{102} + x^{92} + x^{78} + x^{76} + x^{74} + x^{66} + x^{62} + x^{60} + x^{58} \\
& + x^{56} + x^{50} + x^{46} + x^{38} + x^{36} + x^{26} + x^{16} + x^{14} + x^{12} + x^{10} + x^4, \\
p_{12}(x) = & x^{112} + x^{80} + x^{16} + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 1, 0, 0, 0, 1, 0, 0, 0, 0, 0, 1, 0]$.

For $\phi(M^o, 0, 0)$:

$$\begin{aligned}
p_0(x) = & x^{129} + x^{127} + x^{125} + x^{124} + x^{120} + x^{116} + x^{115} + x^{114} + x^{112} + x^{111} \\
& + x^{110} + x^{107} + x^{106} + x^{104} + x^{103} + x^{101} + x^{99} + x^{98} + x^{97} + x^{95} \\
& + x^{94} + x^{93} + x^{92} + x^{91} + x^{89} + x^{88} + x^{86} + x^{85} + x^{84} + x^{83} + x^{81} \\
& + x^{80} + x^{79} + x^{78} + x^{77} + x^{72} + x^{71} + x^{69} + x^{68} + x^{65} + x^{60} + x^{59} \\
& + x^{58} + x^{55} + x^{53} + x^{52} + x^{51} + x^{50} + x^{48} + x^{45} + x^{43} + x^{42} + x^{41} \\
& + x^{37} + x^{35} + x^{33} + x^{30} + x^{25} + x^{24},
\end{aligned}$$

$$\begin{aligned}
p_3(x) = & x^{125} + x^{124} + x^{121} + x^{115} + x^{114} + x^{113} + x^{108} + x^{106} + x^{103} + x^{101} \\
& + x^{100} + x^{96} + x^{93} + x^{91} + x^{87} + x^{85} + x^{84} + x^{83} + x^{82} + x^{79} + x^{78} \\
& + x^{77} + x^{74} + x^{73} + x^{71} + x^{69} + x^{66} + x^{65} + x^{62} + x^{61} + x^{59} + x^{57} \\
& + x^{56} + x^{55} + x^{52} + x^{51} + x^{48} + x^{47} + x^{45} + x^{44} + x^{40} + x^{35} + x^{34} \\
& + x^{32} + x^{31} + x^{30} + x^{28} + x^{27} + x^{24} + x^{22} + x^{17} + x^{16},
\end{aligned}$$

$$\begin{aligned}
p_6(x) = & x^{125} + x^{124} + x^{121} + x^{115} + x^{114} + x^{113} + x^{108} + x^{106} + x^{105} + x^{103} \\
& + x^{101} + x^{99} + x^{98} + x^{95} + x^{94} + x^{92} + x^{91} + x^{90} + x^{86} + x^{85} + x^{84} \\
& + x^{83} + x^{81} + x^{80} + x^{79} + x^{78} + x^{77} + x^{76} + x^{75} + x^{72} + x^{69} + x^{68} \\
& + x^{67} + x^{66} + x^{65} + x^{64} + x^{63} + x^{61} + x^{60} + x^{57} + x^{56} + x^{55} + x^{54} \\
& + x^{50} + x^{48} + x^{46} + x^{45} + x^{43} + x^{40} + x^{37} + x^{32} + x^{31} + x^{30} + x^{29} \\
& + x^{27} + x^{25} + x^{23} + x^{22} + x^{21} + x^{19} + x^{16} + x^{14} + x^{13} + x^{12} + x^5 \\
& + x^4 + x + 1,
\end{aligned}$$

$$\begin{aligned}
p_9(x) = & x^{125} + x^{124} + x^{117} + x^{116} + x^{105} + x^{104} + x^{101} + x^{100} + x^{93} + x^{92} \\
& + x^{89} + x^{88} + x^{29} + x^{28} + x^{21} + x^{20} + x^{13} + x^{12} + x^9 + x^8,
\end{aligned}$$

$$\begin{aligned}
p_{12}(x) = & x^{125} + x^{124} + x^{117} + x^{116} + x^{109} + x^{108} + x^{97} + x^{96} + x^{85} + x^{84} \\
& + x^{81} + x^{80} + x^{29} + x^{28} + x^{21} + x^{20} + x^5 + x^4 + x + 1,
\end{aligned}$$

and the initial terms are $[1, 0, 0, 0, 1, 0, 0, 1, 0, 0, 1, 0, 0, 0, 1]$.

For $\phi(M^o, 0, 1)$:

$$\begin{aligned}
p_0(x) = & x^{138} + x^{136} + x^{125} + x^{124} + x^{123} + x^{117} + x^{116} + x^{114} + x^{113} + x^{110} \\
& + x^{109} + x^{108} + x^{107} + x^{105} + x^{104} + x^{103} + x^{94} + x^{93} + x^{90} + x^{84} \\
& + x^{83} + x^{82} + x^{79} + x^{78} + x^{76} + x^{75} + x^{73} + x^{72} + x^{69} + x^{68} + x^{67} \\
& + x^{65} + x^{63} + x^{61} + x^{60} + x^{58} + x^{57} + x^{53} + x^{50} + x^{45} + x^{44} + x^{43} \\
& + x^{41} + x^{38} + x^{36} + x^{34} + x^{33},
\end{aligned}$$

$$\begin{aligned}
p_3(x) = & x^{110} + x^{109} + x^{106} + x^{102} + x^{101} + x^{98} + x^{94} + x^{93} + x^{90} + x^{86} + x^{85} \\
& + x^{82} + x^{78} + x^{77} + x^{74} + x^{46} + x^{45} + x^{42} + x^{30} + x^{29} + x^{26} + x^{22} \\
& + x^{21} + x^{18},
\end{aligned}$$

$$\begin{aligned}
p_6(x) = & x^{114} + x^{104} + x^{102} + x^{100} + x^{94} + x^{84} + x^{80} + x^{78} + x^{74} + x^{72} + x^{70} \\
& + x^{66} + x^{62} + x^{60} + x^{58} + x^{50} + x^{48} + x^{44} + x^{42} + x^{40} + x^{34} + x^{30} \\
& + x^{24} + x^{16} + x^{14} + x^{12},
\end{aligned}$$

$$\begin{aligned}
p_9(x) = & x^{118} + x^{117} + x^{116} + x^{115} + x^{113} + x^{111} + x^{109} + x^{108} + x^{105} + x^{101} \\
& + x^{100} + x^{99} + x^{97} + x^{95} + x^{93} + x^{92} + x^{89} + x^{86} + x^{22} + x^{21} + x^{20} \\
& + x^{19} + x^{17} + x^{15} + x^{13} + x^{12} + x^9 + x^6,
\end{aligned}$$

$$\begin{aligned}
p_{12}(x) = & x^{122} + x^{120} + x^{118} + x^{116} + x^{106} + x^{104} + x^{102} + x^{100} + x^{98} + x^{96} \\
& + x^{82} + x^{80} + x^{26} + x^{24} + x^{22} + x^{20} + x^2 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 1, 1, 1, 1, 0, 0, 0, 0, 1, 1, 0]$.

For $\phi(M^o, 1, 0)$:

$$\begin{aligned}
p_0(x) &= x^{120} + x^{118} + x^{116} + x^{115} + x^{111} + x^{108} + x^{104} + x^{103} + x^{101} + x^{99} \\
&\quad + x^{97} + x^{95} + x^{94} + x^{91} + x^{90} + x^{89} + x^{86} + x^{85} + x^{84} + x^{82} + x^{80} \\
&\quad + x^{79} + x^{77} + x^{75} + x^{73} + x^{71} + x^{70} + x^{69} + x^{66} + x^{64} + x^{59} + x^{57} \\
&\quad + x^{56} + x^{54} + x^{52} + x^{51} + x^{50} + x^{49} + x^{45} + x^{44} + x^{43} + x^{40} + x^{37} \\
&\quad + x^{36} + x^{33}, \\
p_3(x) &= x^{116} + x^{115} + x^{114} + x^{113} + x^{111} + x^{109} + x^{107} + x^{105} + x^{103} + x^{101} \\
&\quad + x^{99} + x^{97} + x^{95} + x^{93} + x^{91} + x^{89} + x^{87} + x^{85} + x^{83} + x^{81} + x^{79} \\
&\quad + x^{77} + x^{76} + x^{74} + x^{52} + x^{51} + x^{50} + x^{49} + x^{47} + x^{45} + x^{44} + x^{42} \\
&\quad + x^{36} + x^{35} + x^{34} + x^{33} + x^{31} + x^{29} + x^{27} + x^{25} + x^{23} + x^{21} + x^{20} \\
&\quad + x^{18}, \\
p_6(x) &= x^{120} + x^{118} + x^{116} + x^{114} + x^{110} + x^{106} + x^{104} + x^{98} + x^{96} + x^{94} \\
&\quad + x^{90} + x^{88} + x^{84} + x^{80} + x^{78} + x^{76} + x^{74} + x^{72} + x^{66} + x^{64} + x^{62} \\
&\quad + x^{58} + x^{56} + x^{50} + x^{48} + x^{46} + x^{44} + x^{38} + x^{32} + x^{28} + x^{26} + x^{24} \\
&\quad + x^{22} + x^{18} + x^{16} + x^{12}, \\
p_9(x) &= x^{124} + x^{123} + x^{119} + x^{116} + x^{114} + x^{113} + x^{112} + x^{111} + x^{110} + x^{108} \\
&\quad + x^{106} + x^{105} + x^{104} + x^{103} + x^{102} + x^{100} + x^{98} + x^{97} + x^{96} + x^{95} \\
&\quad + x^{94} + x^{91} + x^{90} + x^{89} + x^{88} + x^{86} + x^{28} + x^{27} + x^{23} + x^{20} + x^{18} \\
&\quad + x^{17} + x^{16} + x^{15} + x^{14} + x^{11} + x^{10} + x^9 + x^8 + x^6, \\
p_{12}(x) &= x^{128} + x^{124} + x^{120} + x^{116} + x^{112} + x^{108} + x^{100} + x^{96} + x^{88} + x^{80} \\
&\quad + x^{32} + x^{28} + x^{24} + x^{20} + x^8 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 1, 0, 1, 0, 0, 0, 0, 1, 0, 0, 0, 0, 0]$.

For $\phi(M^o, 1, 1)$:

$$\begin{aligned}
p_0(x) &= x^{129} + x^{127} + x^{117} + x^{108} + x^{107} + x^{105} + x^{104} + x^{97} + x^{92} + x^{91} \\
&\quad + x^{88} + x^{85} + x^{84} + x^{77} + x^{76} + x^{73} + x^{72} + x^{69} + x^{68} + x^{66} + x^{65} \\
&\quad + x^{64} + x^{60} + x^{59} + x^{58} + x^{53} + x^{51} + x^{50} + x^{42}, \\
p_3(x) &= x^{125} + x^{124} + x^{121} + x^{117} + x^{116} + x^{113} + x^{109} + x^{107} + x^{105} + x^{104} \\
&\quad + x^{103} + x^{99} + x^{98} + x^{97} + x^{92} + x^{90} + x^{89} + x^{88} + x^{87} + x^{81} + x^{80} \\
&\quad + x^{79} + x^{78} + x^{77} + x^{74} + x^{72} + x^{71} + x^{69} + x^{66} + x^{65} + x^{62} + x^{61} \\
&\quad + x^{59} + x^{57} + x^{56} + x^{55} + x^{53} + x^{50} + x^{48} + x^{47} + x^{45} + x^{44} + x^{41} \\
&\quad + x^{37} + x^{36} + x^{32} + x^{31} + x^{30} + x^{29} + x^{26} + x^{25} + x^{22}, \\
p_6(x) &= x^{125} + x^{124} + x^{121} + x^{117} + x^{116} + x^{113} + x^{109} + x^{107} + x^{104} + x^{103} \\
&\quad + x^{101} + x^{99} + x^{98} + x^{92} + x^{91} + x^{90} + x^{89} + x^{88} + x^{87} + x^{85} + x^{84} \\
&\quad + x^{82} + x^{75} + x^{72} + x^{71} + x^{70} + x^{69} + x^{68} + x^{65} + x^{64} + x^{63} + x^{61} \\
&\quad + x^{60} + x^{59} + x^{58} + x^{55} + x^{54} + x^{50} + x^{48} + x^{47} + x^{45} + x^{43} + x^{40}
\end{aligned}$$

$$\begin{aligned}
& + x^{37} + x^{35} + x^{34} + x^{33} + x^{28} + x^{26} + x^{24} + x^{23} + x^{22} + x^{21} + x^{18} \\
& + x^{17} + x^{14} + x^{13} + x^{12} + x^9 + x^8 + x^5 + x^4 + x + 1, \\
p_9(x) &= x^{125} + x^{124} + x^{117} + x^{116} + x^{105} + x^{104} + x^{101} + x^{100} + x^{93} + x^{92} \\
& + x^{89} + x^{88} + x^{29} + x^{28} + x^{21} + x^{20} + x^{13} + x^{12} + x^9 + x^8, \\
p_{12}(x) &= x^{125} + x^{124} + x^{117} + x^{116} + x^{109} + x^{108} + x^{97} + x^{96} + x^{85} + x^{84} \\
& + x^{81} + x^{80} + x^{29} + x^{28} + x^{21} + x^{20} + x^5 + x^4 + x + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 0, 0, 1, 1, 0, 0, 0, 0, 0, 1, 1]$.

For $\phi(W^o, 0, 0)$:

$$\begin{aligned}
p_0(x) &= x^{129} + x^{127} + x^{125} + x^{124} + x^{120} + x^{116} + x^{115} + x^{114} + x^{112} + x^{111} \\
& + x^{110} + x^{107} + x^{106} + x^{104} + x^{103} + x^{101} + x^{99} + x^{98} + x^{97} + x^{95} \\
& + x^{94} + x^{93} + x^{92} + x^{91} + x^{89} + x^{88} + x^{86} + x^{85} + x^{84} + x^{83} + x^{81} \\
& + x^{80} + x^{79} + x^{78} + x^{77} + x^{72} + x^{71} + x^{69} + x^{68} + x^{65} + x^{60} + x^{59} \\
& + x^{58} + x^{55} + x^{53} + x^{52} + x^{51} + x^{50} + x^{48} + x^{45} + x^{43} + x^{42} + x^{41} \\
& + x^{37} + x^{35} + x^{33} + x^{30} + x^{25} + x^{24}, \\
p_3(x) &= x^{125} + x^{124} + x^{121} + x^{115} + x^{114} + x^{113} + x^{108} + x^{106} + x^{103} + x^{101} \\
& + x^{100} + x^{96} + x^{93} + x^{91} + x^{87} + x^{85} + x^{84} + x^{83} + x^{82} + x^{79} + x^{78} \\
& + x^{77} + x^{74} + x^{73} + x^{71} + x^{69} + x^{66} + x^{65} + x^{62} + x^{61} + x^{59} + x^{57} \\
& + x^{56} + x^{55} + x^{52} + x^{51} + x^{48} + x^{47} + x^{45} + x^{44} + x^{40} + x^{35} + x^{34} \\
& + x^{32} + x^{31} + x^{30} + x^{28} + x^{27} + x^{24} + x^{22} + x^{17} + x^{16}, \\
p_6(x) &= x^{125} + x^{124} + x^{121} + x^{115} + x^{114} + x^{113} + x^{108} + x^{106} + x^{105} + x^{103} \\
& + x^{101} + x^{99} + x^{98} + x^{95} + x^{94} + x^{92} + x^{91} + x^{90} + x^{86} + x^{85} + x^{84} \\
& + x^{83} + x^{81} + x^{80} + x^{79} + x^{78} + x^{77} + x^{76} + x^{75} + x^{72} + x^{69} + x^{68} \\
& + x^{67} + x^{66} + x^{65} + x^{64} + x^{63} + x^{61} + x^{60} + x^{57} + x^{56} + x^{55} + x^{54} \\
& + x^{50} + x^{48} + x^{46} + x^{45} + x^{43} + x^{40} + x^{37} + x^{32} + x^{31} + x^{30} + x^{29} \\
& + x^{27} + x^{25} + x^{23} + x^{22} + x^{21} + x^{19} + x^{16} + x^{14} + x^{13} + x^{12} + x^5 \\
& + x^4 + x + 1, \\
p_9(x) &= x^{125} + x^{124} + x^{117} + x^{116} + x^{105} + x^{104} + x^{101} + x^{100} + x^{93} + x^{92} \\
& + x^{89} + x^{88} + x^{29} + x^{28} + x^{21} + x^{20} + x^{13} + x^{12} + x^9 + x^8, \\
p_{12}(x) &= x^{125} + x^{124} + x^{117} + x^{116} + x^{109} + x^{108} + x^{97} + x^{96} + x^{85} + x^{84} \\
& + x^{81} + x^{80} + x^{29} + x^{28} + x^{21} + x^{20} + x^5 + x^4 + x + 1,
\end{aligned}$$

and the initial terms are $[1, 0, 0, 0, 1, 0, 0, 1, 0, 0, 1, 0, 0, 0, 0, 1]$.

For $\phi(W^o, 0, 1)$:

$$\begin{aligned}
p_0(x) &= x^{120} + x^{118} + x^{116} + x^{115} + x^{111} + x^{108} + x^{104} + x^{103} + x^{101} + x^{99} \\
& + x^{97} + x^{95} + x^{94} + x^{91} + x^{90} + x^{89} + x^{86} + x^{85} + x^{84} + x^{82} + x^{80} \\
& + x^{79} + x^{77} + x^{75} + x^{73} + x^{71} + x^{70} + x^{69} + x^{66} + x^{64} + x^{59} + x^{57}
\end{aligned}$$

$$\begin{aligned}
& + x^{56} + x^{54} + x^{52} + x^{51} + x^{50} + x^{49} + x^{45} + x^{44} + x^{43} + x^{40} + x^{37} \\
& + x^{36} + x^{33}, \\
p_3(x) &= x^{116} + x^{115} + x^{114} + x^{113} + x^{111} + x^{109} + x^{107} + x^{105} + x^{103} + x^{101} \\
& + x^{99} + x^{97} + x^{95} + x^{93} + x^{91} + x^{89} + x^{87} + x^{85} + x^{83} + x^{81} + x^{79} \\
& + x^{77} + x^{76} + x^{74} + x^{52} + x^{51} + x^{50} + x^{49} + x^{47} + x^{45} + x^{44} + x^{42} \\
& + x^{36} + x^{35} + x^{34} + x^{33} + x^{31} + x^{29} + x^{27} + x^{25} + x^{23} + x^{21} + x^{20} \\
& + x^{18}, \\
p_6(x) &= x^{120} + x^{118} + x^{116} + x^{114} + x^{110} + x^{106} + x^{104} + x^{98} + x^{96} + x^{94} \\
& + x^{90} + x^{88} + x^{84} + x^{80} + x^{78} + x^{76} + x^{74} + x^{72} + x^{66} + x^{64} + x^{62} \\
& + x^{58} + x^{56} + x^{50} + x^{48} + x^{46} + x^{44} + x^{38} + x^{32} + x^{28} + x^{26} + x^{24} \\
& + x^{22} + x^{18} + x^{16} + x^{12}, \\
p_9(x) &= x^{124} + x^{123} + x^{119} + x^{116} + x^{114} + x^{113} + x^{112} + x^{111} + x^{110} + x^{108} \\
& + x^{106} + x^{105} + x^{104} + x^{103} + x^{102} + x^{100} + x^{98} + x^{97} + x^{96} + x^{95} \\
& + x^{94} + x^{91} + x^{90} + x^{89} + x^{88} + x^{86} + x^{28} + x^{27} + x^{23} + x^{20} + x^{18} \\
& + x^{17} + x^{16} + x^{15} + x^{14} + x^{11} + x^{10} + x^9 + x^8 + x^6, \\
p_{12}(x) &= x^{128} + x^{124} + x^{120} + x^{116} + x^{112} + x^{108} + x^{100} + x^{96} + x^{88} + x^{80} \\
& + x^{32} + x^{28} + x^{24} + x^{20} + x^8 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 1, 0, 1, 0, 0, 0, 0, 1, 0, 0, 0, 0, 0]$.

For $\phi(W^o, 1, 0)$:

$$\begin{aligned}
p_0(x) &= x^{138} + x^{136} + x^{125} + x^{124} + x^{123} + x^{117} + x^{116} + x^{114} + x^{113} + x^{110} \\
& + x^{109} + x^{108} + x^{107} + x^{105} + x^{104} + x^{103} + x^{94} + x^{93} + x^{90} + x^{84} \\
& + x^{83} + x^{82} + x^{79} + x^{78} + x^{76} + x^{75} + x^{73} + x^{72} + x^{69} + x^{68} + x^{67} \\
& + x^{65} + x^{63} + x^{61} + x^{60} + x^{58} + x^{57} + x^{53} + x^{50} + x^{45} + x^{44} + x^{43} \\
& + x^{41} + x^{38} + x^{36} + x^{34} + x^{33}, \\
p_3(x) &= x^{110} + x^{109} + x^{106} + x^{102} + x^{101} + x^{98} + x^{94} + x^{93} + x^{90} + x^{86} + x^{85} \\
& + x^{82} + x^{78} + x^{77} + x^{74} + x^{46} + x^{45} + x^{42} + x^{30} + x^{29} + x^{26} + x^{22} \\
& + x^{21} + x^{18}, \\
p_6(x) &= x^{114} + x^{104} + x^{102} + x^{100} + x^{94} + x^{84} + x^{80} + x^{78} + x^{74} + x^{72} + x^{70} \\
& + x^{66} + x^{62} + x^{60} + x^{58} + x^{50} + x^{48} + x^{44} + x^{42} + x^{40} + x^{34} + x^{30} \\
& + x^{24} + x^{16} + x^{14} + x^{12}, \\
p_9(x) &= x^{118} + x^{117} + x^{116} + x^{115} + x^{113} + x^{111} + x^{109} + x^{108} + x^{105} + x^{101} \\
& + x^{100} + x^{99} + x^{97} + x^{95} + x^{93} + x^{92} + x^{89} + x^{86} + x^{22} + x^{21} + x^{20} \\
& + x^{19} + x^{17} + x^{15} + x^{13} + x^{12} + x^9 + x^6, \\
p_{12}(x) &= x^{122} + x^{120} + x^{118} + x^{116} + x^{106} + x^{104} + x^{102} + x^{100} + x^{98} + x^{96} \\
& + x^{82} + x^{80} + x^{26} + x^{24} + x^{22} + x^{20} + x^2 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 1, 1, 1, 1, 0, 0, 0, 0, 1, 1, 0]$.

For $\phi(W^o, 1, 1)$:

$$\begin{aligned}
p_0(x) &= x^{129} + x^{127} + x^{117} + x^{108} + x^{107} + x^{105} + x^{104} + x^{97} + x^{92} + x^{91} \\
&\quad + x^{88} + x^{85} + x^{84} + x^{77} + x^{76} + x^{73} + x^{72} + x^{69} + x^{68} + x^{66} + x^{65} \\
&\quad + x^{64} + x^{60} + x^{59} + x^{58} + x^{53} + x^{51} + x^{50} + x^{42}, \\
p_3(x) &= x^{125} + x^{124} + x^{121} + x^{117} + x^{116} + x^{113} + x^{109} + x^{107} + x^{105} + x^{104} \\
&\quad + x^{103} + x^{99} + x^{98} + x^{97} + x^{92} + x^{90} + x^{89} + x^{88} + x^{87} + x^{81} + x^{80} \\
&\quad + x^{79} + x^{78} + x^{77} + x^{74} + x^{72} + x^{71} + x^{69} + x^{66} + x^{65} + x^{62} + x^{61} \\
&\quad + x^{59} + x^{57} + x^{56} + x^{55} + x^{53} + x^{50} + x^{48} + x^{47} + x^{45} + x^{44} + x^{41} \\
&\quad + x^{37} + x^{36} + x^{32} + x^{31} + x^{30} + x^{29} + x^{26} + x^{25} + x^{22}, \\
p_6(x) &= x^{125} + x^{124} + x^{121} + x^{117} + x^{116} + x^{113} + x^{109} + x^{107} + x^{104} + x^{103} \\
&\quad + x^{101} + x^{99} + x^{98} + x^{92} + x^{91} + x^{90} + x^{89} + x^{88} + x^{87} + x^{85} + x^{84} \\
&\quad + x^{82} + x^{75} + x^{72} + x^{71} + x^{70} + x^{69} + x^{68} + x^{65} + x^{64} + x^{63} + x^{61} \\
&\quad + x^{60} + x^{59} + x^{58} + x^{55} + x^{54} + x^{50} + x^{48} + x^{47} + x^{45} + x^{43} + x^{40} \\
&\quad + x^{37} + x^{35} + x^{34} + x^{33} + x^{28} + x^{26} + x^{24} + x^{23} + x^{22} + x^{21} + x^{18} \\
&\quad + x^{17} + x^{14} + x^{13} + x^{12} + x^9 + x^8 + x^5 + x^4 + x + 1, \\
p_9(x) &= x^{125} + x^{124} + x^{117} + x^{116} + x^{105} + x^{104} + x^{101} + x^{100} + x^{93} + x^{92} \\
&\quad + x^{89} + x^{88} + x^{29} + x^{28} + x^{21} + x^{20} + x^{13} + x^{12} + x^9 + x^8, \\
p_{12}(x) &= x^{125} + x^{124} + x^{117} + x^{116} + x^{109} + x^{108} + x^{97} + x^{96} + x^{85} + x^{84} \\
&\quad + x^{81} + x^{80} + x^{29} + x^{28} + x^{21} + x^{20} + x^5 + x^4 + x + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 0, 0, 1, 1, 0, 0, 0, 0, 0, 1, 1]$.

Below is the transition function and output function of an 2-automaton that generates $T = M_{0,0}^o$:

Transition function $(n, j) \mapsto \delta(n, j)$ ($\Lambda(n) := [\delta(n, 0), \delta(n, 1)]$):

n	$\Lambda(n)$	n	$\Lambda(n)$	n	$\Lambda(n)$	n	$\Lambda(n)$	n	$\Lambda(n)$
0	[1, 2]	58	[36, 96]	116	[177, 173]	174	[206, 207]	232	[97, 40]
1	[3, 4]	59	[97, 33]	117	[178, 42]	175	[208, 209]	233	[258, 85]
2	[5, 6]	60	[41, 98]	118	[179, 36]	176	[210, 211]	234	[259, 11]
3	[7, 8]	61	[82, 99]	119	[180, 181]	177	[212, 145]	235	[172, 96]
4	[9, 2]	62	[100, 2]	120	[33, 172]	178	[213, 214]	236	[39, 39]
5	[10, 11]	63	[12, 76]	121	[182, 183]	179	[215, 158]	237	[260, 181]
6	[12, 11]	64	[12, 88]	122	[172, 184]	180	[123, 216]	238	[175, 192]
7	[13, 5]	65	[101, 102]	123	[185, 77]	181	[217, 218]	239	[12, 86]
8	[14, 15]	66	[33, 36]	124	[29, 99]	182	[219, 63]	240	[85, 11]
9	[16, 17]	67	[103, 104]	125	[8, 28]	183	[220, 221]	241	[104, 104]
10	[18, 19]	68	[105, 106]	126	[110, 11]	184	[222, 223]	242	[261, 104]
11	[20, 21]	69	[104, 106]	127	[186, 88]	185	[224, 57]	243	[195, 196]
12	[22, 23]	70	[107, 33]	128	[187, 40]	186	[225, 226]	244	[106, 106]
13	[24, 12]	71	[108, 42]	129	[108, 98]	187	[227, 228]	245	[10, 88]

14	[25, 26]	72	[82, 30]	130	[176, 184]	188	[58, 229]	246	[85, 88]
15	[27, 28]	73	[109, 28]	131	[39, 38]	189	[230, 231]	247	[260, 189]
16	[29, 30]	74	[110, 88]	132	[97, 179]	190	[232, 147]	248	[262, 36]
17	[31, 2]	75	[87, 11]	133	[101, 184]	191	[233, 234]	249	[11, 263]
18	[32, 33]	76	[111, 112]	134	[95, 39]	192	[235, 236]	250	[195, 105]
19	[34, 35]	77	[113, 114]	135	[33, 188]	193	[237, 238]	251	[261, 196]
20	[36, 37]	78	[115, 116]	136	[180, 189]	194	[239, 240]	252	[177, 95]
21	[38, 39]	79	[117, 118]	137	[190, 188]	195	[241, 242]	253	[29, 26]
22	[32, 40]	80	[119, 120]	138	[191, 178]	196	[243, 244]	254	[93, 194]
23	[41, 42]	81	[121, 122]	139	[192, 102]	197	[245, 246]	255	[264, 265]
24	[43, 20]	82	[54, 123]	140	[193, 81]	198	[247, 248]	256	[266, 267]
25	[44, 45]	83	[124, 125]	141	[88, 79]	199	[216, 249]	257	[268, 133]
26	[46, 44]	84	[126, 127]	142	[82, 92]	200	[250, 251]	258	[269, 270]
27	[47, 48]	85	[128, 129]	143	[9, 194]	201	[122, 252]	259	[271, 272]
28	[49, 50]	86	[130, 131]	144	[103, 105]	202	[253, 254]	260	[273, 274]
29	[51, 52]	87	[132, 117]	145	[195, 104]	203	[176, 102]	261	[275, 276]
30	[53, 54]	88	[133, 134]	146	[196, 105]	204	[173, 39]	262	[277, 278]
31	[55, 56]	89	[19, 135]	147	[34, 197]	205	[106, 105]	263	[23, 279]
32	[57, 58]	90	[136, 137]	148	[40, 36]	206	[25, 99]	264	[104, 196]
33	[59, 60]	91	[138, 139]	149	[94, 38]	207	[31, 194]	265	[280, 196]
34	[61, 62]	92	[140, 141]	150	[85, 76]	208	[187, 175]	266	[281, 178]
35	[63, 64]	93	[142, 143]	151	[186, 11]	209	[183, 42]	267	[36, 184]
36	[65, 43]	94	[144, 67]	152	[198, 91]	210	[176, 37]	268	[38, 95]
37	[60, 66]	95	[145, 146]	153	[86, 79]	211	[255, 95]	269	[95, 95]
38	[67, 68]	96	[147, 148]	154	[199, 30]	212	[105, 196]	270	[201, 102]
39	[68, 69]	97	[149, 65]	155	[100, 194]	213	[25, 30]	271	[107, 175]
40	[70, 71]	98	[150, 151]	156	[101, 37]	214	[8, 84]	272	[282, 197]
41	[72, 73]	99	[152, 153]	157	[94, 173]	215	[32, 175]	273	[283, 77]
42	[74, 75]	100	[154, 155]	158	[178, 98]	216	[11, 89]	274	[86, 263]
43	[38, 38]	101	[156, 157]	159	[175, 188]	217	[193, 256]	275	[195, 106]
44	[76, 77]	102	[158, 159]	160	[105, 104]	218	[86, 77]	276	[280, 104]
45	[78, 79]	103	[160, 161]	161	[103, 196]	219	[257, 12]	277	[187, 179]
46	[80, 81]	104	[162, 163]	162	[104, 105]	220	[199, 92]	278	[282, 35]
47	[82, 26]	105	[105, 105]	163	[196, 106]	221	[83, 28]	279	[40, 188]
48	[83, 84]	106	[161, 162]	164	[200, 38]	222	[174, 197]	280	[284, 243]
49	[85, 86]	107	[164, 165]	165	[201, 184]	223	[179, 188]	281	[285, 245]
50	[87, 88]	108	[166, 167]	166	[29, 92]	224	[36, 102]	282	[286, 4]
51	[76, 89]	109	[168, 169]	167	[27, 84]	225	[107, 179]	283	[287, 288]
52	[86, 89]	110	[170, 171]	168	[199, 99]	226	[183, 98]	284	[103, 106]
53	[90, 91]	111	[172, 37]	169	[109, 84]	227	[200, 95]	285	[257, 85]
54	[11, 77]	112	[173, 38]	170	[187, 33]	228	[192, 37]	286	[199, 26]
55	[25, 92]	113	[174, 35]	171	[202, 197]	229	[95, 38]	287	[172, 102]
56	[93, 2]	114	[175, 36]	172	[203, 204]	230	[80, 256]	288	[255, 173]
57	[94, 95]	115	[176, 96]	173	[69, 205]	231	[11, 79]		

Output function $n \mapsto \tau(n)$:

n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$
0	0	42	1	84	1	126	1	168	0	210	1	252	0
1	0	43	0	85	1	127	1	169	0	211	1	253	1
2	0	44	1	86	1	128	1	170	1	212	0	254	0
3	0	45	1	87	0	129	1	171	1	213	1	255	1
4	1	46	0	88	0	130	1	172	1	214	1	256	0
5	0	47	0	89	0	131	0	173	1	215	0	257	0
6	0	48	1	90	0	132	0	174	1	216	0	258	1
7	0	49	1	91	1	133	0	175	1	217	1	259	1
8	1	50	0	92	1	134	1	176	1	218	1	260	1
9	1	51	1	93	0	135	0	177	0	219	0	261	1
10	0	52	1	94	0	136	0	178	1	220	0	262	1
11	0	53	0	95	1	137	0	179	0	221	1	263	0
12	0	54	0	96	0	138	1	180	0	222	1	264	1
13	0	55	1	97	0	139	1	181	1	223	0	265	0
14	1	56	0	98	1	140	1	182	0	224	0	266	0
15	0	57	0	99	1	141	0	183	0	225	1	267	0
16	1	58	0	100	0	142	0	184	1	226	0	268	0
17	1	59	0	101	0	143	1	185	0	227	1	269	1
18	0	60	0	102	1	144	0	186	1	228	1	270	1
19	0	61	0	103	0	145	1	187	1	229	1	271	1
20	0	62	0	104	1	146	1	188	0	230	0	272	0
21	0	63	0	105	0	147	0	189	0	231	0	273	1
22	0	64	0	106	0	148	1	190	0	232	0	274	1
23	0	65	0	107	1	149	0	191	1	233	1	275	1
24	0	66	0	108	1	150	1	192	1	234	1	276	0
25	1	67	0	109	0	151	1	193	1	235	1	277	1
26	0	68	0	110	1	152	1	194	0	236	0	278	0
27	0	69	1	111	1	153	1	195	1	237	1	279	1
28	1	70	1	112	1	154	0	196	1	238	1	280	0
29	1	71	1	113	1	155	0	197	0	239	0	281	0
30	0	72	0	114	1	156	0	198	1	240	1	282	0
31	1	73	0	115	1	157	0	199	0	241	1	283	1
32	0	74	1	116	0	158	1	200	1	242	1	284	0
33	0	75	0	117	1	159	1	201	1	243	1	285	0
34	0	76	1	118	0	160	0	202	1	244	0	286	0
35	0	77	1	119	0	161	0	203	1	245	0	287	1
36	0	78	1	120	0	162	1	204	1	246	1	288	1
37	0	79	1	121	0	163	1	205	0	247	1		
38	0	80	0	122	1	164	1	206	1	248	1		
39	0	81	0	123	0	165	1	207	1	249	0		
40	1	82	0	124	1	166	1	208	1	250	1		

	41	0		83	1		125	1		167	0		209	0		251	1			
--	----	---	--	----	---	--	-----	---	--	-----	---	--	-----	---	--	-----	---	--	--	--

UNIVERSITÉ DE STRASBOURG, CNRS, IRMA UMR 7501, F-67000 STRASBOURG, FRANCE
Email address: **guoniu.han@unistra.fr**

SCHOOL OF MATHEMATICS AND STATISTICS, HUAZHONG UNIVERSITY OF SCIENCE AND TECHNOLOGY, WUHAN, PR CHINA
Email address: **huyining@hust.edu.cn**