

PROOF OF THE CONTINUED FRACTION CONJECTURE FOR

$$(a, b) = (z^3, z^4 + 1)$$

GUO-NIU HAN* AND YINING HU*

ABSTRACT. The continued fraction conjecture claims that the continued fraction defined by the (a, b) -Thue-Morse sequence is algebraic over $\mathbb{F}_2(x)$ for all pairs of distinct elements (a, b) in $\mathbb{F}_2[x] \setminus \mathbb{F}_2$. In this paper we prove the conjecture for $(a, b) = (z^3, z^4 + 1)$.

1. INTRODUCTION

Let $\mathbb{F}_2$ be the finite field of cardinality 2. Given a sequence of polynomials $\mathbf{a} = (a_0(z), a_1(z), a_2(z), \dots)$ of elements from $\mathbb{F}_2[z] \setminus \mathbb{F}_2$, we define the infinite continued fraction

$$(1.1) \quad \text{CF}(\mathbf{a}(z)) = \frac{1}{a_0(z) + \frac{1}{a_1(z) + \frac{1}{a_2(z) + \frac{1}{\ddots}}}}$$

as the limit of the finite continued fractions

$$(1.2) \quad \text{CF}_n(\mathbf{a}(z)) = \frac{1}{a_0(z) + \frac{1}{a_1(z) + \frac{1}{\ddots + \frac{1}{a_n(z)}}}} \in \mathbb{F}_2((1/z)).$$

Conversely, each power series in $\mathbb{F}_2[[1/z]]$ without constant term

$$(1.3) \quad c_{-1}z^{-1} + c_{-2}z^{-2} + c_{-3}z^{-3} + \dots$$

admits a continued fraction expansion of form (1.1).

Let a and b be two distinct elements from $\mathbb{F}_2[z] \setminus \mathbb{F}_2$. We consider the continued fraction (1.1) associated with the (a, b) -Thue-Morse sequence

$$\mathbf{t} = (t_0(z), t_1(z), t_2(z), \dots) = (a, b, b, a, b, a, a, b, \dots)$$

that we shall denote

$$(1.4) \quad \text{CF}(a, b) := \text{CF}(\mathbf{t}).$$

Date: 9/04/2020 17:26:06 .

2010 Mathematics Subject Classification. 11B85, 11J70, 11B50, 11Y65, 05A15.

Key words and phrases. automatic sequence, continued fraction, Thue-Morse sequence.

* This paper was automatically generated from a computer program developed by the authors, with 11.6 minutes CPU time used.

Theorem 1.1. *When $(a, b) = (z^3, z^4 + 1)$, the two power series $\text{CF}(a, b)$ and $\text{CF}(b, a)$ are algebraic over $\mathbb{F}_2(z)$, with minimal polynomials of the form*

$$p_4(z)y^4 + p_3(z)y^3 + p_2(z)y^2 + p_1(z)y + p_0(z) = 0$$

where, for $\text{CF}(a, b)$,

$$\begin{aligned} p_0(z) &= z^{21} + z^{17} + z^{16} + z^{15} + z^{14} + z^{11} + z^7 + z^6 + z^3 + 1, \\ p_1(z) &= z^{25} + z^{24} + z^{20} + z^{19} + z^{18} + z^{17} + z^{14} + z^{10} + z^6 + z^3, \\ p_2(z) &= z^{28} + z^{22} + z^{20} + z^6, \\ p_3(z) &= z^{25} + z^{24} + z^{20} + z^{19} + z^{18} + z^{17} + z^{14} + z^{10} + z^6 + z^3, \\ p_4(z) &= z^{22} + z^{21} + z^{20} + z^{17} + z^{15} + z^{14} + z^{11} + z^7 + z^3, \end{aligned}$$

and for $\text{CF}(b, a)$,

$$\begin{aligned} p_0(z) &= z^{21} + z^{17} + z^{15} + z^{14} + z^{12} + z^{11} + z^7 + z^6 + z^3, \\ p_1(z) &= z^{25} + z^{24} + z^{20} + z^{19} + z^{18} + z^{17} + z^{14} + z^{10} + z^6 + z^3, \\ p_2(z) &= z^{28} + z^{22} + z^{20} + z^6, \\ p_3(z) &= z^{25} + z^{24} + z^{20} + z^{19} + z^{18} + z^{17} + z^{14} + z^{10} + z^6 + z^3, \\ p_4(z) &= z^{22} + z^{21} + z^{20} + z^{17} + z^{16} + z^{15} + z^{14} + z^{12} + z^{11} + z^7 + z^3 + 1. \end{aligned}$$

2. PROOF

Define the sequences of polynomials $(P_n(z))_{n \geq 0}$ and $(Q_n(z))_{n \geq 0}$ by

$$\begin{pmatrix} P_n(z) & Q_n(z) \\ P_{n-1}(z) & Q_{n-1}(z) \end{pmatrix} := \begin{pmatrix} t_n(z) & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} t_{n-1}(z) & 1 \\ 1 & 0 \end{pmatrix} \cdots \begin{pmatrix} t_0(z) & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}.$$

By the basic properties of continued fractions, we have

$$(2.1) \quad \text{CF}_n(a, b) = \frac{P_n(z)}{Q_n(z)} \in \mathbb{F}_2((1/z)).$$

As formula (2.1) is not efficient for calculating $\text{CF}_n(a, b)$, a fast method is to be derived and described as follows. Define

$$\begin{aligned} M_n(x) &= x^{\deg(t_{2^n-1})} \begin{pmatrix} t_{2^n-1}(1/x) & 1 \\ 1 & 0 \end{pmatrix} \times x^{\deg(t_{2^n-2})} \begin{pmatrix} t_{2^n-2}(1/x) & 1 \\ 1 & 0 \end{pmatrix} \times \\ &\quad \cdots \times x^{\deg(t_0)} \begin{pmatrix} t_0(1/x) & 1 \\ 1 & 0 \end{pmatrix}, \\ W_n(x) &= x^{\deg(\bar{t}_{2^n-1})} \begin{pmatrix} \bar{t}_{2^n-1}(1/x) & 1 \\ 1 & 0 \end{pmatrix} \times x^{\deg(\bar{t}_{2^n-2})} \begin{pmatrix} \bar{t}_{2^n-2}(1/x) & 1 \\ 1 & 0 \end{pmatrix} \times \\ &\quad \cdots \times x^{\deg(\bar{t}_0)} \begin{pmatrix} \bar{t}_0(1/x) & 1 \\ 1 & 0 \end{pmatrix}, \end{aligned}$$

where $\bar{t}$ is the (b, a) -Thue-Morse sequence. By the properties of the Thue-Morse sequence, we have for $n \geq 0$

$$\begin{aligned} M_{n+1}(x) &= W_n(x) \cdot M_n(x), \\ W_{n+1}(x) &= M_n(x) \cdot W_n(x). \end{aligned}$$

Let $x := 1/z$. For each non-zero polynomial $P(z)$, define $\tilde{P}(x)$ to be $P(1/x)$. Then,

$$\text{CF}_n(a, b) = \frac{P_n(z)}{Q_n(z)} = \frac{\tilde{P}_n(x)}{\tilde{Q}_n(x)} \in \mathbb{F}_2((x)) = \mathbb{F}_2((1/z)).$$

Comparing the definition of $M_n(x)$ with the definition of $P_n(x)$ and $Q_n(x)$, we see that

$$\begin{aligned} M_n(x)_{0,1} &= x^{d_n} \tilde{P}_{2^n-1}(x), \\ M_n(x)_{0,0} &= x^{d_n} \tilde{Q}_{2^n-1}(x), \end{aligned}$$

for some positive integer d_n . Hence,

$$(2.2) \quad \text{CF}_{2^{2^n}-1}(a, b) = \frac{\tilde{P}_{2^{2^n}-1}(x)}{\tilde{Q}_{2^{2^n}-1}(x)} = \frac{M_{2^n}(x)_{0,1}}{M_{2^n}(x)_{0,0}}.$$

Therefore, by the convergence theorem of continued fraction, the algebraicity of $\text{CF}(a, b)$ will be established if it is shown that both $M_{2^n}(x)_{0,1}$ and $M_{2^n}(x)_{0,0}$ converge to algebraic series in $\mathbb{F}_2[[x]]$.

Actually, we will prove that for all $0 \leq i, j \leq 1$ the four sequences $(M_{2^n}(x)_{i,j})_n$, $(M_{2^{n+1}}(x)_{i,j})_n$, $(W_{2^n}(x)_{i,j})_n$, and $(W_{2^{n+1}}(x)_{i,j})_n$ converge to algebraic series in $\mathbb{F}_2[[x]]$. For this purpose, we define four 2×2 matrices M^e, M^o, W^e, W^o as follows: For each $T \in \{M^e, M^o, W^e, W^o\}$ and $0 \leq i, j \leq 1$, $T_{i,j}$ is defined to be the unique solution in $\mathbb{F}_2[[x]]$ of the polynomial $\phi(T, i, j)$ under certain initial conditions; the polynomials $\phi(T, i, j)$ and initial conditions are given in Section 3. We will prove that these four matrices, whose components are algebraic by definition, are the limits of $(M_{2^n}(x))_n$, $(M_{2^{n+1}}(x))_n$, $(W_{2^n}(x))_n$, and $(W_{2^{n+1}}(x))_n$.

Let us explain how the polynomials $\phi(T, i, j)$ and initial conditions are found, and why the solutions exist and are unique. For $0 \leq i, j \leq 1$, the coefficients of the polynomial $\phi(M^e, i, j)$ (resp. $\phi(M^o, i, j)$, $\phi(W^e, i, j)$, and $\phi(W^o, i, j)$) are the Padé-Hermite approximants of type

$$(175, 175, 175, 175, 175)$$

of the vector

$$(1, T^3, T^6, T^9, T^{12}),$$

where $T = M_{12,i,j}$ (resp. $M_{11,i,j}$, $W_{12,i,j}$, and $W_{11,i,j}$), found by the Derksen algorithm. We take the first 8 terms of $M_{12,i,j}$ (resp. $M_{11,i,j}$, $W_{12,i,j}$, and $W_{11,i,j}$) as initial conditions for $\phi(M^e, i, j)$ (resp. $\phi(M^o, i, j)$, $\phi(W^e, i, j)$, and $\phi(W^o, i, j)$). The following fact will be used to ensure that the solution exists and is unique: let $P(x, y) \in \mathbb{F}_2[x, y]$ and for each series $f(x) = \sum_{n=0}^{\infty} a_n x^n \in \mathbb{F}_2[[x]]$ denote the partial sum $\sum_{j=0}^{n-1} a_j x^j$ by $f_n(x)$ for $n \geq 0$. If for some $n \geq 0$ and $a_0, a_1, \dots, a_{n-1} \in \mathbb{F}_2$ $P(x, \sum_{j=0}^{n-1} a_j x^j) = O(x^n)$ and $Q(x, y) := P(x, \sum_{j=0}^{n-1} a_j x^j + x^n y)$ can be written as $x^m \sum_{j=0}^{\infty} q_j(x) y^j$ where $q_j(x)$ are polynomials for $j \geq 0$, $q_1(0) = 1$, and $q_j(0) = 0$ for $j > 1$, then there exists a unique solution $f(x) \in \mathbb{F}_2[[x]]$ of $P(x, f(x)) = 0$ that satisfies the initial condition $f_n(x) = \sum_{j=0}^{n-1} a_j x^j$.

We state two lemmas concerning the four matrices M^e, M^o, W^e, W^o . The first one is about relations between them; the second, about the structure of the each matrix.

Lemma 2.1. *We have*

$$(2.3) \quad M^e = W^o \cdot M^o,$$

$$(2.4) \quad M^o = W^e \cdot M^e,$$

$$(2.5) \quad W^e = M^o \cdot W^o,$$

$$(2.6) \quad W^o = M^e \cdot W^e.$$

Proof. We give the proof of the identity

$$M_{0,0}^e = W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o,$$

the proofs of the others being similar.

First, we compute the minimal polynomials of $W_{0,0}^o M_{0,0}^o$ and $W_{0,1}^o M_{1,0}^o$. We know that

$$P(x, y) = \text{Res}_z (\phi(W^o, 0, 0)(x, z), z^{12} \cdot \phi(M^o, 0, 0)(x, y/z))$$

is an annihilating polynomial of $W_{0,0}^o M_{0,0}^o$. We use Padé-Hermite approximation to find a candidate for the minimal polynomial of $W_{0,0}^o M_{0,0}^o$, that will be called $\phi_0(x, y)$. To prove that $\phi_0(x, y)$ is indeed the minimal polynomial, it suffices to prove that it is an irreducible factor of $P(x, y)$ of multiplicity m and that $Q(x, y) := P(x, y)/\phi_0(x, y)^m$ is not an annihilating polynomial of $W_{0,0}^o M_{0,0}^o$. We verify the first point directly. For the second point, we truncate $W_{0,0}^o M_{0,0}^o$ to order 630 and substitute it for y in $Q(x, y)$. We get a series of valuation less than 630, which proves that $Q(x, y)$ is not an annihilating polynomial of $W_{0,0}^o M_{0,0}^o$. We find the minimal polynomial $\phi_1(x, y)$ of $W_{0,1}^o M_{1,0}^o$ in a similar way.

Now we prove that $\phi(M^e, 0, 0)$ is the minimal polynomial of $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$. We know that

$$S(x, y) = \text{Res}_z (\phi_0(x, z), \phi_1(x, y + z))$$

is an annihilating of $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$. We verify that $\phi(M^e, 0, 0)$ is an irreducible factor of $S(x, y)$ of multiplicity μ , and that $Q(x, y) := S(x, y)/\phi(M^e, 0, 0)^\mu$ is not an annihilating polynomial of $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$. To see the last point, we truncate $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$ to order 770 and substitute it for y in $Q(x, y)$. We get a series of valuation less than 770, and therefore $Q(x, y)$ is not an annihilating polynomial of $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$.

Finally, the first 16 terms of $M_{0,0}^e$ and $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$ coincide. As these first terms determine a unique solution of $\phi(M^e, 0, 0)$, we know that the two series are one and the same. $\square$

Define

$$R^e = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \quad \text{and} \quad R^o = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}.$$

Lemma 2.2. *For all integers $k \geq 2$ and $u = 2^{2k-1}$, the following identities hold.*

$$\begin{aligned} M^e[:14u] &= M^e[:7u] + x^{4u} M^e[:3u] + x^{6u} M^e[:u] + x^{4u} M^e[:7u] + x^{8u} M^e[:3u] \\ &\quad + x^{10u} M^e[:u] + x^{6u} M^e[:7u] + x^{10u} M^e[:3u] + x^{12u} M^e[:u] \\ &\quad + x^{10u} M^e[:4u] + (x^{7u} + x^{11u} + x^{13u}) R^e, \\ W^e[:14u] &= W^e[:7u] + x^{4u} W^e[:3u] + x^{6u} W^e[:u] + x^{4u} W^e[:7u] + x^{8u} W^e[:3u] \\ &\quad + x^{10u} W^e[:u] + x^{6u} W^e[:7u] + x^{10u} W^e[:3u] + x^{12u} W^e[:u] \\ &\quad + x^{10u} W^e[:4u] + (x^{7u} + x^{11u} + x^{13u}) R^e. \end{aligned}$$

For all integers $k \geq 2$ and $u = 2^{2k}$, the following identities hold.

$$\begin{aligned}
M^o[:14u] &= M^o[:7u] + x^{4u}M^o[:3u] + x^{6u}M^o[:u] + x^{4u}M^o[:7u] + x^{8u}M^o[:3u] \\
&\quad + x^{10u}M^o[:u] + x^{6u}M^o[:7u] + x^{10u}M^o[:3u] + x^{12u}M^o[:u] \\
&\quad + x^{10u}M^o[:4u] + (x^{7u} + x^{11u} + x^{13u})R^o, \\
W^o[:14u] &= W^o[:7u] + x^{4u}W^o[:3u] + x^{6u}W^o[:u] + x^{4u}W^o[:7u] + x^{8u}W^o[:3u] \\
&\quad + x^{10u}W^o[:u] + x^{6u}W^o[:7u] + x^{10u}W^o[:3u] + x^{12u}W^o[:u] \\
&\quad + x^{10u}W^o[:4u] + (x^{7u} + x^{11u} + x^{13u})R^o.
\end{aligned}$$

Proof. To prove Lemma 2.2 we first construct an automaton for each sequence concerned, and then transform the conditions on infinitely many k 's into finitely many conditions on the states of the automaton. In the following, we will prove that for $T = M_{0,0}^o$, for all integer $k \geq 2$ and $u = 2^{2k}$,

$$\begin{aligned}
T[:14u] &= T[:7u] + x^{4u}T[:3u] + x^{6u}T[:u] + x^{4u}T[:7u] + x^{8u}T[:3u] + x^{10u}T[:u] \\
&\quad + x^{6u}T[:7u] + x^{10u}T[:3u] + x^{12u}T[:u] + x^{10u}T[:4u].
\end{aligned}$$

The proof of the other 15 cases are similar. We break down the above identity into 7 parts:

$$\begin{aligned}
0 &= x^{6u}T[u:2u] + x^{4u}T[3u:4u] + T[7u:8u], \\
0 &= x^{8u}T[:u] + x^{6u}T[2u:3u] + x^{4u}T[4u:5u] + T[8u:9u], \\
0 &= x^{8u}T[u:2u] + x^{6u}T[3u:4u] + x^{4u}T[5u:6u] + T[9u:10u], \\
0 &= x^{10u}T[:u] + x^{8u}T[2u:3u] + x^{6u}T[4u:5u] + x^{4u}T[6u:7u] \\
&\quad + T[10u:11u], \\
0 &= x^{6u}T[5u:6u] + T[11u:12u], \\
0 &= x^{12u}T[:u] + x^{6u}T[6u:7u] + T[12u:13u], \\
0 &= x^{10u}T[3u:4u] + T[13u:14u],
\end{aligned}$$

which can be rewritten as

$$(2.7) \quad 0 = T[[1w]_2] + T[[11w]_2] + T[[111w]_2],$$

$$(2.8) \quad 0 = T[[w]_2] + T[[10w]_2] + T[[100w]_2] + T[[1000w]_2],$$

$$(2.9) \quad 0 = T[[1w]_2] + T[[11w]_2] + T[[101w]_2] + T[[1001w]_2],$$

$$(2.10) \quad 0 = T[[w]_2] + T[[10w]_2] + T[[100w]_2] + T[[110w]_2] + T[[1010w]_2],$$

$$(2.11) \quad 0 = T[[101w]_2] + T[[1011w]_2],$$

$$(2.12) \quad 0 = T[[w]_2] + T[[110w]_2] + T[[1100w]_2],$$

$$(2.13) \quad 0 = T[[11w]_2] + T[[1101w]_2],$$

for all binary word w of length $2k$ and $w \neq 0^{2k}$, and

$$(2.14) \quad 0 = T[[1w]_2] + T[[11w]_2] + T[[111w]_2],$$

$$(2.15) \quad 0 = T[[w]_2] + T[[10w]_2] + T[[100w]_2] + T[[1000w]_2],$$

$$(2.16) \quad 0 = T[[1w]_2] + T[[11w]_2] + T[[101w]_2] + T[[1001w]_2],$$

$$(2.17) \quad 0 = T[[w]_2] + T[[10w]_2] + T[[100w]_2] + T[[110w]_2] + T[[1010w]_2],$$

$$(2.18) \quad 0 = T[[101w]_2] + T[[1011w]_2],$$

$$(2.19) \quad 0 = T[[w]_2] + T[[110w]_2] + T[[1100w]_2],$$

$$(2.20) \quad 0 = T[[11w]_2] + T[[1101w]_2],$$

for $w = 0^{2k}$.

First we calculate an 2-automaton that generates T from its minimal polynomial and its first terms. This automaton has 379 states; its transition function and output function can be found in the annex. Let $A(s, w)$ denote the state reached after reading w from right to left starting from the state s , and τ the output function. Define

$$E_{2k} = \{A(i, w) : |w| = 2k, w \neq 0^{2k}\}.$$

Identities (2.7) through (2.20) can be written as

$$(2.21) \quad 0 = \tau(A(s, 1)) + \tau(A(s, 11)) + \tau(A(s, 111)),$$

$$(2.22) \quad 0 = \tau(A(s, \epsilon)) + \tau(A(s, 10)) + \tau(A(s, 100)) + \tau(A(s, 1000)),$$

$$(2.23) \quad 0 = \tau(A(s, 1)) + \tau(A(s, 11)) + \tau(A(s, 101)) + \tau(A(s, 1001)),$$

$$0 = \tau(A(s, \epsilon)) + \tau(A(s, 10)) + \tau(A(s, 100)) + \tau(A(s, 110))$$

$$(2.24) \quad + \tau(A(s, 1010)),$$

$$(2.25) \quad 0 = \tau(A(s, 101)) + \tau(A(s, 1011)),$$

$$(2.26) \quad 0 = \tau(A(s, \epsilon)) + \tau(A(s, 110)) + \tau(A(s, 1100)),$$

$$(2.27) \quad 0 = \tau(A(s, 11)) + \tau(A(s, 1101)),$$

for all $s \in E_{2k}$, and

$$(2.28) \quad 0 = \tau(A(s, 1)) + \tau(A(s, 11)) + \tau(A(s, 111)),$$

$$(2.29) \quad 0 = \tau(A(s, \epsilon)) + \tau(A(s, 10)) + \tau(A(s, 100)) + \tau(A(s, 1000)),$$

$$(2.30) \quad 0 = \tau(A(s, 1)) + \tau(A(s, 11)) + \tau(A(s, 101)) + \tau(A(s, 1001)),$$

$$0 = \tau(A(s, \epsilon)) + \tau(A(s, 10)) + \tau(A(s, 100)) + \tau(A(s, 110))$$

$$(2.31) \quad + \tau(A(s, 1010)),$$

$$(2.32) \quad 0 = \tau(A(s, 101)) + \tau(A(s, 1011)),$$

$$(2.33) \quad 0 = \tau(A(s, \epsilon)) + \tau(A(s, 110)) + \tau(A(s, 1100)),$$

$$(2.34) \quad 0 = \tau(A(s, 11)) + \tau(A(s, 1101)),$$

for $s = A(i, 0^{2k})$. We find that $(A(i, 0^{30}), E_{30}) = (A(i, 0^{18}), E_{18})$, so that we only have to verify that identities (2.21) through (2.34) hold for $2 \leq k \leq 15$, which turns out to be true. $\square$

In the following lemma, we express M_{2k} , M_{2k+1} , W_{2k} , and W_{2k+1} in terms of M^e , M^o , W^e , and W^o .

Lemma 2.3. *For all integer $k \geq 2$, and $u = 2^{2k-1}$,*

$$M_{2k} = M^e[:7u] + x^{4u}M^e[:3u] + x^{6u}M^e[:u] + x^{7u}R^e,$$

$$W_{2k} = W^e[:7u] + x^{4u}W^e[:3u] + x^{6u}W^e[:u] + x^{7u}R^e.$$

For all integer $k \geq 2$, and $u = 2^{2k}$,

$$M_{2k+1} = M^o[:7u] + x^{4u}M^o[:3u] + x^{6u}M^o[:u] + x^{7u}R^o,$$

$$W_{2k+1} = W^o[:7u] + x^{4u}W^o[:3u] + x^{6u}W^o[:u] + x^{7u}R^o.$$

Proof. We call the four identities in Lemma 2.3 also by the name M_{2k} , W_{2k} , M_{2k+1} , and W_{2k+1} . For $n = 2$, the identities can be verified directly. For $n \geq 2$, we claim that

$$\begin{aligned} M_{2k} \wedge W_{2k} &\Rightarrow M_{2k+1} \wedge W_{2k+1}, \\ M_{2k+1} \wedge W_{2k+1} &\Rightarrow M_{2k+2} \wedge W_{2k+2}. \end{aligned}$$

We give the proof of

$$M_{2k} \wedge W_{2k} \Rightarrow M_{2k+1},$$

the other ones being similar. Set $u = 2^{2k}$ and $v = 2^{2k-1}$. By definition and induction hypothesis, the left side of identity M_{2k+1} is equal to

$$\begin{aligned} (2.35) \quad W_{2k} M_{2k} &= (W^e[:7v] + x^{4v} W^e[:3v] + x^{6v} W^e[:v] + x^{7u} R^e) \times (M^e[:7v] \\ &\quad + x^{4v} M^e[:3v] + x^{6v} M^e[:v] + x^{7u} R^e); \end{aligned}$$

Call this expression *lhs*. Note that both sides of identity M_{2k+1} have the same term of highest degree $x^{14v} R^o$. Therefore we only need to prove that their difference is $O(x^{14v})$. Using Lemma 2.1 it can be seen that the right side of identity M_{2k+1} is congruent, modulo x^{14v} , to

$$W^e[:14v] M^e[:14v] + x^{8v} W^e[:6v] M^e[:6v] + x^{12v} W^e[:2v] M^e[:2v].$$

For all $n \leq 14$, replace the occurrences of $W^e[:n \cdot v]$ and $M^e[:n \cdot v]$ in the above expression by the reduction modulo $x^{n \cdot v}$ of the right side of the corresponding identity in Lemma 2.2 and get a new expression, which we call *rhs*. Define

$$\begin{aligned} X &:= x^v, \\ a_n &:= W^e[n \cdot v : (n+1) \cdot v] / X^n, \\ b_n &:= M^e[n \cdot v : (n+1) \cdot v] / X^n, \\ c &:= R^e. \end{aligned}$$

Using the notation introduced above, we can represent the expressions *lhs* (2.35) and *rhs* as polynomials in $\mathbb{F}_2[a_1, \dots, a_{14}, b_1, \dots, b_{14}, c][X]$. Note that it is not a problem that a_j commutes with b_k while W^e does not commute with M^e , because in the expressions concerned, the products of W^e -terms and M^e -terms are always in the same order. We let the computer do the simplification and check that the difference between these two polynomials is indeed $O(X^{14})$, which completes the proof. $\square$

Proof of Theorem 1.1. We prove the theorem for $\text{CF}(a, b)$; for $\text{CF}(b, a)$, the proof is similar. By Lemma 2.3, we have For all $0 \leq j, k \leq 1$,

$$\begin{aligned} \lim_{n \rightarrow \infty} M_{2n,j,k} &= M_{j,k}^e, \\ \lim_{n \rightarrow \infty} M_{2n+1,j,k} &= M_{j,k}^o, \\ \lim_{n \rightarrow \infty} W_{2n,j,k} &= W_{j,k}^e, \\ \lim_{n \rightarrow \infty} W_{2n+1,j,k} &= W_{j,k}^o. \end{aligned}$$

Let $z = 1/x$. By the convergence theorem and identity (2.2),

$$\text{CF}(a, b) = \frac{M_{0,1}^e(x)}{M_{0,0}^e(x)}.$$

By definition, $\phi(M^e, 0, 1)$ and $\phi(M^e, 0, 0)$ are minimal polynomials of $M_{0,1}^e$ and $M_{0,0}^e$. Therefore

$$P(x, y) = \text{Res}_t (\phi(M^e, 0, 1)(x, t), y^{12}\phi(M^e, 0, 1)(x, t/y))$$

is an annihilating polynomial of $f(x) = M_{0,1}^e/M_{0,0}^e$.

Define

$$Q(x, y) = q_4(x)y^4 + q_3(x)y^3 + q_2(x)y^2 + q_1(x)y + q_0(x),$$

where

$$q_0(x) = x^{28} + x^{25} + x^{22} + x^{21} + x^{17} + x^{14} + x^{13} + x^{12} + x^{11} + x^7,$$

$$q_1(x) = x^{25} + x^{22} + x^{18} + x^{14} + x^{11} + x^{10} + x^9 + x^8 + x^4 + x^3,$$

$$q_2(x) = x^{22} + x^8 + x^6 + 1,$$

$$q_3(x) = x^{25} + x^{22} + x^{18} + x^{14} + x^{11} + x^{10} + x^9 + x^8 + x^4 + x^3,$$

$$q_4(x) = x^{25} + x^{21} + x^{17} + x^{14} + x^{13} + x^{11} + x^8 + x^7 + x^6.$$

The polynomial $Q(x, y)$ is the candidate for the minimal polynomial of $f(x)$ found by Padé-Hermite approximation. To prove that it is indeed the minimal polynomial of $f(x)$, we only need to prove that it is an irreducible factor of $P(x, y)$ of multiplicity m and $R(x, y) := P(x, y)/Q(x, y)^m$ is not an annihilating polynomial of $f(x)$. We verify the first point directly. For the second point, we find that when we truncate $f(x)$ to order 216, and substitute it for y in $R(z, y)$, we get a series with valuation smaller than 216, which proves that $R(z, y)$ is not an annihilating polynomial of $f(x)$. Finally, $z^{12}Q(1/z, y)$ is the minimal polynomial of $\text{CF}(a, b) = f(1/z)$. $\square$

3. ANNEXE

All 16 polynomials are of the form

$$p_0(x) + p_3(x)y^3 + p_6(x)y^6 + p_9(x)y^9 + p_{12}(x)y^{12}.$$

The coefficients $p_j(x)$ and the 16 initial terms to determine the solutions uniquely are given below.

For $\phi(M^e, 0, 0)$:

$$p_0(x) = x^{150} + x^{142} + x^{128} + x^{122} + x^{116} + x^{114} + x^{100} + x^{94} + x^{82} + x^{74} \\ + x^{70} + x^{64} + x^{62} + x^{58} + x^{52} + x^{48} + x^{42} + x^{38} + x^{36},$$

$$p_3(x) = x^{136} + x^{132} + x^{128} + x^{126} + x^{124} + x^{120} + x^{116} + x^{110} + x^{108} + x^{104} \\ + x^{96} + x^{88} + x^{76} + x^{72} + x^{70} + x^{64} + x^{62} + x^{54} + x^{48} + x^{46} + x^{38} \\ + x^{36} + x^{28} + x^{24} + x^{22} + x^{20},$$

$$p_6(x) = x^{138} + x^{136} + x^{132} + x^{130} + x^{128} + x^{124} + x^{122} + x^{120} + x^{114} + x^{112} \\ + x^{108} + x^{104} + x^{102} + x^{100} + x^{94} + x^{82} + x^{76} + x^{72} + x^{64} + x^{60} \\ + x^{56} + x^{52} + x^{46} + x^{44} + x^{42} + x^{40} + x^{34} + x^{30} + x^{26} + x^{24} + x^{20} \\ + x^{14} + x^{12} + x^8 + x^6 + 1,$$

$$p_9(x) = x^{144} + x^{138} + x^{124} + x^{116} + x^{112} + x^{110} + x^{108} + x^{106} + x^{100} + x^{96} \\ + x^{94} + x^{88} + x^{82} + x^{80} + x^{74} + x^{68} + x^{54} + x^{48} + x^{46} + x^{42} + x^{40} \\ + x^{38} + x^{32} + x^{30} + x^{28} + x^{24} + x^{22} + x^{20} + x^{18} + x^{12} + x^8 + x^6,$$

$$p_{12}(x) = x^{144} + x^{112} + x^{80} + x^{48} + x^{32} + 1,$$

and the initial terms are $[1, 0, 0, 0, 0, 0, 0, 0, 1, 0, 0, 0, 1, 0, 1, 0]$.

For $\phi(M^e, 0, 1)$:

$$\begin{aligned} p_0(x) = & x^{147} + x^{144} + x^{140} + x^{139} + x^{138} + x^{136} + x^{134} + x^{129} + x^{128} + x^{125} \\ & + x^{122} + x^{114} + x^{113} + x^{112} + x^{108} + x^{104} + x^{100} + x^{98} + x^{95} + x^{94} \\ & + x^{88} + x^{87} + x^{84} + x^{83} + x^{82} + x^{80} + x^{79} + x^{75} + x^{72} + x^{71} + x^{70} \\ & + x^{68} + x^{67} + x^{66} + x^{65} + x^{64} + x^{63} + x^{62} + x^{61} + x^{59} + x^{58} + x^{55} \\ & + x^{49} + x^{47} + x^{45} + x^{44} + x^{42} + x^{40} + x^{39}, \end{aligned}$$

$$\begin{aligned} p_3(x) = & x^{141} + x^{138} + x^{137} + x^{127} + x^{125} + x^{124} + x^{123} + x^{122} + x^{121} + x^{119} \\ & + x^{117} + x^{116} + x^{115} + x^{114} + x^{113} + x^{103} + x^{101} + x^{100} + x^{99} + x^{98} \\ & + x^{97} + x^{95} + x^{92} + x^{91} + x^{77} + x^{74} + x^{73} + x^{63} + x^{60} + x^{59} + x^{55} \\ & + x^{53} + x^{52} + x^{51} + x^{50} + x^{49} + x^{47} + x^{45} + x^{44} + x^{43} + x^{42} + x^{41} \\ & + x^{37} + x^{34} + x^{33} + x^{31} + x^{28} + x^{27}, \end{aligned}$$

$$\begin{aligned} p_6(x) = & x^{138} + x^{134} + x^{132} + x^{128} + x^{126} + x^{124} + x^{118} + x^{116} + x^{112} + x^{110} \\ & + x^{108} + x^{106} + x^{98} + x^{94} + x^{92} + x^{88} + x^{84} + x^{80} + x^{76} + x^{74} + x^{72} \\ & + x^{66} + x^{64} + x^{62} + x^{60} + x^{52} + x^{50} + x^{46} + x^{40} + x^{32} + x^{30} + x^{26} \\ & + x^{20} + x^{18}, \end{aligned}$$

$$\begin{aligned} p_9(x) = & x^{135} + x^{132} + x^{131} + x^{129} + x^{126} + x^{125} + x^{119} + x^{116} + x^{113} + x^{112} \\ & + x^{111} + x^{110} + x^{109} + x^{107} + x^{104} + x^{103} + x^{101} + x^{98} + x^{97} + x^{87} \\ & + x^{84} + x^{83} + x^{79} + x^{76} + x^{75} + x^{73} + x^{70} + x^{69} + x^{65} + x^{63} + x^{62} \\ & + x^{61} + x^{60} + x^{57} + x^{56} + x^{55} + x^{54} + x^{53} + x^{49} + x^{47} + x^{46} + x^{44} \\ & + x^{42} + x^{40} + x^{38} + x^{36} + x^{35} + x^{34} + x^{33} + x^{27} + x^{25} + x^{24} + x^{23} \\ & + x^{22} + x^{21} + x^{19} + x^{16} + x^{13} + x^{12} + x^{11} + x^{10} + x^9, \end{aligned}$$

$$\begin{aligned} p_{12}(x) = & x^{132} + x^{128} + x^{116} + x^{112} + x^{68} + x^{64} + x^{52} + x^{48} + x^{20} + x^{16} + x^4 \\ & + 1, \end{aligned}$$

and the initial terms are $[0, 0, 0, 1, 0, 0, 0, 0, 0, 0, 1, 1, 0, 0, 1, 1]$.

For $\phi(M^e, 1, 0)$:

$$\begin{aligned} p_0(x) = & x^{147} + x^{144} + x^{140} + x^{139} + x^{138} + x^{136} + x^{134} + x^{129} + x^{128} + x^{125} \\ & + x^{122} + x^{114} + x^{113} + x^{112} + x^{108} + x^{104} + x^{100} + x^{98} + x^{95} + x^{94} \\ & + x^{88} + x^{87} + x^{84} + x^{83} + x^{82} + x^{80} + x^{79} + x^{75} + x^{72} + x^{71} + x^{70} \\ & + x^{68} + x^{67} + x^{66} + x^{65} + x^{64} + x^{63} + x^{62} + x^{61} + x^{59} + x^{58} + x^{55} \\ & + x^{49} + x^{47} + x^{45} + x^{44} + x^{42} + x^{40} + x^{39}, \end{aligned}$$

$$\begin{aligned} p_3(x) = & x^{141} + x^{138} + x^{137} + x^{127} + x^{125} + x^{124} + x^{123} + x^{122} + x^{121} + x^{119} \\ & + x^{117} + x^{116} + x^{115} + x^{114} + x^{113} + x^{103} + x^{101} + x^{100} + x^{99} + x^{98} \\ & + x^{97} + x^{95} + x^{92} + x^{91} + x^{77} + x^{74} + x^{73} + x^{63} + x^{60} + x^{59} + x^{55} \end{aligned}$$

$$\begin{aligned}
& + x^{53} + x^{52} + x^{51} + x^{50} + x^{49} + x^{47} + x^{45} + x^{44} + x^{43} + x^{42} + x^{41} \\
& + x^{37} + x^{34} + x^{33} + x^{31} + x^{28} + x^{27}, \\
p_6(x) &= x^{138} + x^{134} + x^{132} + x^{128} + x^{126} + x^{124} + x^{118} + x^{116} + x^{112} + x^{110} \\
& + x^{108} + x^{106} + x^{98} + x^{94} + x^{92} + x^{88} + x^{84} + x^{80} + x^{76} + x^{74} + x^{72} \\
& + x^{66} + x^{64} + x^{62} + x^{60} + x^{52} + x^{50} + x^{46} + x^{40} + x^{32} + x^{30} + x^{26} \\
& + x^{20} + x^{18}, \\
p_9(x) &= x^{135} + x^{132} + x^{131} + x^{129} + x^{126} + x^{125} + x^{119} + x^{116} + x^{113} + x^{112} \\
& + x^{111} + x^{110} + x^{109} + x^{107} + x^{104} + x^{103} + x^{101} + x^{98} + x^{97} + x^{87} \\
& + x^{84} + x^{83} + x^{79} + x^{76} + x^{75} + x^{73} + x^{70} + x^{69} + x^{65} + x^{63} + x^{62} \\
& + x^{61} + x^{60} + x^{57} + x^{56} + x^{55} + x^{54} + x^{53} + x^{49} + x^{47} + x^{46} + x^{44} \\
& + x^{42} + x^{40} + x^{38} + x^{36} + x^{35} + x^{34} + x^{33} + x^{27} + x^{25} + x^{24} + x^{23} \\
& + x^{22} + x^{21} + x^{19} + x^{16} + x^{13} + x^{12} + x^{11} + x^{10} + x^9, \\
p_{12}(x) &= x^{132} + x^{128} + x^{116} + x^{112} + x^{68} + x^{64} + x^{52} + x^{48} + x^{20} + x^{16} + x^4 \\
& + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 1, 0, 0, 0, 0, 0, 0, 1, 1, 0, 0, 1, 1]$.

For $\phi(M^e, 1, 1)$:

$$\begin{aligned}
p_0(x) &= x^{144} + x^{138} + x^{136} + x^{126} + x^{114} + x^{108} + x^{100} + x^{98} + x^{92} + x^{86} \\
& + x^{84} + x^{82} + x^{80} + x^{76} + x^{74} + x^{72} + x^{68} + x^{66} + x^{64} + x^{62} + x^{60} \\
& + x^{56} + x^{54} + x^{52} + x^{42}, \\
p_3(x) &= x^{126} + x^{120} + x^{118} + x^{114} + x^{112} + x^{106} + x^{102} + x^{94} + x^{88} + x^{80} \\
& + x^{64} + x^{62} + x^{58} + x^{56} + x^{54} + x^{46} + x^{42} + x^{30} + x^{26} + x^{18}, \\
p_6(x) &= x^{126} + x^{118} + x^{108} + x^{106} + x^{100} + x^{92} + x^{86} + x^{84} + x^{80} + x^{76} + x^{70} \\
& + x^{68} + x^{62} + x^{60} + x^{56} + x^{50} + x^{48} + x^{44} + x^{42} + x^{38} + x^{36} + x^{30} \\
& + x^{28} + x^{18} + x^6 + 1, \\
p_9(x) &= x^{120} + x^{114} + x^{112} + x^{106} + x^{100} + x^{86} + x^{78} + x^{72} + x^{70} + x^{62} + x^{58} \\
& + x^{54} + x^{48} + x^{46} + x^{44} + x^{42} + x^{40} + x^{38} + x^{36} + x^{26} + x^{24} + x^{22} \\
& + x^{18} + x^{16} + x^{14} + x^{12} + x^8 + x^6, \\
p_{12}(x) &= x^{120} + x^{112} + x^{56} + x^{48} + x^8 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 0, 1, 0, 0, 0, 0, 0, 0, 0, 1, 0]$.

For $\phi(W^e, 0, 0)$:

$$\begin{aligned}
p_0(x) &= x^{144} + x^{138} + x^{136} + x^{132} + x^{126} + x^{124} + x^{120} + x^{112} + x^{108} + x^{100} \\
& + x^{98} + x^{96} + x^{90} + x^{88} + x^{82} + x^{80} + x^{74} + x^{70} + x^{66} + x^{64} + x^{58} \\
& + x^{50} + x^{46} + x^{44} + x^{42} + x^{38} + x^{36}, \\
p_3(x) &= x^{126} + x^{120} + x^{118} + x^{112} + x^{108} + x^{96} + x^{88} + x^{86} + x^{78} + x^{70} + x^{56} \\
& + x^{52} + x^{48} + x^{46} + x^{44} + x^{40} + x^{38} + x^{24} + x^{22} + x^{20},
\end{aligned}$$

$$\begin{aligned}
p_6(x) &= x^{126} + x^{118} + x^{114} + x^{108} + x^{102} + x^{94} + x^{92} + x^{88} + x^{84} + x^{80} + x^{78} \\
&\quad + x^{76} + x^{72} + x^{68} + x^{60} + x^{58} + x^{54} + x^{50} + x^{48} + x^{40} + x^{32} + x^{28} \\
&\quad + x^{26} + x^{22} + x^{16} + x^{12} + x^6 + 1, \\
p_9(x) &= x^{120} + x^{114} + x^{112} + x^{106} + x^{100} + x^{86} + x^{78} + x^{72} + x^{70} + x^{62} + x^{58} \\
&\quad + x^{54} + x^{48} + x^{46} + x^{44} + x^{42} + x^{40} + x^{38} + x^{36} + x^{26} + x^{24} + x^{22} \\
&\quad + x^{18} + x^{16} + x^{14} + x^{12} + x^8 + x^6, \\
p_{12}(x) &= x^{120} + x^{112} + x^{56} + x^{48} + x^8 + 1,
\end{aligned}$$

and the initial terms are $[1, 0, 0, 0, 0, 0, 1, 0, 0, 0, 0, 0, 1, 0, 1, 0]$.

For $\phi(W^e, 0, 1)$:

$$\begin{aligned}
p_0(x) &= x^{147} + x^{141} + x^{139} + x^{138} + x^{135} + x^{134} + x^{132} + x^{128} + x^{127} + x^{123} \\
&\quad + x^{117} + x^{115} + x^{112} + x^{111} + x^{108} + x^{106} + x^{105} + x^{103} + x^{99} + x^{96} \\
&\quad + x^{91} + x^{90} + x^{89} + x^{88} + x^{86} + x^{83} + x^{82} + x^{78} + x^{75} + x^{73} + x^{72} \\
&\quad + x^{66} + x^{62} + x^{61} + x^{59} + x^{58} + x^{57} + x^{54} + x^{53} + x^{51} + x^{47} + x^{45} \\
&\quad + x^{42} + x^{40} + x^{39}, \\
p_3(x) &= x^{141} + x^{138} + x^{137} + x^{127} + x^{125} + x^{124} + x^{123} + x^{122} + x^{121} + x^{119} \\
&\quad + x^{117} + x^{116} + x^{115} + x^{114} + x^{113} + x^{103} + x^{101} + x^{100} + x^{99} + x^{98} \\
&\quad + x^{97} + x^{95} + x^{92} + x^{91} + x^{77} + x^{74} + x^{73} + x^{63} + x^{60} + x^{59} + x^{55} \\
&\quad + x^{53} + x^{52} + x^{51} + x^{50} + x^{49} + x^{47} + x^{45} + x^{44} + x^{43} + x^{42} + x^{41} \\
&\quad + x^{37} + x^{34} + x^{33} + x^{31} + x^{28} + x^{27}, \\
p_6(x) &= x^{138} + x^{134} + x^{132} + x^{128} + x^{126} + x^{124} + x^{118} + x^{116} + x^{112} + x^{110} \\
&\quad + x^{108} + x^{106} + x^{98} + x^{94} + x^{92} + x^{88} + x^{84} + x^{80} + x^{76} + x^{74} + x^{72} \\
&\quad + x^{66} + x^{64} + x^{62} + x^{60} + x^{52} + x^{50} + x^{46} + x^{40} + x^{32} + x^{30} + x^{26} \\
&\quad + x^{20} + x^{18}, \\
p_9(x) &= x^{135} + x^{132} + x^{131} + x^{129} + x^{126} + x^{125} + x^{119} + x^{116} + x^{113} + x^{112} \\
&\quad + x^{111} + x^{110} + x^{109} + x^{107} + x^{104} + x^{103} + x^{101} + x^{98} + x^{97} + x^{87} \\
&\quad + x^{84} + x^{83} + x^{79} + x^{76} + x^{75} + x^{73} + x^{70} + x^{69} + x^{65} + x^{63} + x^{62} \\
&\quad + x^{61} + x^{60} + x^{57} + x^{56} + x^{55} + x^{54} + x^{53} + x^{49} + x^{47} + x^{46} + x^{44} \\
&\quad + x^{42} + x^{40} + x^{38} + x^{36} + x^{35} + x^{34} + x^{33} + x^{27} + x^{25} + x^{24} + x^{23} \\
&\quad + x^{22} + x^{21} + x^{19} + x^{16} + x^{13} + x^{12} + x^{11} + x^{10} + x^9, \\
p_{12}(x) &= x^{132} + x^{128} + x^{116} + x^{112} + x^{68} + x^{64} + x^{52} + x^{48} + x^{20} + x^{16} + x^4 \\
&\quad + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 1, 0, 0, 0, 1, 0, 1, 1, 1, 0, 1, 0]$.

For $\phi(W^e, 1, 0)$:

$$\begin{aligned}
p_0(x) &= x^{147} + x^{141} + x^{139} + x^{138} + x^{135} + x^{134} + x^{132} + x^{128} + x^{127} + x^{123} \\
&\quad + x^{117} + x^{115} + x^{112} + x^{111} + x^{108} + x^{106} + x^{105} + x^{103} + x^{99} + x^{96}
\end{aligned}$$

$$\begin{aligned}
& + x^{91} + x^{90} + x^{89} + x^{88} + x^{86} + x^{83} + x^{82} + x^{78} + x^{75} + x^{73} + x^{72} \\
& + x^{66} + x^{62} + x^{61} + x^{59} + x^{58} + x^{57} + x^{54} + x^{53} + x^{51} + x^{47} + x^{45} \\
& + x^{42} + x^{40} + x^{39}, \\
p_3(x) &= x^{141} + x^{138} + x^{137} + x^{127} + x^{125} + x^{124} + x^{123} + x^{122} + x^{121} + x^{119} \\
& + x^{117} + x^{116} + x^{115} + x^{114} + x^{113} + x^{103} + x^{101} + x^{100} + x^{99} + x^{98} \\
& + x^{97} + x^{95} + x^{92} + x^{91} + x^{77} + x^{74} + x^{73} + x^{63} + x^{60} + x^{59} + x^{55} \\
& + x^{53} + x^{52} + x^{51} + x^{50} + x^{49} + x^{47} + x^{45} + x^{44} + x^{43} + x^{42} + x^{41} \\
& + x^{37} + x^{34} + x^{33} + x^{31} + x^{28} + x^{27}, \\
p_6(x) &= x^{138} + x^{134} + x^{132} + x^{128} + x^{126} + x^{124} + x^{118} + x^{116} + x^{112} + x^{110} \\
& + x^{108} + x^{106} + x^{98} + x^{94} + x^{92} + x^{88} + x^{84} + x^{80} + x^{76} + x^{74} + x^{72} \\
& + x^{66} + x^{64} + x^{62} + x^{60} + x^{52} + x^{50} + x^{46} + x^{40} + x^{32} + x^{30} + x^{26} \\
& + x^{20} + x^{18}, \\
p_9(x) &= x^{135} + x^{132} + x^{131} + x^{129} + x^{126} + x^{125} + x^{119} + x^{116} + x^{113} + x^{112} \\
& + x^{111} + x^{110} + x^{109} + x^{107} + x^{104} + x^{103} + x^{101} + x^{98} + x^{97} + x^{87} \\
& + x^{84} + x^{83} + x^{79} + x^{76} + x^{75} + x^{73} + x^{70} + x^{69} + x^{65} + x^{63} + x^{62} \\
& + x^{61} + x^{60} + x^{57} + x^{56} + x^{55} + x^{54} + x^{53} + x^{49} + x^{47} + x^{46} + x^{44} \\
& + x^{42} + x^{40} + x^{38} + x^{36} + x^{35} + x^{34} + x^{33} + x^{27} + x^{25} + x^{24} + x^{23} \\
& + x^{22} + x^{21} + x^{19} + x^{16} + x^{13} + x^{12} + x^{11} + x^{10} + x^9, \\
p_{12}(x) &= x^{132} + x^{128} + x^{116} + x^{112} + x^{68} + x^{64} + x^{52} + x^{48} + x^{20} + x^{16} + x^4 \\
& + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 1, 0, 0, 0, 1, 0, 1, 1, 1, 0, 1, 0]$.

For $\phi(W^e, 1, 1)$:

$$\begin{aligned}
p_0(x) &= x^{150} + x^{144} + x^{142} + x^{138} + x^{130} + x^{120} + x^{114} + x^{110} + x^{108} + x^{106} \\
& + x^{102} + x^{98} + x^{96} + x^{94} + x^{92} + x^{90} + x^{86} + x^{80} + x^{78} + x^{76} + x^{74} \\
& + x^{70} + x^{62} + x^{60} + x^{58} + x^{56} + x^{54} + x^{50} + x^{42}, \\
p_3(x) &= x^{138} + x^{136} + x^{128} + x^{112} + x^{106} + x^{104} + x^{96} + x^{88} + x^{82} + x^{74} \\
& + x^{72} + x^{42} + x^{32} + x^{18}, \\
p_6(x) &= x^{136} + x^{132} + x^{130} + x^{128} + x^{126} + x^{122} + x^{120} + x^{116} + x^{114} + x^{110} \\
& + x^{106} + x^{102} + x^{94} + x^{76} + x^{74} + x^{72} + x^{70} + x^{68} + x^{64} + x^{62} + x^{60} \\
& + x^{54} + x^{52} + x^{44} + x^{38} + x^{34} + x^{30} + x^{28} + x^{26} + x^{22} + x^{18} + x^{16} \\
& + x^{14} + x^8 + x^6 + 1, \\
p_9(x) &= x^{144} + x^{138} + x^{124} + x^{116} + x^{112} + x^{110} + x^{108} + x^{106} + x^{100} + x^{96} \\
& + x^{94} + x^{88} + x^{82} + x^{80} + x^{74} + x^{68} + x^{54} + x^{48} + x^{46} + x^{42} + x^{40} \\
& + x^{38} + x^{32} + x^{30} + x^{28} + x^{24} + x^{22} + x^{20} + x^{18} + x^{12} + x^8 + x^6, \\
p_{12}(x) &= x^{144} + x^{112} + x^{80} + x^{48} + x^{32} + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 0, 0, 0, 1, 0, 0, 0, 0, 0, 1, 0]$.

For $\phi(M^o, 0, 0)$:

$$\begin{aligned}
p_0(x) &= x^{147} + x^{144} + x^{141} + x^{140} + x^{139} + x^{138} + x^{136} + x^{135} + x^{134} + x^{133} \\
&\quad + x^{132} + x^{130} + x^{129} + x^{126} + x^{125} + x^{123} + x^{122} + x^{121} + x^{120} \\
&\quad + x^{119} + x^{118} + x^{113} + x^{112} + x^{111} + x^{110} + x^{109} + x^{104} + x^{103} \\
&\quad + x^{102} + x^{99} + x^{98} + x^{97} + x^{93} + x^{92} + x^{91} + x^{90} + x^{86} + x^{83} + x^{79} \\
&\quad + x^{77} + x^{75} + x^{70} + x^{68} + x^{67} + x^{64} + x^{63} + x^{62} + x^{61} + x^{60} + x^{57} \\
&\quad + x^{52} + x^{51} + x^{50} + x^{49} + x^{48} + x^{44} + x^{43} + x^{42} + x^{40} + x^{38} + x^{36}, \\
p_3(x) &= x^{144} + x^{141} + x^{140} + x^{138} + x^{134} + x^{132} + x^{127} + x^{126} + x^{125} + x^{119} \\
&\quad + x^{117} + x^{114} + x^{112} + x^{108} + x^{106} + x^{103} + x^{102} + x^{101} + x^{96} + x^{95} \\
&\quad + x^{92} + x^{90} + x^{88} + x^{86} + x^{82} + x^{80} + x^{78} + x^{77} + x^{76} + x^{70} + x^{68} \\
&\quad + x^{64} + x^{63} + x^{62} + x^{60} + x^{55} + x^{54} + x^{53} + x^{52} + x^{50} + x^{48} + x^{47} \\
&\quad + x^{45} + x^{44} + x^{40} + x^{38} + x^{37} + x^{36} + x^{34} + x^{31} + x^{30} + x^{28} + x^{26} \\
&\quad + x^{24}, \\
p_6(x) &= x^{144} + x^{141} + x^{140} + x^{138} + x^{134} + x^{132} + x^{129} + x^{127} + x^{125} + x^{122} \\
&\quad + x^{119} + x^{115} + x^{114} + x^{113} + x^{110} + x^{108} + x^{107} + x^{106} + x^{101} \\
&\quad + x^{92} + x^{90} + x^{88} + x^{87} + x^{86} + x^{84} + x^{82} + x^{79} + x^{78} + x^{77} + x^{73} \\
&\quad + x^{72} + x^{71} + x^{66} + x^{65} + x^{61} + x^{59} + x^{58} + x^{57} + x^{50} + x^{49} + x^{48} \\
&\quad + x^{47} + x^{45} + x^{43} + x^{41} + x^{40} + x^{39} + x^{38} + x^{37} + x^{30} + x^{29} + x^{28} \\
&\quad + x^{27} + x^{26} + x^{25} + x^{23} + x^{22} + x^{20} + x^{19} + x^{16} + x^{14} + x^{12} + x^4 + 1, \\
p_9(x) &= x^{144} + x^{140} + x^{132} + x^{124} + x^{80} + x^{76} + x^{68} + x^{60} + x^{32} + x^{28} + x^{20} \\
&\quad + x^{12}, \\
p_{12}(x) &= x^{144} + x^{140} + x^{128} + x^{124} + x^{116} + x^{112} + x^{80} + x^{76} + x^{64} + x^{60} \\
&\quad + x^{52} + x^{48} + x^{32} + x^{28} + x^{16} + x^{12} + x^4 + 1,
\end{aligned}$$

and the initial terms are $[1, 0, 0, 0, 0, 0, 1, 1, 1, 0, 0, 1, 0, 1, 1, 0]$.

For $\phi(M^o, 0, 1)$:

$$\begin{aligned}
p_0(x) &= x^{152} + x^{143} + x^{140} + x^{139} + x^{138} + x^{137} + x^{136} + x^{135} + x^{133} + x^{131} \\
&\quad + x^{130} + x^{129} + x^{128} + x^{126} + x^{124} + x^{123} + x^{121} + x^{120} + x^{119} \\
&\quad + x^{118} + x^{116} + x^{114} + x^{111} + x^{109} + x^{107} + x^{106} + x^{105} + x^{100} \\
&\quad + x^{96} + x^{95} + x^{93} + x^{92} + x^{90} + x^{88} + x^{86} + x^{85} + x^{84} + x^{82} + x^{81} \\
&\quad + x^{79} + x^{78} + x^{77} + x^{76} + x^{75} + x^{73} + x^{72} + x^{71} + x^{65} + x^{62} + x^{61} \\
&\quad + x^{58} + x^{55} + x^{52} + x^{49} + x^{47} + x^{46} + x^{43} + x^{42} + x^{40} + x^{39}, \\
p_3(x) &= x^{131} + x^{128} + x^{124} + x^{120} + x^{116} + x^{115} + x^{107} + x^{104} + x^{100} + x^{96} \\
&\quad + x^{92} + x^{91} + x^{67} + x^{64} + x^{60} + x^{56} + x^{52} + x^{48} + x^{44} + x^{43} + x^{35} \\
&\quad + x^{32} + x^{28} + x^{27},
\end{aligned}$$

$$\begin{aligned}
p_6(x) &= x^{134} + x^{128} + x^{122} + x^{120} + x^{118} + x^{116} + x^{108} + x^{106} + x^{94} + x^{88} \\
&\quad + x^{80} + x^{72} + x^{70} + x^{66} + x^{60} + x^{58} + x^{54} + x^{50} + x^{48} + x^{44} + x^{42} \\
&\quad + x^{40} + x^{38} + x^{34} + x^{30} + x^{28} + x^{24} + x^{22} + x^{20} + x^{18}, \\
p_9(x) &= x^{137} + x^{134} + x^{131} + x^{130} + x^{128} + x^{126} + x^{124} + x^{123} + x^{122} + x^{121} \\
&\quad + x^{73} + x^{70} + x^{67} + x^{66} + x^{64} + x^{62} + x^{60} + x^{59} + x^{58} + x^{57} + x^{25} \\
&\quad + x^{22} + x^{19} + x^{18} + x^{16} + x^{14} + x^{12} + x^{11} + x^{10} + x^9, \\
p_{12}(x) &= x^{140} + x^{124} + x^{112} + x^{76} + x^{60} + x^{48} + x^{28} + x^{12} + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 1, 0, 0, 0, 0, 0, 1, 0, 1, 0, 0, 0, 0]$.

For $\phi(M^o, 1, 0)$:

$$\begin{aligned}
p_0(x) &= x^{150} + x^{147} + x^{144} + x^{143} + x^{142} + x^{139} + x^{136} + x^{133} + x^{132} + x^{131} \\
&\quad + x^{130} + x^{128} + x^{127} + x^{124} + x^{122} + x^{117} + x^{115} + x^{109} + x^{107} \\
&\quad + x^{106} + x^{103} + x^{102} + x^{99} + x^{96} + x^{95} + x^{94} + x^{93} + x^{90} + x^{88} + x^{87} \\
&\quad + x^{84} + x^{83} + x^{82} + x^{81} + x^{78} + x^{76} + x^{73} + x^{72} + x^{71} + x^{70} + x^{68} \\
&\quad + x^{67} + x^{66} + x^{65} + x^{63} + x^{62} + x^{60} + x^{57} + x^{56} + x^{55} + x^{52} + x^{51} \\
&\quad + x^{50} + x^{48} + x^{47} + x^{46} + x^{44} + x^{43} + x^{42} + x^{40} + x^{39}, \\
p_3(x) &= x^{147} + x^{144} + x^{140} + x^{136} + x^{132} + x^{128} + x^{124} + x^{123} + x^{115} + x^{112} \\
&\quad + x^{108} + x^{104} + x^{100} + x^{96} + x^{92} + x^{91} + x^{83} + x^{80} + x^{76} + x^{72} + x^{68} \\
&\quad + x^{67} + x^{59} + x^{56} + x^{52} + x^{51} + x^{35} + x^{32} + x^{28} + x^{27}, \\
p_6(x) &= x^{150} + x^{144} + x^{138} + x^{136} + x^{132} + x^{128} + x^{124} + x^{120} + x^{118} + x^{116} \\
&\quad + x^{110} + x^{108} + x^{106} + x^{104} + x^{96} + x^{94} + x^{86} + x^{82} + x^{80} + x^{76} \\
&\quad + x^{74} + x^{72} + x^{64} + x^{56} + x^{48} + x^{46} + x^{42} + x^{36} + x^{30} + x^{28} + x^{24} \\
&\quad + x^{22} + x^{20} + x^{18}, \\
p_9(x) &= x^{153} + x^{150} + x^{147} + x^{146} + x^{144} + x^{142} + x^{140} + x^{139} + x^{138} + x^{134} \\
&\quad + x^{131} + x^{130} + x^{128} + x^{126} + x^{124} + x^{123} + x^{122} + x^{121} + x^{89} + x^{86} \\
&\quad + x^{83} + x^{82} + x^{80} + x^{78} + x^{76} + x^{75} + x^{74} + x^{70} + x^{67} + x^{66} + x^{64} \\
&\quad + x^{62} + x^{60} + x^{59} + x^{58} + x^{57} + x^{41} + x^{38} + x^{35} + x^{34} + x^{32} + x^{30} \\
&\quad + x^{28} + x^{27} + x^{26} + x^{22} + x^{19} + x^{18} + x^{16} + x^{14} + x^{12} + x^{11} + x^{10} \\
&\quad + x^9, \\
p_{12}(x) &= x^{156} + x^{128} + x^{124} + x^{112} + x^{92} + x^{64} + x^{60} + x^{48} + x^{44} + x^{16} + x^{12} \\
&\quad + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 1, 0, 0, 0, 1, 0, 1, 0, 0, 0, 1, 0]$.

For $\phi(M^o, 1, 1)$:

$$\begin{aligned}
p_0(x) &= x^{147} + x^{139} + x^{135} + x^{133} + x^{132} + x^{128} + x^{127} + x^{123} + x^{121} + x^{117} \\
&\quad + x^{115} + x^{111} + x^{109} + x^{108} + x^{107} + x^{105} + x^{102} + x^{100} + x^{99} + x^{98} \\
&\quad + x^{97} + x^{95} + x^{94} + x^{93} + x^{90} + x^{89} + x^{88} + x^{87} + x^{86} + x^{84} + x^{82}
\end{aligned}$$

$$\begin{aligned}
& + x^{80} + x^{78} + x^{77} + x^{75} + x^{74} + x^{73} + x^{71} + x^{70} + x^{67} + x^{66} + x^{65} \\
& + x^{62} + x^{61} + x^{60} + x^{59} + x^{58} + x^{56} + x^{55} + x^{54} + x^{53} + x^{52} + x^{51} \\
& + x^{50} + x^{47} + x^{46} + x^{42}, \\
p_3(x) &= x^{144} + x^{141} + x^{140} + x^{132} + x^{127} + x^{126} + x^{125} + x^{124} + x^{122} + x^{120} \\
& + x^{119} + x^{118} + x^{117} + x^{116} + x^{110} + x^{108} + x^{106} + x^{103} + x^{102} \\
& + x^{101} + x^{100} + x^{98} + x^{95} + x^{94} + x^{90} + x^{86} + x^{82} + x^{80} + x^{78} + x^{77} \\
& + x^{76} + x^{74} + x^{68} + x^{64} + x^{63} + x^{62} + x^{56} + x^{55} + x^{54} + x^{53} + x^{47} \\
& + x^{46} + x^{45} + x^{42} + x^{37} + x^{36} + x^{31} + x^{26}, \\
p_6(x) &= x^{144} + x^{141} + x^{140} + x^{132} + x^{129} + x^{127} + x^{126} + x^{125} + x^{124} + x^{122} \\
& + x^{120} + x^{119} + x^{118} + x^{116} + x^{115} + x^{114} + x^{113} + x^{107} + x^{104} \\
& + x^{101} + x^{100} + x^{98} + x^{94} + x^{92} + x^{90} + x^{88} + x^{87} + x^{86} + x^{82} + x^{80} \\
& + x^{79} + x^{78} + x^{77} + x^{76} + x^{74} + x^{73} + x^{71} + x^{68} + x^{65} + x^{64} + x^{62} \\
& + x^{61} + x^{59} + x^{58} + x^{57} + x^{56} + x^{52} + x^{49} + x^{48} + x^{47} + x^{46} + x^{45} \\
& + x^{44} + x^{43} + x^{41} + x^{40} + x^{39} + x^{38} + x^{37} + x^{32} + x^{29} + x^{28} + x^{27} \\
& + x^{25} + x^{24} + x^{23} + x^{22} + x^{20} + x^{19} + x^{18} + x^{16} + x^{14} + x^4 + 1, \\
p_9(x) &= x^{144} + x^{140} + x^{132} + x^{124} + x^{80} + x^{76} + x^{68} + x^{60} + x^{32} + x^{28} + x^{20} \\
& + x^{12}, \\
p_{12}(x) &= x^{144} + x^{140} + x^{128} + x^{124} + x^{116} + x^{112} + x^{80} + x^{76} + x^{64} + x^{60} \\
& + x^{52} + x^{48} + x^{32} + x^{28} + x^{16} + x^{12} + x^4 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 0, 0, 1, 0, 0, 0, 1, 0, 1, 1, 0]$.

For $\phi(W^\circ, 0, 0)$:

$$\begin{aligned}
p_0(x) &= x^{147} + x^{144} + x^{141} + x^{140} + x^{139} + x^{138} + x^{136} + x^{135} + x^{134} + x^{133} \\
& + x^{132} + x^{130} + x^{129} + x^{126} + x^{125} + x^{123} + x^{122} + x^{121} + x^{120} \\
& + x^{119} + x^{118} + x^{113} + x^{112} + x^{111} + x^{110} + x^{109} + x^{104} + x^{103} \\
& + x^{102} + x^{99} + x^{98} + x^{97} + x^{93} + x^{92} + x^{91} + x^{90} + x^{86} + x^{83} + x^{79} \\
& + x^{77} + x^{75} + x^{70} + x^{68} + x^{67} + x^{64} + x^{63} + x^{62} + x^{61} + x^{60} + x^{57} \\
& + x^{52} + x^{51} + x^{50} + x^{49} + x^{48} + x^{44} + x^{43} + x^{42} + x^{40} + x^{38} + x^{36}, \\
p_3(x) &= x^{144} + x^{141} + x^{140} + x^{138} + x^{134} + x^{132} + x^{127} + x^{126} + x^{125} + x^{119} \\
& + x^{117} + x^{114} + x^{112} + x^{108} + x^{106} + x^{103} + x^{102} + x^{101} + x^{96} + x^{95} \\
& + x^{92} + x^{90} + x^{88} + x^{86} + x^{82} + x^{80} + x^{78} + x^{77} + x^{76} + x^{70} + x^{68} \\
& + x^{64} + x^{63} + x^{62} + x^{60} + x^{55} + x^{54} + x^{53} + x^{52} + x^{50} + x^{48} + x^{47} \\
& + x^{45} + x^{44} + x^{40} + x^{38} + x^{37} + x^{36} + x^{34} + x^{31} + x^{30} + x^{28} + x^{26} \\
& + x^{24}, \\
p_6(x) &= x^{144} + x^{141} + x^{140} + x^{138} + x^{134} + x^{132} + x^{129} + x^{127} + x^{125} + x^{122} \\
& + x^{119} + x^{115} + x^{114} + x^{113} + x^{110} + x^{108} + x^{107} + x^{106} + x^{101}
\end{aligned}$$

$$\begin{aligned}
& + x^{92} + x^{90} + x^{88} + x^{87} + x^{86} + x^{84} + x^{82} + x^{79} + x^{78} + x^{77} + x^{73} \\
& + x^{72} + x^{71} + x^{66} + x^{65} + x^{61} + x^{59} + x^{58} + x^{57} + x^{50} + x^{49} + x^{48} \\
& + x^{47} + x^{45} + x^{43} + x^{41} + x^{40} + x^{39} + x^{38} + x^{37} + x^{30} + x^{29} + x^{28} \\
& + x^{27} + x^{26} + x^{25} + x^{23} + x^{22} + x^{20} + x^{19} + x^{16} + x^{14} + x^{12} + x^4 + 1, \\
p_9(x) &= x^{144} + x^{140} + x^{132} + x^{124} + x^{80} + x^{76} + x^{68} + x^{60} + x^{32} + x^{28} + x^{20} \\
& + x^{12}, \\
p_{12}(x) &= x^{144} + x^{140} + x^{128} + x^{124} + x^{116} + x^{112} + x^{80} + x^{76} + x^{64} + x^{60} \\
& + x^{52} + x^{48} + x^{32} + x^{28} + x^{16} + x^{12} + x^4 + 1,
\end{aligned}$$

and the initial terms are $[1, 0, 0, 0, 0, 0, 1, 1, 1, 0, 0, 1, 0, 1, 1, 0]$.

For $\phi(W^o, 0, 1)$:

$$\begin{aligned}
p_0(x) &= x^{150} + x^{147} + x^{144} + x^{143} + x^{142} + x^{139} + x^{136} + x^{133} + x^{132} + x^{131} \\
& + x^{130} + x^{128} + x^{127} + x^{124} + x^{122} + x^{117} + x^{115} + x^{109} + x^{107} \\
& + x^{106} + x^{103} + x^{102} + x^{99} + x^{96} + x^{95} + x^{94} + x^{93} + x^{90} + x^{88} + x^{87} \\
& + x^{84} + x^{83} + x^{82} + x^{81} + x^{78} + x^{76} + x^{73} + x^{72} + x^{71} + x^{70} + x^{68} \\
& + x^{67} + x^{66} + x^{65} + x^{63} + x^{62} + x^{60} + x^{57} + x^{56} + x^{55} + x^{52} + x^{51} \\
& + x^{50} + x^{48} + x^{47} + x^{46} + x^{44} + x^{43} + x^{42} + x^{40} + x^{39}, \\
p_3(x) &= x^{147} + x^{144} + x^{140} + x^{136} + x^{132} + x^{128} + x^{124} + x^{123} + x^{115} + x^{112} \\
& + x^{108} + x^{104} + x^{100} + x^{96} + x^{92} + x^{91} + x^{83} + x^{80} + x^{76} + x^{72} + x^{68} \\
& + x^{67} + x^{59} + x^{56} + x^{52} + x^{51} + x^{35} + x^{32} + x^{28} + x^{27}, \\
p_6(x) &= x^{150} + x^{144} + x^{138} + x^{136} + x^{132} + x^{128} + x^{124} + x^{120} + x^{118} + x^{116} \\
& + x^{110} + x^{108} + x^{106} + x^{104} + x^{96} + x^{94} + x^{86} + x^{82} + x^{80} + x^{76} \\
& + x^{74} + x^{72} + x^{64} + x^{56} + x^{48} + x^{46} + x^{42} + x^{36} + x^{30} + x^{28} + x^{24} \\
& + x^{22} + x^{20} + x^{18}, \\
p_9(x) &= x^{153} + x^{150} + x^{147} + x^{146} + x^{144} + x^{142} + x^{140} + x^{139} + x^{138} + x^{134} \\
& + x^{131} + x^{130} + x^{128} + x^{126} + x^{124} + x^{123} + x^{122} + x^{121} + x^{89} + x^{86} \\
& + x^{83} + x^{82} + x^{80} + x^{78} + x^{76} + x^{75} + x^{74} + x^{70} + x^{67} + x^{66} + x^{64} \\
& + x^{62} + x^{60} + x^{59} + x^{58} + x^{57} + x^{41} + x^{38} + x^{35} + x^{34} + x^{32} + x^{30} \\
& + x^{28} + x^{27} + x^{26} + x^{22} + x^{19} + x^{18} + x^{16} + x^{14} + x^{12} + x^{11} + x^{10} \\
& + x^9, \\
p_{12}(x) &= x^{156} + x^{128} + x^{124} + x^{112} + x^{92} + x^{64} + x^{60} + x^{48} + x^{44} + x^{16} + x^{12} \\
& + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 1, 0, 0, 0, 1, 0, 1, 0, 0, 0, 1, 0]$.

For $\phi(W^o, 1, 0)$:

$$\begin{aligned}
p_0(x) &= x^{152} + x^{143} + x^{140} + x^{139} + x^{138} + x^{137} + x^{136} + x^{135} + x^{133} + x^{131} \\
& + x^{130} + x^{129} + x^{128} + x^{126} + x^{124} + x^{123} + x^{121} + x^{120} + x^{119}
\end{aligned}$$

$$\begin{aligned}
& + x^{118} + x^{116} + x^{114} + x^{111} + x^{109} + x^{107} + x^{106} + x^{105} + x^{100} \\
& + x^{96} + x^{95} + x^{93} + x^{92} + x^{90} + x^{88} + x^{86} + x^{85} + x^{84} + x^{82} + x^{81} \\
& + x^{79} + x^{78} + x^{77} + x^{76} + x^{75} + x^{73} + x^{72} + x^{71} + x^{65} + x^{62} + x^{61} \\
& + x^{58} + x^{55} + x^{52} + x^{49} + x^{47} + x^{46} + x^{43} + x^{42} + x^{40} + x^{39}, \\
p_3(x) &= x^{131} + x^{128} + x^{124} + x^{120} + x^{116} + x^{115} + x^{107} + x^{104} + x^{100} + x^{96} \\
& + x^{92} + x^{91} + x^{67} + x^{64} + x^{60} + x^{56} + x^{52} + x^{48} + x^{44} + x^{43} + x^{35} \\
& + x^{32} + x^{28} + x^{27}, \\
p_6(x) &= x^{134} + x^{128} + x^{122} + x^{120} + x^{118} + x^{116} + x^{108} + x^{106} + x^{94} + x^{88} \\
& + x^{80} + x^{72} + x^{70} + x^{66} + x^{60} + x^{58} + x^{54} + x^{50} + x^{48} + x^{44} + x^{42} \\
& + x^{40} + x^{38} + x^{34} + x^{30} + x^{28} + x^{24} + x^{22} + x^{20} + x^{18}, \\
p_9(x) &= x^{137} + x^{134} + x^{131} + x^{130} + x^{128} + x^{126} + x^{124} + x^{123} + x^{122} + x^{121} \\
& + x^{73} + x^{70} + x^{67} + x^{66} + x^{64} + x^{62} + x^{60} + x^{59} + x^{58} + x^{57} + x^{25} \\
& + x^{22} + x^{19} + x^{18} + x^{16} + x^{14} + x^{12} + x^{11} + x^{10} + x^9, \\
p_{12}(x) &= x^{140} + x^{124} + x^{112} + x^{76} + x^{60} + x^{48} + x^{28} + x^{12} + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 1, 0, 0, 0, 0, 0, 1, 0, 1, 0, 0, 0, 0]$.

For $\phi(W^o, 1, 1)$:

$$\begin{aligned}
p_0(x) &= x^{147} + x^{139} + x^{135} + x^{133} + x^{132} + x^{128} + x^{127} + x^{123} + x^{121} + x^{117} \\
& + x^{115} + x^{111} + x^{109} + x^{108} + x^{107} + x^{105} + x^{102} + x^{100} + x^{99} + x^{98} \\
& + x^{97} + x^{95} + x^{94} + x^{93} + x^{90} + x^{89} + x^{88} + x^{87} + x^{86} + x^{84} + x^{82} \\
& + x^{80} + x^{78} + x^{77} + x^{75} + x^{74} + x^{73} + x^{71} + x^{70} + x^{67} + x^{66} + x^{65} \\
& + x^{62} + x^{61} + x^{60} + x^{59} + x^{58} + x^{56} + x^{55} + x^{54} + x^{53} + x^{52} + x^{51} \\
& + x^{50} + x^{47} + x^{46} + x^{42}, \\
p_3(x) &= x^{144} + x^{141} + x^{140} + x^{132} + x^{127} + x^{126} + x^{125} + x^{124} + x^{122} + x^{120} \\
& + x^{119} + x^{118} + x^{117} + x^{116} + x^{110} + x^{108} + x^{106} + x^{103} + x^{102} \\
& + x^{101} + x^{100} + x^{98} + x^{95} + x^{94} + x^{90} + x^{86} + x^{82} + x^{80} + x^{78} + x^{77} \\
& + x^{76} + x^{74} + x^{68} + x^{64} + x^{63} + x^{62} + x^{56} + x^{55} + x^{54} + x^{53} + x^{47} \\
& + x^{46} + x^{45} + x^{42} + x^{37} + x^{36} + x^{31} + x^{26}, \\
p_6(x) &= x^{144} + x^{141} + x^{140} + x^{132} + x^{129} + x^{127} + x^{126} + x^{125} + x^{124} + x^{122} \\
& + x^{120} + x^{119} + x^{118} + x^{116} + x^{115} + x^{114} + x^{113} + x^{107} + x^{104} \\
& + x^{101} + x^{100} + x^{98} + x^{94} + x^{92} + x^{90} + x^{88} + x^{87} + x^{86} + x^{82} + x^{80} \\
& + x^{79} + x^{78} + x^{77} + x^{76} + x^{74} + x^{73} + x^{71} + x^{68} + x^{65} + x^{64} + x^{62} \\
& + x^{61} + x^{59} + x^{58} + x^{57} + x^{56} + x^{52} + x^{49} + x^{48} + x^{47} + x^{46} + x^{45} \\
& + x^{44} + x^{43} + x^{41} + x^{40} + x^{39} + x^{38} + x^{37} + x^{32} + x^{29} + x^{28} + x^{27} \\
& + x^{25} + x^{24} + x^{23} + x^{22} + x^{20} + x^{19} + x^{18} + x^{16} + x^{14} + x^4 + 1, \\
p_9(x) &= x^{144} + x^{140} + x^{132} + x^{124} + x^{80} + x^{76} + x^{68} + x^{60} + x^{32} + x^{28} + x^{20}
\end{aligned}$$

$$\begin{aligned}
& + x^{12}, \\
p_{12}(x) = & x^{144} + x^{140} + x^{128} + x^{124} + x^{116} + x^{112} + x^{80} + x^{76} + x^{64} + x^{60} \\
& + x^{52} + x^{48} + x^{32} + x^{28} + x^{16} + x^{12} + x^4 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 0, 0, 1, 0, 0, 0, 1, 0, 1, 1, 0]$.

Below is the transition function and output function of an 2-automaton that generates $T = M_{0,0}^o$:

Transition function $(n, j) \mapsto \delta(n, j)$ ($\Lambda(n) := [\delta(n, 0), \delta(n, 1)]$):

n	$\Lambda(n)$	n	$\Lambda(n)$	n	$\Lambda(n)$	n	$\Lambda(n)$	n	$\Lambda(n)$
0	[1, 2]	76	[118, 95]	152	[210, 211]	228	[282, 30]	304	[145, 139]
1	[3, 4]	77	[48, 119]	153	[212, 213]	229	[84, 147]	305	[263, 341]
2	[5, 6]	78	[120, 53]	154	[214, 193]	230	[24, 162]	306	[183, 164]
3	[7, 8]	79	[121, 122]	155	[95, 215]	231	[283, 86]	307	[79, 77]
4	[9, 9]	80	[123, 124]	156	[216, 217]	232	[284, 174]	308	[343, 158]
5	[10, 11]	81	[125, 126]	157	[218, 187]	233	[285, 8]	309	[159, 83]
6	[12, 2]	82	[127, 128]	158	[219, 204]	234	[286, 174]	310	[69, 140]
7	[13, 14]	83	[129, 130]	159	[220, 88]	235	[142, 155]	311	[285, 63]
8	[15, 16]	84	[131, 132]	160	[221, 222]	236	[287, 32]	312	[272, 179]
9	[17, 18]	85	[133, 45]	161	[39, 223]	237	[266, 61]	313	[344, 65]
10	[19, 20]	86	[134, 135]	162	[37, 134]	238	[280, 138]	314	[7, 76]
11	[11, 11]	87	[136, 65]	163	[224, 195]	239	[274, 60]	315	[345, 142]
12	[21, 22]	88	[75, 63]	164	[101, 225]	240	[288, 35]	316	[346, 174]
13	[23, 24]	89	[137, 61]	165	[226, 121]	241	[175, 179]	317	[282, 32]
14	[25, 26]	90	[59, 138]	166	[227, 53]	242	[137, 185]	318	[141, 63]
15	[27, 26]	91	[139, 140]	167	[228, 123]	243	[65, 35]	319	[347, 85]
16	[28, 26]	92	[141, 142]	168	[229, 117]	244	[273, 86]	320	[348, 164]
17	[29, 29]	93	[142, 143]	169	[14, 230]	245	[162, 255]	321	[158, 60]
18	[30, 30]	94	[144, 145]	170	[231, 188]	246	[289, 164]	322	[156, 29]
19	[31, 32]	95	[146, 147]	171	[232, 233]	247	[31, 30]	323	[9, 33]
20	[33, 32]	96	[148, 147]	172	[234, 92]	248	[173, 290]	324	[269, 65]
21	[34, 35]	97	[146, 149]	173	[235, 217]	249	[291, 153]	325	[349, 155]
22	[33, 9]	98	[150, 85]	174	[236, 44]	250	[275, 151]	326	[350, 24]
23	[36, 37]	99	[84, 151]	175	[237, 238]	251	[275, 81]	327	[25, 261]
24	[38, 39]	100	[152, 81]	176	[239, 222]	252	[285, 142]	328	[257, 35]
25	[40, 41]	101	[82, 153]	177	[240, 119]	253	[142, 169]	329	[287, 30]
26	[42, 43]	102	[154, 155]	178	[241, 242]	254	[292, 134]	330	[171, 81]
27	[44, 45]	103	[156, 11]	179	[88, 243]	255	[41, 293]	331	[28, 153]
28	[46, 47]	104	[12, 67]	180	[244, 245]	256	[251, 294]	332	[78, 32]
29	[20, 48]	105	[157, 158]	181	[246, 49]	257	[295, 251]	333	[72, 351]
30	[18, 49]	106	[159, 153]	182	[247, 248]	258	[106, 296]	334	[352, 73]
31	[50, 51]	107	[160, 161]	183	[249, 250]	259	[297, 91]	335	[353, 252]
32	[51, 52]	108	[162, 161]	184	[126, 251]	260	[104, 111]	336	[193, 354]
33	[53, 17]	109	[85, 59]	185	[252, 253]	261	[298, 299]	337	[355, 39]
34	[54, 55]	110	[163, 63]	186	[254, 85]	262	[300, 211]	338	[356, 190]

35	[56, 57]	111	[64, 34]	187	[86, 161]	263	[198, 301]	339	[357, 305]
36	[58, 59]	112	[65, 164]	188	[86, 255]	264	[130, 97]	340	[358, 101]
37	[60, 61]	113	[165, 29]	189	[65, 256]	265	[302, 18]	341	[211, 359]
38	[62, 63]	114	[166, 167]	190	[136, 257]	266	[303, 245]	342	[360, 14]
39	[64, 65]	115	[168, 169]	191	[257, 258]	267	[304, 305]	343	[361, 301]
40	[66, 67]	116	[170, 145]	192	[259, 145]	268	[306, 307]	344	[202, 55]
41	[68, 65]	117	[171, 149]	193	[171, 147]	269	[308, 309]	345	[299, 16]
42	[69, 61]	118	[172, 147]	194	[260, 26]	270	[310, 198]	346	[362, 48]
43	[70, 71]	119	[32, 9]	195	[28, 261]	271	[311, 128]	347	[363, 242]
44	[72, 73]	120	[10, 29]	196	[24, 59]	272	[312, 89]	348	[364, 106]
45	[7, 63]	121	[77, 11]	197	[262, 139]	273	[313, 314]	349	[365, 327]
46	[74, 73]	122	[11, 79]	198	[263, 140]	274	[315, 92]	350	[366, 90]
47	[75, 76]	123	[173, 174]	199	[264, 142]	275	[316, 252]	351	[367, 210]
48	[32, 32]	124	[141, 8]	200	[265, 182]	276	[317, 22]	352	[368, 323]
49	[29, 77]	125	[175, 161]	201	[266, 140]	277	[318, 41]	353	[7, 142]
50	[78, 30]	126	[176, 85]	202	[267, 158]	278	[319, 250]	354	[145, 60]
51	[79, 29]	127	[177, 67]	203	[268, 182]	279	[320, 17]	355	[369, 67]
52	[30, 33]	128	[68, 34]	204	[269, 183]	280	[222, 37]	356	[68, 257]
53	[9, 30]	129	[178, 179]	205	[270, 140]	281	[321, 238]	357	[339, 140]
54	[80, 81]	130	[180, 145]	206	[267, 262]	282	[322, 323]	358	[370, 158]
55	[82, 83]	131	[181, 182]	207	[271, 59]	283	[324, 47]	359	[257, 164]
56	[84, 81]	132	[136, 183]	208	[272, 161]	284	[325, 122]	360	[159, 26]
57	[85, 86]	133	[184, 65]	209	[176, 273]	285	[326, 327]	361	[371, 139]
58	[87, 88]	134	[59, 179]	210	[166, 182]	286	[328, 20]	362	[372, 35]
59	[89, 90]	135	[60, 185]	211	[269, 257]	287	[329, 307]	363	[71, 59]
60	[91, 91]	136	[186, 56]	212	[160, 179]	288	[330, 309]	364	[146, 81]
61	[92, 93]	137	[187, 188]	213	[180, 274]	289	[331, 99]	365	[82, 26]
62	[94, 95]	138	[45, 189]	214	[275, 147]	290	[332, 333]	366	[373, 60]
63	[96, 97]	139	[108, 187]	215	[145, 263]	291	[334, 314]	367	[276, 29]
64	[98, 99]	140	[190, 191]	216	[276, 11]	292	[335, 60]	368	[63, 169]
65	[100, 101]	141	[192, 193]	217	[77, 79]	293	[63, 336]	369	[374, 121]
66	[102, 11]	142	[194, 195]	218	[277, 86]	294	[85, 280]	370	[375, 108]
67	[103, 104]	143	[16, 196]	219	[278, 257]	295	[337, 153]	371	[376, 233]
68	[105, 106]	144	[197, 198]	220	[279, 73]	296	[158, 137]	372	[377, 56]
69	[107, 108]	145	[199, 124]	221	[270, 61]	297	[338, 139]	373	[378, 132]
70	[109, 89]	146	[200, 190]	222	[280, 179]	298	[339, 61]	374	[76, 155]
71	[110, 111]	147	[201, 202]	223	[63, 155]	299	[70, 24]	375	[24, 86]
72	[112, 51]	148	[203, 204]	224	[281, 24]	300	[340, 257]	376	[75, 142]
73	[113, 114]	149	[205, 206]	225	[158, 139]	301	[139, 341]	377	[25, 153]
74	[115, 52]	150	[207, 135]	226	[165, 11]	302	[342, 169]	378	[64, 257]
75	[116, 117]	151	[208, 209]	227	[8, 169]	303	[178, 161]		

Output function $n \mapsto \tau(n)$:

n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$
0	0	55	0	110	1	165	0	220	1	275	1	330	0
1	0	56	0	111	1	166	1	221	1	276	1	331	0
2	0	57	0	112	1	167	1	222	1	277	0	332	0
3	0	58	0	113	0	168	0	223	0	278	1	333	1
4	0	59	0	114	1	169	1	224	1	279	1	334	0
5	0	60	1	115	0	170	1	225	1	280	1	335	0
6	0	61	0	116	1	171	0	226	0	281	1	336	0
7	0	62	0	117	0	172	1	227	1	282	1	337	1
8	1	63	0	118	1	173	0	228	1	283	1	338	0
9	0	64	1	119	1	174	1	229	0	284	0	339	0
10	0	65	1	120	0	175	0	230	0	285	1	340	0
11	0	66	1	121	1	176	0	231	1	286	1	341	1
12	0	67	1	122	0	177	0	232	0	287	1	342	1
13	0	68	0	123	0	178	0	233	1	288	0	343	1
14	1	69	1	124	0	179	1	234	1	289	0	344	1
15	1	70	0	125	0	180	1	235	0	290	0	345	0
16	0	71	1	126	0	181	0	236	1	291	0	346	1
17	0	72	1	127	0	182	0	237	0	292	0	347	1
18	1	73	0	128	0	183	0	238	1	293	0	348	1
19	0	74	0	129	0	184	0	239	0	294	0	349	0
20	0	75	1	130	1	185	1	240	0	295	1	350	1
21	0	76	1	131	0	186	0	241	0	296	1	351	1
22	0	77	1	132	0	187	0	242	0	297	0	352	0
23	0	78	0	133	0	188	0	243	1	298	0	353	0
24	0	79	1	134	0	189	1	244	1	299	0	354	1
25	1	80	0	135	1	190	0	245	1	300	0	355	1
26	1	81	0	136	0	191	1	246	0	301	1	356	0
27	1	82	0	137	0	192	0	247	0	302	1	357	0
28	0	83	0	138	0	193	0	248	0	303	0	358	0
29	0	84	0	139	1	194	0	249	0	304	1	359	1
30	1	85	0	140	0	195	0	250	1	305	0	360	1
31	0	86	0	141	0	196	0	251	1	306	0	361	1
32	1	87	0	142	0	197	0	252	1	307	1	362	1
33	0	88	1	143	0	198	0	253	0	308	1	363	1
34	0	89	0	144	0	199	1	254	0	309	1	364	1
35	0	90	0	145	1	200	1	255	0	310	1	365	0
36	0	91	1	146	1	201	0	256	1	311	1	366	1
37	1	92	0	147	0	202	1	257	1	312	1	367	1
38	0	93	0	148	0	203	0	258	1	313	1	368	0
39	1	94	0	149	1	204	1	259	0	314	0	369	1
40	1	95	1	150	1	205	1	260	0	315	0	370	0
41	0	96	0	151	1	206	1	261	0	316	1	371	1
42	1	97	1	152	1	207	1	262	0	317	1	372	1

43	0	98	1	153	1	208	1	263	0	318	0	373	1
44	1	99	0	154	1	209	0	264	1	319	1	374	1
45	0	100	1	155	1	210	1	265	1	320	1	375	0
46	0	101	0	156	1	211	1	266	0	321	1	376	1
47	1	102	1	157	0	212	1	267	1	322	1	377	1
48	1	103	1	158	1	213	1	268	0	323	0	378	1
49	0	104	0	159	1	214	1	269	1	324	1		
50	0	105	0	160	1	215	1	270	1	325	0		
51	1	106	1	161	1	216	1	271	1	326	1		
52	1	107	1	162	1	217	1	272	1	327	1		
53	0	108	1	163	1	218	0	273	1	328	1		
54	0	109	0	164	0	219	1	274	0	329	1		

UNIVERSITÉ DE STRASBOURG, CNRS, IRMA UMR 7501, F-67000 STRASBOURG, FRANCE
Email address: `guoniu.han@unistra.fr`

SCHOOL OF MATHEMATICS AND STATISTICS, HUAZHONG UNIVERSITY OF SCIENCE AND TECHNOLOGY, WUHAN, PR CHINA
Email address: `huyining@hust.edu.cn`