

PROOF OF THE CONTINUED FRACTION CONJECTURE FOR

$$(a, b) = (z^2, z^5 + z^2)$$

GUO-NIU HAN* AND YINING HU*

ABSTRACT. The continued fraction conjecture claims that the continued fraction defined by the (a, b) -Thue-Morse sequence is algebraic over $\mathbb{F}_2(x)$ for all pairs of distinct elements (a, b) in $\mathbb{F}_2[x] \setminus \mathbb{F}_2$. In this paper we prove the conjecture for $(a, b) = (z^2, z^5 + z^2)$.

1. INTRODUCTION

Let $\mathbb{F}_2$ be the finite field of cardinality 2. Given a sequence of polynomials $\mathbf{a} = (a_0(z), a_1(z), a_2(z), \dots)$ of elements from $\mathbb{F}_2[z] \setminus \mathbb{F}_2$, we define the infinite continued fraction

$$(1.1) \quad \text{CF}(\mathbf{a}(z)) = \frac{1}{a_0(z) + \frac{1}{a_1(z) + \frac{1}{a_2(z) + \frac{1}{\ddots}}}}$$

as the limit of the finite continued fractions

$$(1.2) \quad \text{CF}_n(\mathbf{a}(z)) = \frac{1}{a_0(z) + \frac{1}{a_1(z) + \frac{1}{\ddots + \frac{1}{a_n(z)}}}} \in \mathbb{F}_2((1/z)).$$

Conversely, each power series in $\mathbb{F}_2[[1/z]]$ without constant term

$$(1.3) \quad c_{-1}z^{-1} + c_{-2}z^{-2} + c_{-3}z^{-3} + \dots$$

admits a continued fraction expansion of form (1.1).

Let a and b be two distinct elements from $\mathbb{F}_2[z] \setminus \mathbb{F}_2$. We consider the continued fraction (1.1) associated with the (a, b) -Thue-Morse sequence

$$\mathbf{t} = (t_0(z), t_1(z), t_2(z), \dots) = (a, b, b, a, b, a, a, b, \dots)$$

that we shall denote

$$(1.4) \quad \text{CF}(a, b) := \text{CF}(\mathbf{t}).$$

Date: 9/04/2020 17:30:32 .

2010 Mathematics Subject Classification. 11B85, 11J70, 11B50, 11Y65, 05A15.

Key words and phrases. automatic sequence, continued fraction, Thue-Morse sequence.

* This paper was automatically generated from a computer program developed by the authors, with 5.3 minutes CPU time used.

Theorem 1.1. *When $(a, b) = (z^2, z^5 + z^2)$, the two power series $\text{CF}(a, b)$ and $\text{CF}(b, a)$ are algebraic over $\mathbb{F}_2(z)$, with minimal polynomials of the form*

$$p_4(z)y^4 + p_3(z)y^3 + p_2(z)y^2 + p_1(z)y + p_0(z) = 0$$

where, for $\text{CF}(a, b)$,

$$\begin{aligned} p_0(z) &= z^9 + z^8 + z^6 + z^5 + z^3 + 1, \\ p_1(z) &= z^{14} + z^{11} + z^{10} + z^8 + z^7 + z^5, \\ p_2(z) &= z^{16} + z^{12} + z^6 + z^4, \\ p_3(z) &= z^{14} + z^{11} + z^{10} + z^8 + z^7 + z^5, \\ p_4(z) &= z^{12} + z^9 + z^5 + z^3 + 1, \end{aligned}$$

and for $\text{CF}(b, a)$,

$$\begin{aligned} p_0(z) &= z^9 + z^6 + z^5 + z^3 + 1, \\ p_1(z) &= z^{14} + z^{11} + z^{10} + z^8 + z^7 + z^5, \\ p_2(z) &= z^{16} + z^{12} + z^6 + z^4, \\ p_3(z) &= z^{14} + z^{11} + z^{10} + z^8 + z^7 + z^5, \\ p_4(z) &= z^{12} + z^9 + z^8 + z^5 + z^3 + 1. \end{aligned}$$

2. PROOF

Define the sequences of polynomials $(P_n(z))_{n \geq 0}$ and $(Q_n(z))_{n \geq 0}$ by

$$\begin{pmatrix} P_n(z) & Q_n(z) \\ P_{n-1}(z) & Q_{n-1}(z) \end{pmatrix} := \begin{pmatrix} t_n(z) & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} t_{n-1}(z) & 1 \\ 1 & 0 \end{pmatrix} \cdots \begin{pmatrix} t_0(z) & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}.$$

By the basic properties of continued fractions, we have

$$(2.1) \quad \text{CF}_n(a, b) = \frac{P_n(z)}{Q_n(z)} \in \mathbb{F}_2((1/z)).$$

As formula (2.1) is not efficient for calculating $\text{CF}_n(a, b)$, a fast method is to be derived and described as follows. Define

$$\begin{aligned} M_n(x) &= x^{\deg(t_{2^n-1})} \begin{pmatrix} t_{2^n-1}(1/x) & 1 \\ 1 & 0 \end{pmatrix} \times x^{\deg(t_{2^n-2})} \begin{pmatrix} t_{2^n-2}(1/x) & 1 \\ 1 & 0 \end{pmatrix} \times \\ &\quad \cdots \times x^{\deg(t_0)} \begin{pmatrix} t_0(1/x) & 1 \\ 1 & 0 \end{pmatrix}, \\ W_n(x) &= x^{\deg(\bar{t}_{2^n-1})} \begin{pmatrix} \bar{t}_{2^n-1}(1/x) & 1 \\ 1 & 0 \end{pmatrix} \times x^{\deg(\bar{t}_{2^n-2})} \begin{pmatrix} \bar{t}_{2^n-2}(1/x) & 1 \\ 1 & 0 \end{pmatrix} \times \\ &\quad \cdots \times x^{\deg(\bar{t}_0)} \begin{pmatrix} \bar{t}_0(1/x) & 1 \\ 1 & 0 \end{pmatrix}, \end{aligned}$$

where $\bar{t}$ is the (b, a) -Thue-Morse sequence. By the properties of the Thue-Morse sequence, we have for $n \geq 0$

$$\begin{aligned} M_{n+1}(x) &= W_n(x) \cdot M_n(x), \\ W_{n+1}(x) &= M_n(x) \cdot W_n(x). \end{aligned}$$

Let $x := 1/z$. For each non-zero polynomial $P(z)$, define $\tilde{P}(x)$ to be $P(1/x)$. Then,

$$\text{CF}_n(a, b) = \frac{P_n(z)}{Q_n(z)} = \frac{\tilde{P}_n(x)}{\tilde{Q}_n(x)} \in \mathbb{F}_2((x)) = \mathbb{F}_2((1/z)).$$

Comparing the definition of $M_n(x)$ with the definition of $P_n(x)$ and $Q_n(x)$, we see that

$$\begin{aligned} M_n(x)_{0,1} &= x^{d_n} \tilde{P}_{2^n-1}(x), \\ M_n(x)_{0,0} &= x^{d_n} \tilde{Q}_{2^n-1}(x), \end{aligned}$$

for some positive integer d_n . Hence,

$$(2.2) \quad \text{CF}_{2^{2^n}-1}(a, b) = \frac{\tilde{P}_{2^{2^n}-1}(x)}{\tilde{Q}_{2^{2^n}-1}(x)} = \frac{M_{2^n}(x)_{0,1}}{M_{2^n}(x)_{0,0}}.$$

Therefore, by the convergence theorem of continued fraction, the algebraicity of $\text{CF}(a, b)$ will be established if it is shown that both $M_{2^n}(x)_{0,1}$ and $M_{2^n}(x)_{0,0}$ converge to algebraic series in $\mathbb{F}_2[[x]]$.

Actually, we will prove that for all $0 \leq i, j \leq 1$ the four sequences $(M_{2^n}(x)_{i,j})_n$, $(M_{2^{n+1}}(x)_{i,j})_n$, $(W_{2^n}(x)_{i,j})_n$, and $(W_{2^{n+1}}(x)_{i,j})_n$ converge to algebraic series in $\mathbb{F}_2[[x]]$. For this purpose, we define four 2×2 matrices M^e, M^o, W^e, W^o as follows: For each $T \in \{M^e, M^o, W^e, W^o\}$ and $0 \leq i, j \leq 1$, $T_{i,j}$ is defined to be the unique solution in $\mathbb{F}_2[[x]]$ of the polynomial $\phi(T, i, j)$ under certain initial conditions; the polynomials $\phi(T, i, j)$ and initial conditions are given in Section 3. We will prove that these four matrices, whose components are algebraic by definition, are the limits of $(M_{2^n}(x))_n$, $(M_{2^{n+1}}(x))_n$, $(W_{2^n}(x))_n$, and $(W_{2^{n+1}}(x))_n$.

Let us explain how the polynomials $\phi(T, i, j)$ and initial conditions are found, and why the solutions exist and are unique. For $0 \leq i, j \leq 1$, the coefficients of the polynomial $\phi(M^e, i, j)$ (resp. $\phi(M^o, i, j)$, $\phi(W^e, i, j)$, and $\phi(W^o, i, j)$) are the Padé-Hermite approximants of type

$$(175, 175, 175, 175, 175)$$

of the vector

$$(1, T^3, T^6, T^9, T^{12}),$$

where $T = M_{12,i,j}$ (resp. $M_{11,i,j}$, $W_{12,i,j}$, and $W_{11,i,j}$), found by the Derksen algorithm. We take the first 8 terms of $M_{12,i,j}$ (resp. $M_{11,i,j}$, $W_{12,i,j}$, and $W_{11,i,j}$) as initial conditions for $\phi(M^e, i, j)$ (resp. $\phi(M^o, i, j)$, $\phi(W^e, i, j)$, and $\phi(W^o, i, j)$). The following fact will be used to ensure that the solution exists and is unique: let $P(x, y) \in \mathbb{F}_2[x, y]$ and for each series $f(x) = \sum_{n=0}^{\infty} a_n x^n \in \mathbb{F}_2[[x]]$ denote the partial sum $\sum_{j=0}^{n-1} a_j x^j$ by $f_n(x)$ for $n \geq 0$. If for some $n \geq 0$ and $a_0, a_1, \dots, a_{n-1} \in \mathbb{F}_2$ $P(x, \sum_{j=0}^{n-1} a_j x^j) = O(x^n)$ and $Q(x, y) := P(x, \sum_{j=0}^{n-1} a_j x^j + x^n y)$ can be written as $x^m \sum_{j=0}^{\infty} q_j(x) y^j$ where $q_j(x)$ are polynomials for $j \geq 0$, $q_1(0) = 1$, and $q_j(0) = 0$ for $j > 1$, then there exists a unique solution $f(x) \in \mathbb{F}_2[[x]]$ of $P(x, f(x)) = 0$ that satisfies the initial condition $f_n(x) = \sum_{j=0}^{n-1} a_j x^j$.

We state two lemmas concerning the four matrices M^e, M^o, W^e, W^o . The first one is about relations between them; the second, about the structure of the each matrix.

Lemma 2.1. *We have*

$$(2.3) \quad M^e = W^o \cdot M^o,$$

$$(2.4) \quad M^o = W^e \cdot M^e,$$

$$(2.5) \quad W^e = M^o \cdot W^o,$$

$$(2.6) \quad W^o = M^e \cdot W^e.$$

Proof. We give the proof of the identity

$$M_{0,0}^e = W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o,$$

the proofs of the others being similar.

First, we compute the minimal polynomials of $W_{0,0}^o M_{0,0}^o$ and $W_{0,1}^o M_{1,0}^o$. We know that

$$P(x, y) = \text{Res}_z (\phi(W^o, 0, 0)(x, z), z^{12} \cdot \phi(M^o, 0, 0)(x, y/z))$$

is an annihilating polynomial of $W_{0,0}^o M_{0,0}^o$. We use Padé-Hermite approximation to find a candidate for the minimal polynomial of $W_{0,0}^o M_{0,0}^o$, that will be called $\phi_0(x, y)$. To prove that $\phi_0(x, y)$ is indeed the minimal polynomial, it suffices to prove that it is an irreducible factor of $P(x, y)$ of multiplicity m and that $Q(x, y) := P(x, y)/\phi_0(x, y)^m$ is not an annihilating polynomial of $W_{0,0}^o M_{0,0}^o$. We verify the first point directly. For the second point, we truncate $W_{0,0}^o M_{0,0}^o$ to order 630 and substitute it for y in $Q(x, y)$. We get a series of valuation less than 630, which proves that $Q(x, y)$ is not an annihilating polynomial of $W_{0,0}^o M_{0,0}^o$. We find the minimal polynomial $\phi_1(x, y)$ of $W_{0,1}^o M_{1,0}^o$ in a similar way.

Now we prove that $\phi(M^e, 0, 0)$ is the minimal polynomial of $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$. We know that

$$S(x, y) = \text{Res}_z (\phi_0(x, z), \phi_1(x, y + z))$$

is an annihilating of $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$. We verify that $\phi(M^e, 0, 0)$ is an irreducible factor of $S(x, y)$ of multiplicity μ , and that $Q(x, y) := S(x, y)/\phi(M^e, 0, 0)^\mu$ is not an annihilating polynomial of $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$. To see the last point, we truncate $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$ to order 770 and substitute it for y in $Q(x, y)$. We get a series of valuation less than 770, and therefore $Q(x, y)$ is not an annihilating polynomial of $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$.

Finally, the first 16 terms of $M_{0,0}^e$ and $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$ coincide. As these first terms determine a unique solution of $\phi(M^e, 0, 0)$, we know that the two series are one and the same. $\square$

Define

$$R^e = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \quad \text{and} \quad R^o = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}.$$

Lemma 2.2. *For all integers $k \geq 2$ and $u = 2^{2k-1}$, the following identities hold.*

$$\begin{aligned} M^e[:14u] &= M^e[:7u] + x^{3u} M^e[:4u] + x^{4u} M^e[:3u] + x^{3u} M^e[:7u] + x^{6u} M^e[:4u] \\ &\quad + x^{7u} M^e[:3u] + x^{4u} M^e[:7u] + x^{7u} M^e[:4u] + x^{8u} M^e[:3u] \\ &\quad + x^{6u} M^e[:7u] + x^{9u} M^e[:4u] + x^{10u} M^e[:3u] + x^{7u} M^e[:7u] \\ &\quad + x^{10u} M^e[:4u] + x^{11u} M^e[:3u] + x^{9u} M^e[:5u] + x^{13u} M^e[:u] \\ &\quad + (x^{7u} + x^{10u} + x^{11u} + x^{13u}) R^e, \\ W^e[:14u] &= W^e[:7u] + x^{3u} W^e[:4u] + x^{4u} W^e[:3u] + x^{3u} W^e[:7u] + x^{6u} W^e[:4u] \end{aligned}$$

$$\begin{aligned}
& + x^{7u}W^e[:3u] + x^{4u}W^e[:7u] + x^{7u}W^e[:4u] + x^{8u}W^e[:3u] \\
& + x^{6u}W^e[:7u] + x^{9u}W^e[:4u] + x^{10u}W^e[:3u] + x^{7u}W^e[:7u] \\
& + x^{10u}W^e[:4u] + x^{11u}W^e[:3u] + x^{9u}W^e[:5u] + x^{13u}W^e[:u] \\
& + (x^{7u} + x^{10u} + x^{11u} + x^{13u})R^e.
\end{aligned}$$

For all integers $k \geq 2$ and $u = 2^{2k}$, the following identities hold.

$$\begin{aligned}
M^o[:14u] &= M^o[:7u] + x^{3u}M^o[:4u] + x^{4u}M^o[:3u] + x^{3u}M^o[:7u] + x^{6u}M^o[:4u] \\
&+ x^{7u}M^o[:3u] + x^{4u}M^o[:7u] + x^{7u}M^o[:4u] + x^{8u}M^o[:3u] \\
&+ x^{6u}M^o[:7u] + x^{9u}M^o[:4u] + x^{10u}M^o[:3u] + x^{7u}M^o[:7u] \\
&+ x^{10u}M^o[:4u] + x^{11u}M^o[:3u] + x^{9u}M^o[:5u] + x^{13u}M^o[:u] \\
&+ (x^{7u} + x^{10u} + x^{11u} + x^{13u})R^o, \\
W^o[:14u] &= W^o[:7u] + x^{3u}W^o[:4u] + x^{4u}W^o[:3u] + x^{3u}W^o[:7u] + x^{6u}W^o[:4u] \\
&+ x^{7u}W^o[:3u] + x^{4u}W^o[:7u] + x^{7u}W^o[:4u] + x^{8u}W^o[:3u] \\
&+ x^{6u}W^o[:7u] + x^{9u}W^o[:4u] + x^{10u}W^o[:3u] + x^{7u}W^o[:7u] \\
&+ x^{10u}W^o[:4u] + x^{11u}W^o[:3u] + x^{9u}W^o[:5u] + x^{13u}W^o[:u] \\
&+ (x^{7u} + x^{10u} + x^{11u} + x^{13u})R^o.
\end{aligned}$$

Proof. To prove Lemma 2.2 we first construct an automaton for each sequence concerned, and then transform the conditions on infinitely many k 's into finitely many conditions on the states of the automaton. In the following, we will prove that for $T = M_{0,0}^o$, for all integer $k \geq 2$ and $u = 2^{2k}$,

$$\begin{aligned}
T[:14u] &= T[:7u] + x^{3u}T[:4u] + x^{4u}T[:3u] + x^{3u}T[:7u] + x^{6u}T[:4u] + x^{7u}T[:3u] \\
&+ x^{4u}T[:7u] + x^{7u}T[:4u] + x^{8u}T[:3u] + x^{6u}T[:7u] + x^{9u}T[:4u] \\
&+ x^{10u}T[:3u] + x^{7u}T[:7u] + x^{10u}T[:4u] + x^{11u}T[:3u] + x^{9u}T[:5u] \\
&+ x^{13u}T[:u].
\end{aligned}$$

The proof of the other 15 cases are similar. We break down the above identity into 7 parts:

$$\begin{aligned}
0 &= x^{7u}T[:u] + x^{4u}T[3u:4u] + x^{3u}T[4u:5u] + T[7u:8u], \\
0 &= x^{8u}T[:u] + x^{7u}T[u:2u] + x^{4u}T[4u:5u] + x^{3u}T[5u:6u] + T[8u:9u], \\
0 &= x^{8u}T[u:2u] + x^{7u}T[2u:3u] + x^{4u}T[5u:6u] + x^{3u}T[6u:7u] \\
&+ T[9u:10u], \\
0 &= x^{8u}T[2u:3u] + x^{6u}T[4u:5u] + x^{4u}T[6u:7u] + T[10u:11u], \\
0 &= x^{11u}T[:u] + x^{7u}T[4u:5u] + x^{6u}T[5u:6u] + T[11u:12u], \\
0 &= x^{11u}T[u:2u] + x^{7u}T[5u:6u] + x^{6u}T[6u:7u] + T[12u:13u], \\
0 &= x^{13u}T[:u] + x^{11u}T[2u:3u] + x^{10u}T[3u:4u] + x^{9u}T[4u:5u] \\
&+ x^{7u}T[6u:7u] + T[13u:14u],
\end{aligned}$$

which can be rewritten as

$$(2.7) \quad 0 = T[[w]_2] + T[[11w]_2] + T[[100w]_2] + T[[111w]_2],$$

$$\begin{aligned}
(2.8) \quad & 0 = T[[w]_2] + T[[1w]_2] + T[[100w]_2] + T[[101w]_2] + T[[1000w]_2], \\
(2.9) \quad & 0 = T[[1w]_2] + T[[10w]_2] + T[[101w]_2] + T[[110w]_2] + T[[1001w]_2], \\
(2.10) \quad & 0 = T[[10w]_2] + T[[100w]_2] + T[[110w]_2] + T[[1010w]_2], \\
(2.11) \quad & 0 = T[[w]_2] + T[[100w]_2] + T[[101w]_2] + T[[1011w]_2], \\
(2.12) \quad & 0 = T[[1w]_2] + T[[101w]_2] + T[[110w]_2] + T[[1100w]_2], \\
(2.13) \quad & 0 = T[[w]_2] + T[[10w]_2] + T[[11w]_2] + T[[100w]_2] + T[[110w]_2] \\
& \quad + T[[1101w]_2],
\end{aligned}$$

for all binary word w of length $2k$ and $w \neq 0^{2k}$, and

$$\begin{aligned}
(2.14) \quad & 0 = T[[w]_2] + T[[11w]_2] + T[[100w]_2] + T[[111w]_2], \\
(2.15) \quad & 0 = T[[w]_2] + T[[1w]_2] + T[[100w]_2] + T[[101w]_2] + T[[1000w]_2], \\
(2.16) \quad & 0 = T[[1w]_2] + T[[10w]_2] + T[[101w]_2] + T[[110w]_2] + T[[1001w]_2], \\
(2.17) \quad & 0 = T[[10w]_2] + T[[100w]_2] + T[[110w]_2] + T[[1010w]_2], \\
(2.18) \quad & 0 = T[[w]_2] + T[[100w]_2] + T[[101w]_2] + T[[1011w]_2], \\
(2.19) \quad & 0 = T[[1w]_2] + T[[101w]_2] + T[[110w]_2] + T[[1100w]_2], \\
(2.20) \quad & 0 = T[[w]_2] + T[[10w]_2] + T[[11w]_2] + T[[100w]_2] + T[[110w]_2] \\
& \quad + T[[1101w]_2],
\end{aligned}$$

for $w = 0^{2k}$.

First we calculate an 2-automaton that generates T from its minimal polynomial and its first terms. This automaton has 214 states; its transition function and output function can be found in the annex. Let $A(s, w)$ denote the state reached after reading w from right to left starting from the state s , and τ the output function. Define

$$E_{2k} = \{A(i, w) : |w| = 2k, w \neq 0^{2k}\}.$$

Identities (2.7) through (2.20) can be written as

$$\begin{aligned}
(2.21) \quad & 0 = \tau(A(s, \epsilon)) + \tau(A(s, 11)) + \tau(A(s, 100)) + \tau(A(s, 111)), \\
& 0 = \tau(A(s, \epsilon)) + \tau(A(s, 1)) + \tau(A(s, 100)) + \tau(A(s, 101)) \\
(2.22) \quad & + \tau(A(s, 1000)), \\
& 0 = \tau(A(s, 1)) + \tau(A(s, 10)) + \tau(A(s, 101)) + \tau(A(s, 110)) \\
(2.23) \quad & + \tau(A(s, 1001)), \\
(2.24) \quad & 0 = \tau(A(s, 10)) + \tau(A(s, 100)) + \tau(A(s, 110)) + \tau(A(s, 1010)), \\
(2.25) \quad & 0 = \tau(A(s, \epsilon)) + \tau(A(s, 100)) + \tau(A(s, 101)) + \tau(A(s, 1011)), \\
(2.26) \quad & 0 = \tau(A(s, 1)) + \tau(A(s, 101)) + \tau(A(s, 110)) + \tau(A(s, 1100)), \\
& 0 = \tau(A(s, \epsilon)) + \tau(A(s, 10)) + \tau(A(s, 11)) + \tau(A(s, 100)) \\
(2.27) \quad & + \tau(A(s, 110)) + \tau(A(s, 1101)),
\end{aligned}$$

for all $s \in E_{2k}$, and

$$\begin{aligned}
(2.28) \quad & 0 = \tau(A(s, \epsilon)) + \tau(A(s, 11)) + \tau(A(s, 100)) + \tau(A(s, 111)), \\
& 0 = \tau(A(s, \epsilon)) + \tau(A(s, 1)) + \tau(A(s, 100)) + \tau(A(s, 101)) \\
(2.29) \quad & + \tau(A(s, 1000)),
\end{aligned}$$

$$(2.30) \quad 0 = \tau(A(s, 1)) + \tau(A(s, 10)) + \tau(A(s, 101)) + \tau(A(s, 110)) \\ + \tau(A(s, 1001)),$$

$$(2.31) \quad 0 = \tau(A(s, 10)) + \tau(A(s, 100)) + \tau(A(s, 110)) + \tau(A(s, 1010)),$$

$$(2.32) \quad 0 = \tau(A(s, \epsilon)) + \tau(A(s, 100)) + \tau(A(s, 101)) + \tau(A(s, 1011)),$$

$$(2.33) \quad 0 = \tau(A(s, 1)) + \tau(A(s, 101)) + \tau(A(s, 110)) + \tau(A(s, 1100)),$$

$$(2.34) \quad 0 = \tau(A(s, \epsilon)) + \tau(A(s, 10)) + \tau(A(s, 11)) + \tau(A(s, 100)) \\ + \tau(A(s, 110)) + \tau(A(s, 1101)),$$

for $s = A(i, 0^{2k})$. We find that $(A(i, 0^{20}), E_{20}) = (A(i, 0^{12}), E_{12})$, so that we only have to verify that identities (2.21) through (2.34) hold for $2 \leq k \leq 10$, which turns out to be true. $\square$

In the following lemma, we express M_{2k} , M_{2k+1} , W_{2k} , and W_{2k+1} in terms of M^e , M^o , W^e , and W^o .

Lemma 2.3. *For all integer $k \geq 2$, and $u = 2^{2k-1}$,*

$$M_{2k} = M^e[:7u] + x^{3u}M^e[:4u] + x^{4u}M^e[:3u] + x^{7u}R^e,$$

$$W_{2k} = W^e[:7u] + x^{3u}W^e[:4u] + x^{4u}W^e[:3u] + x^{7u}R^e.$$

For all integer $k \geq 2$, and $u = 2^{2k}$,

$$M_{2k+1} = M^o[:7u] + x^{3u}M^o[:4u] + x^{4u}M^o[:3u] + x^{7u}R^o,$$

$$W_{2k+1} = W^o[:7u] + x^{3u}W^o[:4u] + x^{4u}W^o[:3u] + x^{7u}R^o.$$

Proof. We call the four identities in Lemma 2.3 also by the name M_{2k} , W_{2k} , M_{2k+1} , and W_{2k+1} . For $n = 2$, the identities can be verified directly. For $n \geq 2$, we claim that

$$M_{2k} \wedge W_{2k} \Rightarrow M_{2k+1} \wedge W_{2k+1}, \\ M_{2k+1} \wedge W_{2k+1} \Rightarrow M_{2k+2} \wedge W_{2k+2}.$$

We give the proof of

$$M_{2k} \wedge W_{2k} \Rightarrow M_{2k+1},$$

the other ones being similar. Set $u = 2^{2k}$ and $v = 2^{2k-1}$. By definition and induction hypothesis, the left side of identity M_{2k+1} is equal to

$$(2.35) \quad W_{2k}M_{2k} = (W^e[:7v] + x^{3v}W^e[:4v] + x^{4v}W^e[:3v] + x^{7v}R^e) \times (M^e[:7v] \\ + x^{3v}M^e[:4v] + x^{4v}M^e[:3v] + x^{7v}R^e);$$

Call this expression *lhs*. Note that both sides of identity M_{2k+1} have the same term of highest degree $x^{14v}R^o$. Therefore we only need to prove that their difference is $O(x^{14v})$. Using Lemma 2.1 it can be seen that the right side of identity M_{2k+1} is congruent, modulo x^{14v} , to

$$W^e[:14v]M^e[:14v] + x^{6v}W^e[:8v]M^e[:8v] + x^{8v}W^e[:6v]M^e[:6v].$$

For all $n \leq 14$, replace the occurrences of $W^e[:n \cdot v]$ and $M^e[:n \cdot v]$ in the above expression by the reduction modulo $x^{n \cdot v}$ of the right side of the corresponding identity in Lemma 2.2 and get a new expression, which we call *rhs*. Define

$$X := x^v, \\ a_n := W^e[n \cdot v : (n+1) \cdot v] / X^n,$$

$$\begin{aligned} b_n &:= M^e[n \cdot v : (n+1) \cdot v]/X^n, \\ c &:= R^e. \end{aligned}$$

Using the notation introduced above, we can represent the expressions *lhs* (2.35) and *rhs* as polynomials in $\mathbb{F}_2[a_1, \dots, a_{14}, b_1, \dots, b_{14}, c][X]$. Note that it is not a problem that a_j commutes with b_k while W^e does not commute with M^e , because in the expressions concerned, the products of W^e -terms and M^e -terms are always in the same order. We let the computer do the simplification and check that the difference between these two polynomials is indeed $O(X^{14})$, which completes the proof. $\square$

Proof of Theorem 1.1. We prove the theorem for $\text{CF}(a, b)$; for $\text{CF}(b, a)$, the proof is similar. By Lemma 2.3, we have For all $0 \leq j, k \leq 1$,

$$\begin{aligned} \lim_{n \rightarrow \infty} M_{2n,j,k} &= M_{j,k}^e, \\ \lim_{n \rightarrow \infty} M_{2n+1,j,k} &= M_{j,k}^o, \\ \lim_{n \rightarrow \infty} W_{2n,j,k} &= W_{j,k}^e, \\ \lim_{n \rightarrow \infty} W_{2n+1,j,k} &= W_{j,k}^o. \end{aligned}$$

Let $z = 1/x$. By the convergence theorem and identity (2.2),

$$\text{CF}(a, b) = \frac{M_{0,1}^e(x)}{M_{0,0}^e(x)}.$$

By definition, $\phi(M^e, 0, 1)$ and $\phi(M^e, 0, 0)$ are minimal polynomials of $M_{0,1}^e$ and $M_{0,0}^e$. Therefore

$$P(x, y) = \text{Res}_t (\phi(M^e, 0, 1)(x, t), y^{12} \phi(M^e, 0, 1)(x, t/y))$$

is an annihilating polynomial of $f(x) = M_{0,1}^e/M_{0,0}^e$.

Define

$$Q(x, y) = q_4(x)y^4 + q_3(x)y^3 + q_2(x)y^2 + q_1(x)y + q_0(x),$$

where

$$\begin{aligned} q_0(x) &= x^{16} + x^{13} + x^{11} + x^{10} + x^8 + x^7, \\ q_1(x) &= x^{11} + x^9 + x^8 + x^6 + x^5 + x^2, \\ q_2(x) &= x^{12} + x^{10} + x^4 + 1, \\ q_3(x) &= x^{11} + x^9 + x^8 + x^6 + x^5 + x^2, \\ q_4(x) &= x^{16} + x^{13} + x^{11} + x^7 + x^4. \end{aligned}$$

The polynomial $Q(x, y)$ is the candidate for the minimal polynomial of $f(x)$ found by Padé-Hermite approximation. To prove that it is indeed the minimal polynomial of $f(x)$, we only need to prove that it is an irreducible factor of $P(x, y)$ of multiplicity m and $R(x, y) := P(x, y)/Q(x, y)^m$ is not an annihilating polynomial of $f(x)$. We verify the first point directly. For the second point, we find that when we truncate $f(x)$ to order 216, and substitute it for y in $R(z, y)$, we get a series with valuation smaller than 216, which proves that $R(z, y)$ is not an annihilating polynomial of $f(x)$. Finally, $z^{12}Q(1/z, y)$ is the minimal polynomial of $\text{CF}(a, b) = f(1/z)$. $\square$

3. ANNEXE

All 16 polynomials are of the form

$$p_0(x) + p_3(x)y^3 + p_6(x)y^6 + p_9(x)y^9 + p_{12}(x)y^{12}.$$

The coefficients $p_j(x)$ and the 16 initial terms to determine the solutions uniquely are given below.

For $\phi(M^e, 0, 0)$:

$$\begin{aligned} p_0(x) &= x^{96} + x^{90} + x^{86} + x^{84} + x^{76} + x^{74} + x^{72} + x^{70} + x^{58} + x^{54} + x^{52} \\ &\quad + x^{50} + x^{48} + x^{38} + x^{36} + x^{30} + x^{24}, \\ p_3(x) &= x^{78} + x^{70} + x^{68} + x^{66} + x^{64} + x^{62} + x^{60} + x^{58} + x^{54} + x^{50} + x^{48} \\ &\quad + x^{46} + x^{44} + x^{42} + x^{38} + x^{34} + x^{30} + x^{26} + x^{18} + x^{16}, \\ p_6(x) &= x^{80} + x^{78} + x^{74} + x^{70} + x^{68} + x^{64} + x^{60} + x^{52} + x^{46} + x^{38} + x^{34} \\ &\quad + x^{26} + x^{20} + x^{18} + x^{16} + x^{14} + x^{10} + x^8 + x^4 + 1, \\ p_9(x) &= x^{70} + x^{68} + x^{66} + x^{62} + x^{56} + x^{50} + x^{48} + x^{44} + x^{38} + x^{36} + x^{34} \\ &\quad + x^{28} + x^{26} + x^{24} + x^{22} + x^{18} + x^{12} + x^{10} + x^8 + x^4, \\ p_{12}(x) &= x^{72} + x^{56} + x^{48} + x^{32} + x^{24} + 1, \end{aligned}$$

and the initial terms are $[1, 0, 0, 0, 0, 0, 0, 0, 1, 0, 1, 0, 0, 0, 1, 0]$.

For $\phi(M^e, 0, 1)$:

$$\begin{aligned} p_0(x) &= x^{87} + x^{84} + x^{79} + x^{78} + x^{77} + x^{76} + x^{75} + x^{69} + x^{68} + x^{67} + x^{66} \\ &\quad + x^{65} + x^{64} + x^{62} + x^{61} + x^{60} + x^{58} + x^{56} + x^{55} + x^{54} + x^{51} + x^{50} \\ &\quad + x^{45} + x^{44} + x^{43} + x^{42} + x^{41} + x^{39} + x^{38} + x^{37} + x^{35} + x^{34} + x^{33}, \\ p_3(x) &= x^{54} + x^{52} + x^{44} + x^{42} + x^{38} + x^{34} + x^{30} + x^{28} + x^{22} + x^{18}, \\ p_6(x) &= x^{57} + x^{54} + x^{53} + x^{51} + x^{50} + x^{49} + x^{48} + x^{47} + x^{46} + x^{44} + x^{43} \\ &\quad + x^{40} + x^{37} + x^{34} + x^{33} + x^{31} + x^{30} + x^{29} + x^{28} + x^{27} + x^{26} + x^{24} \\ &\quad + x^{23} + x^{21} + x^{20} + x^{18} + x^{15} + x^{12}, \\ p_9(x) &= x^{60} + x^{58} + x^{56} + x^{52} + x^{48} + x^{44} + x^{38} + x^{36} + x^{34} + x^{32} + x^{28} \\ &\quad + x^{22} + x^{20} + x^{18} + x^{14} + x^{12} + x^{10} + x^6, \\ p_{12}(x) &= x^{63} + x^{60} + x^{51} + x^{48} + x^{47} + x^{44} + x^{35} + x^{32} + x^{15} + x^{12} + x^3 + 1, \end{aligned}$$

and the initial terms are $[0, 0, 1, 0, 0, 0, 0, 0, 0, 1, 1, 0, 0, 0, 0, 1]$.

For $\phi(M^e, 1, 0)$:

$$\begin{aligned} p_0(x) &= x^{87} + x^{84} + x^{79} + x^{78} + x^{77} + x^{76} + x^{75} + x^{69} + x^{68} + x^{67} + x^{66} \\ &\quad + x^{65} + x^{64} + x^{62} + x^{61} + x^{60} + x^{58} + x^{56} + x^{55} + x^{54} + x^{51} + x^{50} \\ &\quad + x^{45} + x^{44} + x^{43} + x^{42} + x^{41} + x^{39} + x^{38} + x^{37} + x^{35} + x^{34} + x^{33}, \\ p_3(x) &= x^{54} + x^{52} + x^{44} + x^{42} + x^{38} + x^{34} + x^{30} + x^{28} + x^{22} + x^{18}, \\ p_6(x) &= x^{57} + x^{54} + x^{53} + x^{51} + x^{50} + x^{49} + x^{48} + x^{47} + x^{46} + x^{44} + x^{43} \\ &\quad + x^{40} + x^{37} + x^{34} + x^{33} + x^{31} + x^{30} + x^{29} + x^{28} + x^{27} + x^{26} + x^{24} \\ &\quad + x^{23} + x^{21} + x^{20} + x^{18} + x^{15} + x^{12}, \end{aligned}$$

$$p_9(x) = x^{60} + x^{58} + x^{56} + x^{52} + x^{48} + x^{44} + x^{38} + x^{36} + x^{34} + x^{32} + x^{28} \\ + x^{22} + x^{20} + x^{18} + x^{14} + x^{12} + x^{10} + x^6,$$

$$p_{12}(x) = x^{63} + x^{60} + x^{51} + x^{48} + x^{47} + x^{44} + x^{35} + x^{32} + x^{15} + x^{12} + x^3 + 1,$$

and the initial terms are $[0, 0, 1, 0, 0, 0, 0, 0, 0, 1, 1, 0, 0, 0, 0, 1]$.

For $\phi(M^e, 1, 1)$:

$$p_0(x) = x^{78} + x^{68} + x^{66} + x^{58} + x^{54} + x^{48} + x^{46} + x^{44} + x^{42},$$

$$p_3(x) = x^{60} + x^{54} + x^{52} + x^{50} + x^{44} + x^{40} + x^{38} + x^{36} + x^{34} + x^{32} + x^{30} \\ + x^{28} + x^{26} + x^{22} + x^{20} + x^{18} + x^{16} + x^{12},$$

$$p_6(x) = x^{62} + x^{60} + x^{54} + x^{52} + x^{50} + x^{48} + x^{46} + x^{44} + x^{40} + x^{38} + x^{36} \\ + x^{34} + x^{26} + x^{18} + x^{16} + x^{14} + x^{12} + x^6 + x^4 + 1,$$

$$p_9(x) = x^{52} + x^{50} + x^{48} + x^{46} + x^{42} + x^{30} + x^{28} + x^{26} + x^{22} + x^{16} + x^{14} \\ + x^{12} + x^8 + x^4,$$

$$p_{12}(x) = x^{54} + x^{48} + x^{38} + x^{32} + x^6 + 1,$$

and the initial terms are $[0, 0, 0, 0, 1, 0, 0, 0, 0, 0, 0, 0, 1, 0, 1, 0]$.

For $\phi(W^e, 0, 0)$:

$$p_0(x) = x^{78} + x^{68} + x^{60} + x^{58} + x^{48} + x^{42} + x^{40} + x^{32} + x^{24},$$

$$p_3(x) = x^{60} + x^{54} + x^{52} + x^{50} + x^{48} + x^{44} + x^{38} + x^{34} + x^{32} + x^{30} + x^{26} \\ + x^{24} + x^{22} + x^{20} + x^{18} + x^{16},$$

$$p_6(x) = x^{62} + x^{60} + x^{54} + x^{52} + x^{46} + x^{42} + x^{36} + x^{34} + x^{32} + x^{30} + x^{28} \\ + x^{20} + x^{18} + x^{16} + x^8 + x^6 + x^4 + 1,$$

$$p_9(x) = x^{52} + x^{50} + x^{48} + x^{46} + x^{42} + x^{30} + x^{28} + x^{26} + x^{22} + x^{16} + x^{14} \\ + x^{12} + x^8 + x^4,$$

$$p_{12}(x) = x^{54} + x^{48} + x^{38} + x^{32} + x^6 + 1,$$

and the initial terms are $[1, 0, 0, 0, 1, 0, 0, 0, 1, 0, 0, 0, 1, 0, 1, 0]$.

For $\phi(W^e, 0, 1)$:

$$p_0(x) = x^{87} + x^{84} + x^{79} + x^{78} + x^{77} + x^{76} + x^{75} + x^{73} + x^{70} + x^{69} + x^{68} \\ + x^{66} + x^{65} + x^{62} + x^{60} + x^{59} + x^{57} + x^{55} + x^{53} + x^{51} + x^{50} + x^{46} \\ + x^{45} + x^{43} + x^{42} + x^{41} + x^{40} + x^{36} + x^{33},$$

$$p_3(x) = x^{54} + x^{52} + x^{44} + x^{42} + x^{38} + x^{34} + x^{30} + x^{28} + x^{22} + x^{18},$$

$$p_6(x) = x^{57} + x^{54} + x^{53} + x^{51} + x^{50} + x^{49} + x^{48} + x^{47} + x^{46} + x^{44} + x^{43} \\ + x^{40} + x^{37} + x^{34} + x^{33} + x^{31} + x^{30} + x^{29} + x^{28} + x^{27} + x^{26} + x^{24} \\ + x^{23} + x^{21} + x^{20} + x^{18} + x^{15} + x^{12},$$

$$p_9(x) = x^{60} + x^{58} + x^{56} + x^{52} + x^{48} + x^{44} + x^{38} + x^{36} + x^{34} + x^{32} + x^{28} \\ + x^{22} + x^{20} + x^{18} + x^{14} + x^{12} + x^{10} + x^6,$$

$$p_{12}(x) = x^{63} + x^{60} + x^{51} + x^{48} + x^{47} + x^{44} + x^{35} + x^{32} + x^{15} + x^{12} + x^3 + 1,$$

and the initial terms are $[0, 0, 0, 0, 0, 1, 0, 0, 1, 1, 0, 1, 0, 1, 1, 1]$.

For $\phi(W^e, 1, 0)$:

$$\begin{aligned}
p_0(x) &= x^{87} + x^{84} + x^{79} + x^{78} + x^{77} + x^{76} + x^{75} + x^{73} + x^{70} + x^{69} + x^{68} \\
&\quad + x^{66} + x^{65} + x^{62} + x^{60} + x^{59} + x^{57} + x^{55} + x^{53} + x^{51} + x^{50} + x^{46} \\
&\quad + x^{45} + x^{43} + x^{42} + x^{41} + x^{40} + x^{36} + x^{33}, \\
p_3(x) &= x^{54} + x^{52} + x^{44} + x^{42} + x^{38} + x^{34} + x^{30} + x^{28} + x^{22} + x^{18}, \\
p_6(x) &= x^{57} + x^{54} + x^{53} + x^{51} + x^{50} + x^{49} + x^{48} + x^{47} + x^{46} + x^{44} + x^{43} \\
&\quad + x^{40} + x^{37} + x^{34} + x^{33} + x^{31} + x^{30} + x^{29} + x^{28} + x^{27} + x^{26} + x^{24} \\
&\quad + x^{23} + x^{21} + x^{20} + x^{18} + x^{15} + x^{12}, \\
p_9(x) &= x^{60} + x^{58} + x^{56} + x^{52} + x^{48} + x^{44} + x^{38} + x^{36} + x^{34} + x^{32} + x^{28} \\
&\quad + x^{22} + x^{20} + x^{18} + x^{14} + x^{12} + x^{10} + x^6, \\
p_{12}(x) &= x^{63} + x^{60} + x^{51} + x^{48} + x^{47} + x^{44} + x^{35} + x^{32} + x^{15} + x^{12} + x^3 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 1, 0, 0, 1, 1, 0, 1, 0, 1, 1, 1]$.

For $\phi(W^e, 1, 1)$:

$$\begin{aligned}
p_0(x) &= x^{96} + x^{90} + x^{86} + x^{76} + x^{74} + x^{70} + x^{66} + x^{64} + x^{62} + x^{58} + x^{50} \\
&\quad + x^{48} + x^{42}, \\
p_3(x) &= x^{78} + x^{70} + x^{68} + x^{64} + x^{62} + x^{54} + x^{52} + x^{50} + x^{48} + x^{46} + x^{44} \\
&\quad + x^{42} + x^{38} + x^{30} + x^{28} + x^{26} + x^{16} + x^{12}, \\
p_6(x) &= x^{80} + x^{78} + x^{74} + x^{70} + x^{66} + x^{64} + x^{60} + x^{58} + x^{56} + x^{48} + x^{46} \\
&\quad + x^{38} + x^{36} + x^{28} + x^{26} + x^{24} + x^{16} + x^{14} + x^{12} + x^{10} + x^4 + 1, \\
p_9(x) &= x^{70} + x^{68} + x^{66} + x^{62} + x^{56} + x^{50} + x^{48} + x^{44} + x^{38} + x^{36} + x^{34} \\
&\quad + x^{28} + x^{26} + x^{24} + x^{22} + x^{18} + x^{12} + x^{10} + x^8 + x^4, \\
p_{12}(x) &= x^{72} + x^{56} + x^{48} + x^{32} + x^{24} + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 1, 0, 0, 0, 1, 0]$.

For $\phi(M^o, 0, 0)$:

$$\begin{aligned}
p_0(x) &= x^{87} + x^{84} + x^{81} + x^{79} + x^{78} + x^{77} + x^{73} + x^{70} + x^{69} + x^{68} + x^{67} \\
&\quad + x^{66} + x^{61} + x^{60} + x^{59} + x^{57} + x^{54} + x^{51} + x^{50} + x^{49} + x^{48} + x^{46} \\
&\quad + x^{42} + x^{41} + x^{37} + x^{36} + x^{34} + x^{33} + x^{32} + x^{30} + x^{28} + x^{27} + x^{24}, \\
p_3(x) &= x^{67} + x^{64} + x^{61} + x^{58} + x^{57} + x^{56} + x^{51} + x^{48} + x^{46} + x^{45} + x^{44} \\
&\quad + x^{42} + x^{41} + x^{40} + x^{38} + x^{37} + x^{36} + x^{34} + x^{33} + x^{32} + x^{29} + x^{26} \\
&\quad + x^{25} + x^{24} + x^{22} + x^{20} + x^{19} + x^{16}, \\
p_6(x) &= x^{71} + x^{68} + x^{67} + x^{65} + x^{64} + x^{63} + x^{62} + x^{59} + x^{58} + x^{57} + x^{55} \\
&\quad + x^{53} + x^{52} + x^{47} + x^{44} + x^{43} + x^{40} + x^{38} + x^{37} + x^{36} + x^{35} + x^{33} \\
&\quad + x^{32} + x^{29} + x^{26} + x^{24} + x^{23} + x^{22} + x^{21} + x^{19} + x^{18} + x^{17} + x^{15} \\
&\quad + x^{14} + x^{11} + x^8 + x^3 + 1, \\
p_9(x) &= x^{59} + x^{56} + x^{43} + x^{40} + x^{11} + x^8,
\end{aligned}$$

$$p_{12}(x) = x^{63} + x^{60} + x^{59} + x^{56} + x^{51} + x^{48} + x^{47} + x^{44} + x^{43} + x^{40} + x^{35} \\ + x^{32} + x^{15} + x^{12} + x^{11} + x^8 + x^3 + 1,$$

and the initial terms are $[1, 0, 0, 0, 1, 0, 0, 1, 0, 0, 0, 1, 0, 1, 0, 0]$.

For $\phi(M^o, 0, 1)$:

$$p_0(x) = x^{84} + x^{75} + x^{74} + x^{73} + x^{70} + x^{68} + x^{67} + x^{66} + x^{61} + x^{60} + x^{58} \\ + x^{57} + x^{56} + x^{55} + x^{54} + x^{50} + x^{49} + x^{46} + x^{44} + x^{42} + x^{41} + x^{39} \\ + x^{38} + x^{37} + x^{36} + x^{35} + x^{34} + x^{33}, \\ p_3(x) = x^{51} + x^{48} + x^{45} + x^{43} + x^{42} + x^{40} + x^{37} + x^{34} + x^{27} + x^{24} + x^{21} \\ + x^{18}, \\ p_6(x) = x^{54} + x^{50} + x^{48} + x^{46} + x^{44} + x^{40} + x^{34} + x^{30} + x^{28} + x^{26} + x^{24} \\ + x^{20} + x^{18} + x^{12}, \\ p_9(x) = x^{57} + x^{54} + x^{41} + x^{38} + x^9 + x^6, \\ p_{12}(x) = x^{60} + x^{56} + x^{48} + x^{44} + x^{40} + x^{32} + x^{12} + x^8 + 1,$$

and the initial terms are $[0, 0, 1, 0, 0, 0, 1, 0, 0, 0, 0, 0, 1, 0, 0, 0]$.

For $\phi(M^o, 1, 0)$:

$$p_0(x) = x^{96} + x^{87} + x^{86} + x^{85} + x^{84} + x^{79} + x^{78} + x^{76} + x^{75} + x^{74} + x^{72} \\ + x^{69} + x^{68} + x^{62} + x^{60} + x^{59} + x^{54} + x^{52} + x^{50} + x^{44} + x^{43} + x^{42} \\ + x^{41} + x^{37} + x^{33}, \\ p_3(x) = x^{63} + x^{60} + x^{57} + x^{55} + x^{54} + x^{52} + x^{51} + x^{49} + x^{48} + x^{46} + x^{45} \\ + x^{43} + x^{42} + x^{40} + x^{39} + x^{37} + x^{36} + x^{34} + x^{33} + x^{30} + x^{27} + x^{24} \\ + x^{21} + x^{18}, \\ p_6(x) = x^{66} + x^{62} + x^{60} + x^{58} + x^{56} + x^{54} + x^{52} + x^{50} + x^{48} + x^{44} + x^{42} \\ + x^{38} + x^{36} + x^{34} + x^{32} + x^{28} + x^{26} + x^{20} + x^{18} + x^{12}, \\ p_9(x) = x^{69} + x^{66} + x^{57} + x^{54} + x^{53} + x^{50} + x^{41} + x^{38} + x^{21} + x^{18} + x^9 + x^6, \\ p_{12}(x) = x^{72} + x^{68} + x^{52} + x^{48} + x^{40} + x^{32} + x^{24} + x^{20} + x^8 + 1,$$

and the initial terms are $[0, 0, 0, 0, 0, 1, 0, 0, 1, 1, 0, 1, 1, 0, 1, 0]$.

For $\phi(M^o, 1, 1)$:

$$p_0(x) = x^{87} + x^{84} + x^{81} + x^{79} + x^{78} + x^{77} + x^{75} + x^{72} + x^{69} + x^{68} + x^{66} \\ + x^{64} + x^{61} + x^{60} + x^{59} + x^{57} + x^{54} + x^{52} + x^{49} + x^{48} + x^{44} + x^{43} \\ + x^{42}, \\ p_3(x) = x^{67} + x^{64} + x^{61} + x^{58} + x^{57} + x^{56} + x^{55} + x^{53} + x^{52} + x^{51} + x^{50} \\ + x^{48} + x^{46} + x^{44} + x^{43} + x^{41} + x^{39} + x^{38} + x^{37} + x^{35} + x^{34} + x^{33} \\ + x^{31} + x^{28} + x^{25} + x^{24} + x^{23} + x^{22}, \\ p_6(x) = x^{71} + x^{68} + x^{67} + x^{65} + x^{64} + x^{63} + x^{62} + x^{58} + x^{56} + x^{54} + x^{50} \\ + x^{49} + x^{47} + x^{46} + x^{44} + x^{43} + x^{40} + x^{38} + x^{36} + x^{34} + x^{30} + x^{24} \\ + x^{22} + x^{20} + x^{17} + x^{16} + x^{14} + x^{12} + x^3 + 1,$$

$$\begin{aligned}
p_9(x) &= x^{59} + x^{56} + x^{43} + x^{40} + x^{11} + x^8, \\
p_{12}(x) &= x^{63} + x^{60} + x^{59} + x^{56} + x^{51} + x^{48} + x^{47} + x^{44} + x^{43} + x^{40} + x^{35} \\
&\quad + x^{32} + x^{15} + x^{12} + x^{11} + x^8 + x^3 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 0, 0, 1, 0, 0, 1, 1, 0, 1, 0, 0]$.

For $\phi(W^o, 0, 0)$:

$$\begin{aligned}
p_0(x) &= x^{87} + x^{84} + x^{81} + x^{79} + x^{78} + x^{77} + x^{73} + x^{70} + x^{69} + x^{68} + x^{67} \\
&\quad + x^{66} + x^{61} + x^{60} + x^{59} + x^{57} + x^{54} + x^{51} + x^{50} + x^{49} + x^{48} + x^{46} \\
&\quad + x^{42} + x^{41} + x^{37} + x^{36} + x^{34} + x^{33} + x^{32} + x^{30} + x^{28} + x^{27} + x^{24}, \\
p_3(x) &= x^{67} + x^{64} + x^{61} + x^{58} + x^{57} + x^{56} + x^{51} + x^{48} + x^{46} + x^{45} + x^{44} \\
&\quad + x^{42} + x^{41} + x^{40} + x^{38} + x^{37} + x^{36} + x^{34} + x^{33} + x^{32} + x^{29} + x^{26} \\
&\quad + x^{25} + x^{24} + x^{22} + x^{20} + x^{19} + x^{16}, \\
p_6(x) &= x^{71} + x^{68} + x^{67} + x^{65} + x^{64} + x^{63} + x^{62} + x^{59} + x^{58} + x^{57} + x^{55} \\
&\quad + x^{53} + x^{52} + x^{47} + x^{44} + x^{43} + x^{40} + x^{38} + x^{37} + x^{36} + x^{35} + x^{33} \\
&\quad + x^{32} + x^{29} + x^{26} + x^{24} + x^{23} + x^{22} + x^{21} + x^{19} + x^{18} + x^{17} + x^{15} \\
&\quad + x^{14} + x^{11} + x^8 + x^3 + 1, \\
p_9(x) &= x^{59} + x^{56} + x^{43} + x^{40} + x^{11} + x^8, \\
p_{12}(x) &= x^{63} + x^{60} + x^{59} + x^{56} + x^{51} + x^{48} + x^{47} + x^{44} + x^{43} + x^{40} + x^{35} \\
&\quad + x^{32} + x^{15} + x^{12} + x^{11} + x^8 + x^3 + 1,
\end{aligned}$$

and the initial terms are $[1, 0, 0, 0, 1, 0, 0, 1, 0, 0, 0, 1, 0, 1, 0, 0]$.

For $\phi(W^o, 0, 1)$:

$$\begin{aligned}
p_0(x) &= x^{96} + x^{87} + x^{86} + x^{85} + x^{84} + x^{79} + x^{78} + x^{76} + x^{75} + x^{74} + x^{72} \\
&\quad + x^{69} + x^{68} + x^{62} + x^{60} + x^{59} + x^{54} + x^{52} + x^{50} + x^{44} + x^{43} + x^{42} \\
&\quad + x^{41} + x^{37} + x^{33}, \\
p_3(x) &= x^{63} + x^{60} + x^{57} + x^{55} + x^{54} + x^{52} + x^{51} + x^{49} + x^{48} + x^{46} + x^{45} \\
&\quad + x^{43} + x^{42} + x^{40} + x^{39} + x^{37} + x^{36} + x^{34} + x^{33} + x^{30} + x^{27} + x^{24} \\
&\quad + x^{21} + x^{18}, \\
p_6(x) &= x^{66} + x^{62} + x^{60} + x^{58} + x^{56} + x^{54} + x^{52} + x^{50} + x^{48} + x^{44} + x^{42} \\
&\quad + x^{38} + x^{36} + x^{34} + x^{32} + x^{28} + x^{26} + x^{20} + x^{18} + x^{12}, \\
p_9(x) &= x^{69} + x^{66} + x^{57} + x^{54} + x^{53} + x^{50} + x^{41} + x^{38} + x^{21} + x^{18} + x^9 + x^6, \\
p_{12}(x) &= x^{72} + x^{68} + x^{52} + x^{48} + x^{40} + x^{32} + x^{24} + x^{20} + x^8 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 1, 0, 0, 1, 1, 0, 1, 1, 0, 1, 0]$.

For $\phi(W^o, 1, 0)$:

$$\begin{aligned}
p_0(x) &= x^{84} + x^{75} + x^{74} + x^{73} + x^{70} + x^{68} + x^{67} + x^{66} + x^{61} + x^{60} + x^{58} \\
&\quad + x^{57} + x^{56} + x^{55} + x^{54} + x^{50} + x^{49} + x^{46} + x^{44} + x^{42} + x^{41} + x^{39} \\
&\quad + x^{38} + x^{37} + x^{36} + x^{35} + x^{34} + x^{33}, \\
p_3(x) &= x^{51} + x^{48} + x^{45} + x^{43} + x^{42} + x^{40} + x^{37} + x^{34} + x^{27} + x^{24} + x^{21}
\end{aligned}$$

$$\begin{aligned}
& + x^{18}, \\
p_6(x) &= x^{54} + x^{50} + x^{48} + x^{46} + x^{44} + x^{40} + x^{34} + x^{30} + x^{28} + x^{26} + x^{24} \\
& + x^{20} + x^{18} + x^{12}, \\
p_9(x) &= x^{57} + x^{54} + x^{41} + x^{38} + x^9 + x^6, \\
p_{12}(x) &= x^{60} + x^{56} + x^{48} + x^{44} + x^{40} + x^{32} + x^{12} + x^8 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 1, 0, 0, 0, 1, 0, 0, 0, 0, 1, 0, 0, 0]$.

For $\phi(W^o, 1, 1)$:

$$\begin{aligned}
p_0(x) &= x^{87} + x^{84} + x^{81} + x^{79} + x^{78} + x^{77} + x^{75} + x^{72} + x^{69} + x^{68} + x^{66} \\
& + x^{64} + x^{61} + x^{60} + x^{59} + x^{57} + x^{54} + x^{52} + x^{49} + x^{48} + x^{44} + x^{43} \\
& + x^{42}, \\
p_3(x) &= x^{67} + x^{64} + x^{61} + x^{58} + x^{57} + x^{56} + x^{55} + x^{53} + x^{52} + x^{51} + x^{50} \\
& + x^{48} + x^{46} + x^{44} + x^{43} + x^{41} + x^{39} + x^{38} + x^{37} + x^{35} + x^{34} + x^{33} \\
& + x^{31} + x^{28} + x^{25} + x^{24} + x^{23} + x^{22}, \\
p_6(x) &= x^{71} + x^{68} + x^{67} + x^{65} + x^{64} + x^{63} + x^{62} + x^{58} + x^{56} + x^{54} + x^{50} \\
& + x^{49} + x^{47} + x^{46} + x^{44} + x^{43} + x^{40} + x^{38} + x^{36} + x^{34} + x^{30} + x^{24} \\
& + x^{22} + x^{20} + x^{17} + x^{16} + x^{14} + x^{12} + x^3 + 1, \\
p_9(x) &= x^{59} + x^{56} + x^{43} + x^{40} + x^{11} + x^8, \\
p_{12}(x) &= x^{63} + x^{60} + x^{59} + x^{56} + x^{51} + x^{48} + x^{47} + x^{44} + x^{43} + x^{40} + x^{35} \\
& + x^{32} + x^{15} + x^{12} + x^{11} + x^8 + x^3 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 0, 0, 1, 0, 0, 1, 1, 0, 1, 0, 0]$.

Below is the transition function and output function of an 2-automaton that generates $T = M_{0,0}^o$:

Transition function $(n, j) \mapsto \delta(n, j)$ ($\Lambda(n) := [\delta(n, 0), \delta(n, 1)]$):

n	$\Lambda(n)$	n	$\Lambda(n)$	n	$\Lambda(n)$	n	$\Lambda(n)$	n	$\Lambda(n)$
0	[1, 2]	43	[71, 72]	86	[129, 130]	129	[165, 85]	172	[191, 75]
1	[3, 4]	44	[73, 74]	87	[79, 131]	130	[42, 12]	173	[192, 149]
2	[5, 6]	45	[75, 76]	88	[132, 63]	131	[166, 167]	174	[193, 146]
3	[7, 8]	46	[77, 78]	89	[133, 130]	132	[107, 168]	175	[158, 160]
4	[9, 10]	47	[78, 66]	90	[3, 134]	133	[103, 169]	176	[194, 128]
5	[11, 12]	48	[16, 79]	91	[63, 75]	134	[33, 170]	177	[195, 145]
6	[13, 14]	49	[80, 81]	92	[135, 64]	135	[125, 102]	178	[196, 65]
7	[15, 16]	50	[82, 6]	93	[74, 136]	136	[114, 171]	179	[21, 17]
8	[17, 18]	51	[83, 35]	94	[18, 134]	137	[172, 122]	180	[197, 140]
9	[19, 20]	52	[84, 85]	95	[137, 138]	138	[173, 45]	181	[198, 66]
10	[6, 21]	53	[86, 87]	96	[72, 81]	139	[85, 167]	182	[199, 142]
11	[22, 23]	54	[88, 89]	97	[139, 24]	140	[174, 175]	183	[200, 67]
12	[24, 20]	55	[90, 91]	98	[4, 27]	141	[176, 123]	184	[143, 72]
13	[25, 26]	56	[92, 37]	99	[128, 140]	142	[121, 111]	185	[201, 82]
14	[27, 28]	57	[45, 93]	100	[141, 136]	143	[167, 89]	186	[202, 17]

15	[29, 30]	58	[94, 37]	101	[142, 55]	144	[177, 107]	187	[203, 53]
16	[31, 32]	59	[95, 96]	102	[130, 23]	145	[178, 168]	188	[204, 93]
17	[14, 33]	60	[35, 84]	103	[59, 143]	146	[48, 94]	189	[87, 125]
18	[34, 35]	61	[97, 8]	104	[144, 15]	147	[169, 30]	190	[32, 42]
19	[36, 32]	62	[98, 99]	105	[140, 26]	148	[179, 99]	191	[89, 36]
20	[37, 38]	63	[100, 101]	106	[145, 62]	149	[171, 123]	192	[163, 50]
21	[12, 39]	64	[40, 10]	107	[146, 69]	150	[96, 114]	193	[105, 126]
22	[2, 40]	65	[102, 103]	108	[81, 142]	151	[10, 175]	194	[99, 96]
23	[41, 42]	66	[104, 105]	109	[147, 127]	152	[180, 83]	195	[38, 118]
24	[43, 7]	67	[106, 107]	110	[148, 62]	153	[181, 126]	196	[205, 170]
25	[44, 45]	68	[39, 91]	111	[149, 82]	154	[182, 169]	197	[206, 179]
26	[46, 47]	69	[50, 108]	112	[150, 21]	155	[175, 179]	198	[170, 106]
27	[8, 48]	70	[109, 102]	113	[151, 152]	156	[183, 39]	199	[168, 83]
28	[49, 50]	71	[110, 111]	114	[136, 153]	157	[164, 184]	200	[184, 47]
29	[51, 52]	72	[112, 106]	115	[154, 60]	158	[185, 36]	201	[207, 44]
30	[53, 53]	73	[113, 114]	116	[155, 56]	159	[47, 14]	202	[208, 94]
31	[54, 55]	74	[108, 44]	117	[156, 69]	160	[186, 184]	203	[93, 116]
32	[56, 24]	75	[115, 116]	118	[67, 78]	161	[187, 164]	204	[209, 160]
33	[26, 57]	76	[117, 118]	119	[157, 158]	162	[188, 131]	205	[210, 79]
34	[58, 59]	77	[119, 38]	120	[159, 74]	163	[138, 152]	206	[211, 76]
35	[60, 61]	78	[120, 121]	121	[160, 145]	164	[134, 27]	207	[212, 143]
36	[20, 15]	79	[30, 122]	122	[52, 60]	165	[189, 155]	208	[213, 57]
37	[62, 63]	80	[123, 121]	123	[153, 149]	166	[190, 133]	209	[122, 101]
38	[64, 65]	81	[124, 125]	124	[161, 56]	167	[76, 158]	210	[118, 7]
39	[23, 64]	82	[126, 40]	125	[131, 59]	168	[57, 138]	211	[111, 87]
40	[66, 67]	83	[55, 127]	126	[65, 73]	169	[70, 133]	212	[116, 163]
41	[68, 16]	84	[127, 128]	127	[162, 163]	170	[28, 146]	213	[101, 105]
42	[69, 70]	85	[61, 52]	128	[91, 164]	171	[152, 155]		

n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$
0	0	31	1	62	0	93	0	124	0	155	1	186	0
1	0	32	0	63	1	94	1	125	0	156	0	187	0
2	0	33	1	64	1	95	0	126	1	157	1	188	0
3	0	34	1	65	1	96	1	127	0	158	1	189	1
4	0	35	0	66	1	97	1	128	1	159	1	190	0
5	0	36	0	67	0	98	0	129	1	160	0	191	0
6	0	37	0	68	1	99	1	130	1	161	0	192	1
7	0	38	1	69	1	100	1	131	0	162	0	193	1
8	0	39	1	70	1	101	0	132	1	163	1	194	1
9	0	40	1	71	1	102	1	133	0	164	1	195	1
10	0	41	1	72	1	103	0	134	1	165	1	196	0
11	0	42	1	73	0	104	1	135	0	166	0	197	0
12	1	43	1	74	0	105	1	136	0	167	0	198	0

13	0	44	0	75	1	106	0	137	0	168	1	199	1
14	0	45	1	76	0	107	1	138	1	169	1	200	0
15	0	46	1	77	1	108	0	139	1	170	0	201	1
16	1	47	1	78	1	109	1	140	1	171	0	202	0
17	0	48	1	79	1	110	1	141	1	172	0	203	0
18	1	49	0	80	0	111	0	142	0	173	1	204	0
19	0	50	1	81	0	112	1	143	0	174	1	205	0
20	0	51	0	82	1	113	0	144	1	175	1	206	0
21	1	52	0	83	0	114	0	145	0	176	1	207	1
22	0	53	1	84	0	115	1	146	1	177	1	208	0
23	1	54	1	85	1	116	1	147	1	178	0	209	0
24	1	55	0	86	1	117	0	148	1	179	1	210	0
25	0	56	0	87	1	118	0	149	0	180	0	211	0
26	1	57	1	88	1	119	1	150	1	181	0	212	1
27	0	58	1	89	0	120	1	151	0	182	1	213	0
28	0	59	0	90	0	121	0	152	0	183	0		
29	0	60	0	91	1	122	0	153	0	184	0		
30	1	61	1	92	0	123	0	154	1	185	1		

UNIVERSITÉ DE STRASBOURG, CNRS, IRMA UMR 7501, F-67000 STRASBOURG, FRANCE
Email address: `guoniu.han@unistra.fr`

SCHOOL OF MATHEMATICS AND STATISTICS, HUAZHONG UNIVERSITY OF SCIENCE AND TECHNOLOGY, WUHAN, PR CHINA
Email address: `huyining@hust.edu.cn`