

PROOF OF THE CONTINUED FRACTION CONJECTURE FOR

$$(a, b) = (z^3 + z^2, z^4 + z)$$

GUO-NIU HAN* AND YINING HU*

ABSTRACT. The continued fraction conjecture claims that the continued fraction defined by the (a, b) -Thue-Morse sequence is algebraic over $\mathbb{F}_2(x)$ for all pairs of distinct elements (a, b) in $\mathbb{F}_2[x] \setminus \mathbb{F}_2$. In this paper we prove the conjecture for $(a, b) = (z^3 + z^2, z^4 + z)$.

1. INTRODUCTION

Let $\mathbb{F}_2$ be the finite field of cardinality 2. Given a sequence of polynomials $\mathbf{a} = (a_0(z), a_1(z), a_2(z), \dots)$ of elements from $\mathbb{F}_2[z] \setminus \mathbb{F}_2$, we define the infinite continued fraction

$$(1.1) \quad \text{CF}(\mathbf{a}(z)) = \frac{1}{a_0(z) + \frac{1}{a_1(z) + \frac{1}{a_2(z) + \frac{1}{\ddots}}}}$$

as the limit of the finite continued fractions

$$(1.2) \quad \text{CF}_n(\mathbf{a}(z)) = \frac{1}{a_0(z) + \frac{1}{a_1(z) + \frac{1}{\ddots + \frac{1}{a_n(z)}}}} \in \mathbb{F}_2((1/z)).$$

Conversely, each power series in $\mathbb{F}_2[[1/z]]$ without constant term

$$(1.3) \quad c_{-1}z^{-1} + c_{-2}z^{-2} + c_{-3}z^{-3} + \dots$$

admits a continued fraction expansion of form (1.1).

Let a and b be two distinct elements from $\mathbb{F}_2[z] \setminus \mathbb{F}_2$. We consider the continued fraction (1.1) associated with the (a, b) -Thue-Morse sequence

$$\mathbf{t} = (t_0(z), t_1(z), t_2(z), \dots) = (a, b, b, a, b, a, a, b, \dots)$$

that we shall denote

$$(1.4) \quad \text{CF}(a, b) := \text{CF}(\mathbf{t}).$$

Date: 9/04/2020 19:08:42 .

2010 Mathematics Subject Classification. 11B85, 11J70, 11B50, 11Y65, 05A15.

Key words and phrases. automatic sequence, continued fraction, Thue-Morse sequence.

* This paper was automatically generated from a computer program developed by the authors, with 8.4 minutes CPU time used.

Theorem 1.1. *When $(a, b) = (z^3 + z^2, z^4 + z)$, the two power series $\text{CF}(a, b)$ and $\text{CF}(b, a)$ are algebraic over $\mathbb{F}_2(z)$, with minimal polynomials of the form*

$$p_4(z)y^4 + p_3(z)y^3 + p_2(z)y^2 + p_1(z)y + p_0(z) = 0$$

where, for $\text{CF}(a, b)$,

$$\begin{aligned} p_0(z) &= z^{11} + z^{10} + z^8 + z^4 + z^2 + z + 1, \\ p_1(z) &= z^{15} + z^{12} + z^5 + z^2, \\ p_2(z) &= z^{18} + z^{16} + z^{14} + z^8 + z^6 + z^4, \\ p_3(z) &= z^{15} + z^{12} + z^5 + z^2, \\ p_4(z) &= z^{12} + z^{11} + z^8 + z^4 + z, \end{aligned}$$

and for $\text{CF}(b, a)$,

$$\begin{aligned} p_0(z) &= z^{11} + z^{10} + z^8 + z^6 + z, \\ p_1(z) &= z^{15} + z^{12} + z^5 + z^2, \\ p_2(z) &= z^{18} + z^{16} + z^{14} + z^8 + z^6 + z^4, \\ p_3(z) &= z^{15} + z^{12} + z^5 + z^2, \\ p_4(z) &= z^{12} + z^{11} + z^8 + z^6 + z^2 + z + 1. \end{aligned}$$

2. PROOF

Define the sequences of polynomials $(P_n(z))_{n \geq 0}$ and $(Q_n(z))_{n \geq 0}$ by

$$\begin{pmatrix} P_n(z) & Q_n(z) \\ P_{n-1}(z) & Q_{n-1}(z) \end{pmatrix} := \begin{pmatrix} t_n(z) & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} t_{n-1}(z) & 1 \\ 1 & 0 \end{pmatrix} \cdots \begin{pmatrix} t_0(z) & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}.$$

By the basic properties of continued fractions, we have

$$(2.1) \quad \text{CF}_n(a, b) = \frac{P_n(z)}{Q_n(z)} \in \mathbb{F}_2((1/z)).$$

As formula (2.1) is not efficient for calculating $\text{CF}_n(a, b)$, a fast method is to be derived and described as follows. Define

$$\begin{aligned} M_n(x) &= x^{\deg(t_{2^n-1})} \begin{pmatrix} t_{2^n-1}(1/x) & 1 \\ 1 & 0 \end{pmatrix} \times x^{\deg(t_{2^n-2})} \begin{pmatrix} t_{2^n-2}(1/x) & 1 \\ 1 & 0 \end{pmatrix} \times \\ &\quad \cdots \times x^{\deg(t_0)} \begin{pmatrix} t_0(1/x) & 1 \\ 1 & 0 \end{pmatrix}, \\ W_n(x) &= x^{\deg(\bar{t}_{2^n-1})} \begin{pmatrix} \bar{t}_{2^n-1}(1/x) & 1 \\ 1 & 0 \end{pmatrix} \times x^{\deg(\bar{t}_{2^n-2})} \begin{pmatrix} \bar{t}_{2^n-2}(1/x) & 1 \\ 1 & 0 \end{pmatrix} \times \\ &\quad \cdots \times x^{\deg(\bar{t}_0)} \begin{pmatrix} \bar{t}_0(1/x) & 1 \\ 1 & 0 \end{pmatrix}, \end{aligned}$$

where $\bar{t}$ is the (b, a) -Thue-Morse sequence. By the properties of the Thue-Morse sequence, we have for $n \geq 0$

$$\begin{aligned} M_{n+1}(x) &= W_n(x) \cdot M_n(x), \\ W_{n+1}(x) &= M_n(x) \cdot W_n(x). \end{aligned}$$

Let $x := 1/z$. For each non-zero polynomial $P(z)$, define $\tilde{P}(x)$ to be $P(1/x)$. Then,

$$\text{CF}_n(a, b) = \frac{P_n(z)}{Q_n(z)} = \frac{\tilde{P}_n(x)}{\tilde{Q}_n(x)} \in \mathbb{F}_2((x)) = \mathbb{F}_2((1/z)).$$

Comparing the definition of $M_n(x)$ with the definition of $P_n(x)$ and $Q_n(x)$, we see that

$$\begin{aligned} M_n(x)_{0,1} &= x^{d_n} \tilde{P}_{2^n-1}(x), \\ M_n(x)_{0,0} &= x^{d_n} \tilde{Q}_{2^n-1}(x), \end{aligned}$$

for some positive integer d_n . Hence,

$$(2.2) \quad \text{CF}_{2^{2^n}-1}(a, b) = \frac{\tilde{P}_{2^{2^n}-1}(x)}{\tilde{Q}_{2^{2^n}-1}(x)} = \frac{M_{2^n}(x)_{0,1}}{M_{2^n}(x)_{0,0}}.$$

Therefore, by the convergence theorem of continued fraction, the algebraicity of $\text{CF}(a, b)$ will be established if it is shown that both $M_{2^n}(x)_{0,1}$ and $M_{2^n}(x)_{0,0}$ converge to algebraic series in $\mathbb{F}_2[[x]]$.

Actually, we will prove that for all $0 \leq i, j \leq 1$ the four sequences $(M_{2^n}(x)_{i,j})_n$, $(M_{2^{n+1}}(x)_{i,j})_n$, $(W_{2^n}(x)_{i,j})_n$, and $(W_{2^{n+1}}(x)_{i,j})_n$ converge to algebraic series in $\mathbb{F}_2[[x]]$. For this purpose, we define four 2×2 matrices M^e, M^o, W^e, W^o as follows: For each $T \in \{M^e, M^o, W^e, W^o\}$ and $0 \leq i, j \leq 1$, $T_{i,j}$ is defined to be the unique solution in $\mathbb{F}_2[[x]]$ of the polynomial $\phi(T, i, j)$ under certain initial conditions; the polynomials $\phi(T, i, j)$ and initial conditions are given in Section 3. We will prove that these four matrices, whose components are algebraic by definition, are the limits of $(M_{2^n}(x))_n$, $(M_{2^{n+1}}(x))_n$, $(W_{2^n}(x))_n$, and $(W_{2^{n+1}}(x))_n$.

Let us explain how the polynomials $\phi(T, i, j)$ and initial conditions are found, and why the solutions exist and are unique. For $0 \leq i, j \leq 1$, the coefficients of the polynomial $\phi(M^e, i, j)$ (resp. $\phi(M^o, i, j)$, $\phi(W^e, i, j)$, and $\phi(W^o, i, j)$) are the Padé-Hermite approximants of type

$$(175, 175, 175, 175, 175)$$

of the vector

$$(1, T^3, T^6, T^9, T^{12}),$$

where $T = M_{12,i,j}$ (resp. $M_{11,i,j}$, $W_{12,i,j}$, and $W_{11,i,j}$), found by the Derksen algorithm. We take the first 8 terms of $M_{12,i,j}$ (resp. $M_{11,i,j}$, $W_{12,i,j}$, and $W_{11,i,j}$) as initial conditions for $\phi(M^e, i, j)$ (resp. $\phi(M^o, i, j)$, $\phi(W^e, i, j)$, and $\phi(W^o, i, j)$). The following fact will be used to ensure that the solution exists and is unique: let $P(x, y) \in \mathbb{F}_2[x, y]$ and for each series $f(x) = \sum_{n=0}^{\infty} a_n x^n \in \mathbb{F}_2[[x]]$ denote the partial sum $\sum_{j=0}^{n-1} a_j x^j$ by $f_n(x)$ for $n \geq 0$. If for some $n \geq 0$ and $a_0, a_1, \dots, a_{n-1} \in \mathbb{F}_2$ $P(x, \sum_{j=0}^{n-1} a_j x^j) = O(x^n)$ and $Q(x, y) := P(x, \sum_{j=0}^{n-1} a_j x^j + x^n y)$ can be written as $x^m \sum_{j=0}^{\infty} q_j(x) y^j$ where $q_j(x)$ are polynomials for $j \geq 0$, $q_1(0) = 1$, and $q_j(0) = 0$ for $j > 1$, then there exists a unique solution $f(x) \in \mathbb{F}_2[[x]]$ of $P(x, f(x)) = 0$ that satisfies the initial condition $f_n(x) = \sum_{j=0}^{n-1} a_j x^j$.

We state two lemmas concerning the four matrices M^e, M^o, W^e, W^o . The first one is about relations between them; the second, about the structure of the each matrix.

Lemma 2.1. *We have*

$$(2.3) \quad M^e = W^o \cdot M^o,$$

$$(2.4) \quad M^o = W^e \cdot M^e,$$

$$(2.5) \quad W^e = M^o \cdot W^o,$$

$$(2.6) \quad W^o = M^e \cdot W^e.$$

Proof. We give the proof of the identity

$$M_{0,0}^e = W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o,$$

the proofs of the others being similar.

First, we compute the minimal polynomials of $W_{0,0}^o M_{0,0}^o$ and $W_{0,1}^o M_{1,0}^o$. We know that

$$P(x, y) = \text{Res}_z (\phi(W^o, 0, 0)(x, z), z^{12} \cdot \phi(M^o, 0, 0)(x, y/z))$$

is an annihilating polynomial of $W_{0,0}^o M_{0,0}^o$. We use Padé-Hermite approximation to find a candidate for the minimal polynomial of $W_{0,0}^o M_{0,0}^o$, that will be called $\phi_0(x, y)$. To prove that $\phi_0(x, y)$ is indeed the minimal polynomial, it suffices to prove that it is an irreducible factor of $P(x, y)$ of multiplicity m and that $Q(x, y) := P(x, y)/\phi_0(x, y)^m$ is not an annihilating polynomial of $W_{0,0}^o M_{0,0}^o$. We verify the first point directly. For the second point, we truncate $W_{0,0}^o M_{0,0}^o$ to order 630 and substitute it for y in $Q(x, y)$. We get a series of valuation less than 630, which proves that $Q(x, y)$ is not an annihilating polynomial of $W_{0,0}^o M_{0,0}^o$. We find the minimal polynomial $\phi_1(x, y)$ of $W_{0,1}^o M_{1,0}^o$ in a similar way.

Now we prove that $\phi(M^e, 0, 0)$ is the minimal polynomial of $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$. We know that

$$S(x, y) = \text{Res}_z (\phi_0(x, z), \phi_1(x, y + z))$$

is an annihilating of $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$. We verify that $\phi(M^e, 0, 0)$ is an irreducible factor of $S(x, y)$ of multiplicity μ , and that $Q(x, y) := S(x, y)/\phi(M^e, 0, 0)^\mu$ is not an annihilating polynomial of $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$. To see the last point, we truncate $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$ to order 770 and substitute it for y in $Q(x, y)$. We get a series of valuation less than 770, and therefore $Q(x, y)$ is not an annihilating polynomial of $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$.

Finally, the first 16 terms of $M_{0,0}^e$ and $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$ coincide. As these first terms determine a unique solution of $\phi(M^e, 0, 0)$, we know that the two series are one and the same. $\square$

Define

$$R^e = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \quad \text{and} \quad R^o = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}.$$

Lemma 2.2. *For all integers $k \geq 2$ and $u = 2^{2k-1}$, the following identities hold.*

$$\begin{aligned} M^e[:14u] &= M^e[:7u] + x^u M^e[:6u] + x^{3u} M^e[:4u] + x^{4u} M^e[:3u] + x^{6u} M^e[:u] \\ &\quad + x^u M^e[:7u] + x^{2u} M^e[:6u] + x^{4u} M^e[:4u] + x^{5u} M^e[:3u] \\ &\quad + x^{7u} M^e[:u] + x^{2u} M^e[:7u] + x^{3u} M^e[:6u] + x^{5u} M^e[:4u] \\ &\quad + x^{6u} M^e[:3u] + x^{8u} M^e[:u] + x^{10u} M^e[:4u] + x^{12u} M^e[:2u] \\ &\quad + (x^{7u} + x^{8u} + x^{9u}) R^e, \\ W^e[:14u] &= W^e[:7u] + x^u W^e[:6u] + x^{3u} W^e[:4u] + x^{4u} W^e[:3u] + x^{6u} W^e[:u] \end{aligned}$$

$$\begin{aligned}
& + x^u W^e[:7u] + x^{2u} W^e[:6u] + x^{4u} W^e[:4u] + x^{5u} W^e[:3u] \\
& + x^{7u} W^e[:u] + x^{2u} W^e[:7u] + x^{3u} W^e[:6u] + x^{5u} W^e[:4u] \\
& + x^{6u} W^e[:3u] + x^{8u} W^e[:u] + x^{10u} W^e[:4u] + x^{12u} W^e[:2u] \\
& + (x^{7u} + x^{8u} + x^{9u}) R^e.
\end{aligned}$$

For all integers $k \geq 2$ and $u = 2^{2k}$, the following identities hold.

$$\begin{aligned}
M^o[:14u] &= M^o[:7u] + x^u M^o[:6u] + x^{3u} M^o[:4u] + x^{4u} M^o[:3u] + x^{6u} M^o[:u] \\
&+ x^u M^o[:7u] + x^{2u} M^o[:6u] + x^{4u} M^o[:4u] + x^{5u} M^o[:3u] \\
&+ x^{7u} M^o[:u] + x^{2u} M^o[:7u] + x^{3u} M^o[:6u] + x^{5u} M^o[:4u] \\
&+ x^{6u} M^o[:3u] + x^{8u} M^o[:u] + x^{10u} M^o[:4u] + x^{12u} M^o[:2u] \\
&+ (x^{7u} + x^{8u} + x^{9u}) R^o, \\
W^o[:14u] &= W^o[:7u] + x^u W^o[:6u] + x^{3u} W^o[:4u] + x^{4u} W^o[:3u] + x^{6u} W^o[:u] \\
&+ x^u W^o[:7u] + x^{2u} W^o[:6u] + x^{4u} W^o[:4u] + x^{5u} W^o[:3u] \\
&+ x^{7u} W^o[:u] + x^{2u} W^o[:7u] + x^{3u} W^o[:6u] + x^{5u} W^o[:4u] \\
&+ x^{6u} W^o[:3u] + x^{8u} W^o[:u] + x^{10u} W^o[:4u] + x^{12u} W^o[:2u] \\
&+ (x^{7u} + x^{8u} + x^{9u}) R^o.
\end{aligned}$$

Proof. To prove Lemma 2.2 we first construct an automaton for each sequence concerned, and then transform the conditions on infinitely many k 's into finitely many conditions on the states of the automaton. In the following, we will prove that for $T = M_{0,0}^o$, for all integer $k \geq 2$ and $u = 2^{2k}$,

$$\begin{aligned}
T[:14u] &= T[:7u] + x^u T[:6u] + x^{3u} T[:4u] + x^{4u} T[:3u] + x^{6u} T[:u] + x^u T[:7u] \\
&+ x^{2u} T[:6u] + x^{4u} T[:4u] + x^{5u} T[:3u] + x^{7u} T[:u] + x^{2u} T[:7u] \\
&+ x^{3u} T[:6u] + x^{5u} T[:4u] + x^{6u} T[:3u] + x^{8u} T[:u] + x^{10u} T[:4u] \\
&+ x^{12u} T[:2u].
\end{aligned}$$

The proof of the other 15 cases are similar. We break down the above identity into 7 parts:

$$\begin{aligned}
0 &= x^{7u} T[:u] + x^{6u} T[u:2u] + x^{4u} T[3u:4u] + x^{3u} T[4u:5u] + x^u T[6u:7u] \\
&+ T[7u:8u], \\
0 &= x^{8u} T[:u] + x^{6u} T[2u:3u] + x^{5u} T[3u:4u] + x^{3u} T[5u:6u] \\
&+ x^{2u} T[6u:7u] + T[8u:9u], \\
0 &= T[9u:10u], \\
0 &= x^{10u} T[:u] + T[10u:11u], \\
0 &= x^{10u} T[u:2u] + T[11u:12u], \\
0 &= x^{12u} T[:u] + x^{10u} T[2u:3u] + T[12u:13u], \\
0 &= x^{12u} T[u:2u] + x^{10u} T[3u:4u] + T[13u:14u],
\end{aligned}$$

which can be rewritten as

$$0 = T[[w]_2] + T[[1w]_2] + T[[11w]_2] + T[[100w]_2] + T[[110w]_2]$$

$$\begin{aligned}
(2.7) \quad & + T[[111w]_2], \\
& 0 = T[[w]_2] + T[[10w]_2] + T[[11w]_2] + T[[101w]_2] + T[[110w]_2] \\
(2.8) \quad & + T[[1000w]_2], \\
(2.9) \quad & 0 = T[[1001w]_2], \\
(2.10) \quad & 0 = T[[w]_2] + T[[1010w]_2], \\
(2.11) \quad & 0 = T[[1w]_2] + T[[1011w]_2], \\
(2.12) \quad & 0 = T[[w]_2] + T[[10w]_2] + T[[1100w]_2], \\
(2.13) \quad & 0 = T[[1w]_2] + T[[11w]_2] + T[[1101w]_2],
\end{aligned}$$

for all binary word w of length $2k$ and $w \neq 0^{2k}$, and

$$\begin{aligned}
& 0 = T[[w]_2] + T[[1w]_2] + T[[11w]_2] + T[[100w]_2] + T[[110w]_2] \\
(2.14) \quad & + T[[111w]_2], \\
& 0 = T[[w]_2] + T[[10w]_2] + T[[11w]_2] + T[[101w]_2] + T[[110w]_2] \\
(2.15) \quad & + T[[1000w]_2], \\
(2.16) \quad & 0 = T[[1001w]_2], \\
(2.17) \quad & 0 = T[[w]_2] + T[[1010w]_2], \\
(2.18) \quad & 0 = T[[1w]_2] + T[[1011w]_2], \\
(2.19) \quad & 0 = T[[w]_2] + T[[10w]_2] + T[[1100w]_2], \\
(2.20) \quad & 0 = T[[1w]_2] + T[[11w]_2] + T[[1101w]_2],
\end{aligned}$$

for $w = 0^{2k}$.

First we calculate an 2-automaton that generates T from its minimal polynomial and its first terms. This automaton has 382 states; its transition function and output function can be found in the annex. Let $A(s, w)$ denote the state reached after reading w from right to left starting from the state s , and τ the output function. Define

$$E_{2k} = \{A(i, w) : |w| = 2k, w \neq 0^{2k}\}.$$

Identities (2.7) through (2.20) can be written as

$$\begin{aligned}
& 0 = \tau(A(s, \epsilon)) + \tau(A(s, 1)) + \tau(A(s, 11)) + \tau(A(s, 100)) \\
(2.21) \quad & + \tau(A(s, 110)) + \tau(A(s, 111)), \\
& 0 = \tau(A(s, \epsilon)) + \tau(A(s, 10)) + \tau(A(s, 11)) + \tau(A(s, 101)) \\
(2.22) \quad & + \tau(A(s, 110)) + \tau(A(s, 1000)), \\
(2.23) \quad & 0 = \tau(A(s, 1001)), \\
(2.24) \quad & 0 = \tau(A(s, \epsilon)) + \tau(A(s, 1010)), \\
(2.25) \quad & 0 = \tau(A(s, 1)) + \tau(A(s, 1011)), \\
(2.26) \quad & 0 = \tau(A(s, \epsilon)) + \tau(A(s, 10)) + \tau(A(s, 1100)), \\
(2.27) \quad & 0 = \tau(A(s, 1)) + \tau(A(s, 11)) + \tau(A(s, 1101)),
\end{aligned}$$

for all $s \in E_{2k}$, and

$$\begin{aligned}
& 0 = \tau(A(s, \epsilon)) + \tau(A(s, 1)) + \tau(A(s, 11)) + \tau(A(s, 100)) \\
(2.28) \quad & + \tau(A(s, 110)) + \tau(A(s, 111)),
\end{aligned}$$

$$(2.29) \quad 0 = \tau(A(s, \epsilon)) + \tau(A(s, 10)) + \tau(A(s, 11)) + \tau(A(s, 101)) \\ + \tau(A(s, 110)) + \tau(A(s, 1000)),$$

$$(2.30) \quad 0 = \tau(A(s, 1001)),$$

$$(2.31) \quad 0 = \tau(A(s, \epsilon)) + \tau(A(s, 1010)),$$

$$(2.32) \quad 0 = \tau(A(s, 1)) + \tau(A(s, 1011)),$$

$$(2.33) \quad 0 = \tau(A(s, \epsilon)) + \tau(A(s, 10)) + \tau(A(s, 1100)),$$

$$(2.34) \quad 0 = \tau(A(s, 1)) + \tau(A(s, 11)) + \tau(A(s, 1101)),$$

for $s = A(i, 0^{2k})$. We find that $(A(i, 0^{16}), E_{16}) = (A(i, 0^{14}), E_{14})$, so that we only have to verify that identities (2.21) through (2.34) hold for $2 \leq k \leq 8$, which turns out to be true. $\square$

In the following lemma, we express M_{2k} , M_{2k+1} , W_{2k} , and W_{2k+1} in terms of M^e , M^o , W^e , and W^o .

Lemma 2.3. *For all integer $k \geq 2$, and $u = 2^{2k-1}$,*

$$M_{2k} = M^e[:7u] + x^u M^e[:6u] + x^{3u} M^e[:4u] + x^{4u} M^e[:3u] + x^{6u} M^e[:u] \\ + x^{7u} R^e,$$

$$W_{2k} = W^e[:7u] + x^u W^e[:6u] + x^{3u} W^e[:4u] + x^{4u} W^e[:3u] + x^{6u} W^e[:u] \\ + x^{7u} R^e.$$

For all integer $k \geq 2$, and $u = 2^{2k}$,

$$M_{2k+1} = M^o[:7u] + x^u M^o[:6u] + x^{3u} M^o[:4u] + x^{4u} M^o[:3u] + x^{6u} M^o[:u] \\ + x^{7u} R^o,$$

$$W_{2k+1} = W^o[:7u] + x^u W^o[:6u] + x^{3u} W^o[:4u] + x^{4u} W^o[:3u] + x^{6u} W^o[:u] \\ + x^{7u} R^o.$$

Proof. We call the four identities in Lemma 2.3 also by the name M_{2k} , W_{2k} , M_{2k+1} , and W_{2k+1} . For $n = 2$, the identities can be verified directly. For $n \geq 2$, we claim that

$$M_{2k} \wedge W_{2k} \Rightarrow M_{2k+1} \wedge W_{2k+1}, \\ M_{2k+1} \wedge W_{2k+1} \Rightarrow M_{2k+2} \wedge W_{2k+2}.$$

We give the proof of

$$M_{2k} \wedge W_{2k} \Rightarrow M_{2k+1},$$

the other ones being similar. Set $u = 2^{2k}$ and $v = 2^{2k-1}$. By definition and induction hypothesis, the left side of identity M_{2k+1} is equal to

$$(2.35) \quad W_{2k} M_{2k} = (W^e[:7v] + x^v W^e[:6v] + x^{3v} W^e[:4v] + x^{4v} W^e[:3v] + x^{6v} W^e[:v] \\ + x^{7v} R^e) \times (M^e[:7v] + x^v M^e[:6v] + x^{3v} M^e[:4v] + x^{4v} M^e[:3v] \\ + x^{6v} M^e[:v] + x^{7v} R^e);$$

Call this expression lhs . Note that both sides of identity M_{2k+1} have the same term of highest degree $x^{14v} R^o$. Therefore we only need to prove that their difference is $O(x^{14v})$. Using Lemma 2.1 it can be seen that the right side of identity M_{2k+1} is congruent, modulo x^{14v} , to

$$W^e[:14v]M^e[:14v] + x^{2v}W^e[:12v]M^e[:12v] + x^{6v}W^e[:8v]M^e[:8v] \\ + x^{8v}W^e[:6v]M^e[:6v] + x^{12v}W^e[:2v]M^e[:2v].$$

For all $n \leq 14$, replace the occurrences of $W^e[:n \cdot v]$ and $M^e[:n \cdot v]$ in the above expression by the reduction modulo $x^{n \cdot v}$ of the right side of the corresponding identity in Lemma 2.2 and get a new expression, which we call *rhs*. Define

$$\begin{aligned} X &:= x^v, \\ a_n &:= W^e[n \cdot v : (n+1) \cdot v] / X^n, \\ b_n &:= M^e[n \cdot v : (n+1) \cdot v] / X^n, \\ c &:= R^e. \end{aligned}$$

Using the notation introduced above, we can represent the expressions *lhs* (2.35) and *rhs* as polynomials in $\mathbb{F}_2[a_1, \dots, a_{14}, b_1, \dots, b_{14}, c][X]$. Note that it is not a problem that a_j commutes with b_k while W^e does not commute with M^e , because in the expressions concerned, the products of W^e -terms and M^e -terms are always in the same order. We let the computer do the simplification and check that the difference between these two polynomials is indeed $O(X^{14})$, which completes the proof. $\square$

Proof of Theorem 1.1. We prove the theorem for $\text{CF}(a, b)$; for $\text{CF}(b, a)$, the proof is similar. By Lemma 2.3, we have For all $0 \leq j, k \leq 1$,

$$\begin{aligned} \lim_{n \rightarrow \infty} M_{2n,j,k} &= M_{j,k}^e, \\ \lim_{n \rightarrow \infty} M_{2n+1,j,k} &= M_{j,k}^o, \\ \lim_{n \rightarrow \infty} W_{2n,j,k} &= W_{j,k}^e, \\ \lim_{n \rightarrow \infty} W_{2n+1,j,k} &= W_{j,k}^o. \end{aligned}$$

Let $z = 1/x$. By the convergence theorem and identity (2.2),

$$\text{CF}(a, b) = \frac{M_{0,1}^e(x)}{M_{0,0}^e(x)}.$$

By definition, $\phi(M^e, 0, 1)$ and $\phi(M^e, 0, 0)$ are minimal polynomials of $M_{0,1}^e$ and $M_{0,0}^e$. Therefore

$$P(x, y) = \text{Res}_t (\phi(M^e, 0, 1)(x, t), y^{12}\phi(M^e, 0, 1)(x, t/y))$$

is an annihilating polynomial of $f(x) = M_{0,1}^e/M_{0,0}^e$.

Define

$$Q(x, y) = q_4(x)y^4 + q_3(x)y^3 + q_2(x)y^2 + q_1(x)y + q_0(x),$$

where

$$\begin{aligned} q_0(x) &= x^{18} + x^{17} + x^{16} + x^{14} + x^{10} + x^8 + x^7, \\ q_1(x) &= x^{16} + x^{13} + x^6 + x^3, \\ q_2(x) &= x^{14} + x^{12} + x^{10} + x^4 + x^2 + 1, \\ q_3(x) &= x^{16} + x^{13} + x^6 + x^3, \\ q_4(x) &= x^{17} + x^{14} + x^{10} + x^7 + x^6. \end{aligned}$$

The polynomial $Q(x, y)$ is the candidate for the minimal polynomial of $f(x)$ found by Padé-Hermite approximation. To prove that it is indeed the minimal polynomial

of $f(x)$, we only need to prove that it is an irreducible factor of $P(x, y)$ of multiplicity m and $R(x, y) := P(x, y)/Q(x, y)^m$ is not an annihilating polynomial of $f(x)$. We verify the first point directly. For the second point, we find that when we truncate $f(x)$ to order 216, and substitute it for y in $R(z, y)$, we get a series with valuation smaller than 216, which proves that $R(z, y)$ is not an annihilating polynomial of $f(x)$. Finally, $z^{12}Q(1/z, y)$ is the minimal polynomial of $\text{CF}(a, b) = f(1/z)$. $\square$

3. ANNEXE

All 16 polynomials are of the form

$$p_0(x) + p_3(x)y^3 + p_6(x)y^6 + p_9(x)y^9 + p_{12}(x)y^{12}.$$

The coefficients $p_j(x)$ and the 16 initial terms to determine the solutions uniquely are given below.

For $\phi(M^e, 0, 0)$:

$$\begin{aligned} p_0(x) &= x^{102} + x^{96} + x^{90} + x^{88} + x^{84} + x^{82} + x^{80} + x^{76} + x^{74} + x^{70} + x^{62} \\ &\quad + x^{60} + x^{58} + x^{56} + x^{54} + x^{48} + x^{44} + x^{42} + x^{40} + x^{38} + x^{36}, \\ p_3(x) &= x^{94} + x^{92} + x^{90} + x^{86} + x^{80} + x^{78} + x^{76} + x^{74} + x^{68} + x^{66} + x^{62} \\ &\quad + x^{60} + x^{54} + x^{52} + x^{50} + x^{46} + x^{40} + x^{38} + x^{36} + x^{34} + x^{28} + x^{26} \\ &\quad + x^{22} + x^{20}, \\ p_6(x) &= x^{94} + x^{92} + x^{86} + x^{84} + x^{82} + x^{78} + x^{74} + x^{70} + x^{68} + x^{64} + x^{62} \\ &\quad + x^{60} + x^{58} + x^{56} + x^{54} + x^{52} + x^{42} + x^{40} + x^{38} + x^{34} + x^{30} + x^{28} \\ &\quad + x^{24} + x^{22} + x^{20} + x^{18} + x^{16} + x^6 + x^4 + 1, \\ p_9(x) &= x^{92} + x^{84} + x^{82} + x^{76} + x^{74} + x^{72} + x^{66} + x^{64} + x^{62} + x^{56} + x^{54} \\ &\quad + x^{52} + x^{46} + x^{44} + x^{42} + x^{36} + x^{34} + x^{32} + x^{26} + x^{24} + x^{22} + x^{16} \\ &\quad + x^{14} + x^6, \\ p_{12}(x) &= x^{92} + x^{88} + x^{76} + x^{72} + x^{68} + x^{64} + x^{60} + x^{56} + x^{52} + x^{48} + x^{44} \\ &\quad + x^{40} + x^{36} + x^{32} + x^{28} + x^{24} + x^{20} + x^{16} + x^4 + 1, \end{aligned}$$

and the initial terms are $[1, 0, 0, 0, 0, 0, 0, 0, 1, 0, 0, 0, 1, 0, 0, 0]$.

For $\phi(M^e, 0, 1)$:

$$\begin{aligned} p_0(x) &= x^{99} + x^{96} + x^{95} + x^{94} + x^{93} + x^{92} + x^{87} + x^{86} + x^{85} + x^{84} + x^{78} \\ &\quad + x^{73} + x^{72} + x^{70} + x^{69} + x^{68} + x^{67} + x^{65} + x^{63} + x^{61} + x^{58} + x^{54} \\ &\quad + x^{53} + x^{51} + x^{50} + x^{48} + x^{47} + x^{40} + x^{39}, \\ p_3(x) &= x^{92} + x^{91} + x^{86} + x^{85} + x^{84} + x^{83} + x^{82} + x^{81} + x^{78} + x^{77} + x^{76} \\ &\quad + x^{75} + x^{74} + x^{73} + x^{68} + x^{67} + x^{52} + x^{51} + x^{46} + x^{45} + x^{44} + x^{43} \\ &\quad + x^{42} + x^{41} + x^{38} + x^{37} + x^{36} + x^{35} + x^{34} + x^{33} + x^{28} + x^{27}, \\ p_6(x) &= x^{90} + x^{89} + x^{88} + x^{86} + x^{85} + x^{83} + x^{82} + x^{80} + x^{79} + x^{78} + x^{70} \\ &\quad + x^{69} + x^{68} + x^{66} + x^{65} + x^{63} + x^{62} + x^{60} + x^{59} + x^{58} + x^{50} + x^{49} \\ &\quad + x^{48} + x^{46} + x^{45} + x^{43} + x^{42} + x^{40} + x^{39} + x^{38} + x^{30} + x^{29} + x^{28} \\ &\quad + x^{26} + x^{25} + x^{23} + x^{22} + x^{20} + x^{19} + x^{18}, \end{aligned}$$

$$\begin{aligned}
p_9(x) &= x^{88} + x^{87} + x^{84} + x^{83} + x^{80} + x^{79} + x^{78} + x^{77} + x^{74} + x^{73} + x^{70} \\
&\quad + x^{69} + x^{68} + x^{67} + x^{64} + x^{63} + x^{60} + x^{59} + x^{58} + x^{57} + x^{54} + x^{53} \\
&\quad + x^{50} + x^{49} + x^{48} + x^{47} + x^{44} + x^{43} + x^{40} + x^{39} + x^{38} + x^{37} + x^{34} \\
&\quad + x^{33} + x^{30} + x^{29} + x^{28} + x^{27} + x^{24} + x^{23} + x^{20} + x^{19} + x^{18} + x^{17} \\
&\quad + x^{14} + x^{13} + x^{10} + x^9, \\
p_{12}(x) &= x^{86} + x^{85} + x^{84} + x^{78} + x^{77} + x^{76} + x^{74} + x^{73} + x^{72} + x^{70} + x^{69} \\
&\quad + x^{68} + x^{66} + x^{65} + x^{64} + x^{62} + x^{61} + x^{60} + x^{58} + x^{57} + x^{56} + x^{54} \\
&\quad + x^{53} + x^{52} + x^{50} + x^{49} + x^{48} + x^{46} + x^{45} + x^{44} + x^{42} + x^{41} + x^{40} \\
&\quad + x^{38} + x^{37} + x^{36} + x^{34} + x^{33} + x^{32} + x^{30} + x^{29} + x^{28} + x^{26} + x^{25} \\
&\quad + x^{24} + x^{22} + x^{21} + x^{20} + x^{18} + x^{17} + x^{16} + x^{14} + x^{13} + x^{12} + x^{10} \\
&\quad + x^9 + x^8 + x^2 + x + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 1, 1, 1, 1, 1, 1, 0, 0, 1, 1, 1, 0]$.

For $\phi(M^e, 1, 0)$:

$$\begin{aligned}
p_0(x) &= x^{99} + x^{96} + x^{95} + x^{94} + x^{93} + x^{92} + x^{87} + x^{86} + x^{85} + x^{84} + x^{78} \\
&\quad + x^{73} + x^{72} + x^{70} + x^{69} + x^{68} + x^{67} + x^{65} + x^{63} + x^{61} + x^{58} + x^{54} \\
&\quad + x^{53} + x^{51} + x^{50} + x^{48} + x^{47} + x^{40} + x^{39}, \\
p_3(x) &= x^{92} + x^{91} + x^{86} + x^{85} + x^{84} + x^{83} + x^{82} + x^{81} + x^{78} + x^{77} + x^{76} \\
&\quad + x^{75} + x^{74} + x^{73} + x^{68} + x^{67} + x^{52} + x^{51} + x^{46} + x^{45} + x^{44} + x^{43} \\
&\quad + x^{42} + x^{41} + x^{38} + x^{37} + x^{36} + x^{35} + x^{34} + x^{33} + x^{28} + x^{27}, \\
p_6(x) &= x^{90} + x^{89} + x^{88} + x^{86} + x^{85} + x^{83} + x^{82} + x^{80} + x^{79} + x^{78} + x^{70} \\
&\quad + x^{69} + x^{68} + x^{66} + x^{65} + x^{63} + x^{62} + x^{60} + x^{59} + x^{58} + x^{50} + x^{49} \\
&\quad + x^{48} + x^{46} + x^{45} + x^{43} + x^{42} + x^{40} + x^{39} + x^{38} + x^{30} + x^{29} + x^{28} \\
&\quad + x^{26} + x^{25} + x^{23} + x^{22} + x^{20} + x^{19} + x^{18}, \\
p_9(x) &= x^{88} + x^{87} + x^{84} + x^{83} + x^{80} + x^{79} + x^{78} + x^{77} + x^{74} + x^{73} + x^{70} \\
&\quad + x^{69} + x^{68} + x^{67} + x^{64} + x^{63} + x^{60} + x^{59} + x^{58} + x^{57} + x^{54} + x^{53} \\
&\quad + x^{50} + x^{49} + x^{48} + x^{47} + x^{44} + x^{43} + x^{40} + x^{39} + x^{38} + x^{37} + x^{34} \\
&\quad + x^{33} + x^{30} + x^{29} + x^{28} + x^{27} + x^{24} + x^{23} + x^{20} + x^{19} + x^{18} + x^{17} \\
&\quad + x^{14} + x^{13} + x^{10} + x^9, \\
p_{12}(x) &= x^{86} + x^{85} + x^{84} + x^{78} + x^{77} + x^{76} + x^{74} + x^{73} + x^{72} + x^{70} + x^{69} \\
&\quad + x^{68} + x^{66} + x^{65} + x^{64} + x^{62} + x^{61} + x^{60} + x^{58} + x^{57} + x^{56} + x^{54} \\
&\quad + x^{53} + x^{52} + x^{50} + x^{49} + x^{48} + x^{46} + x^{45} + x^{44} + x^{42} + x^{41} + x^{40} \\
&\quad + x^{38} + x^{37} + x^{36} + x^{34} + x^{33} + x^{32} + x^{30} + x^{29} + x^{28} + x^{26} + x^{25} \\
&\quad + x^{24} + x^{22} + x^{21} + x^{20} + x^{18} + x^{17} + x^{16} + x^{14} + x^{13} + x^{12} + x^{10} \\
&\quad + x^9 + x^8 + x^2 + x + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 1, 1, 1, 1, 1, 1, 0, 0, 1, 1, 1, 0]$.

For $\phi(M^e, 1, 1)$:

$$\begin{aligned}
p_0(x) &= x^{96} + x^{88} + x^{86} + x^{82} + x^{80} + x^{78} + x^{76} + x^{72} + x^{70} + x^{54} + x^{52} \\
&\quad + x^{46} + x^{42}, \\
p_3(x) &= x^{84} + x^{80} + x^{74} + x^{72} + x^{68} + x^{64} + x^{62} + x^{58} + x^{44} + x^{40} + x^{34} \\
&\quad + x^{32} + x^{28} + x^{24} + x^{22} + x^{18}, \\
p_6(x) &= x^{84} + x^{76} + x^{74} + x^{72} + x^{70} + x^{68} + x^{66} + x^{60} + x^{58} + x^{56} + x^{54} \\
&\quad + x^{52} + x^{48} + x^{44} + x^{42} + x^{40} + x^{36} + x^{34} + x^{32} + x^{30} + x^{28} + x^{26} \\
&\quad + x^{20} + x^{18} + x^{16} + x^{14} + x^{12} + x^8 + x^2 + 1, \\
p_9(x) &= x^{80} + x^{78} + x^{76} + x^{70} + x^{68} + x^{66} + x^{60} + x^{58} + x^{56} + x^{50} + x^{48} \\
&\quad + x^{46} + x^{40} + x^{38} + x^{36} + x^{30} + x^{28} + x^{26} + x^{20} + x^{18} + x^{16} + x^{10} \\
&\quad + x^8 + x^6, \\
p_{12}(x) &= x^{80} + x^{78} + x^{74} + x^{70} + x^{66} + x^{62} + x^{58} + x^{54} + x^{50} + x^{46} + x^{42} \\
&\quad + x^{38} + x^{34} + x^{30} + x^{26} + x^{22} + x^{18} + x^{14} + x^{10} + x^6 + x^2 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 0, 1, 0, 1, 0, 1, 0, 1, 0, 0, 0]$.

For $\phi(W^e, 0, 0)$:

$$\begin{aligned}
p_0(x) &= x^{96} + x^{84} + x^{82} + x^{80} + x^{76} + x^{72} + x^{68} + x^{64} + x^{60} + x^{58} + x^{56} \\
&\quad + x^{54} + x^{52} + x^{50} + x^{42} + x^{40} + x^{36}, \\
p_3(x) &= x^{84} + x^{80} + x^{76} + x^{70} + x^{68} + x^{60} + x^{44} + x^{40} + x^{36} + x^{30} + x^{28} \\
&\quad + x^{20}, \\
p_6(x) &= x^{84} + x^{72} + x^{70} + x^{68} + x^{62} + x^{60} + x^{58} + x^{56} + x^{48} + x^{44} + x^{42} \\
&\quad + x^{40} + x^{32} + x^{30} + x^{28} + x^{22} + x^{20} + x^{18} + x^{16} + x^8 + x^2 + 1, \\
p_9(x) &= x^{80} + x^{78} + x^{76} + x^{70} + x^{68} + x^{66} + x^{60} + x^{58} + x^{56} + x^{50} + x^{48} \\
&\quad + x^{46} + x^{40} + x^{38} + x^{36} + x^{30} + x^{28} + x^{26} + x^{20} + x^{18} + x^{16} + x^{10} \\
&\quad + x^8 + x^6, \\
p_{12}(x) &= x^{80} + x^{78} + x^{74} + x^{70} + x^{66} + x^{62} + x^{58} + x^{54} + x^{50} + x^{46} + x^{42} \\
&\quad + x^{38} + x^{34} + x^{30} + x^{26} + x^{22} + x^{18} + x^{14} + x^{10} + x^6 + x^2 + 1,
\end{aligned}$$

and the initial terms are $[1, 0, 0, 0, 0, 0, 1, 0, 1, 0, 1, 0, 0, 0, 0, 0]$.

For $\phi(W^e, 0, 1)$:

$$\begin{aligned}
p_0(x) &= x^{99} + x^{94} + x^{93} + x^{91} + x^{90} + x^{89} + x^{88} + x^{85} + x^{78} + x^{77} + x^{75} \\
&\quad + x^{72} + x^{69} + x^{68} + x^{67} + x^{66} + x^{63} + x^{62} + x^{60} + x^{54} + x^{51} + x^{49} \\
&\quad + x^{48} + x^{47} + x^{46} + x^{45} + x^{44} + x^{40} + x^{39}, \\
p_3(x) &= x^{92} + x^{91} + x^{86} + x^{85} + x^{84} + x^{83} + x^{82} + x^{81} + x^{78} + x^{77} + x^{76} \\
&\quad + x^{75} + x^{74} + x^{73} + x^{68} + x^{67} + x^{52} + x^{51} + x^{46} + x^{45} + x^{44} + x^{43} \\
&\quad + x^{42} + x^{41} + x^{38} + x^{37} + x^{36} + x^{35} + x^{34} + x^{33} + x^{28} + x^{27}, \\
p_6(x) &= x^{90} + x^{89} + x^{88} + x^{86} + x^{85} + x^{83} + x^{82} + x^{80} + x^{79} + x^{78} + x^{70}
\end{aligned}$$

$$\begin{aligned}
& + x^{69} + x^{68} + x^{66} + x^{65} + x^{63} + x^{62} + x^{60} + x^{59} + x^{58} + x^{50} + x^{49} \\
& + x^{48} + x^{46} + x^{45} + x^{43} + x^{42} + x^{40} + x^{39} + x^{38} + x^{30} + x^{29} + x^{28} \\
& + x^{26} + x^{25} + x^{23} + x^{22} + x^{20} + x^{19} + x^{18}, \\
p_9(x) &= x^{88} + x^{87} + x^{84} + x^{83} + x^{80} + x^{79} + x^{78} + x^{77} + x^{74} + x^{73} + x^{70} \\
& + x^{69} + x^{68} + x^{67} + x^{64} + x^{63} + x^{60} + x^{59} + x^{58} + x^{57} + x^{54} + x^{53} \\
& + x^{50} + x^{49} + x^{48} + x^{47} + x^{44} + x^{43} + x^{40} + x^{39} + x^{38} + x^{37} + x^{34} \\
& + x^{33} + x^{30} + x^{29} + x^{28} + x^{27} + x^{24} + x^{23} + x^{20} + x^{19} + x^{18} + x^{17} \\
& + x^{14} + x^{13} + x^{10} + x^9, \\
p_{12}(x) &= x^{86} + x^{85} + x^{84} + x^{78} + x^{77} + x^{76} + x^{74} + x^{73} + x^{72} + x^{70} + x^{69} \\
& + x^{68} + x^{66} + x^{65} + x^{64} + x^{62} + x^{61} + x^{60} + x^{58} + x^{57} + x^{56} + x^{54} \\
& + x^{53} + x^{52} + x^{50} + x^{49} + x^{48} + x^{46} + x^{45} + x^{44} + x^{42} + x^{41} + x^{40} \\
& + x^{38} + x^{37} + x^{36} + x^{34} + x^{33} + x^{32} + x^{30} + x^{29} + x^{28} + x^{26} + x^{25} \\
& + x^{24} + x^{22} + x^{21} + x^{20} + x^{18} + x^{17} + x^{16} + x^{14} + x^{13} + x^{12} + x^{10} \\
& + x^9 + x^8 + x^2 + x + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 1, 0, 0, 1, 0, 0, 0, 1, 0, 1, 0, 0]$.

For $\phi(W^e, 1, 0)$:

$$\begin{aligned}
p_0(x) &= x^{99} + x^{94} + x^{93} + x^{91} + x^{90} + x^{89} + x^{88} + x^{85} + x^{78} + x^{77} + x^{75} \\
& + x^{72} + x^{69} + x^{68} + x^{67} + x^{66} + x^{63} + x^{62} + x^{60} + x^{54} + x^{51} + x^{49} \\
& + x^{48} + x^{47} + x^{46} + x^{45} + x^{44} + x^{40} + x^{39}, \\
p_3(x) &= x^{92} + x^{91} + x^{86} + x^{85} + x^{84} + x^{83} + x^{82} + x^{81} + x^{78} + x^{77} + x^{76} \\
& + x^{75} + x^{74} + x^{73} + x^{68} + x^{67} + x^{52} + x^{51} + x^{46} + x^{45} + x^{44} + x^{43} \\
& + x^{42} + x^{41} + x^{38} + x^{37} + x^{36} + x^{35} + x^{34} + x^{33} + x^{28} + x^{27}, \\
p_6(x) &= x^{90} + x^{89} + x^{88} + x^{86} + x^{85} + x^{83} + x^{82} + x^{80} + x^{79} + x^{78} + x^{70} \\
& + x^{69} + x^{68} + x^{66} + x^{65} + x^{63} + x^{62} + x^{60} + x^{59} + x^{58} + x^{50} + x^{49} \\
& + x^{48} + x^{46} + x^{45} + x^{43} + x^{42} + x^{40} + x^{39} + x^{38} + x^{30} + x^{29} + x^{28} \\
& + x^{26} + x^{25} + x^{23} + x^{22} + x^{20} + x^{19} + x^{18}, \\
p_9(x) &= x^{88} + x^{87} + x^{84} + x^{83} + x^{80} + x^{79} + x^{78} + x^{77} + x^{74} + x^{73} + x^{70} \\
& + x^{69} + x^{68} + x^{67} + x^{64} + x^{63} + x^{60} + x^{59} + x^{58} + x^{57} + x^{54} + x^{53} \\
& + x^{50} + x^{49} + x^{48} + x^{47} + x^{44} + x^{43} + x^{40} + x^{39} + x^{38} + x^{37} + x^{34} \\
& + x^{33} + x^{30} + x^{29} + x^{28} + x^{27} + x^{24} + x^{23} + x^{20} + x^{19} + x^{18} + x^{17} \\
& + x^{14} + x^{13} + x^{10} + x^9, \\
p_{12}(x) &= x^{86} + x^{85} + x^{84} + x^{78} + x^{77} + x^{76} + x^{74} + x^{73} + x^{72} + x^{70} + x^{69} \\
& + x^{68} + x^{66} + x^{65} + x^{64} + x^{62} + x^{61} + x^{60} + x^{58} + x^{57} + x^{56} + x^{54} \\
& + x^{53} + x^{52} + x^{50} + x^{49} + x^{48} + x^{46} + x^{45} + x^{44} + x^{42} + x^{41} + x^{40} \\
& + x^{38} + x^{37} + x^{36} + x^{34} + x^{33} + x^{32} + x^{30} + x^{29} + x^{28} + x^{26} + x^{25}
\end{aligned}$$

$$\begin{aligned}
& + x^{24} + x^{22} + x^{21} + x^{20} + x^{18} + x^{17} + x^{16} + x^{14} + x^{13} + x^{12} + x^{10} \\
& + x^9 + x^8 + x^2 + x + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 1, 0, 0, 1, 0, 0, 0, 1, 0, 1, 0, 0]$.

For $\phi(W^e, 1, 1)$:

$$\begin{aligned}
p_0(x) &= x^{102} + x^{92} + x^{88} + x^{84} + x^{74} + x^{72} + x^{68} + x^{66} + x^{60} + x^{54} + x^{46} \\
&\quad + x^{44} + x^{42}, \\
p_3(x) &= x^{94} + x^{92} + x^{90} + x^{88} + x^{86} + x^{82} + x^{80} + x^{78} + x^{76} + x^{74} + x^{68} \\
&\quad + x^{66} + x^{64} + x^{62} + x^{60} + x^{58} + x^{54} + x^{52} + x^{50} + x^{48} + x^{46} + x^{42} \\
&\quad + x^{40} + x^{38} + x^{36} + x^{34} + x^{28} + x^{26} + x^{24} + x^{22} + x^{20} + x^{18}, \\
p_6(x) &= x^{94} + x^{92} + x^{88} + x^{86} + x^{80} + x^{78} + x^{76} + x^{74} + x^{70} + x^{68} + x^{62} \\
&\quad + x^{54} + x^{48} + x^{44} + x^{38} + x^{36} + x^{34} + x^{30} + x^{28} + x^{22} + x^{12} + x^6 \\
&\quad + x^4 + 1, \\
p_9(x) &= x^{92} + x^{84} + x^{82} + x^{76} + x^{74} + x^{72} + x^{66} + x^{64} + x^{62} + x^{56} + x^{54} \\
&\quad + x^{52} + x^{46} + x^{44} + x^{42} + x^{36} + x^{34} + x^{32} + x^{26} + x^{24} + x^{22} + x^{16} \\
&\quad + x^{14} + x^6, \\
p_{12}(x) &= x^{92} + x^{88} + x^{76} + x^{72} + x^{68} + x^{64} + x^{60} + x^{56} + x^{52} + x^{48} + x^{44} \\
&\quad + x^{40} + x^{36} + x^{32} + x^{28} + x^{24} + x^{20} + x^{16} + x^4 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 0, 0, 0, 1, 0, 0, 0, 0, 0, 0, 0]$.

For $\phi(M^o, 0, 0)$:

$$\begin{aligned}
p_0(x) &= x^{99} + x^{96} + x^{91} + x^{89} + x^{84} + x^{83} + x^{82} + x^{77} + x^{73} + x^{72} + x^{69} \\
&\quad + x^{68} + x^{62} + x^{61} + x^{60} + x^{59} + x^{58} + x^{56} + x^{54} + x^{50} + x^{47} + x^{46} \\
&\quad + x^{45} + x^{44} + x^{42} + x^{40} + x^{39} + x^{37} + x^{36}, \\
p_3(x) &= x^{94} + x^{92} + x^{91} + x^{90} + x^{88} + x^{85} + x^{79} + x^{78} + x^{77} + x^{72} + x^{71} \\
&\quad + x^{70} + x^{69} + x^{67} + x^{65} + x^{64} + x^{54} + x^{52} + x^{51} + x^{50} + x^{48} + x^{45} \\
&\quad + x^{39} + x^{38} + x^{37} + x^{32} + x^{31} + x^{30} + x^{29} + x^{27} + x^{25} + x^{24}, \\
p_6(x) &= x^{94} + x^{92} + x^{91} + x^{90} + x^{88} + x^{83} + x^{82} + x^{80} + x^{79} + x^{78} + x^{77} \\
&\quad + x^{75} + x^{74} + x^{73} + x^{71} + x^{70} + x^{68} + x^{66} + x^{65} + x^{64} + x^{63} + x^{62} \\
&\quad + x^{60} + x^{59} + x^{56} + x^{55} + x^{52} + x^{51} + x^{49} + x^{43} + x^{41} + x^{39} + x^{38} \\
&\quad + x^{37} + x^{35} + x^{34} + x^{33} + x^{31} + x^{30} + x^{28} + x^{26} + x^{25} + x^{24} + x^{23} \\
&\quad + x^{22} + x^{20} + x^{19} + x^{16} + x^{15} + x^{14} + x^{10} + x^9 + x^8 + x^2 + x + 1, \\
p_9(x) &= x^{94} + x^{93} + x^{92} + x^{14} + x^{13} + x^{12}, \\
p_{12}(x) &= x^{94} + x^{93} + x^{92} + x^{86} + x^{85} + x^{84} + x^{78} + x^{77} + x^{76} + x^{74} + x^{73} \\
&\quad + x^{72} + x^{70} + x^{69} + x^{68} + x^{66} + x^{65} + x^{64} + x^{62} + x^{61} + x^{60} + x^{58} \\
&\quad + x^{57} + x^{56} + x^{54} + x^{53} + x^{52} + x^{50} + x^{49} + x^{48} + x^{46} + x^{45} + x^{44} \\
&\quad + x^{42} + x^{41} + x^{40} + x^{38} + x^{37} + x^{36} + x^{34} + x^{33} + x^{32} + x^{30} + x^{29}
\end{aligned}$$

$$\begin{aligned}
& + x^{28} + x^{26} + x^{25} + x^{24} + x^{22} + x^{21} + x^{20} + x^{18} + x^{17} + x^{16} + x^{10} \\
& + x^9 + x^8 + x^2 + x + 1,
\end{aligned}$$

and the initial terms are $[1, 0, 0, 0, 0, 0, 1, 1, 1, 1, 1, 0, 1, 0, 1, 0]$.

For $\phi(M^o, 0, 1)$:

$$\begin{aligned}
p_0(x) &= x^{100} + x^{98} + x^{95} + x^{93} + x^{89} + x^{88} + x^{86} + x^{85} + x^{84} + x^{83} + x^{78} \\
&+ x^{76} + x^{74} + x^{73} + x^{66} + x^{65} + x^{63} + x^{62} + x^{61} + x^{58} + x^{57} + x^{56} \\
&+ x^{55} + x^{50} + x^{48} + x^{47} + x^{46} + x^{44} + x^{42} + x^{41} + x^{39}, \\
p_3(x) &= x^{86} + x^{84} + x^{83} + x^{81} + x^{80} + x^{77} + x^{76} + x^{73} + x^{72} + x^{70} + x^{69} \\
&+ x^{67} + x^{46} + x^{44} + x^{43} + x^{41} + x^{40} + x^{37} + x^{36} + x^{33} + x^{32} + x^{30} \\
&+ x^{29} + x^{27}, \\
p_6(x) &= x^{88} + x^{84} + x^{82} + x^{78} + x^{68} + x^{64} + x^{62} + x^{58} + x^{48} + x^{44} + x^{42} \\
&+ x^{38} + x^{28} + x^{24} + x^{22} + x^{18}, \\
p_9(x) &= x^{90} + x^{88} + x^{87} + x^{86} + x^{85} + x^{84} + x^{83} + x^{82} + x^{81} + x^{80} + x^{79} \\
&+ x^{78} + x^{77} + x^{76} + x^{75} + x^{74} + x^{73} + x^{72} + x^{71} + x^{70} + x^{69} + x^{68} \\
&+ x^{67} + x^{66} + x^{65} + x^{64} + x^{63} + x^{62} + x^{61} + x^{60} + x^{59} + x^{58} + x^{57} \\
&+ x^{56} + x^{55} + x^{54} + x^{53} + x^{52} + x^{51} + x^{50} + x^{49} + x^{48} + x^{47} + x^{46} \\
&+ x^{45} + x^{44} + x^{43} + x^{42} + x^{41} + x^{40} + x^{39} + x^{38} + x^{37} + x^{36} + x^{35} \\
&+ x^{34} + x^{33} + x^{32} + x^{31} + x^{30} + x^{29} + x^{28} + x^{27} + x^{26} + x^{25} + x^{24} \\
&+ x^{23} + x^{22} + x^{21} + x^{20} + x^{19} + x^{18} + x^{17} + x^{16} + x^{15} + x^{14} + x^{13} \\
&+ x^{12} + x^{11} + x^9, \\
p_{12}(x) &= x^{92} + x^{84} + x^{76} + x^{72} + x^{68} + x^{64} + x^{60} + x^{56} + x^{52} + x^{48} + x^{44} \\
&+ x^{40} + x^{36} + x^{32} + x^{28} + x^{24} + x^{20} + x^{16} + x^8 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 1, 1, 1, 1, 1, 1, 0, 0, 0, 0, 1, 1, 0]$.

For $\phi(M^o, 1, 0)$:

$$\begin{aligned}
p_0(x) &= x^{102} + x^{99} + x^{97} + x^{96} + x^{94} + x^{87} + x^{85} + x^{84} + x^{82} + x^{81} + x^{80} \\
&+ x^{79} + x^{78} + x^{77} + x^{76} + x^{73} + x^{71} + x^{69} + x^{68} + x^{67} + x^{66} + x^{65} \\
&+ x^{62} + x^{61} + x^{58} + x^{56} + x^{55} + x^{48} + x^{45} + x^{43} + x^{42} + x^{41} + x^{39}, \\
p_3(x) &= x^{94} + x^{92} + x^{91} + x^{90} + x^{89} + x^{87} + x^{86} + x^{84} + x^{83} + x^{81} + x^{80} \\
&+ x^{78} + x^{77} + x^{75} + x^{74} + x^{72} + x^{71} + x^{70} + x^{69} + x^{67} + x^{54} + x^{52} \\
&+ x^{51} + x^{50} + x^{49} + x^{47} + x^{46} + x^{44} + x^{43} + x^{41} + x^{40} + x^{38} + x^{37} \\
&+ x^{35} + x^{34} + x^{32} + x^{31} + x^{30} + x^{29} + x^{27}, \\
p_6(x) &= x^{96} + x^{90} + x^{84} + x^{78} + x^{76} + x^{70} + x^{64} + x^{58} + x^{56} + x^{50} + x^{44} \\
&+ x^{38} + x^{36} + x^{30} + x^{24} + x^{18}, \\
p_9(x) &= x^{98} + x^{96} + x^{95} + x^{93} + x^{90} + x^{88} + x^{87} + x^{86} + x^{85} + x^{84} + x^{83} \\
&+ x^{82} + x^{81} + x^{80} + x^{79} + x^{78} + x^{77} + x^{76} + x^{75} + x^{74} + x^{73} + x^{72}
\end{aligned}$$

$$\begin{aligned}
& + x^{71} + x^{70} + x^{69} + x^{68} + x^{67} + x^{66} + x^{65} + x^{64} + x^{63} + x^{62} + x^{61} \\
& + x^{60} + x^{59} + x^{58} + x^{57} + x^{56} + x^{55} + x^{54} + x^{53} + x^{52} + x^{51} + x^{50} \\
& + x^{49} + x^{48} + x^{47} + x^{46} + x^{45} + x^{44} + x^{43} + x^{42} + x^{41} + x^{40} + x^{39} \\
& + x^{38} + x^{37} + x^{36} + x^{35} + x^{34} + x^{33} + x^{32} + x^{31} + x^{30} + x^{29} + x^{28} \\
& + x^{27} + x^{26} + x^{25} + x^{24} + x^{23} + x^{22} + x^{21} + x^{20} + x^{19} + x^{17} + x^{14} \\
& + x^{12} + x^{11} + x^9, \\
p_{12}(x) &= x^{100} + x^{96} + x^{88} + x^{76} + x^{72} + x^{68} + x^{64} + x^{60} + x^{56} + x^{52} + x^{48} \\
& + x^{44} + x^{40} + x^{36} + x^{32} + x^{28} + x^{24} + x^{12} + x^4 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 1, 0, 0, 1, 0, 0, 0, 0, 0, 1, 0]$.

For $\phi(M^o, 1, 1)$:

$$\begin{aligned}
p_0(x) &= x^{99} + x^{95} + x^{91} + x^{90} + x^{88} + x^{87} + x^{86} + x^{84} + x^{78} + x^{76} + x^{74} \\
& + x^{73} + x^{70} + x^{64} + x^{62} + x^{60} + x^{58} + x^{57} + x^{56} + x^{55} + x^{51} + x^{48} \\
& + x^{47} + x^{46} + x^{45} + x^{43} + x^{42}, \\
p_3(x) &= x^{94} + x^{92} + x^{91} + x^{89} + x^{85} + x^{84} + x^{83} + x^{81} + x^{77} + x^{76} + x^{75} \\
& + x^{73} + x^{72} + x^{69} + x^{67} + x^{66} + x^{54} + x^{52} + x^{51} + x^{49} + x^{45} + x^{44} \\
& + x^{43} + x^{41} + x^{37} + x^{36} + x^{35} + x^{33} + x^{32} + x^{29} + x^{27} + x^{26}, \\
p_6(x) &= x^{94} + x^{92} + x^{91} + x^{89} + x^{84} + x^{77} + x^{75} + x^{72} + x^{71} + x^{69} + x^{66} \\
& + x^{65} + x^{58} + x^{56} + x^{55} + x^{54} + x^{53} + x^{51} + x^{50} + x^{48} + x^{44} + x^{42} \\
& + x^{41} + x^{40} + x^{37} + x^{35} + x^{32} + x^{31} + x^{29} + x^{26} + x^{25} + x^{18} + x^{16} \\
& + x^{15} + x^{13} + x^{12} + x^{10} + x^9 + x^8 + x^2 + x + 1, \\
p_9(x) &= x^{94} + x^{93} + x^{92} + x^{14} + x^{13} + x^{12}, \\
p_{12}(x) &= x^{94} + x^{93} + x^{92} + x^{86} + x^{85} + x^{84} + x^{78} + x^{77} + x^{76} + x^{74} + x^{73} \\
& + x^{72} + x^{70} + x^{69} + x^{68} + x^{66} + x^{65} + x^{64} + x^{62} + x^{61} + x^{60} + x^{58} \\
& + x^{57} + x^{56} + x^{54} + x^{53} + x^{52} + x^{50} + x^{49} + x^{48} + x^{46} + x^{45} + x^{44} \\
& + x^{42} + x^{41} + x^{40} + x^{38} + x^{37} + x^{36} + x^{34} + x^{33} + x^{32} + x^{30} + x^{29} \\
& + x^{28} + x^{26} + x^{25} + x^{24} + x^{22} + x^{21} + x^{20} + x^{18} + x^{17} + x^{16} + x^{10} \\
& + x^9 + x^8 + x^2 + x + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 0, 0, 1, 1, 1, 0, 0, 0, 0, 1, 0]$.

For $\phi(W^o, 0, 0)$:

$$\begin{aligned}
p_0(x) &= x^{99} + x^{96} + x^{91} + x^{89} + x^{84} + x^{83} + x^{82} + x^{77} + x^{73} + x^{72} + x^{69} \\
& + x^{68} + x^{62} + x^{61} + x^{60} + x^{59} + x^{58} + x^{56} + x^{54} + x^{50} + x^{47} + x^{46} \\
& + x^{45} + x^{44} + x^{42} + x^{40} + x^{39} + x^{37} + x^{36}, \\
p_3(x) &= x^{94} + x^{92} + x^{91} + x^{90} + x^{88} + x^{85} + x^{79} + x^{78} + x^{77} + x^{72} + x^{71} \\
& + x^{70} + x^{69} + x^{67} + x^{65} + x^{64} + x^{54} + x^{52} + x^{51} + x^{50} + x^{48} + x^{45} \\
& + x^{39} + x^{38} + x^{37} + x^{32} + x^{31} + x^{30} + x^{29} + x^{27} + x^{25} + x^{24},
\end{aligned}$$

$$\begin{aligned}
p_6(x) &= x^{94} + x^{92} + x^{91} + x^{90} + x^{88} + x^{83} + x^{82} + x^{80} + x^{79} + x^{78} + x^{77} \\
&\quad + x^{75} + x^{74} + x^{73} + x^{71} + x^{70} + x^{68} + x^{66} + x^{65} + x^{64} + x^{63} + x^{62} \\
&\quad + x^{60} + x^{59} + x^{56} + x^{55} + x^{52} + x^{51} + x^{49} + x^{43} + x^{41} + x^{39} + x^{38} \\
&\quad + x^{37} + x^{35} + x^{34} + x^{33} + x^{31} + x^{30} + x^{28} + x^{26} + x^{25} + x^{24} + x^{23} \\
&\quad + x^{22} + x^{20} + x^{19} + x^{16} + x^{15} + x^{14} + x^{10} + x^9 + x^8 + x^2 + x + 1, \\
p_9(x) &= x^{94} + x^{93} + x^{92} + x^{14} + x^{13} + x^{12}, \\
p_{12}(x) &= x^{94} + x^{93} + x^{92} + x^{86} + x^{85} + x^{84} + x^{78} + x^{77} + x^{76} + x^{74} + x^{73} \\
&\quad + x^{72} + x^{70} + x^{69} + x^{68} + x^{66} + x^{65} + x^{64} + x^{62} + x^{61} + x^{60} + x^{58} \\
&\quad + x^{57} + x^{56} + x^{54} + x^{53} + x^{52} + x^{50} + x^{49} + x^{48} + x^{46} + x^{45} + x^{44} \\
&\quad + x^{42} + x^{41} + x^{40} + x^{38} + x^{37} + x^{36} + x^{34} + x^{33} + x^{32} + x^{30} + x^{29} \\
&\quad + x^{28} + x^{26} + x^{25} + x^{24} + x^{22} + x^{21} + x^{20} + x^{18} + x^{17} + x^{16} + x^{10} \\
&\quad + x^9 + x^8 + x^2 + x + 1,
\end{aligned}$$

and the initial terms are $[1, 0, 0, 0, 0, 0, 1, 1, 1, 1, 1, 0, 1, 0, 1, 0]$.

For $\phi(W^o, 0, 1)$:

$$\begin{aligned}
p_0(x) &= x^{102} + x^{99} + x^{97} + x^{96} + x^{94} + x^{87} + x^{85} + x^{84} + x^{82} + x^{81} + x^{80} \\
&\quad + x^{79} + x^{78} + x^{77} + x^{76} + x^{73} + x^{71} + x^{69} + x^{68} + x^{67} + x^{66} + x^{65} \\
&\quad + x^{62} + x^{61} + x^{58} + x^{56} + x^{55} + x^{48} + x^{45} + x^{43} + x^{42} + x^{41} + x^{39}, \\
p_3(x) &= x^{94} + x^{92} + x^{91} + x^{90} + x^{89} + x^{87} + x^{86} + x^{84} + x^{83} + x^{81} + x^{80} \\
&\quad + x^{78} + x^{77} + x^{75} + x^{74} + x^{72} + x^{71} + x^{70} + x^{69} + x^{67} + x^{54} + x^{52} \\
&\quad + x^{51} + x^{50} + x^{49} + x^{47} + x^{46} + x^{44} + x^{43} + x^{41} + x^{40} + x^{38} + x^{37} \\
&\quad + x^{35} + x^{34} + x^{32} + x^{31} + x^{30} + x^{29} + x^{27}, \\
p_6(x) &= x^{96} + x^{90} + x^{84} + x^{78} + x^{76} + x^{70} + x^{64} + x^{58} + x^{56} + x^{50} + x^{44} \\
&\quad + x^{38} + x^{36} + x^{30} + x^{24} + x^{18}, \\
p_9(x) &= x^{98} + x^{96} + x^{95} + x^{93} + x^{90} + x^{88} + x^{87} + x^{86} + x^{85} + x^{84} + x^{83} \\
&\quad + x^{82} + x^{81} + x^{80} + x^{79} + x^{78} + x^{77} + x^{76} + x^{75} + x^{74} + x^{73} + x^{72} \\
&\quad + x^{71} + x^{70} + x^{69} + x^{68} + x^{67} + x^{66} + x^{65} + x^{64} + x^{63} + x^{62} + x^{61} \\
&\quad + x^{60} + x^{59} + x^{58} + x^{57} + x^{56} + x^{55} + x^{54} + x^{53} + x^{52} + x^{51} + x^{50} \\
&\quad + x^{49} + x^{48} + x^{47} + x^{46} + x^{45} + x^{44} + x^{43} + x^{42} + x^{41} + x^{40} + x^{39} \\
&\quad + x^{38} + x^{37} + x^{36} + x^{35} + x^{34} + x^{33} + x^{32} + x^{31} + x^{30} + x^{29} + x^{28} \\
&\quad + x^{27} + x^{26} + x^{25} + x^{24} + x^{23} + x^{22} + x^{21} + x^{20} + x^{19} + x^{17} + x^{14} \\
&\quad + x^{12} + x^{11} + x^9, \\
p_{12}(x) &= x^{100} + x^{96} + x^{88} + x^{76} + x^{72} + x^{68} + x^{64} + x^{60} + x^{56} + x^{52} + x^{48} \\
&\quad + x^{44} + x^{40} + x^{36} + x^{32} + x^{28} + x^{24} + x^{12} + x^4 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 1, 0, 0, 1, 0, 0, 0, 0, 0, 0, 1, 0]$.

For $\phi(W^o, 1, 0)$:

$$\begin{aligned}
p_0(x) &= x^{100} + x^{98} + x^{95} + x^{93} + x^{89} + x^{88} + x^{86} + x^{85} + x^{84} + x^{83} + x^{78} \\
&\quad + x^{76} + x^{74} + x^{73} + x^{66} + x^{65} + x^{63} + x^{62} + x^{61} + x^{58} + x^{57} + x^{56} \\
&\quad + x^{55} + x^{50} + x^{48} + x^{47} + x^{46} + x^{44} + x^{42} + x^{41} + x^{39}, \\
p_3(x) &= x^{86} + x^{84} + x^{83} + x^{81} + x^{80} + x^{77} + x^{76} + x^{73} + x^{72} + x^{70} + x^{69} \\
&\quad + x^{67} + x^{46} + x^{44} + x^{43} + x^{41} + x^{40} + x^{37} + x^{36} + x^{33} + x^{32} + x^{30} \\
&\quad + x^{29} + x^{27}, \\
p_6(x) &= x^{88} + x^{84} + x^{82} + x^{78} + x^{68} + x^{64} + x^{62} + x^{58} + x^{48} + x^{44} + x^{42} \\
&\quad + x^{38} + x^{28} + x^{24} + x^{22} + x^{18}, \\
p_9(x) &= x^{90} + x^{88} + x^{87} + x^{86} + x^{85} + x^{84} + x^{83} + x^{82} + x^{81} + x^{80} + x^{79} \\
&\quad + x^{78} + x^{77} + x^{76} + x^{75} + x^{74} + x^{73} + x^{72} + x^{71} + x^{70} + x^{69} + x^{68} \\
&\quad + x^{67} + x^{66} + x^{65} + x^{64} + x^{63} + x^{62} + x^{61} + x^{60} + x^{59} + x^{58} + x^{57} \\
&\quad + x^{56} + x^{55} + x^{54} + x^{53} + x^{52} + x^{51} + x^{50} + x^{49} + x^{48} + x^{47} + x^{46} \\
&\quad + x^{45} + x^{44} + x^{43} + x^{42} + x^{41} + x^{40} + x^{39} + x^{38} + x^{37} + x^{36} + x^{35} \\
&\quad + x^{34} + x^{33} + x^{32} + x^{31} + x^{30} + x^{29} + x^{28} + x^{27} + x^{26} + x^{25} + x^{24} \\
&\quad + x^{23} + x^{22} + x^{21} + x^{20} + x^{19} + x^{18} + x^{17} + x^{16} + x^{15} + x^{14} + x^{13} \\
&\quad + x^{12} + x^{11} + x^9, \\
p_{12}(x) &= x^{92} + x^{84} + x^{76} + x^{72} + x^{68} + x^{64} + x^{60} + x^{56} + x^{52} + x^{48} + x^{44} \\
&\quad + x^{40} + x^{36} + x^{32} + x^{28} + x^{24} + x^{20} + x^{16} + x^8 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 1, 1, 1, 1, 1, 1, 0, 0, 0, 0, 1, 1, 0]$.

For $\phi(W^o, 1, 1)$:

$$\begin{aligned}
p_0(x) &= x^{99} + x^{95} + x^{91} + x^{90} + x^{88} + x^{87} + x^{86} + x^{84} + x^{78} + x^{76} + x^{74} \\
&\quad + x^{73} + x^{70} + x^{64} + x^{62} + x^{60} + x^{58} + x^{57} + x^{56} + x^{55} + x^{51} + x^{48} \\
&\quad + x^{47} + x^{46} + x^{45} + x^{43} + x^{42}, \\
p_3(x) &= x^{94} + x^{92} + x^{91} + x^{89} + x^{85} + x^{84} + x^{83} + x^{81} + x^{77} + x^{76} + x^{75} \\
&\quad + x^{73} + x^{72} + x^{69} + x^{67} + x^{66} + x^{54} + x^{52} + x^{51} + x^{49} + x^{45} + x^{44} \\
&\quad + x^{43} + x^{41} + x^{37} + x^{36} + x^{35} + x^{33} + x^{32} + x^{29} + x^{27} + x^{26}, \\
p_6(x) &= x^{94} + x^{92} + x^{91} + x^{89} + x^{84} + x^{77} + x^{75} + x^{72} + x^{71} + x^{69} + x^{66} \\
&\quad + x^{65} + x^{58} + x^{56} + x^{55} + x^{54} + x^{53} + x^{51} + x^{50} + x^{48} + x^{44} + x^{42} \\
&\quad + x^{41} + x^{40} + x^{37} + x^{35} + x^{32} + x^{31} + x^{29} + x^{26} + x^{25} + x^{18} + x^{16} \\
&\quad + x^{15} + x^{13} + x^{12} + x^{10} + x^9 + x^8 + x^2 + x + 1, \\
p_9(x) &= x^{94} + x^{93} + x^{92} + x^{14} + x^{13} + x^{12}, \\
p_{12}(x) &= x^{94} + x^{93} + x^{92} + x^{86} + x^{85} + x^{84} + x^{78} + x^{77} + x^{76} + x^{74} + x^{73} \\
&\quad + x^{72} + x^{70} + x^{69} + x^{68} + x^{66} + x^{65} + x^{64} + x^{62} + x^{61} + x^{60} + x^{58} \\
&\quad + x^{57} + x^{56} + x^{54} + x^{53} + x^{52} + x^{50} + x^{49} + x^{48} + x^{46} + x^{45} + x^{44}
\end{aligned}$$

$$\begin{aligned}
& + x^{42} + x^{41} + x^{40} + x^{38} + x^{37} + x^{36} + x^{34} + x^{33} + x^{32} + x^{30} + x^{29} \\
& + x^{28} + x^{26} + x^{25} + x^{24} + x^{22} + x^{21} + x^{20} + x^{18} + x^{17} + x^{16} + x^{10} \\
& + x^9 + x^8 + x^2 + x + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 0, 0, 1, 1, 1, 0, 0, 0, 0, 1, 0]$.

Below is the transition function and output function of an 2-automaton that generates $T = M_{0,0}^o$:

Transition function $(n, j) \mapsto \delta(n, j)$ ($\Lambda(n) := [\delta(n, 0), \delta(n, 1)]$):

n	$\Lambda(n)$	n	$\Lambda(n)$	n	$\Lambda(n)$	n	$\Lambda(n)$	n	$\Lambda(n)$
0	[1, 2]	77	[120, 121]	154	[118, 49]	231	[300, 301]	308	[30, 130]
1	[3, 4]	78	[107, 122]	155	[135, 210]	232	[175, 302]	309	[337, 199]
2	[5, 6]	79	[123, 11]	156	[211, 212]	233	[56, 26]	310	[338, 204]
3	[7, 8]	80	[12, 124]	157	[213, 214]	234	[139, 303]	311	[266, 339]
4	[9, 10]	81	[125, 126]	158	[215, 216]	235	[304, 305]	312	[340, 270]
5	[11, 12]	82	[127, 128]	159	[217, 113]	236	[158, 306]	313	[341, 342]
6	[13, 14]	83	[38, 105]	160	[218, 219]	237	[307, 286]	314	[343, 344]
7	[15, 16]	84	[96, 129]	161	[109, 220]	238	[150, 177]	315	[345, 276]
8	[17, 18]	85	[1, 40]	162	[220, 127]	239	[173, 154]	316	[180, 346]
9	[19, 20]	86	[130, 31]	163	[121, 128]	240	[170, 21]	317	[347, 348]
10	[21, 22]	87	[49, 131]	164	[35, 46]	241	[308, 93]	318	[292, 243]
11	[23, 24]	88	[95, 131]	165	[221, 222]	242	[309, 310]	319	[148, 349]
12	[25, 26]	89	[132, 52]	166	[223, 224]	243	[113, 234]	320	[350, 80]
13	[6, 27]	90	[133, 13]	167	[125, 225]	244	[217, 203]	321	[351, 352]
14	[28, 29]	91	[134, 35]	168	[226, 227]	245	[39, 311]	322	[299, 353]
15	[30, 31]	92	[135, 47]	169	[112, 119]	246	[223, 196]	323	[354, 355]
16	[32, 33]	93	[48, 136]	170	[11, 42]	247	[209, 136]	324	[356, 167]
17	[34, 35]	94	[87, 137]	171	[228, 229]	248	[131, 118]	325	[305, 357]
18	[36, 37]	95	[88, 54]	172	[230, 231]	249	[203, 114]	326	[358, 359]
19	[7, 38]	96	[138, 139]	173	[118, 95]	250	[312, 46]	327	[279, 291]
20	[1, 39]	97	[20, 59]	174	[101, 197]	251	[110, 127]	328	[360, 361]
21	[40, 41]	98	[140, 141]	175	[232, 205]	252	[238, 236]	329	[80, 5]
22	[42, 43]	99	[16, 142]	176	[233, 49]	253	[313, 216]	330	[362, 184]
23	[44, 45]	100	[143, 144]	177	[203, 234]	254	[35, 135]	331	[83, 363]
24	[46, 47]	101	[145, 146]	178	[235, 212]	255	[314, 211]	332	[303, 296]
25	[48, 2]	102	[147, 148]	179	[213, 236]	256	[315, 238]	333	[364, 365]
26	[49, 50]	103	[149, 150]	180	[237, 201]	257	[204, 316]	334	[301, 366]
27	[12, 43]	104	[151, 152]	181	[238, 214]	258	[123, 317]	335	[363, 367]
28	[51, 10]	105	[153, 154]	182	[127, 109]	259	[242, 122]	336	[368, 290]
29	[52, 53]	106	[155, 156]	183	[128, 220]	260	[318, 319]	337	[160, 18]
30	[54, 55]	107	[157, 158]	184	[51, 98]	261	[46, 210]	338	[369, 262]
31	[55, 56]	108	[159, 160]	185	[222, 195]	262	[313, 318]	339	[326, 195]
32	[57, 58]	109	[161, 162]	186	[131, 94]	263	[211, 36]	340	[336, 222]
33	[59, 60]	110	[163, 143]	187	[239, 50]	264	[94, 49]	341	[314, 235]
34	[61, 62]	111	[92, 164]	188	[207, 240]	265	[320, 229]	342	[338, 232]

35	[63, 64]	112	[165, 166]	189	[241, 52]	266	[321, 40]	343	[370, 110]
36	[65, 66]	113	[167, 145]	190	[120, 127]	267	[231, 322]	344	[371, 372]
37	[67, 68]	114	[22, 168]	191	[242, 108]	268	[228, 322]	345	[373, 235]
38	[69, 70]	115	[70, 169]	192	[220, 121]	269	[133, 194]	346	[112, 328]
39	[60, 71]	116	[85, 170]	193	[243, 244]	270	[227, 193]	347	[34, 312]
40	[72, 67]	117	[171, 172]	194	[245, 246]	271	[222, 223]	348	[119, 334]
41	[4, 73]	118	[142, 173]	195	[247, 248]	272	[104, 317]	349	[312, 135]
42	[74, 75]	119	[78, 163]	196	[249, 250]	273	[195, 224]	350	[10, 129]
43	[24, 76]	120	[162, 82]	197	[146, 251]	274	[323, 126]	351	[132, 226]
44	[77, 78]	121	[66, 174]	198	[252, 70]	275	[197, 101]	352	[230, 335]
45	[79, 80]	122	[76, 175]	199	[253, 254]	276	[200, 218]	353	[226, 53]
46	[81, 82]	123	[176, 25]	200	[255, 256]	277	[235, 36]	354	[337, 219]
47	[27, 83]	124	[177, 178]	201	[257, 155]	278	[324, 202]	355	[315, 374]
48	[84, 85]	125	[179, 180]	202	[258, 139]	279	[215, 318]	356	[375, 128]
49	[86, 87]	126	[169, 181]	203	[259, 144]	280	[232, 316]	357	[326, 223]
50	[26, 88]	127	[182, 183]	204	[260, 261]	281	[36, 325]	358	[333, 45]
51	[89, 90]	128	[82, 182]	205	[262, 263]	282	[221, 326]	359	[212, 37]
52	[91, 92]	129	[184, 185]	206	[186, 264]	283	[99, 106]	360	[323, 225]
53	[93, 86]	130	[130, 130]	207	[265, 266]	284	[216, 327]	361	[110, 121]
54	[50, 94]	131	[137, 186]	208	[68, 267]	285	[328, 329]	362	[320, 322]
55	[95, 50]	132	[187, 188]	209	[268, 269]	286	[237, 218]	363	[194, 14]
56	[94, 95]	133	[189, 28]	210	[270, 271]	287	[206, 118]	364	[100, 197]
57	[96, 97]	134	[190, 81]	211	[272, 273]	288	[330, 331]	365	[371, 376]
58	[9, 98]	135	[191, 192]	212	[274, 275]	289	[134, 312]	366	[52, 227]
59	[38, 99]	136	[73, 138]	213	[181, 24]	290	[32, 332]	367	[105, 115]
60	[39, 41]	137	[31, 130]	214	[276, 277]	291	[202, 113]	368	[233, 95]
61	[100, 101]	138	[10, 97]	215	[278, 279]	292	[318, 327]	369	[373, 211]
62	[102, 103]	139	[53, 193]	216	[280, 281]	293	[333, 112]	370	[251, 275]
63	[104, 11]	140	[194, 116]	217	[282, 283]	294	[328, 334]	371	[377, 253]
64	[105, 106]	141	[195, 196]	218	[284, 285]	295	[117, 332]	372	[306, 284]
65	[107, 108]	142	[31, 31]	219	[286, 159]	296	[335, 322]	373	[378, 65]
66	[109, 110]	143	[197, 197]	220	[183, 161]	297	[317, 42]	374	[379, 380]
67	[53, 111]	144	[101, 101]	221	[287, 288]	298	[241, 226]	375	[174, 163]
68	[8, 105]	145	[128, 110]	222	[289, 249]	299	[321, 39]	376	[244, 381]
69	[44, 112]	146	[121, 109]	223	[290, 55]	300	[40, 311]	377	[198, 219]
70	[113, 114]	147	[198, 199]	224	[261, 149]	301	[227, 111]	378	[375, 109]
71	[99, 115]	148	[200, 201]	225	[291, 292]	302	[119, 329]	379	[216, 319]
72	[13, 116]	149	[202, 203]	226	[293, 294]	303	[317, 12]	380	[212, 325]
73	[8, 99]	150	[204, 205]	227	[295, 264]	304	[336, 326]	381	[374, 214]
74	[117, 33]	151	[206, 94]	228	[296, 166]	305	[42, 124]		
75	[50, 118]	152	[207, 208]	229	[58, 297]	306	[45, 328]		
76	[45, 119]	153	[209, 2]	230	[298, 299]	307	[324, 217]		

Output function $n \mapsto \tau(n)$:

n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$
0	0	55	1	110	1	165	1	220	0	275	1	330	0
1	0	56	0	111	1	166	0	221	1	276	1	331	0
2	0	57	0	112	1	167	0	222	0	277	0	332	1
3	0	58	0	113	0	168	1	223	0	278	0	333	1
4	0	59	0	114	1	169	1	224	0	279	0	334	1
5	0	60	1	115	0	170	0	225	0	280	1	335	1
6	0	61	1	116	0	171	1	226	1	281	0	336	0
7	0	62	0	117	1	172	0	227	1	282	1	337	1
8	1	63	1	118	1	173	1	228	1	283	0	338	0
9	0	64	1	119	0	174	0	229	0	284	1	339	1
10	0	65	0	120	0	175	1	230	0	285	1	340	0
11	0	66	1	121	1	176	0	231	0	286	0	341	1
12	0	67	0	122	0	177	1	232	1	287	1	342	0
13	0	68	1	123	0	178	0	233	0	288	0	343	1
14	1	69	0	124	1	179	0	234	0	289	0	344	0
15	0	70	0	125	0	180	0	235	0	290	0	345	0
16	0	71	0	126	1	181	0	236	0	291	0	346	1
17	1	72	0	127	0	182	0	237	0	292	0	347	1
18	0	73	1	128	0	183	0	238	0	293	1	348	0
19	0	74	1	129	1	184	1	239	1	294	1	349	0
20	0	75	0	130	0	185	0	240	0	295	1	350	0
21	0	76	0	131	1	186	1	241	0	296	1	351	1
22	1	77	0	132	1	187	1	242	1	297	1	352	0
23	0	78	0	133	0	188	0	243	0	298	0	353	1
24	0	79	0	134	0	189	0	244	1	299	1	354	1
25	0	80	0	135	1	190	0	245	1	300	0	355	0
26	0	81	0	136	1	191	1	246	0	301	1	356	0
27	0	82	0	137	1	192	0	247	1	302	0	357	1
28	1	83	0	138	0	193	0	248	1	303	1	358	1
29	0	84	0	139	0	194	1	249	1	304	0	359	1
30	0	85	0	140	1	195	1	250	0	305	1	360	1
31	1	86	0	141	1	196	1	251	1	306	0	361	1
32	0	87	0	142	1	197	1	252	0	307	0	362	0
33	0	88	1	143	1	198	0	253	1	308	0	363	1
34	1	89	1	144	0	199	1	254	1	309	1	364	1
35	1	90	0	145	0	200	1	255	1	310	0	365	0
36	0	91	0	146	1	201	0	256	0	311	1	366	0
37	0	92	1	147	0	202	0	257	0	312	0	367	1
38	0	93	0	148	1	203	1	258	0	313	1	368	0
39	1	94	0	149	0	204	0	259	1	314	1	369	0
40	0	95	1	150	0	205	1	260	0	315	0	370	1
41	0	96	0	151	1	206	1	261	0	316	0	371	0
42	1	97	0	152	0	207	0	262	1	317	1	372	0

43	0	98	1	153	1	208	1	263	1	318	0	373	0
44	0	99	0	154	1	209	1	264	0	319	1	374	1
45	0	100	1	155	1	210	1	265	0	320	0	375	0
46	0	101	0	156	1	211	1	266	1	321	1	376	1
47	0	102	0	157	0	212	1	267	0	322	1	377	0
48	0	103	0	158	0	213	0	268	1	323	1	378	0
49	0	104	1	159	1	214	1	269	0	324	0	379	1
50	0	105	1	160	1	215	0	270	1	325	1	380	1
51	1	106	1	161	1	216	1	271	0	326	1	381	1
52	0	107	0	162	0	217	1	272	1	327	0		
53	0	108	1	163	1	218	1	273	1	328	1		
54	0	109	1	164	1	219	0	274	1	329	0		

UNIVERSITÉ DE STRASBOURG, CNRS, IRMA UMR 7501, F-67000 STRASBOURG, FRANCE
Email address: `guoniu.han@unistra.fr`

SCHOOL OF MATHEMATICS AND STATISTICS, HUAZHONG UNIVERSITY OF SCIENCE AND TECHNOLOGY, WUHAN, PR CHINA
Email address: `huyining@hust.edu.cn`