

PROOF OF THE CONTINUED FRACTION CONJECTURE FOR

$$(a, b) = (z^4 + z, z^3 + z)$$

GUO-NIU HAN* AND YINING HU*

ABSTRACT. The continued fraction conjecture claims that the continued fraction defined by the (a, b) -Thue-Morse sequence is algebraic over $\mathbb{F}_2(x)$ for all pairs of distinct elements (a, b) in $\mathbb{F}_2[x] \setminus \mathbb{F}_2$. In this paper we prove the conjecture for $(a, b) = (z^4 + z, z^3 + z)$.

1. INTRODUCTION

Let $\mathbb{F}_2$ be the finite field of cardinality 2. Given a sequence of polynomials $\mathbf{a} = (a_0(z), a_1(z), a_2(z), \dots)$ of elements from $\mathbb{F}_2[z] \setminus \mathbb{F}_2$, we define the infinite continued fraction

$$(1.1) \quad \text{CF}(\mathbf{a}(z)) = \frac{1}{a_0(z) + \frac{1}{a_1(z) + \frac{1}{a_2(z) + \frac{1}{\ddots}}}}$$

as the limit of the finite continued fractions

$$(1.2) \quad \text{CF}_n(\mathbf{a}(z)) = \frac{1}{a_0(z) + \frac{1}{a_1(z) + \frac{1}{\ddots + \frac{1}{a_n(z)}}}} \in \mathbb{F}_2((1/z)).$$

Conversely, each power series in $\mathbb{F}_2[[1/z]]$ without constant term

$$(1.3) \quad c_{-1}z^{-1} + c_{-2}z^{-2} + c_{-3}z^{-3} + \dots$$

admits a continued fraction expansion of form (1.1).

Let a and b be two distinct elements from $\mathbb{F}_2[z] \setminus \mathbb{F}_2$. We consider the continued fraction (1.1) associated with the (a, b) -Thue-Morse sequence

$$\mathbf{t} = (t_0(z), t_1(z), t_2(z), \dots) = (a, b, b, a, b, a, a, b, \dots)$$

that we shall denote

$$(1.4) \quad \text{CF}(a, b) := \text{CF}(\mathbf{t}).$$

Date: 9/04/2020 20:47:34 .

2010 Mathematics Subject Classification. 11B85, 11J70, 11B50, 11Y65, 05A15.

Key words and phrases. automatic sequence, continued fraction, Thue-Morse sequence.

* This paper was automatically generated from a computer program developed by the authors, with 10.3 minutes CPU time used.

Theorem 1.1. *When $(a, b) = (z^4 + z, z^3 + z)$, the two power series $\text{CF}(a, b)$ and $\text{CF}(b, a)$ are algebraic over $\mathbb{F}_2(z)$, with minimal polynomials of the form*

$$p_4(z)y^4 + p_3(z)y^3 + p_2(z)y^2 + p_1(z)y + p_0(z) = 0$$

where, for $\text{CF}(a, b)$,

$$\begin{aligned} p_0(z) &= z^{11} + z^9 + z^8 + z^6 + z^4 + z^3 + z^2 + 1, \\ p_1(z) &= z^{15} + z^{14} + z^{13} + z^{11} + z^{10} + z^9 + z^7 + z^6 + z^4 + z^3, \\ p_2(z) &= z^{18} + z^{14} + z^{12} + z^{10} + z^4 + z^2, \\ p_3(z) &= z^{15} + z^{14} + z^{13} + z^{11} + z^{10} + z^9 + z^7 + z^6 + z^4 + z^3, \\ p_4(z) &= z^{12} + z^{11} + z^{10} + z^9 + z^8 + z^6 + z^3 + z^2 + 1, \end{aligned}$$

and for $\text{CF}(b, a)$,

$$\begin{aligned} p_0(z) &= z^{11} + z^9 + z^8 + z^4 + z^3 + z^2 + 1, \\ p_1(z) &= z^{15} + z^{14} + z^{13} + z^{11} + z^{10} + z^9 + z^7 + z^6 + z^4 + z^3, \\ p_2(z) &= z^{18} + z^{14} + z^{12} + z^{10} + z^4 + z^2, \\ p_3(z) &= z^{15} + z^{14} + z^{13} + z^{11} + z^{10} + z^9 + z^7 + z^6 + z^4 + z^3, \\ p_4(z) &= z^{12} + z^{11} + z^{10} + z^9 + z^8 + z^3 + z^2 + 1. \end{aligned}$$

2. PROOF

Define the sequences of polynomials $(P_n(z))_{n \geq 0}$ and $(Q_n(z))_{n \geq 0}$ by

$$\begin{pmatrix} P_n(z) & Q_n(z) \\ P_{n-1}(z) & Q_{n-1}(z) \end{pmatrix} := \begin{pmatrix} t_n(z) & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} t_{n-1}(z) & 1 \\ 1 & 0 \end{pmatrix} \cdots \begin{pmatrix} t_0(z) & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}.$$

By the basic properties of continued fractions, we have

$$(2.1) \quad \text{CF}_n(a, b) = \frac{P_n(z)}{Q_n(z)} \in \mathbb{F}_2((1/z)).$$

As formula (2.1) is not efficient for calculating $\text{CF}_n(a, b)$, a fast method is to be derived and described as follows. Define

$$\begin{aligned} M_n(x) &= x^{\deg(t_{2^n-1})} \begin{pmatrix} t_{2^n-1}(1/x) & 1 \\ 1 & 0 \end{pmatrix} \times x^{\deg(t_{2^n-2})} \begin{pmatrix} t_{2^n-2}(1/x) & 1 \\ 1 & 0 \end{pmatrix} \times \\ &\quad \cdots \times x^{\deg(t_0)} \begin{pmatrix} t_0(1/x) & 1 \\ 1 & 0 \end{pmatrix}, \\ W_n(x) &= x^{\deg(\bar{t}_{2^n-1})} \begin{pmatrix} \bar{t}_{2^n-1}(1/x) & 1 \\ 1 & 0 \end{pmatrix} \times x^{\deg(\bar{t}_{2^n-2})} \begin{pmatrix} \bar{t}_{2^n-2}(1/x) & 1 \\ 1 & 0 \end{pmatrix} \times \\ &\quad \cdots \times x^{\deg(\bar{t}_0)} \begin{pmatrix} \bar{t}_0(1/x) & 1 \\ 1 & 0 \end{pmatrix}, \end{aligned}$$

where $\bar{t}$ is the (b, a) -Thue-Morse sequence. By the properties of the Thue-Morse sequence, we have for $n \geq 0$

$$\begin{aligned} M_{n+1}(x) &= W_n(x) \cdot M_n(x), \\ W_{n+1}(x) &= M_n(x) \cdot W_n(x). \end{aligned}$$

Let $x := 1/z$. For each non-zero polynomial $P(z)$, define $\tilde{P}(x)$ to be $P(1/x)$. Then,

$$\text{CF}_n(a, b) = \frac{P_n(z)}{Q_n(z)} = \frac{\tilde{P}_n(x)}{\tilde{Q}_n(x)} \in \mathbb{F}_2((x)) = \mathbb{F}_2((1/z)).$$

Comparing the definition of $M_n(x)$ with the definition of $P_n(x)$ and $Q_n(x)$, we see that

$$\begin{aligned} M_n(x)_{0,1} &= x^{d_n} \tilde{P}_{2^n-1}(x), \\ M_n(x)_{0,0} &= x^{d_n} \tilde{Q}_{2^n-1}(x), \end{aligned}$$

for some positive integer d_n . Hence,

$$(2.2) \quad \text{CF}_{2^{2^n}-1}(a, b) = \frac{\tilde{P}_{2^{2^n}-1}(x)}{\tilde{Q}_{2^{2^n}-1}(x)} = \frac{M_{2^n}(x)_{0,1}}{M_{2^n}(x)_{0,0}}.$$

Therefore, by the convergence theorem of continued fraction, the algebraicity of $\text{CF}(a, b)$ will be established if it is shown that both $M_{2^n}(x)_{0,1}$ and $M_{2^n}(x)_{0,0}$ converge to algebraic series in $\mathbb{F}_2[[x]]$.

Actually, we will prove that for all $0 \leq i, j \leq 1$ the four sequences $(M_{2^n}(x)_{i,j})_n$, $(M_{2^{n+1}}(x)_{i,j})_n$, $(W_{2^n}(x)_{i,j})_n$, and $(W_{2^{n+1}}(x)_{i,j})_n$ converge to algebraic series in $\mathbb{F}_2[[x]]$. For this purpose, we define four 2×2 matrices M^e, M^o, W^e, W^o as follows: For each $T \in \{M^e, M^o, W^e, W^o\}$ and $0 \leq i, j \leq 1$, $T_{i,j}$ is defined to be the unique solution in $\mathbb{F}_2[[x]]$ of the polynomial $\phi(T, i, j)$ under certain initial conditions; the polynomials $\phi(T, i, j)$ and initial conditions are given in Section 3. We will prove that these four matrices, whose components are algebraic by definition, are the limits of $(M_{2^n}(x))_n$, $(M_{2^{n+1}}(x))_n$, $(W_{2^n}(x))_n$, and $(W_{2^{n+1}}(x))_n$.

Let us explain how the polynomials $\phi(T, i, j)$ and initial conditions are found, and why the solutions exist and are unique. For $0 \leq i, j \leq 1$, the coefficients of the polynomial $\phi(M^e, i, j)$ (resp. $\phi(M^o, i, j)$, $\phi(W^e, i, j)$, and $\phi(W^o, i, j)$) are the Padé-Hermite approximants of type

$$(175, 175, 175, 175, 175)$$

of the vector

$$(1, T^3, T^6, T^9, T^{12}),$$

where $T = M_{12,i,j}$ (resp. $M_{11,i,j}$, $W_{12,i,j}$, and $W_{11,i,j}$), found by the Derksen algorithm. We take the first 8 terms of $M_{12,i,j}$ (resp. $M_{11,i,j}$, $W_{12,i,j}$, and $W_{11,i,j}$) as initial conditions for $\phi(M^e, i, j)$ (resp. $\phi(M^o, i, j)$, $\phi(W^e, i, j)$, and $\phi(W^o, i, j)$). The following fact will be used to ensure that the solution exists and is unique: let $P(x, y) \in \mathbb{F}_2[x, y]$ and for each series $f(x) = \sum_{n=0}^{\infty} a_n x^n \in \mathbb{F}_2[[x]]$ denote the partial sum $\sum_{j=0}^{n-1} a_j x^j$ by $f_n(x)$ for $n \geq 0$. If for some $n \geq 0$ and $a_0, a_1, \dots, a_{n-1} \in \mathbb{F}_2$ $P(x, \sum_{j=0}^{n-1} a_j x^j) = O(x^n)$ and $Q(x, y) := P(x, \sum_{j=0}^{n-1} a_j x^j + x^n y)$ can be written as $x^m \sum_{j=0}^{\infty} q_j(x) y^j$ where $q_j(x)$ are polynomials for $j \geq 0$, $q_1(0) = 1$, and $q_j(0) = 0$ for $j > 1$, then there exists a unique solution $f(x) \in \mathbb{F}_2[[x]]$ of $P(x, f(x)) = 0$ that satisfies the initial condition $f_n(x) = \sum_{j=0}^{n-1} a_j x^j$.

We state two lemmas concerning the four matrices M^e, M^o, W^e, W^o . The first one is about relations between them; the second, about the structure of the each matrix.

Lemma 2.1. *We have*

$$(2.3) \quad M^e = W^o \cdot M^o,$$

$$(2.4) \quad M^o = W^e \cdot M^e,$$

$$(2.5) \quad W^e = M^o \cdot W^o,$$

$$(2.6) \quad W^o = M^e \cdot W^e.$$

Proof. We give the proof of the identity

$$M_{0,0}^e = W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o,$$

the proofs of the others being similar.

First, we compute the minimal polynomials of $W_{0,0}^o M_{0,0}^o$ and $W_{0,1}^o M_{1,0}^o$. We know that

$$P(x, y) = \text{Res}_z (\phi(W^o, 0, 0)(x, z), z^{12} \cdot \phi(M^o, 0, 0)(x, y/z))$$

is an annihilating polynomial of $W_{0,0}^o M_{0,0}^o$. We use Padé-Hermite approximation to find a candidate for the minimal polynomial of $W_{0,0}^o M_{0,0}^o$, that will be called $\phi_0(x, y)$. To prove that $\phi_0(x, y)$ is indeed the minimal polynomial, it suffices to prove that it is an irreducible factor of $P(x, y)$ of multiplicity m and that $Q(x, y) := P(x, y)/\phi_0(x, y)^m$ is not an annihilating polynomial of $W_{0,0}^o M_{0,0}^o$. We verify the first point directly. For the second point, we truncate $W_{0,0}^o M_{0,0}^o$ to order 630 and substitute it for y in $Q(x, y)$. We get a series of valuation less than 630, which proves that $Q(x, y)$ is not an annihilating polynomial of $W_{0,0}^o M_{0,0}^o$. We find the minimal polynomial $\phi_1(x, y)$ of $W_{0,1}^o M_{1,0}^o$ in a similar way.

Now we prove that $\phi(M^e, 0, 0)$ is the minimal polynomial of $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$. We know that

$$S(x, y) = \text{Res}_z (\phi_0(x, z), \phi_1(x, y + z))$$

is an annihilating of $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$. We verify that $\phi(M^e, 0, 0)$ is an irreducible factor of $S(x, y)$ of multiplicity μ , and that $Q(x, y) := S(x, y)/\phi(M^e, 0, 0)^\mu$ is not an annihilating polynomial of $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$. To see the last point, we truncate $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$ to order 770 and substitute it for y in $Q(x, y)$. We get a series of valuation less than 770, and therefore $Q(x, y)$ is not an annihilating polynomial of $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$.

Finally, the first 16 terms of $M_{0,0}^e$ and $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$ coincide. As these first terms determine a unique solution of $\phi(M^e, 0, 0)$, we know that the two series are one and the same. $\square$

Define

$$R^e = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \quad \text{and} \quad R^o = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}.$$

Lemma 2.2. *For all integers $k \geq 2$ and $u = 2^{2k-1}$, the following identities hold.*

$$\begin{aligned} M^e[:14u] &= M^e[:7u] + x^{2u} M^e[:5u] + x^{3u} M^e[:4u] + x^{5u} M^e[:2u] + x^{6u} M^e[:u] \\ &\quad + x^{2u} M^e[:7u] + x^{4u} M^e[:5u] + x^{5u} M^e[:4u] + x^{7u} M^e[:2u] \\ &\quad + x^{8u} M^e[:u] + x^{3u} M^e[:7u] + x^{5u} M^e[:5u] + x^{6u} M^e[:4u] \\ &\quad + x^{8u} M^e[:2u] + x^{9u} M^e[:u] + x^{4u} M^e[:7u] + x^{6u} M^e[:5u] \\ &\quad + x^{7u} M^e[:4u] + x^{9u} M^e[:2u] + x^{10u} M^e[:u] + x^{5u} M^e[:7u] \\ &\quad + x^{7u} M^e[:5u] + x^{8u} M^e[:4u] + x^{10u} M^e[:2u] + x^{11u} M^e[:u] \end{aligned}$$

$$\begin{aligned}
& + x^{6u}M^e[:7u] + x^{8u}M^e[:5u] + x^{9u}M^e[:4u] + x^{11u}M^e[:2u] \\
& + x^{12u}M^e[:u] + x^{7u}M^e[:7u] + x^{11u}M^e[:3u] + x^{13u}M^e[:u] \\
& + (x^{7u} + x^{9u} + x^{10u} + x^{11u} + x^{12u} + x^{13u})R^e, \\
W^e[:14u] = & W^e[:7u] + x^{2u}W^e[:5u] + x^{3u}W^e[:4u] + x^{5u}W^e[:2u] + x^{6u}W^e[:u] \\
& + x^{2u}W^e[:7u] + x^{4u}W^e[:5u] + x^{5u}W^e[:4u] + x^{7u}W^e[:2u] \\
& + x^{8u}W^e[:u] + x^{3u}W^e[:7u] + x^{5u}W^e[:5u] + x^{6u}W^e[:4u] \\
& + x^{8u}W^e[:2u] + x^{9u}W^e[:u] + x^{4u}W^e[:7u] + x^{6u}W^e[:5u] \\
& + x^{7u}W^e[:4u] + x^{9u}W^e[:2u] + x^{10u}W^e[:u] + x^{5u}W^e[:7u] \\
& + x^{7u}W^e[:5u] + x^{8u}W^e[:4u] + x^{10u}W^e[:2u] + x^{11u}W^e[:u] \\
& + x^{6u}W^e[:7u] + x^{8u}W^e[:5u] + x^{9u}W^e[:4u] + x^{11u}W^e[:2u] \\
& + x^{12u}W^e[:u] + x^{7u}W^e[:7u] + x^{11u}W^e[:3u] + x^{13u}W^e[:u] \\
& + (x^{7u} + x^{9u} + x^{10u} + x^{11u} + x^{12u} + x^{13u})R^e.
\end{aligned}$$

For all integers $k \geq 2$ and $u = 2^{2k}$, the following identities hold.

$$\begin{aligned}
M^o[:14u] = & M^o[:7u] + x^{2u}M^o[:5u] + x^{3u}M^o[:4u] + x^{5u}M^o[:2u] + x^{6u}M^o[:u] \\
& + x^{2u}M^o[:7u] + x^{4u}M^o[:5u] + x^{5u}M^o[:4u] + x^{7u}M^o[:2u] \\
& + x^{8u}M^o[:u] + x^{3u}M^o[:7u] + x^{5u}M^o[:5u] + x^{6u}M^o[:4u] \\
& + x^{8u}M^o[:2u] + x^{9u}M^o[:u] + x^{4u}M^o[:7u] + x^{6u}M^o[:5u] \\
& + x^{7u}M^o[:4u] + x^{9u}M^o[:2u] + x^{10u}M^o[:u] + x^{5u}M^o[:7u] \\
& + x^{7u}M^o[:5u] + x^{8u}M^o[:4u] + x^{10u}M^o[:2u] + x^{11u}M^o[:u] \\
& + x^{6u}M^o[:7u] + x^{8u}M^o[:5u] + x^{9u}M^o[:4u] + x^{11u}M^o[:2u] \\
& + x^{12u}M^o[:u] + x^{7u}M^o[:7u] + x^{11u}M^o[:3u] + x^{13u}M^o[:u] \\
& + (x^{7u} + x^{9u} + x^{10u} + x^{11u} + x^{12u} + x^{13u})R^o, \\
W^o[:14u] = & W^o[:7u] + x^{2u}W^o[:5u] + x^{3u}W^o[:4u] + x^{5u}W^o[:2u] + x^{6u}W^o[:u] \\
& + x^{2u}W^o[:7u] + x^{4u}W^o[:5u] + x^{5u}W^o[:4u] + x^{7u}W^o[:2u] \\
& + x^{8u}W^o[:u] + x^{3u}W^o[:7u] + x^{5u}W^o[:5u] + x^{6u}W^o[:4u] \\
& + x^{8u}W^o[:2u] + x^{9u}W^o[:u] + x^{4u}W^o[:7u] + x^{6u}W^o[:5u] \\
& + x^{7u}W^o[:4u] + x^{9u}W^o[:2u] + x^{10u}W^o[:u] + x^{5u}W^o[:7u] \\
& + x^{7u}W^o[:5u] + x^{8u}W^o[:4u] + x^{10u}W^o[:2u] + x^{11u}W^o[:u] \\
& + x^{6u}W^o[:7u] + x^{8u}W^o[:5u] + x^{9u}W^o[:4u] + x^{11u}W^o[:2u] \\
& + x^{12u}W^o[:u] + x^{7u}W^o[:7u] + x^{11u}W^o[:3u] + x^{13u}W^o[:u] \\
& + (x^{7u} + x^{9u} + x^{10u} + x^{11u} + x^{12u} + x^{13u})R^o.
\end{aligned}$$

Proof. To prove Lemma 2.2 we first construct an automaton for each sequence concerned, and then transform the conditions on infinitely many k 's into finitely many conditions on the states of the automaton. In the following, we will prove that for $T = M_{0,0}^o$, for all integer $k \geq 2$ and $u = 2^{2k}$,

$$T[:14u] = T[:7u] + x^{2u}T[:5u] + x^{3u}T[:4u] + x^{5u}T[:2u] + x^{6u}T[:u] + x^{2u}T[:7u]$$

$$\begin{aligned}
& + x^{4u}T[:5u] + x^{5u}T[:4u] + x^{7u}T[:2u] + x^{8u}T[:u] + x^{3u}T[:7u] \\
& + x^{5u}T[:5u] + x^{6u}T[:4u] + x^{8u}T[:2u] + x^{9u}T[:u] + x^{4u}T[:7u] \\
& + x^{6u}T[:5u] + x^{7u}T[:4u] + x^{9u}T[:2u] + x^{10u}T[:u] + x^{5u}T[:7u] \\
& + x^{7u}T[:5u] + x^{8u}T[:4u] + x^{10u}T[:2u] + x^{11u}T[:u] + x^{6u}T[:7u] \\
& + x^{8u}T[:5u] + x^{9u}T[:4u] + x^{11u}T[:2u] + x^{12u}T[:u] + x^{7u}T[:7u] \\
& + x^{11u}T[:3u] + x^{13u}T[:u].
\end{aligned}$$

The proof of the other 15 cases are similar. We break down the above identity into 7 parts:

$$\begin{aligned}
0 &= x^{6u}T[u:2u] + x^{5u}T[2u:3u] + x^{3u}T[4u:5u] + x^{2u}T[5u:6u] \\
&\quad + T[7u:8u], \\
0 &= x^{6u}T[2u:3u] + x^{5u}T[3u:4u] + x^{3u}T[5u:6u] + x^{2u}T[6u:7u] \\
&\quad + T[8u:9u], \\
0 &= x^{9u}T[:u] + x^{8u}T[u:2u] + x^{7u}T[2u:3u] + x^{6u}T[3u:4u] \\
&\quad + x^{4u}T[5u:6u] + x^{3u}T[6u:7u] + T[9u:10u], \\
0 &= x^{7u}T[3u:4u] + x^{5u}T[5u:6u] + x^{4u}T[6u:7u] + T[10u:11u], \\
0 &= x^{11u}T[:u] + x^{10u}T[u:2u] + x^{9u}T[2u:3u] + x^{6u}T[5u:6u] \\
&\quad + x^{5u}T[6u:7u] + T[11u:12u], \\
0 &= x^{12u}T[:u] + x^{9u}T[3u:4u] + x^{8u}T[4u:5u] + x^{7u}T[5u:6u] \\
&\quad + x^{6u}T[6u:7u] + T[12u:13u], \\
0 &= x^{13u}T[:u] + x^{11u}T[2u:3u] + x^{7u}T[6u:7u] + T[13u:14u],
\end{aligned}$$

which can be rewritten as

$$(2.7) \quad 0 = T[[1w]_2] + T[[10w]_2] + T[[100w]_2] + T[[101w]_2] + T[[111w]_2],$$

$$(2.8) \quad 0 = T[[10w]_2] + T[[11w]_2] + T[[101w]_2] + T[[110w]_2] + T[[1000w]_2],$$

$$0 = T[[w]_2] + T[[1w]_2] + T[[10w]_2] + T[[11w]_2] + T[[101w]_2]$$

$$(2.9) \quad + T[[110w]_2] + T[[1001w]_2],$$

$$(2.10) \quad 0 = T[[11w]_2] + T[[101w]_2] + T[[110w]_2] + T[[1010w]_2],$$

$$(2.11) \quad 0 = T[[w]_2] + T[[1w]_2] + T[[10w]_2] + T[[101w]_2] + T[[110w]_2]$$

$$+ T[[1011w]_2],$$

$$0 = T[[w]_2] + T[[11w]_2] + T[[100w]_2] + T[[101w]_2] + T[[110w]_2]$$

$$(2.12) \quad + T[[1100w]_2],$$

$$(2.13) \quad 0 = T[[w]_2] + T[[10w]_2] + T[[110w]_2] + T[[1101w]_2],$$

for all binary word w of length $2k$ and $w \neq 0^{2k}$, and

$$(2.14) \quad 0 = T[[1w]_2] + T[[10w]_2] + T[[100w]_2] + T[[101w]_2] + T[[111w]_2],$$

$$(2.15) \quad 0 = T[[10w]_2] + T[[11w]_2] + T[[101w]_2] + T[[110w]_2] + T[[1000w]_2],$$

$$(2.16) \quad 0 = T[[w]_2] + T[[1w]_2] + T[[10w]_2] + T[[11w]_2] + T[[101w]_2]$$

$$+ T[[110w]_2] + T[[1001w]_2],$$

$$\begin{aligned}
(2.17) \quad 0 &= T[[11w]_2] + T[[101w]_2] + T[[110w]_2] + T[[1010w]_2], \\
&0 = T[[w]_2] + T[[1w]_2] + T[[10w]_2] + T[[101w]_2] + T[[110w]_2] \\
(2.18) \quad &+ T[[1011w]_2], \\
&0 = T[[w]_2] + T[[11w]_2] + T[[100w]_2] + T[[101w]_2] + T[[110w]_2] \\
(2.19) \quad &+ T[[1100w]_2], \\
(2.20) \quad 0 &= T[[w]_2] + T[[10w]_2] + T[[110w]_2] + T[[1101w]_2],
\end{aligned}$$

for $w = 0^{2k}$.

First we calculate an 2-automaton that generates T from its minimal polynomial and its first terms. This automaton has 531 states; its transition function and output function can be found in the annex. Let $A(s, w)$ denote the state reached after reading w from right to left starting from the state s , and τ the output function. Define

$$E_{2k} = \{A(i, w) : |w| = 2k, w \neq 0^{2k}\}.$$

Identities (2.7) through (2.20) can be written as

$$\begin{aligned}
(2.21) \quad 0 &= \tau(A(s, 1)) + \tau(A(s, 10)) + \tau(A(s, 100)) + \tau(A(s, 101)) \\
&+ \tau(A(s, 111)), \\
(2.22) \quad 0 &= \tau(A(s, 10)) + \tau(A(s, 11)) + \tau(A(s, 101)) + \tau(A(s, 110)) \\
&+ \tau(A(s, 1000)), \\
(2.23) \quad 0 &= \tau(A(s, \epsilon)) + \tau(A(s, 1)) + \tau(A(s, 10)) + \tau(A(s, 11)) + \tau(A(s, 101)) \\
&+ \tau(A(s, 110)) + \tau(A(s, 1001)), \\
(2.24) \quad 0 &= \tau(A(s, 11)) + \tau(A(s, 101)) + \tau(A(s, 110)) + \tau(A(s, 1010)), \\
&0 = \tau(A(s, \epsilon)) + \tau(A(s, 1)) + \tau(A(s, 10)) + \tau(A(s, 101)) \\
(2.25) \quad &+ \tau(A(s, 110)) + \tau(A(s, 1011)), \\
&0 = \tau(A(s, \epsilon)) + \tau(A(s, 11)) + \tau(A(s, 100)) + \tau(A(s, 101)) \\
(2.26) \quad &+ \tau(A(s, 110)) + \tau(A(s, 1100)), \\
(2.27) \quad 0 &= \tau(A(s, \epsilon)) + \tau(A(s, 10)) + \tau(A(s, 110)) + \tau(A(s, 1101)),
\end{aligned}$$

for all $s \in E_{2k}$, and

$$\begin{aligned}
(2.28) \quad 0 &= \tau(A(s, 1)) + \tau(A(s, 10)) + \tau(A(s, 100)) + \tau(A(s, 101)) \\
&+ \tau(A(s, 111)), \\
(2.29) \quad 0 &= \tau(A(s, 10)) + \tau(A(s, 11)) + \tau(A(s, 101)) + \tau(A(s, 110)) \\
&+ \tau(A(s, 1000)), \\
(2.30) \quad 0 &= \tau(A(s, \epsilon)) + \tau(A(s, 1)) + \tau(A(s, 10)) + \tau(A(s, 11)) + \tau(A(s, 101)) \\
&+ \tau(A(s, 110)) + \tau(A(s, 1001)), \\
(2.31) \quad 0 &= \tau(A(s, 11)) + \tau(A(s, 101)) + \tau(A(s, 110)) + \tau(A(s, 1010)), \\
&0 = \tau(A(s, \epsilon)) + \tau(A(s, 1)) + \tau(A(s, 10)) + \tau(A(s, 101)) \\
(2.32) \quad &+ \tau(A(s, 110)) + \tau(A(s, 1011)), \\
&0 = \tau(A(s, \epsilon)) + \tau(A(s, 11)) + \tau(A(s, 100)) + \tau(A(s, 101)) \\
(2.33) \quad &+ \tau(A(s, 110)) + \tau(A(s, 1100)), \\
(2.34) \quad 0 &= \tau(A(s, \epsilon)) + \tau(A(s, 10)) + \tau(A(s, 110)) + \tau(A(s, 1101)),
\end{aligned}$$

for $s = A(i, 0^{2k})$. We find that $(A(i, 0^{20}), E_{20}) = (A(i, 0^{18}), E_{18})$, so that we only have to verify that identities (2.21) through (2.34) hold for $2 \leq k \leq 10$, which turns out to be true. $\square$

In the following lemma, we express M_{2k} , M_{2k+1} , W_{2k} , and W_{2k+1} in terms of M^e , M^o , W^e , and W^o .

Lemma 2.3. *For all integer $k \geq 2$, and $u = 2^{2k-1}$,*

$$\begin{aligned} M_{2k} &= M^e[:7u] + x^{2u}M^e[:5u] + x^{3u}M^e[:4u] + x^{5u}M^e[:2u] + x^{6u}M^e[:u] \\ &\quad + x^{7u}R^e, \\ W_{2k} &= W^e[:7u] + x^{2u}W^e[:5u] + x^{3u}W^e[:4u] + x^{5u}W^e[:2u] + x^{6u}W^e[:u] \\ &\quad + x^{7u}R^e. \end{aligned}$$

For all integer $k \geq 2$, and $u = 2^{2k}$,

$$\begin{aligned} M_{2k+1} &= M^o[:7u] + x^{2u}M^o[:5u] + x^{3u}M^o[:4u] + x^{5u}M^o[:2u] + x^{6u}M^o[:u] \\ &\quad + x^{7u}R^o, \\ W_{2k+1} &= W^o[:7u] + x^{2u}W^o[:5u] + x^{3u}W^o[:4u] + x^{5u}W^o[:2u] + x^{6u}W^o[:u] \\ &\quad + x^{7u}R^o. \end{aligned}$$

Proof. We call the four identities in Lemma 2.3 also by the name M_{2k} , W_{2k} , M_{2k+1} , and W_{2k+1} . For $n = 2$, the identities can be verified directly. For $n \geq 2$, we claim that

$$\begin{aligned} M_{2k} \wedge W_{2k} &\Rightarrow M_{2k+1} \wedge W_{2k+1}, \\ M_{2k+1} \wedge W_{2k+1} &\Rightarrow M_{2k+2} \wedge W_{2k+2}. \end{aligned}$$

We give the proof of

$$M_{2k} \wedge W_{2k} \Rightarrow M_{2k+1},$$

the other ones being similar. Set $u = 2^{2k}$ and $v = 2^{2k-1}$. By definition and induction hypothesis, the left side of identity M_{2k+1} is equal to

$$\begin{aligned} (2.35) \quad W_{2k}M_{2k} &= (W^e[:7v] + x^{2v}W^e[:5v] + x^{3v}W^e[:4v] + x^{5v}W^e[:2v] + x^{6v}W^e[:v] \\ &\quad + x^{7v}R^e) \times (M^e[:7v] + x^{2v}M^e[:5v] + x^{3v}M^e[:4v] + x^{5v}M^e[:2v] \\ &\quad + x^{6v}M^e[:v] + x^{7v}R^e); \end{aligned}$$

Call this expression *lhs*. Note that both sides of identity M_{2k+1} have the same term of highest degree $x^{14v}R^o$. Therefore we only need to prove that their difference is $O(x^{14v})$. Using Lemma 2.1 it can be seen that the right side of identity M_{2k+1} is congruent, modulo x^{14v} , to

$$\begin{aligned} W^e[:14v]M^e[:14v] &+ x^{4v}W^e[:10v]M^e[:10v] + x^{6v}W^e[:8v]M^e[:8v] \\ &+ x^{10v}W^e[:4v]M^e[:4v] + x^{12v}W^e[:2v]M^e[:2v]. \end{aligned}$$

For all $n \leq 14$, replace the occurrences of $W^e[:n \cdot v]$ and $M^e[:n \cdot v]$ in the above expression by the reduction modulo $x^{n \cdot v}$ of the right side of the corresponding identity in Lemma 2.2 and get a new expression, which we call *rhs*. Define

$$\begin{aligned} X &:= x^v, \\ a_n &:= W^e[n \cdot v : (n+1) \cdot v] / X^n, \end{aligned}$$

$$b_n := M^e[n \cdot v : (n+1) \cdot v]/X^n,$$

$$c := R^e.$$

Using the notation introduced above, we can represent the expressions *lhs* (2.35) and *rhs* as polynomials in $\mathbb{F}_2[a_1, \dots, a_{14}, b_1, \dots, b_{14}, c][X]$. Note that it is not a problem that a_j commutes with b_k while W^e does not commute with M^e , because in the expressions concerned, the products of W^e -terms and M^e -terms are always in the same order. We let the computer do the simplification and check that the difference between these two polynomials is indeed $O(X^{14})$, which completes the proof. $\square$

Proof of Theorem 1.1. We prove the theorem for $\text{CF}(a, b)$; for $\text{CF}(b, a)$, the proof is similar. By Lemma 2.3, we have For all $0 \leq j, k \leq 1$,

$$\begin{aligned} \lim_{n \rightarrow \infty} M_{2n,j,k} &= M_{j,k}^e, \\ \lim_{n \rightarrow \infty} M_{2n+1,j,k} &= M_{j,k}^o, \\ \lim_{n \rightarrow \infty} W_{2n,j,k} &= W_{j,k}^e, \\ \lim_{n \rightarrow \infty} W_{2n+1,j,k} &= W_{j,k}^o. \end{aligned}$$

Let $z = 1/x$. By the convergence theorem and identity (2.2),

$$\text{CF}(a, b) = \frac{M_{0,1}^e(x)}{M_{0,0}^e(x)}.$$

By definition, $\phi(M^e, 0, 1)$ and $\phi(M^e, 0, 0)$ are minimal polynomials of $M_{0,1}^e$ and $M_{0,0}^e$. Therefore

$$P(x, y) = \text{Res}_t (\phi(M^e, 0, 1)(x, t), y^{12} \phi(M^e, 0, 1)(x, t/y))$$

is an annihilating polynomial of $f(x) = M_{0,1}^e/M_{0,0}^e$.

Define

$$Q(x, y) = q_4(x)y^4 + q_3(x)y^3 + q_2(x)y^2 + q_1(x)y + q_0(x),$$

where

$$\begin{aligned} q_0(x) &= x^{18} + x^{16} + x^{15} + x^{14} + x^{12} + x^{10} + x^9 + x^7, \\ q_1(x) &= x^{15} + x^{14} + x^{12} + x^{11} + x^9 + x^8 + x^7 + x^5 + x^4 + x^3, \\ q_2(x) &= x^{16} + x^{14} + x^8 + x^6 + x^4 + 1, \\ q_3(x) &= x^{15} + x^{14} + x^{12} + x^{11} + x^9 + x^8 + x^7 + x^5 + x^4 + x^3, \\ q_4(x) &= x^{18} + x^{16} + x^{15} + x^{12} + x^{10} + x^9 + x^8 + x^7 + x^6. \end{aligned}$$

The polynomial $Q(x, y)$ is the candidate for the minimal polynomial of $f(x)$ found by Padé-Hermite approximation. To prove that it is indeed the minimal polynomial of $f(x)$, we only need to prove that it is an irreducible factor of $P(x, y)$ of multiplicity m and $R(x, y) := P(x, y)/Q(x, y)^m$ is not an annihilating polynomial of $f(x)$. We verify the first point directly. For the second point, we find that when we truncate $f(x)$ to order 216, and substitute it for y in $R(z, y)$, we get a series with valuation smaller than 216, which proves that $R(z, y)$ is not an annihilating polynomial of $f(x)$. Finally, $z^{12}Q(1/z, y)$ is the minimal polynomial of $\text{CF}(a, b) = f(1/z)$. $\square$

3. ANNEXE

All 16 polynomials are of the form

$$p_0(x) + p_3(x)y^3 + p_6(x)y^6 + p_9(x)y^9 + p_{12}(x)y^{12}.$$

The coefficients $p_j(x)$ and the 16 initial terms to determine the solutions uniquely are given below.

For $\phi(M^e, 0, 0)$:

$$\begin{aligned} p_0(x) &= x^{96} + x^{94} + x^{86} + x^{80} + x^{78} + x^{74} + x^{58} + x^{52} + x^{50} + x^{48} + x^{46} \\ &\quad + x^{42} + x^{36}, \\ p_3(x) &= x^{86} + x^{80} + x^{78} + x^{70} + x^{60} + x^{52} + x^{48} + x^{36} + x^{32} + x^{30} + x^{24} \\ &\quad + x^{20}, \\ p_6(x) &= x^{88} + x^{84} + x^{78} + x^{66} + x^{64} + x^{62} + x^{46} + x^{42} + x^{40} + x^{24} + x^{22} \\ &\quad + x^{20} + x^{16} + x^{14} + x^{12} + x^{10} + x^6 + x^4 + x^2 + 1, \\ p_9(x) &= x^{82} + x^{80} + x^{78} + x^{76} + x^{68} + x^{66} + x^{64} + x^{56} + x^{50} + x^{48} + x^{44} \\ &\quad + x^{40} + x^{38} + x^{36} + x^{32} + x^{30} + x^{28} + x^{26} + x^{18} + x^{12} + x^{10} + x^6, \\ p_{12}(x) &= x^{84} + x^{82} + x^{80} + x^{68} + x^{66} + x^{64} + x^{36} + x^{34} + x^{32} + x^4 + x^2 + 1, \end{aligned}$$

and the initial terms are $[1, 0, 0, 0, 0, 0, 1, 0, 0, 0, 1, 0, 1, 0, 0, 0]$.

For $\phi(M^e, 0, 1)$:

$$\begin{aligned} p_0(x) &= x^{99} + x^{96} + x^{92} + x^{89} + x^{87} + x^{85} + x^{84} + x^{83} + x^{82} + x^{81} + x^{80} \\ &\quad + x^{79} + x^{77} + x^{74} + x^{72} + x^{71} + x^{70} + x^{68} + x^{67} + x^{63} + x^{62} + x^{61} \\ &\quad + x^{59} + x^{57} + x^{56} + x^{54} + x^{51} + x^{49} + x^{48} + x^{47} + x^{44} + x^{42} + x^{41} \\ &\quad + x^{40} + x^{39}, \\ p_3(x) &= x^{78} + x^{77} + x^{74} + x^{73} + x^{70} + x^{69} + x^{64} + x^{63} + x^{62} + x^{61} + x^{60} \\ &\quad + x^{59} + x^{54} + x^{53} + x^{50} + x^{49} + x^{42} + x^{41} + x^{40} + x^{39} + x^{36} + x^{35} \\ &\quad + x^{32} + x^{31} + x^{30} + x^{29} + x^{28} + x^{27}, \\ p_6(x) &= x^{81} + x^{79} + x^{78} + x^{77} + x^{76} + x^{74} + x^{73} + x^{71} + x^{70} + x^{68} + x^{67} \\ &\quad + x^{64} + x^{63} + x^{60} + x^{59} + x^{57} + x^{56} + x^{54} + x^{45} + x^{43} + x^{42} + x^{41} \\ &\quad + x^{40} + x^{38} + x^{29} + x^{27} + x^{26} + x^{24} + x^{23} + x^{21} + x^{20} + x^{18}, \\ p_9(x) &= x^{84} + x^{83} + x^{82} + x^{81} + x^{80} + x^{79} + x^{70} + x^{69} + x^{68} + x^{67} + x^{66} \\ &\quad + x^{65} + x^{64} + x^{63} + x^{56} + x^{55} + x^{50} + x^{49} + x^{48} + x^{47} + x^{44} + x^{43} \\ &\quad + x^{40} + x^{39} + x^{38} + x^{37} + x^{32} + x^{31} + x^{26} + x^{25} + x^{22} + x^{21} + x^{20} \\ &\quad + x^{19} + x^{18} + x^{17} + x^{14} + x^{13} + x^{10} + x^9, \\ p_{12}(x) &= x^{87} + x^{84} + x^{79} + x^{76} + x^{75} + x^{72} + x^{67} + x^{64} + x^{39} + x^{36} + x^{31} \\ &\quad + x^{28} + x^{27} + x^{24} + x^{23} + x^{20} + x^{19} + x^{16} + x^{15} + x^{12} + x^{11} + x^8 \\ &\quad + x^3 + 1, \end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 1, 0, 0, 1, 0, 0, 0, 1, 0, 1, 1, 1]$.

For $\phi(M^e, 1, 0)$:

$$\begin{aligned}
p_0(x) &= x^{99} + x^{96} + x^{92} + x^{89} + x^{87} + x^{85} + x^{84} + x^{83} + x^{82} + x^{81} + x^{80} \\
&\quad + x^{79} + x^{77} + x^{74} + x^{72} + x^{71} + x^{70} + x^{68} + x^{67} + x^{63} + x^{62} + x^{61} \\
&\quad + x^{59} + x^{57} + x^{56} + x^{54} + x^{51} + x^{49} + x^{48} + x^{47} + x^{44} + x^{42} + x^{41} \\
&\quad + x^{40} + x^{39}, \\
p_3(x) &= x^{78} + x^{77} + x^{74} + x^{73} + x^{70} + x^{69} + x^{64} + x^{63} + x^{62} + x^{61} + x^{60} \\
&\quad + x^{59} + x^{54} + x^{53} + x^{50} + x^{49} + x^{42} + x^{41} + x^{40} + x^{39} + x^{36} + x^{35} \\
&\quad + x^{32} + x^{31} + x^{30} + x^{29} + x^{28} + x^{27}, \\
p_6(x) &= x^{81} + x^{79} + x^{78} + x^{77} + x^{76} + x^{74} + x^{73} + x^{71} + x^{70} + x^{68} + x^{67} \\
&\quad + x^{64} + x^{63} + x^{60} + x^{59} + x^{57} + x^{56} + x^{54} + x^{45} + x^{43} + x^{42} + x^{41} \\
&\quad + x^{40} + x^{38} + x^{29} + x^{27} + x^{26} + x^{24} + x^{23} + x^{21} + x^{20} + x^{18}, \\
p_9(x) &= x^{84} + x^{83} + x^{82} + x^{81} + x^{80} + x^{79} + x^{70} + x^{69} + x^{68} + x^{67} + x^{66} \\
&\quad + x^{65} + x^{64} + x^{63} + x^{56} + x^{55} + x^{50} + x^{49} + x^{48} + x^{47} + x^{44} + x^{43} \\
&\quad + x^{40} + x^{39} + x^{38} + x^{37} + x^{32} + x^{31} + x^{26} + x^{25} + x^{22} + x^{21} + x^{20} \\
&\quad + x^{19} + x^{18} + x^{17} + x^{14} + x^{13} + x^{10} + x^9, \\
p_{12}(x) &= x^{87} + x^{84} + x^{79} + x^{76} + x^{75} + x^{72} + x^{67} + x^{64} + x^{39} + x^{36} + x^{31} \\
&\quad + x^{28} + x^{27} + x^{24} + x^{23} + x^{20} + x^{19} + x^{16} + x^{15} + x^{12} + x^{11} + x^8 \\
&\quad + x^3 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 1, 0, 0, 1, 0, 0, 0, 1, 0, 1, 1, 1]$.

For $\phi(M^e, 1, 1)$:

$$\begin{aligned}
p_0(x) &= x^{102} + x^{100} + x^{98} + x^{92} + x^{86} + x^{84} + x^{82} + x^{80} + x^{78} + x^{74} + x^{72} \\
&\quad + x^{70} + x^{68} + x^{58} + x^{52} + x^{50} + x^{46} + x^{44} + x^{42}, \\
p_3(x) &= x^{92} + x^{88} + x^{80} + x^{74} + x^{72} + x^{70} + x^{66} + x^{64} + x^{58} + x^{56} + x^{54} \\
&\quad + x^{52} + x^{46} + x^{44} + x^{40} + x^{38} + x^{32} + x^{30} + x^{24} + x^{22} + x^{20} + x^{18}, \\
p_6(x) &= x^{94} + x^{88} + x^{84} + x^{82} + x^{78} + x^{76} + x^{74} + x^{70} + x^{68} + x^{60} + x^{54} \\
&\quad + x^{50} + x^{48} + x^{46} + x^{44} + x^{36} + x^{32} + x^{30} + x^{20} + x^{14} + x^{12} + x^6 \\
&\quad + x^2 + 1, \\
p_9(x) &= x^{88} + x^{86} + x^{82} + x^{78} + x^{76} + x^{74} + x^{72} + x^{62} + x^{56} + x^{54} + x^{50} \\
&\quad + x^{46} + x^{44} + x^{42} + x^{38} + x^{34} + x^{32} + x^{30} + x^{28} + x^{24} + x^{22} + x^{20} \\
&\quad + x^{18} + x^{16} + x^{14} + x^6, \\
p_{12}(x) &= x^{90} + x^{88} + x^{66} + x^{64} + x^{42} + x^{40} + x^{26} + x^{24} + x^{18} + x^{16} + x^2 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 0, 0, 0, 1, 0, 0, 0, 0, 0, 0, 0]$.

For $\phi(W^e, 0, 0)$:

$$\begin{aligned}
p_0(x) &= x^{102} + x^{100} + x^{98} + x^{94} + x^{90} + x^{88} + x^{80} + x^{78} + x^{76} + x^{74} + x^{72} \\
&\quad + x^{68} + x^{66} + x^{64} + x^{62} + x^{54} + x^{52} + x^{50} + x^{46} + x^{40} + x^{36},
\end{aligned}$$

$$\begin{aligned}
p_3(x) &= x^{92} + x^{88} + x^{84} + x^{78} + x^{72} + x^{70} + x^{66} + x^{64} + x^{58} + x^{56} + x^{54} \\
&\quad + x^{48} + x^{44} + x^{42} + x^{40} + x^{38} + x^{36} + x^{28} + x^{26} + x^{20}, \\
p_6(x) &= x^{94} + x^{88} + x^{86} + x^{84} + x^{76} + x^{74} + x^{72} + x^{60} + x^{48} + x^{44} + x^{40} \\
&\quad + x^{34} + x^{32} + x^{30} + x^{26} + x^{24} + x^{16} + x^6 + x^2 + 1, \\
p_9(x) &= x^{88} + x^{86} + x^{82} + x^{78} + x^{76} + x^{74} + x^{72} + x^{62} + x^{56} + x^{54} + x^{50} \\
&\quad + x^{46} + x^{44} + x^{42} + x^{38} + x^{34} + x^{32} + x^{30} + x^{28} + x^{24} + x^{22} + x^{20} \\
&\quad + x^{18} + x^{16} + x^{14} + x^6, \\
p_{12}(x) &= x^{90} + x^{88} + x^{66} + x^{64} + x^{42} + x^{40} + x^{26} + x^{24} + x^{18} + x^{16} + x^2 + 1,
\end{aligned}$$

and the initial terms are $[1, 0, 0, 0, 0, 0, 0, 0, 1, 0, 0, 0, 1, 0, 0, 0]$.

For $\phi(W^e, 0, 1)$:

$$\begin{aligned}
p_0(x) &= x^{99} + x^{96} + x^{92} + x^{86} + x^{85} + x^{82} + x^{78} + x^{76} + x^{75} + x^{73} + x^{72} \\
&\quad + x^{68} + x^{67} + x^{66} + x^{65} + x^{62} + x^{59} + x^{58} + x^{56} + x^{55} + x^{54} + x^{53} \\
&\quad + x^{50} + x^{42} + x^{41} + x^{40} + x^{39}, \\
p_3(x) &= x^{78} + x^{77} + x^{74} + x^{73} + x^{70} + x^{69} + x^{64} + x^{63} + x^{62} + x^{61} + x^{60} \\
&\quad + x^{59} + x^{54} + x^{53} + x^{50} + x^{49} + x^{42} + x^{41} + x^{40} + x^{39} + x^{36} + x^{35} \\
&\quad + x^{32} + x^{31} + x^{30} + x^{29} + x^{28} + x^{27}, \\
p_6(x) &= x^{81} + x^{79} + x^{78} + x^{77} + x^{76} + x^{74} + x^{73} + x^{71} + x^{70} + x^{68} + x^{67} \\
&\quad + x^{64} + x^{63} + x^{60} + x^{59} + x^{57} + x^{56} + x^{54} + x^{45} + x^{43} + x^{42} + x^{41} \\
&\quad + x^{40} + x^{38} + x^{29} + x^{27} + x^{26} + x^{24} + x^{23} + x^{21} + x^{20} + x^{18}, \\
p_9(x) &= x^{84} + x^{83} + x^{82} + x^{81} + x^{80} + x^{79} + x^{70} + x^{69} + x^{68} + x^{67} + x^{66} \\
&\quad + x^{65} + x^{64} + x^{63} + x^{56} + x^{55} + x^{50} + x^{49} + x^{48} + x^{47} + x^{44} + x^{43} \\
&\quad + x^{40} + x^{39} + x^{38} + x^{37} + x^{32} + x^{31} + x^{26} + x^{25} + x^{22} + x^{21} + x^{20} \\
&\quad + x^{19} + x^{18} + x^{17} + x^{14} + x^{13} + x^{10} + x^9, \\
p_{12}(x) &= x^{87} + x^{84} + x^{79} + x^{76} + x^{75} + x^{72} + x^{67} + x^{64} + x^{39} + x^{36} + x^{31} \\
&\quad + x^{28} + x^{27} + x^{24} + x^{23} + x^{20} + x^{19} + x^{16} + x^{15} + x^{12} + x^{11} + x^8 \\
&\quad + x^3 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 1, 0, 1, 0, 1, 0, 1, 1, 0, 0, 1, 1, 1]$.

For $\phi(W^e, 1, 0)$:

$$\begin{aligned}
p_0(x) &= x^{99} + x^{96} + x^{92} + x^{86} + x^{85} + x^{82} + x^{78} + x^{76} + x^{75} + x^{73} + x^{72} \\
&\quad + x^{68} + x^{67} + x^{66} + x^{65} + x^{62} + x^{59} + x^{58} + x^{56} + x^{55} + x^{54} + x^{53} \\
&\quad + x^{50} + x^{42} + x^{41} + x^{40} + x^{39}, \\
p_3(x) &= x^{78} + x^{77} + x^{74} + x^{73} + x^{70} + x^{69} + x^{64} + x^{63} + x^{62} + x^{61} + x^{60} \\
&\quad + x^{59} + x^{54} + x^{53} + x^{50} + x^{49} + x^{42} + x^{41} + x^{40} + x^{39} + x^{36} + x^{35} \\
&\quad + x^{32} + x^{31} + x^{30} + x^{29} + x^{28} + x^{27}, \\
p_6(x) &= x^{81} + x^{79} + x^{78} + x^{77} + x^{76} + x^{74} + x^{73} + x^{71} + x^{70} + x^{68} + x^{67}
\end{aligned}$$

$$\begin{aligned}
& + x^{64} + x^{63} + x^{60} + x^{59} + x^{57} + x^{56} + x^{54} + x^{45} + x^{43} + x^{42} + x^{41} \\
& + x^{40} + x^{38} + x^{29} + x^{27} + x^{26} + x^{24} + x^{23} + x^{21} + x^{20} + x^{18}, \\
p_9(x) & = x^{84} + x^{83} + x^{82} + x^{81} + x^{80} + x^{79} + x^{70} + x^{69} + x^{68} + x^{67} + x^{66} \\
& + x^{65} + x^{64} + x^{63} + x^{56} + x^{55} + x^{50} + x^{49} + x^{48} + x^{47} + x^{44} + x^{43} \\
& + x^{40} + x^{39} + x^{38} + x^{37} + x^{32} + x^{31} + x^{26} + x^{25} + x^{22} + x^{21} + x^{20} \\
& + x^{19} + x^{18} + x^{17} + x^{14} + x^{13} + x^{10} + x^9, \\
p_{12}(x) & = x^{87} + x^{84} + x^{79} + x^{76} + x^{75} + x^{72} + x^{67} + x^{64} + x^{39} + x^{36} + x^{31} \\
& + x^{28} + x^{27} + x^{24} + x^{23} + x^{20} + x^{19} + x^{16} + x^{15} + x^{12} + x^{11} + x^8 \\
& + x^3 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 1, 0, 1, 0, 1, 0, 1, 1, 0, 0, 1, 1, 1]$.

For $\phi(W^e, 1, 1)$:

$$\begin{aligned}
p_0(x) & = x^{96} + x^{94} + x^{88} + x^{82} + x^{80} + x^{74} + x^{70} + x^{62} + x^{58} + x^{54} + x^{52} \\
& + x^{44} + x^{42}, \\
p_3(x) & = x^{86} + x^{80} + x^{70} + x^{66} + x^{62} + x^{60} + x^{54} + x^{52} + x^{50} + x^{48} + x^{46} \\
& + x^{42} + x^{38} + x^{36} + x^{34} + x^{32} + x^{26} + x^{24} + x^{20} + x^{18}, \\
p_6(x) & = x^{88} + x^{84} + x^{80} + x^{78} + x^{74} + x^{72} + x^{68} + x^{66} + x^{60} + x^{58} + x^{52} \\
& + x^{50} + x^{40} + x^{38} + x^{34} + x^{30} + x^{26} + x^{20} + x^{16} + x^{10} + x^6 + x^4 + x^2 \\
& + 1, \\
p_9(x) & = x^{82} + x^{80} + x^{78} + x^{76} + x^{68} + x^{66} + x^{64} + x^{56} + x^{50} + x^{48} + x^{44} \\
& + x^{40} + x^{38} + x^{36} + x^{32} + x^{30} + x^{28} + x^{26} + x^{18} + x^{12} + x^{10} + x^6, \\
p_{12}(x) & = x^{84} + x^{82} + x^{80} + x^{68} + x^{66} + x^{64} + x^{36} + x^{34} + x^{32} + x^4 + x^2 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 0, 1, 0, 0, 0, 1, 0, 0, 0, 0, 0]$.

For $\phi(M^o, 0, 0)$:

$$\begin{aligned}
p_0(x) & = x^{99} + x^{96} + x^{95} + x^{92} + x^{90} + x^{87} + x^{85} + x^{84} + x^{83} + x^{82} + x^{80} \\
& + x^{77} + x^{76} + x^{75} + x^{74} + x^{73} + x^{71} + x^{67} + x^{66} + x^{64} + x^{63} + x^{60} \\
& + x^{59} + x^{56} + x^{52} + x^{49} + x^{48} + x^{47} + x^{46} + x^{43} + x^{42} + x^{41} + x^{39} \\
& + x^{38} + x^{36}, \\
p_3(x) & = x^{87} + x^{84} + x^{78} + x^{75} + x^{74} + x^{73} + x^{72} + x^{64} + x^{62} + x^{61} + x^{60} \\
& + x^{59} + x^{58} + x^{56} + x^{55} + x^{54} + x^{52} + x^{51} + x^{50} + x^{49} + x^{48} + x^{46} \\
& + x^{43} + x^{42} + x^{41} + x^{39} + x^{38} + x^{37} + x^{35} + x^{34} + x^{31} + x^{30} + x^{29} \\
& + x^{27} + x^{26} + x^{24}, \\
p_6(x) & = x^{91} + x^{88} + x^{87} + x^{84} + x^{82} + x^{79} + x^{78} + x^{77} + x^{76} + x^{70} + x^{68} \\
& + x^{66} + x^{64} + x^{61} + x^{59} + x^{58} + x^{55} + x^{52} + x^{51} + x^{50} + x^{49} + x^{47} \\
& + x^{46} + x^{45} + x^{43} + x^{42} + x^{39} + x^{38} + x^{37} + x^{34} + x^{30} + x^{27} + x^{25} \\
& + x^{24} + x^{23} + x^{21} + x^{20} + x^{19} + x^{17} + x^{14} + x^{11} + x^8 + x^3 + 1,
\end{aligned}$$

$$\begin{aligned}
p_9(x) &= x^{83} + x^{80} + x^{79} + x^{76} + x^{35} + x^{32} + x^{31} + x^{28} + x^{19} + x^{16} + x^{15} \\
&\quad + x^{12}, \\
p_{12}(x) &= x^{87} + x^{84} + x^{83} + x^{80} + x^{75} + x^{72} + x^{67} + x^{64} + x^{39} + x^{36} + x^{35} \\
&\quad + x^{32} + x^{27} + x^{24} + x^{23} + x^{20} + x^{11} + x^8 + x^3 + 1,
\end{aligned}$$

and the initial terms are $[1, 0, 0, 0, 0, 0, 1, 1, 1, 1, 0, 1, 1, 1, 0, 0]$.

For $\phi(M^o, 0, 1)$:

$$\begin{aligned}
p_0(x) &= x^{102} + x^{100} + x^{96} + x^{95} + x^{94} + x^{92} + x^{91} + x^{90} + x^{88} + x^{86} + x^{80} \\
&\quad + x^{79} + x^{77} + x^{75} + x^{74} + x^{69} + x^{68} + x^{66} + x^{59} + x^{57} + x^{56} + x^{52} \\
&\quad + x^{47} + x^{46} + x^{42} + x^{40} + x^{39}, \\
p_3(x) &= x^{81} + x^{80} + x^{78} + x^{77} + x^{75} + x^{74} + x^{72} + x^{71} + x^{69} + x^{68} + x^{66} \\
&\quad + x^{65} + x^{63} + x^{62} + x^{60} + x^{59} + x^{57} + x^{56} + x^{54} + x^{53} + x^{51} + x^{50} \\
&\quad + x^{49} + x^{47} + x^{46} + x^{44} + x^{43} + x^{41} + x^{40} + x^{38} + x^{37} + x^{35} + x^{34} \\
&\quad + x^{33} + x^{31} + x^{30} + x^{28} + x^{27}, \\
p_6(x) &= x^{84} + x^{78} + x^{68} + x^{62} + x^{56} + x^{52} + x^{50} + x^{46} + x^{44} + x^{38} + x^{32} \\
&\quad + x^{28} + x^{26} + x^{24} + x^{22} + x^{18}, \\
p_9(x) &= x^{87} + x^{86} + x^{85} + x^{75} + x^{74} + x^{73} + x^{39} + x^{38} + x^{37} + x^{27} + x^{26} \\
&\quad + x^{25} + x^{23} + x^{22} + x^{21} + x^{11} + x^{10} + x^9, \\
p_{12}(x) &= x^{90} + x^{88} + x^{86} + x^{84} + x^{82} + x^{80} + x^{78} + x^{76} + x^{66} + x^{64} + x^{42} \\
&\quad + x^{40} + x^{38} + x^{36} + x^{34} + x^{32} + x^{30} + x^{28} + x^{26} + x^{24} + x^{22} + x^{20} \\
&\quad + x^{14} + x^{12} + x^2 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 1, 0, 0, 1, 0, 0, 0, 0, 1, 0, 1, 1]$.

For $\phi(M^o, 1, 0)$:

$$\begin{aligned}
p_0(x) &= x^{100} + x^{96} + x^{94} + x^{93} + x^{92} + x^{91} + x^{90} + x^{89} + x^{87} + x^{85} + x^{83} \\
&\quad + x^{82} + x^{80} + x^{76} + x^{75} + x^{72} + x^{71} + x^{68} + x^{64} + x^{63} + x^{62} + x^{61} \\
&\quad + x^{56} + x^{51} + x^{49} + x^{46} + x^{41} + x^{40} + x^{39}, \\
p_3(x) &= x^{79} + x^{78} + x^{77} + x^{75} + x^{74} + x^{72} + x^{70} + x^{68} + x^{66} + x^{64} + x^{63} \\
&\quad + x^{61} + x^{60} + x^{59} + x^{55} + x^{54} + x^{53} + x^{51} + x^{50} + x^{48} + x^{47} + x^{45} \\
&\quad + x^{44} + x^{43} + x^{39} + x^{38} + x^{37} + x^{35} + x^{34} + x^{32} + x^{31} + x^{29} + x^{28} \\
&\quad + x^{27}, \\
p_6(x) &= x^{82} + x^{80} + x^{76} + x^{72} + x^{70} + x^{58} + x^{56} + x^{54} + x^{46} + x^{44} + x^{40} \\
&\quad + x^{38} + x^{30} + x^{28} + x^{26} + x^{22} + x^{20} + x^{18}, \\
p_9(x) &= x^{85} + x^{84} + x^{82} + x^{81} + x^{77} + x^{76} + x^{74} + x^{73} + x^{37} + x^{36} + x^{34} \\
&\quad + x^{33} + x^{29} + x^{28} + x^{26} + x^{25} + x^{21} + x^{20} + x^{18} + x^{17} + x^{13} + x^{12} \\
&\quad + x^{10} + x^9, \\
p_{12}(x) &= x^{88} + x^{80} + x^{76} + x^{72} + x^{68} + x^{64} + x^{40} + x^{32} + x^{28} + x^{20} + x^{12} + x^8
\end{aligned}$$

$$+ x^4 + 1,$$

and the initial terms are $[0, 0, 0, 1, 0, 1, 0, 1, 0, 0, 0, 1, 0, 0, 0, 0]$.

For $\phi(M^o, 1, 1)$:

$$\begin{aligned} p_0(x) &= x^{99} + x^{96} + x^{95} + x^{92} + x^{91} + x^{90} + x^{89} + x^{88} + x^{87} + x^{86} + x^{85} \\ &\quad + x^{84} + x^{82} + x^{76} + x^{75} + x^{74} + x^{73} + x^{71} + x^{70} + x^{62} + x^{60} + x^{58} \\ &\quad + x^{56} + x^{55} + x^{51} + x^{50} + x^{48} + x^{47} + x^{46} + x^{45} + x^{42}, \\ p_3(x) &= x^{87} + x^{84} + x^{79} + x^{78} + x^{77} + x^{76} + x^{71} + x^{70} + x^{69} + x^{68} + x^{66} \\ &\quad + x^{65} + x^{64} + x^{61} + x^{60} + x^{58} + x^{54} + x^{53} + x^{42} + x^{41} + x^{40} + x^{38} \\ &\quad + x^{36} + x^{33} + x^{32} + x^{31} + x^{29} + x^{26}, \\ p_6(x) &= x^{91} + x^{88} + x^{87} + x^{84} + x^{83} + x^{82} + x^{81} + x^{80} + x^{75} + x^{74} + x^{73} \\ &\quad + x^{72} + x^{71} + x^{66} + x^{64} + x^{63} + x^{61} + x^{60} + x^{59} + x^{58} + x^{57} + x^{54} \\ &\quad + x^{50} + x^{49} + x^{48} + x^{46} + x^{44} + x^{41} + x^{40} + x^{39} + x^{37} + x^{35} + x^{34} \\ &\quad + x^{32} + x^{31} + x^{30} + x^{29} + x^{28} + x^{27} + x^{26} + x^{25} + x^{24} + x^{18} + x^{17} \\ &\quad + x^{16} + x^{15} + x^{14} + x^{12} + x^{11} + x^8 + x^3 + 1, \\ p_9(x) &= x^{83} + x^{80} + x^{79} + x^{76} + x^{35} + x^{32} + x^{31} + x^{28} + x^{19} + x^{16} + x^{15} \\ &\quad + x^{12}, \\ p_{12}(x) &= x^{87} + x^{84} + x^{83} + x^{80} + x^{75} + x^{72} + x^{67} + x^{64} + x^{39} + x^{36} + x^{35} \\ &\quad + x^{32} + x^{27} + x^{24} + x^{23} + x^{20} + x^{11} + x^8 + x^3 + 1, \end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 0, 0, 1, 0, 1, 1, 1, 1, 1, 0, 0]$.

For $\phi(W^o, 0, 0)$:

$$\begin{aligned} p_0(x) &= x^{99} + x^{96} + x^{95} + x^{92} + x^{90} + x^{87} + x^{85} + x^{84} + x^{83} + x^{82} + x^{80} \\ &\quad + x^{77} + x^{76} + x^{75} + x^{74} + x^{73} + x^{71} + x^{67} + x^{66} + x^{64} + x^{63} + x^{60} \\ &\quad + x^{59} + x^{56} + x^{52} + x^{49} + x^{48} + x^{47} + x^{46} + x^{43} + x^{42} + x^{41} + x^{39} \\ &\quad + x^{38} + x^{36}, \\ p_3(x) &= x^{87} + x^{84} + x^{78} + x^{75} + x^{74} + x^{73} + x^{72} + x^{64} + x^{62} + x^{61} + x^{60} \\ &\quad + x^{59} + x^{58} + x^{56} + x^{55} + x^{54} + x^{52} + x^{51} + x^{50} + x^{49} + x^{48} + x^{46} \\ &\quad + x^{43} + x^{42} + x^{41} + x^{39} + x^{38} + x^{37} + x^{35} + x^{34} + x^{31} + x^{30} + x^{29} \\ &\quad + x^{27} + x^{26} + x^{24}, \\ p_6(x) &= x^{91} + x^{88} + x^{87} + x^{84} + x^{82} + x^{79} + x^{78} + x^{77} + x^{76} + x^{70} + x^{68} \\ &\quad + x^{66} + x^{64} + x^{61} + x^{59} + x^{58} + x^{55} + x^{52} + x^{51} + x^{50} + x^{49} + x^{47} \\ &\quad + x^{46} + x^{45} + x^{43} + x^{42} + x^{39} + x^{38} + x^{37} + x^{34} + x^{30} + x^{27} + x^{25} \\ &\quad + x^{24} + x^{23} + x^{21} + x^{20} + x^{19} + x^{17} + x^{14} + x^{11} + x^8 + x^3 + 1, \\ p_9(x) &= x^{83} + x^{80} + x^{79} + x^{76} + x^{35} + x^{32} + x^{31} + x^{28} + x^{19} + x^{16} + x^{15} \\ &\quad + x^{12}, \\ p_{12}(x) &= x^{87} + x^{84} + x^{83} + x^{80} + x^{75} + x^{72} + x^{67} + x^{64} + x^{39} + x^{36} + x^{35} \end{aligned}$$

$$+ x^{32} + x^{27} + x^{24} + x^{23} + x^{20} + x^{11} + x^8 + x^3 + 1,$$

and the initial terms are $[1, 0, 0, 0, 0, 0, 1, 1, 1, 1, 0, 1, 1, 1, 0, 0]$.

For $\phi(W^o, 0, 1)$:

$$\begin{aligned} p_0(x) &= x^{100} + x^{96} + x^{94} + x^{93} + x^{92} + x^{91} + x^{90} + x^{89} + x^{87} + x^{85} + x^{83} \\ &\quad + x^{82} + x^{80} + x^{76} + x^{75} + x^{72} + x^{71} + x^{68} + x^{64} + x^{63} + x^{62} + x^{61} \\ &\quad + x^{56} + x^{51} + x^{49} + x^{46} + x^{41} + x^{40} + x^{39}, \\ p_3(x) &= x^{79} + x^{78} + x^{77} + x^{75} + x^{74} + x^{72} + x^{70} + x^{68} + x^{66} + x^{64} + x^{63} \\ &\quad + x^{61} + x^{60} + x^{59} + x^{55} + x^{54} + x^{53} + x^{51} + x^{50} + x^{48} + x^{47} + x^{45} \\ &\quad + x^{44} + x^{43} + x^{39} + x^{38} + x^{37} + x^{35} + x^{34} + x^{32} + x^{31} + x^{29} + x^{28} \\ &\quad + x^{27}, \\ p_6(x) &= x^{82} + x^{80} + x^{76} + x^{72} + x^{70} + x^{58} + x^{56} + x^{54} + x^{46} + x^{44} + x^{40} \\ &\quad + x^{38} + x^{30} + x^{28} + x^{26} + x^{22} + x^{20} + x^{18}, \\ p_9(x) &= x^{85} + x^{84} + x^{82} + x^{81} + x^{77} + x^{76} + x^{74} + x^{73} + x^{37} + x^{36} + x^{34} \\ &\quad + x^{33} + x^{29} + x^{28} + x^{26} + x^{25} + x^{21} + x^{20} + x^{18} + x^{17} + x^{13} + x^{12} \\ &\quad + x^{10} + x^9, \\ p_{12}(x) &= x^{88} + x^{80} + x^{76} + x^{72} + x^{68} + x^{64} + x^{40} + x^{32} + x^{28} + x^{20} + x^{12} + x^8 \\ &\quad + x^4 + 1, \end{aligned}$$

and the initial terms are $[0, 0, 0, 1, 0, 1, 0, 1, 0, 0, 0, 1, 0, 0, 0, 0]$.

For $\phi(W^o, 1, 0)$:

$$\begin{aligned} p_0(x) &= x^{102} + x^{100} + x^{96} + x^{95} + x^{94} + x^{92} + x^{91} + x^{90} + x^{88} + x^{86} + x^{80} \\ &\quad + x^{79} + x^{77} + x^{75} + x^{74} + x^{69} + x^{68} + x^{66} + x^{59} + x^{57} + x^{56} + x^{52} \\ &\quad + x^{47} + x^{46} + x^{42} + x^{40} + x^{39}, \\ p_3(x) &= x^{81} + x^{80} + x^{78} + x^{77} + x^{75} + x^{74} + x^{72} + x^{71} + x^{69} + x^{68} + x^{66} \\ &\quad + x^{65} + x^{63} + x^{62} + x^{60} + x^{59} + x^{57} + x^{56} + x^{54} + x^{53} + x^{51} + x^{50} \\ &\quad + x^{49} + x^{47} + x^{46} + x^{44} + x^{43} + x^{41} + x^{40} + x^{38} + x^{37} + x^{35} + x^{34} \\ &\quad + x^{33} + x^{31} + x^{30} + x^{28} + x^{27}, \\ p_6(x) &= x^{84} + x^{78} + x^{68} + x^{62} + x^{56} + x^{52} + x^{50} + x^{46} + x^{44} + x^{38} + x^{32} \\ &\quad + x^{28} + x^{26} + x^{24} + x^{22} + x^{18}, \\ p_9(x) &= x^{87} + x^{86} + x^{85} + x^{75} + x^{74} + x^{73} + x^{39} + x^{38} + x^{37} + x^{27} + x^{26} \\ &\quad + x^{25} + x^{23} + x^{22} + x^{21} + x^{11} + x^{10} + x^9, \\ p_{12}(x) &= x^{90} + x^{88} + x^{86} + x^{84} + x^{82} + x^{80} + x^{78} + x^{76} + x^{66} + x^{64} + x^{42} \\ &\quad + x^{40} + x^{38} + x^{36} + x^{34} + x^{32} + x^{30} + x^{28} + x^{26} + x^{24} + x^{22} + x^{20} \\ &\quad + x^{14} + x^{12} + x^2 + 1, \end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 1, 0, 0, 1, 0, 0, 0, 0, 1, 0, 1, 1]$.

For $\phi(W^o, 1, 1)$:

$$p_0(x) = x^{99} + x^{96} + x^{95} + x^{92} + x^{91} + x^{90} + x^{89} + x^{88} + x^{87} + x^{86} + x^{85}$$

$$\begin{aligned}
& + x^{84} + x^{82} + x^{76} + x^{75} + x^{74} + x^{73} + x^{71} + x^{70} + x^{62} + x^{60} + x^{58} \\
& + x^{56} + x^{55} + x^{51} + x^{50} + x^{48} + x^{47} + x^{46} + x^{45} + x^{42}, \\
p_3(x) = & x^{87} + x^{84} + x^{79} + x^{78} + x^{77} + x^{76} + x^{71} + x^{70} + x^{69} + x^{68} + x^{66} \\
& + x^{65} + x^{64} + x^{61} + x^{60} + x^{58} + x^{54} + x^{53} + x^{42} + x^{41} + x^{40} + x^{38} \\
& + x^{36} + x^{33} + x^{32} + x^{31} + x^{29} + x^{26}, \\
p_6(x) = & x^{91} + x^{88} + x^{87} + x^{84} + x^{83} + x^{82} + x^{81} + x^{80} + x^{75} + x^{74} + x^{73} \\
& + x^{72} + x^{71} + x^{66} + x^{64} + x^{63} + x^{61} + x^{60} + x^{59} + x^{58} + x^{57} + x^{54} \\
& + x^{50} + x^{49} + x^{48} + x^{46} + x^{44} + x^{41} + x^{40} + x^{39} + x^{37} + x^{35} + x^{34} \\
& + x^{32} + x^{31} + x^{30} + x^{29} + x^{28} + x^{27} + x^{26} + x^{25} + x^{24} + x^{18} + x^{17} \\
& + x^{16} + x^{15} + x^{14} + x^{12} + x^{11} + x^8 + x^3 + 1, \\
p_9(x) = & x^{83} + x^{80} + x^{79} + x^{76} + x^{35} + x^{32} + x^{31} + x^{28} + x^{19} + x^{16} + x^{15} \\
& + x^{12}, \\
p_{12}(x) = & x^{87} + x^{84} + x^{83} + x^{80} + x^{75} + x^{72} + x^{67} + x^{64} + x^{39} + x^{36} + x^{35} \\
& + x^{32} + x^{27} + x^{24} + x^{23} + x^{20} + x^{11} + x^8 + x^3 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 0, 0, 1, 0, 1, 1, 1, 1, 1, 0, 0]$.

Below is the transition function and output function of an 2-automaton that generates $T = M_{0,0}^o$:

Transition function $(n, j) \mapsto \delta(n, j)$ ($\Lambda(n) := [\delta(n, 0), \delta(n, 1)]$):

n	$\Lambda(n)$	n	$\Lambda(n)$	n	$\Lambda(n)$	n	$\Lambda(n)$	n	$\Lambda(n)$
0	[1, 2]	107	[177, 178]	214	[321, 322]	321	[394, 285]	428	[115, 320]
1	[3, 4]	108	[179, 42]	215	[55, 323]	322	[395, 80]	429	[458, 382]
2	[5, 6]	109	[180, 181]	216	[165, 147]	323	[396, 266]	430	[459, 460]
3	[7, 8]	110	[182, 180]	217	[136, 324]	324	[220, 70]	431	[461, 198]
4	[9, 10]	111	[183, 75]	218	[134, 131]	325	[325, 325]	432	[462, 427]
5	[11, 12]	112	[184, 185]	219	[136, 131]	326	[397, 224]	433	[463, 99]
6	[13, 14]	113	[186, 187]	220	[137, 325]	327	[398, 68]	434	[464, 465]
7	[15, 16]	114	[188, 189]	221	[132, 136]	328	[399, 86]	435	[267, 466]
8	[17, 18]	115	[190, 191]	222	[324, 132]	329	[400, 401]	436	[426, 467]
9	[19, 20]	116	[192, 193]	223	[326, 132]	330	[402, 325]	437	[241, 271]
10	[21, 22]	117	[194, 195]	224	[325, 137]	331	[403, 227]	438	[468, 392]
11	[23, 24]	118	[196, 197]	225	[134, 324]	332	[404, 357]	439	[469, 470]
12	[25, 26]	119	[198, 199]	226	[10, 301]	333	[405, 197]	440	[262, 471]
13	[27, 28]	120	[20, 200]	227	[316, 298]	334	[406, 407]	441	[472, 473]
14	[29, 30]	121	[22, 201]	228	[305, 40]	335	[393, 189]	442	[209, 474]
15	[31, 32]	122	[202, 203]	229	[116, 178]	336	[81, 408]	443	[475, 217]
16	[33, 34]	123	[204, 205]	230	[176, 291]	337	[409, 219]	444	[476, 26]
17	[35, 36]	124	[206, 204]	231	[327, 328]	338	[410, 411]	445	[247, 274]
18	[37, 38]	125	[207, 93]	232	[329, 328]	339	[200, 60]	446	[87, 477]
19	[39, 40]	126	[208, 209]	233	[330, 305]	340	[412, 413]	447	[478, 72]
20	[41, 42]	127	[210, 211]	234	[41, 306]	341	[205, 414]	448	[479, 235]

21	[31, 43]	128	[212, 213]	235	[114, 299]	342	[415, 406]	449	[480, 481]
22	[44, 45]	129	[214, 215]	236	[331, 46]	343	[416, 261]	450	[482, 386]
23	[9, 46]	130	[216, 23]	237	[166, 48]	344	[89, 17]	451	[483, 244]
24	[47, 48]	131	[217, 218]	238	[302, 150]	345	[417, 418]	452	[83, 361]
25	[49, 50]	132	[219, 220]	239	[332, 302]	346	[419, 78]	453	[107, 484]
26	[51, 52]	133	[76, 221]	240	[333, 43]	347	[420, 421]	454	[485, 486]
27	[53, 54]	134	[218, 222]	241	[296, 38]	348	[422, 15]	455	[373, 487]
28	[55, 54]	135	[223, 76]	242	[143, 127]	349	[423, 283]	456	[488, 85]
29	[56, 57]	136	[224, 219]	243	[309, 334]	350	[24, 424]	457	[489, 66]
30	[58, 7]	137	[225, 75]	244	[332, 149]	351	[425, 380]	458	[141, 436]
31	[59, 60]	138	[193, 226]	245	[8, 310]	352	[64, 426]	459	[173, 445]
32	[61, 62]	139	[227, 228]	246	[335, 7]	353	[427, 403]	460	[127, 453]
33	[63, 64]	140	[229, 230]	247	[120, 336]	354	[62, 428]	461	[454, 171]
34	[65, 66]	141	[231, 232]	248	[337, 37]	355	[326, 133]	462	[490, 180]
35	[67, 68]	142	[233, 234]	249	[338, 36]	356	[312, 50]	463	[457, 334]
36	[69, 61]	143	[235, 212]	250	[154, 319]	357	[315, 314]	464	[112, 429]
37	[70, 71]	144	[236, 237]	251	[339, 119]	358	[345, 429]	465	[181, 180]
38	[72, 73]	145	[238, 239]	252	[149, 303]	359	[321, 348]	466	[153, 291]
39	[74, 70]	146	[240, 229]	253	[162, 12]	360	[430, 431]	467	[491, 56]
40	[75, 76]	147	[234, 241]	254	[340, 341]	361	[146, 166]	468	[492, 150]
41	[77, 78]	148	[191, 242]	255	[12, 123]	362	[432, 146]	469	[346, 348]
42	[79, 21]	149	[243, 244]	256	[342, 160]	363	[330, 38]	470	[493, 328]
43	[80, 81]	150	[245, 3]	257	[142, 159]	364	[288, 325]	471	[494, 165]
44	[82, 83]	151	[246, 247]	258	[343, 158]	365	[433, 138]	472	[495, 448]
45	[84, 85]	152	[248, 249]	259	[160, 334]	366	[434, 435]	473	[496, 323]
46	[86, 87]	153	[103, 250]	260	[151, 52]	367	[129, 436]	474	[497, 146]
47	[88, 89]	154	[251, 94]	261	[57, 163]	368	[172, 159]	475	[443, 134]
48	[90, 91]	155	[252, 253]	262	[119, 47]	369	[177, 117]	476	[498, 334]
49	[92, 93]	156	[254, 106]	263	[335, 56]	370	[38, 157]	477	[32, 353]
50	[94, 95]	157	[71, 72]	264	[32, 308]	371	[49, 313]	478	[499, 134]
51	[96, 97]	158	[255, 256]	265	[157, 296]	372	[124, 122]	479	[170, 455]
52	[98, 99]	159	[257, 258]	266	[333, 32]	373	[146, 47]	480	[444, 174]
53	[100, 101]	160	[259, 260]	267	[318, 155]	374	[437, 311]	481	[434, 500]
54	[102, 103]	161	[261, 262]	268	[305, 157]	375	[301, 336]	482	[182, 181]
55	[104, 105]	162	[263, 208]	269	[153, 139]	376	[292, 341]	483	[349, 50]
56	[106, 107]	163	[16, 264]	270	[344, 7]	377	[111, 305]	484	[43, 353]
57	[108, 109]	164	[265, 234]	271	[345, 113]	378	[38, 40]	485	[495, 328]
58	[110, 108]	165	[266, 267]	272	[346, 322]	379	[52, 313]	486	[501, 431]
59	[111, 38]	166	[197, 268]	273	[347, 348]	380	[342, 309]	487	[502, 46]
60	[112, 113]	167	[199, 269]	274	[311, 315]	381	[115, 434]	488	[503, 350]
61	[114, 115]	168	[270, 245]	275	[58, 56]	382	[307, 176]	489	[456, 303]
62	[116, 117]	169	[271, 196]	276	[295, 37]	383	[343, 45]	490	[504, 220]
63	[118, 119]	170	[272, 273]	277	[349, 313]	384	[438, 123]	491	[505, 464]
64	[120, 121]	171	[274, 275]	278	[315, 148]	385	[176, 139]	492	[506, 256]

65	[122, 123]	172	[276, 271]	279	[46, 301]	386	[338, 156]	493	[507, 508]
66	[124, 12]	173	[277, 203]	280	[143, 169]	387	[181, 181]	494	[509, 210]
67	[125, 126]	174	[278, 279]	281	[339, 146]	388	[289, 352]	495	[510, 411]
68	[127, 128]	175	[250, 280]	282	[122, 350]	389	[340, 293]	496	[511, 512]
69	[129, 130]	176	[187, 90]	283	[51, 49]	390	[300, 46]	497	[513, 88]
70	[131, 132]	177	[281, 24]	284	[351, 352]	391	[12, 350]	498	[514, 239]
71	[131, 133]	178	[282, 283]	285	[353, 354]	392	[168, 302]	499	[515, 286]
72	[132, 134]	179	[284, 285]	286	[324, 133]	393	[296, 305]	500	[369, 516]
73	[133, 134]	180	[286, 217]	287	[135, 324]	394	[178, 126]	501	[517, 518]
74	[135, 131]	181	[221, 286]	288	[355, 225]	395	[439, 440]	502	[519, 358]
75	[133, 136]	182	[287, 221]	289	[356, 260]	396	[441, 442]	503	[520, 253]
76	[137, 137]	183	[288, 137]	290	[357, 29]	397	[443, 136]	504	[521, 325]
77	[125, 138]	184	[289, 290]	291	[358, 265]	398	[145, 352]	505	[522, 157]
78	[139, 140]	185	[291, 140]	292	[359, 360]	399	[441, 440]	506	[492, 303]
79	[141, 130]	186	[292, 293]	293	[361, 362]	400	[444, 445]	507	[451, 523]
80	[142, 143]	187	[294, 115]	294	[363, 17]	401	[308, 446]	508	[291, 175]
81	[144, 145]	188	[295, 296]	295	[364, 222]	402	[447, 131]	509	[522, 40]
82	[118, 146]	189	[297, 298]	296	[73, 225]	403	[40, 296]	510	[178, 138]
83	[147, 148]	190	[294, 299]	297	[365, 366]	404	[437, 165]	511	[524, 452]
84	[149, 150]	191	[33, 145]	298	[367, 368]	405	[337, 296]	512	[353, 446]
85	[151, 49]	192	[300, 10]	299	[368, 369]	406	[47, 167]	513	[39, 157]
86	[152, 153]	193	[301, 121]	300	[18, 249]	407	[56, 8]	514	[438, 350]
87	[154, 155]	194	[302, 303]	301	[249, 370]	408	[299, 434]	515	[525, 132]
88	[35, 156]	195	[304, 160]	302	[371, 372]	409	[447, 324]	516	[159, 169]
89	[157, 37]	196	[37, 305]	303	[237, 373]	410	[351, 290]	517	[524, 523]
90	[44, 158]	197	[179, 306]	304	[374, 375]	411	[320, 435]	518	[320, 500]
91	[159, 127]	198	[307, 153]	305	[222, 224]	412	[327, 448]	519	[490, 181]
92	[160, 161]	199	[144, 34]	306	[376, 257]	413	[449, 348]	520	[503, 123]
93	[162, 122]	200	[40, 37]	307	[377, 227]	414	[450, 165]	521	[526, 71]
94	[8, 163]	201	[43, 308]	308	[60, 378]	415	[344, 56]	522	[527, 218]
95	[10, 120]	202	[309, 161]	309	[379, 380]	416	[331, 10]	523	[375, 528]
96	[164, 165]	203	[304, 309]	310	[258, 381]	417	[433, 126]	524	[529, 372]
97	[166, 167]	204	[57, 310]	311	[382, 383]	418	[308, 354]	525	[530, 73]
98	[52, 50]	205	[311, 147]	312	[384, 195]	419	[145, 290]	526	[499, 136]
99	[168, 149]	206	[164, 311]	313	[97, 216]	420	[451, 452]	527	[521, 137]
100	[117, 126]	207	[312, 313]	314	[383, 385]	421	[169, 453]	528	[7, 57]
101	[169, 128]	208	[147, 314]	315	[386, 387]	422	[454, 455]	529	[498, 161]
102	[170, 171]	209	[165, 315]	316	[388, 101]	423	[456, 150]	530	[525, 133]
103	[172, 143]	210	[316, 317]	317	[389, 240]	424	[46, 120]		
104	[173, 174]	211	[180, 180]	318	[390, 278]	425	[457, 161]		
105	[139, 175]	212	[318, 319]	319	[391, 392]	426	[119, 166]		
106	[152, 176]	213	[299, 320]	320	[189, 393]	427	[297, 317]		

Output function $n \mapsto \tau(n)$:

n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$
0	0	76	1	152	1	228	1	304	1	380	0	456	1
1	0	77	0	153	0	229	0	305	1	381	1	457	1
2	0	78	0	154	1	230	1	306	0	382	0	458	1
3	0	79	1	155	0	231	1	307	0	383	1	459	1
4	0	80	1	156	1	232	0	308	0	384	0	460	0
5	0	81	1	157	0	233	1	309	0	385	1	461	0
6	1	82	0	158	1	234	0	310	1	386	1	462	1
7	0	83	0	159	1	235	0	311	0	387	0	463	1
8	0	84	0	160	1	236	1	312	0	388	0	464	0
9	0	85	1	161	1	237	1	313	1	389	1	465	0
10	0	86	1	162	1	238	1	314	1	390	0	466	0
11	0	87	1	163	0	239	1	315	1	391	1	467	1
12	1	88	0	164	0	240	1	316	0	392	0	468	0
13	1	89	0	165	1	241	1	317	1	393	1	469	1
14	1	90	0	166	1	242	0	318	0	394	0	470	1
15	0	91	1	167	1	243	0	319	1	395	1	471	1
16	0	92	1	168	0	244	1	320	1	396	0	472	0
17	0	93	1	169	1	245	0	321	0	397	0	473	0
18	0	94	0	170	1	246	1	322	1	398	1	474	0
19	0	95	0	171	0	247	0	323	0	399	0	475	0
20	0	96	0	172	0	248	1	324	1	400	0	476	0
21	0	97	1	173	1	249	1	325	0	401	0	477	0
22	0	98	0	174	1	250	1	326	0	402	1	478	1
23	0	99	0	175	1	251	1	327	1	403	1	479	1
24	0	100	1	176	1	252	0	328	0	404	1	480	0
25	1	101	1	177	1	253	1	329	0	405	1	481	0
26	0	102	1	178	0	254	1	330	1	406	0	482	0
27	1	103	0	179	1	255	1	331	1	407	1	483	1
28	1	104	1	180	1	256	0	332	1	408	0	484	1
29	1	105	0	181	0	257	1	333	1	409	1	485	0
30	0	106	1	182	0	258	1	334	0	410	1	486	0
31	0	107	1	183	0	259	1	335	1	411	1	487	1
32	0	108	1	184	0	260	1	336	1	412	1	488	1
33	0	109	1	185	1	261	1	337	1	413	0	489	1
34	0	110	0	186	0	262	0	338	1	414	0	490	1
35	0	111	0	187	1	263	1	339	1	415	0	491	1
36	0	112	0	188	0	264	0	340	1	416	1	492	0
37	0	113	0	189	1	265	0	341	0	417	1	493	1
38	0	114	0	190	1	266	1	342	0	418	0	494	1
39	0	115	1	191	0	267	0	343	1	419	1	495	0
40	1	116	0	192	0	268	1	344	0	420	1	496	0
41	0	117	1	193	1	269	0	345	1	421	1	497	0
42	1	118	0	194	1	270	0	346	1	422	0	498	0

43	1	119	0	195	1	271	1	347	1	423	1	499	1
44	0	120	0	196	0	272	1	348	0	424	1	500	1
45	0	121	0	197	1	273	1	349	1	425	1	501	0
46	1	122	0	198	0	274	0	350	0	426	0	502	1
47	0	123	1	199	1	275	0	351	1	427	1	503	1
48	0	124	0	200	1	276	0	352	0	428	1	504	1
49	1	125	0	201	1	277	1	353	1	429	1	505	1
50	0	126	0	202	0	278	1	354	0	430	1	506	0
51	0	127	0	203	1	279	1	355	0	431	0	507	1
52	0	128	0	204	1	280	0	356	0	432	1	508	1
53	1	129	0	205	0	281	1	357	1	433	1	509	1
54	1	130	1	206	0	282	0	358	1	434	0	510	0
55	1	131	0	207	0	283	0	359	0	435	0	511	0
56	1	132	0	208	0	284	1	360	1	436	0	512	1
57	1	133	1	209	1	285	1	361	1	437	1	513	0
58	0	134	1	210	0	286	1	362	1	438	0	514	0
59	0	135	0	211	1	287	0	363	1	439	1	515	1
60	0	136	0	212	0	288	0	364	0	440	0	516	1
61	0	137	1	213	0	289	0	365	1	441	0	517	0
62	0	138	1	214	0	290	1	366	0	442	1	518	1
63	0	139	0	215	1	291	1	367	0	443	0	519	1
64	0	140	0	216	1	292	0	368	0	444	0	520	1
65	0	141	1	217	0	293	1	369	1	445	0	521	1
66	0	142	1	218	1	294	1	370	0	446	1	522	1
67	0	143	0	219	0	295	0	371	1	447	1	523	1
68	0	144	1	220	1	296	1	372	0	448	1	524	0
69	0	145	1	221	0	297	1	373	1	449	0	525	1
70	0	146	1	222	1	298	0	374	1	450	0	526	1
71	0	147	0	223	0	299	0	375	1	451	1	527	1
72	0	148	0	224	0	300	0	376	0	452	0	528	0
73	1	149	0	225	1	301	1	377	0	453	1	529	0
74	0	150	0	226	0	302	1	378	0	454	0	530	1
75	1	151	1	227	0	303	1	379	0	455	1		

UNIVERSITÉ DE STRASBOURG, CNRS, IRMA UMR 7501, F-67000 STRASBOURG, FRANCE
Email address: `guoniu.han@unistra.fr`

SCHOOL OF MATHEMATICS AND STATISTICS, HUAZHONG UNIVERSITY OF SCIENCE AND TECHNOLOGY, WUHAN, PR CHINA
Email address: `huyining@hust.edu.cn`