

PROOF OF THE CONTINUED FRACTION CONJECTURE FOR

$$(a, b) = (z^3 + z, z^4 + 1)$$

GUO-NIU HAN* AND YINING HU*

ABSTRACT. The continued fraction conjecture claims that the continued fraction defined by the (a, b) -Thue-Morse sequence is algebraic over $\mathbb{F}_2(x)$ for all pairs of distinct elements (a, b) in $\mathbb{F}_2[x] \setminus \mathbb{F}_2$. In this paper we prove the conjecture for $(a, b) = (z^3 + z, z^4 + 1)$.

1. INTRODUCTION

Let $\mathbb{F}_2$ be the finite field of cardinality 2. Given a sequence of polynomials $\mathbf{a} = (a_0(z), a_1(z), a_2(z), \dots)$ of elements from $\mathbb{F}_2[z] \setminus \mathbb{F}_2$, we define the infinite continued fraction

$$(1.1) \quad \text{CF}(\mathbf{a}(z)) = \frac{1}{a_0(z) + \frac{1}{a_1(z) + \frac{1}{a_2(z) + \frac{1}{\ddots}}}}$$

as the limit of the finite continued fractions

$$(1.2) \quad \text{CF}_n(\mathbf{a}(z)) = \frac{1}{a_0(z) + \frac{1}{a_1(z) + \frac{1}{\ddots + \frac{1}{a_n(z)}}}} \in \mathbb{F}_2((1/z)).$$

Conversely, each power series in $\mathbb{F}_2[[1/z]]$ without constant term

$$(1.3) \quad c_{-1}z^{-1} + c_{-2}z^{-2} + c_{-3}z^{-3} + \dots$$

admits a continued fraction expansion of form (1.1).

Let a and b be two distinct elements from $\mathbb{F}_2[z] \setminus \mathbb{F}_2$. We consider the continued fraction (1.1) associated with the (a, b) -Thue-Morse sequence

$$\mathbf{t} = (t_0(z), t_1(z), t_2(z), \dots) = (a, b, b, a, b, a, a, b, \dots)$$

that we shall denote

$$(1.4) \quad \text{CF}(a, b) := \text{CF}(\mathbf{t}).$$

Date: 9/04/2020 21:40:18 .

2010 Mathematics Subject Classification. 11B85, 11J70, 11B50, 11Y65, 05A15.

Key words and phrases. automatic sequence, continued fraction, Thue-Morse sequence.

* This paper was automatically generated from a computer program developed by the authors, with 5.9 minutes CPU time used.

Theorem 1.1. *When $(a, b) = (z^3 + z, z^4 + 1)$, the two power series $\text{CF}(a, b)$ and $\text{CF}(b, a)$ are algebraic over $\mathbb{F}_2(z)$, with minimal polynomials of the form*

$$p_4(z)y^4 + p_3(z)y^3 + p_2(z)y^2 + p_1(z)y + p_0(z) = 0$$

where, for $\text{CF}(a, b)$,

$$\begin{aligned} p_0(z) &= z^{13} + z^{11} + z^8 + z^7 + z^6 + z^5 + z^3 + z^2 + z + 1, \\ p_1(z) &= z^{17} + z^{16} + z^{15} + z^{13} + z^{12} + z^{10} + z^9 + z^3 + z^2 + z, \\ p_2(z) &= z^{20} + z^{14} + z^{12} + z^{10} + z^6 + z^2, \\ p_3(z) &= z^{17} + z^{16} + z^{15} + z^{13} + z^{12} + z^{10} + z^9 + z^3 + z^2 + z, \\ p_4(z) &= z^{14} + z^{13} + z^{12} + z^{11} + z^{10} + z^7 + z^5 + z^3 + z, \end{aligned}$$

and for $\text{CF}(b, a)$,

$$\begin{aligned} p_0(z) &= z^{13} + z^{11} + z^7 + z^6 + z^5 + z^4 + z^3 + z^2 + z, \\ p_1(z) &= z^{17} + z^{16} + z^{15} + z^{13} + z^{12} + z^{10} + z^9 + z^3 + z^2 + z, \\ p_2(z) &= z^{20} + z^{14} + z^{12} + z^{10} + z^6 + z^2, \\ p_3(z) &= z^{17} + z^{16} + z^{15} + z^{13} + z^{12} + z^{10} + z^9 + z^3 + z^2 + z, \\ p_4(z) &= z^{14} + z^{13} + z^{12} + z^{11} + z^{10} + z^8 + z^7 + z^5 + z^4 + z^3 + z + 1. \end{aligned}$$

2. PROOF

Define the sequences of polynomials $(P_n(z))_{n \geq 0}$ and $(Q_n(z))_{n \geq 0}$ by

$$\begin{pmatrix} P_n(z) & Q_n(z) \\ P_{n-1}(z) & Q_{n-1}(z) \end{pmatrix} := \begin{pmatrix} t_n(z) & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} t_{n-1}(z) & 1 \\ 1 & 0 \end{pmatrix} \cdots \begin{pmatrix} t_0(z) & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}.$$

By the basic properties of continued fractions, we have

$$(2.1) \quad \text{CF}_n(a, b) = \frac{P_n(z)}{Q_n(z)} \in \mathbb{F}_2((1/z)).$$

As formula (2.1) is not efficient for calculating $\text{CF}_n(a, b)$, a fast method is to be derived and described as follows. Define

$$\begin{aligned} M_n(x) &= x^{\deg(t_{2^n-1})} \begin{pmatrix} t_{2^n-1}(1/x) & 1 \\ 1 & 0 \end{pmatrix} \times x^{\deg(t_{2^n-2})} \begin{pmatrix} t_{2^n-2}(1/x) & 1 \\ 1 & 0 \end{pmatrix} \times \\ &\quad \cdots \times x^{\deg(t_0)} \begin{pmatrix} t_0(1/x) & 1 \\ 1 & 0 \end{pmatrix}, \\ W_n(x) &= x^{\deg(\bar{t}_{2^n-1})} \begin{pmatrix} \bar{t}_{2^n-1}(1/x) & 1 \\ 1 & 0 \end{pmatrix} \times x^{\deg(\bar{t}_{2^n-2})} \begin{pmatrix} \bar{t}_{2^n-2}(1/x) & 1 \\ 1 & 0 \end{pmatrix} \times \\ &\quad \cdots \times x^{\deg(\bar{t}_0)} \begin{pmatrix} \bar{t}_0(1/x) & 1 \\ 1 & 0 \end{pmatrix}, \end{aligned}$$

where $\bar{t}$ is the (b, a) -Thue-Morse sequence. By the properties of the Thue-Morse sequence, we have for $n \geq 0$

$$\begin{aligned} M_{n+1}(x) &= W_n(x) \cdot M_n(x), \\ W_{n+1}(x) &= M_n(x) \cdot W_n(x). \end{aligned}$$

Let $x := 1/z$. For each non-zero polynomial $P(z)$, define $\tilde{P}(x)$ to be $P(1/x)$. Then,

$$\text{CF}_n(a, b) = \frac{P_n(z)}{Q_n(z)} = \frac{\tilde{P}_n(x)}{\tilde{Q}_n(x)} \in \mathbb{F}_2((x)) = \mathbb{F}_2((1/z)).$$

Comparing the definition of $M_n(x)$ with the definition of $P_n(x)$ and $Q_n(x)$, we see that

$$\begin{aligned} M_n(x)_{0,1} &= x^{d_n} \tilde{P}_{2^n-1}(x), \\ M_n(x)_{0,0} &= x^{d_n} \tilde{Q}_{2^n-1}(x), \end{aligned}$$

for some positive integer d_n . Hence,

$$(2.2) \quad \text{CF}_{2^{2^n}-1}(a, b) = \frac{\tilde{P}_{2^{2^n}-1}(x)}{\tilde{Q}_{2^{2^n}-1}(x)} = \frac{M_{2^n}(x)_{0,1}}{M_{2^n}(x)_{0,0}}.$$

Therefore, by the convergence theorem of continued fraction, the algebraicity of $\text{CF}(a, b)$ will be established if it is shown that both $M_{2^n}(x)_{0,1}$ and $M_{2^n}(x)_{0,0}$ converge to algebraic series in $\mathbb{F}_2[[x]]$.

Actually, we will prove that for all $0 \leq i, j \leq 1$ the four sequences $(M_{2^n}(x)_{i,j})_n$, $(M_{2^{n+1}}(x)_{i,j})_n$, $(W_{2^n}(x)_{i,j})_n$, and $(W_{2^{n+1}}(x)_{i,j})_n$ converge to algebraic series in $\mathbb{F}_2[[x]]$. For this purpose, we define four 2×2 matrices M^e, M^o, W^e, W^o as follows: For each $T \in \{M^e, M^o, W^e, W^o\}$ and $0 \leq i, j \leq 1$, $T_{i,j}$ is defined to be the unique solution in $\mathbb{F}_2[[x]]$ of the polynomial $\phi(T, i, j)$ under certain initial conditions; the polynomials $\phi(T, i, j)$ and initial conditions are given in Section 3. We will prove that these four matrices, whose components are algebraic by definition, are the limits of $(M_{2^n}(x))_n$, $(M_{2^{n+1}}(x))_n$, $(W_{2^n}(x))_n$, and $(W_{2^{n+1}}(x))_n$.

Let us explain how the polynomials $\phi(T, i, j)$ and initial conditions are found, and why the solutions exist and are unique. For $0 \leq i, j \leq 1$, the coefficients of the polynomial $\phi(M^e, i, j)$ (resp. $\phi(M^o, i, j)$, $\phi(W^e, i, j)$, and $\phi(W^o, i, j)$) are the Padé-Hermite approximants of type

$$(175, 175, 175, 175, 175)$$

of the vector

$$(1, T^3, T^6, T^9, T^{12}),$$

where $T = M_{12,i,j}$ (resp. $M_{11,i,j}$, $W_{12,i,j}$, and $W_{11,i,j}$), found by the Derksen algorithm. We take the first 8 terms of $M_{12,i,j}$ (resp. $M_{11,i,j}$, $W_{12,i,j}$, and $W_{11,i,j}$) as initial conditions for $\phi(M^e, i, j)$ (resp. $\phi(M^o, i, j)$, $\phi(W^e, i, j)$, and $\phi(W^o, i, j)$). The following fact will be used to ensure that the solution exists and is unique: let $P(x, y) \in \mathbb{F}_2[x, y]$ and for each series $f(x) = \sum_{n=0}^{\infty} a_n x^n \in \mathbb{F}_2[[x]]$ denote the partial sum $\sum_{j=0}^{n-1} a_j x^j$ by $f_n(x)$ for $n \geq 0$. If for some $n \geq 0$ and $a_0, a_1, \dots, a_{n-1} \in \mathbb{F}_2$ $P(x, \sum_{j=0}^{n-1} a_j x^j) = O(x^n)$ and $Q(x, y) := P(x, \sum_{j=0}^{n-1} a_j x^j + x^n y)$ can be written as $x^m \sum_{j=0}^{\infty} q_j(x) y^j$ where $q_j(x)$ are polynomials for $j \geq 0$, $q_1(0) = 1$, and $q_j(0) = 0$ for $j > 1$, then there exists a unique solution $f(x) \in \mathbb{F}_2[[x]]$ of $P(x, f(x)) = 0$ that satisfies the initial condition $f_n(x) = \sum_{j=0}^{n-1} a_j x^j$.

We state two lemmas concerning the four matrices M^e, M^o, W^e, W^o . The first one is about relations between them; the second, about the structure of the each matrix.

Lemma 2.1. *We have*

$$(2.3) \quad M^e = W^o \cdot M^o,$$

$$(2.4) \quad M^o = W^e \cdot M^e,$$

$$(2.5) \quad W^e = M^o \cdot W^o,$$

$$(2.6) \quad W^o = M^e \cdot W^e.$$

Proof. We give the proof of the identity

$$M_{0,0}^e = W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o,$$

the proofs of the others being similar.

First, we compute the minimal polynomials of $W_{0,0}^o M_{0,0}^o$ and $W_{0,1}^o M_{1,0}^o$. We know that

$$P(x, y) = \text{Res}_z (\phi(W^o, 0, 0)(x, z), z^{12} \cdot \phi(M^o, 0, 0)(x, y/z))$$

is an annihilating polynomial of $W_{0,0}^o M_{0,0}^o$. We use Padé-Hermite approximation to find a candidate for the minimal polynomial of $W_{0,0}^o M_{0,0}^o$, that will be called $\phi_0(x, y)$. To prove that $\phi_0(x, y)$ is indeed the minimal polynomial, it suffices to prove that it is an irreducible factor of $P(x, y)$ of multiplicity m and that $Q(x, y) := P(x, y)/\phi_0(x, y)^m$ is not an annihilating polynomial of $W_{0,0}^o M_{0,0}^o$. We verify the first point directly. For the second point, we truncate $W_{0,0}^o M_{0,0}^o$ to order 630 and substitute it for y in $Q(x, y)$. We get a series of valuation less than 630, which proves that $Q(x, y)$ is not an annihilating polynomial of $W_{0,0}^o M_{0,0}^o$. We find the minimal polynomial $\phi_1(x, y)$ of $W_{0,1}^o M_{1,0}^o$ in a similar way.

Now we prove that $\phi(M^e, 0, 0)$ is the minimal polynomial of $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$. We know that

$$S(x, y) = \text{Res}_z (\phi_0(x, z), \phi_1(x, y + z))$$

is an annihilating of $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$. We verify that $\phi(M^e, 0, 0)$ is an irreducible factor of $S(x, y)$ of multiplicity μ , and that $Q(x, y) := S(x, y)/\phi(M^e, 0, 0)^\mu$ is not an annihilating polynomial of $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$. To see the last point, we truncate $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$ to order 770 and substitute it for y in $Q(x, y)$. We get a series of valuation less than 770, and therefore $Q(x, y)$ is not an annihilating polynomial of $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$.

Finally, the first 16 terms of $M_{0,0}^e$ and $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$ coincide. As these first terms determine a unique solution of $\phi(M^e, 0, 0)$, we know that the two series are one and the same. $\square$

Define

$$R^e = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \quad \text{and} \quad R^o = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}.$$

Lemma 2.2. *For all integers $k \geq 2$ and $u = 2^{2k-1}$, the following identities hold.*

$$\begin{aligned} M^e[:14u] &= M^e[:7u] + x^{2u} M^e[:5u] + x^{4u} M^e[:3u] + x^{2u} M^e[:7u] + x^{4u} M^e[:5u] \\ &\quad + x^{6u} M^e[:3u] + x^{6u} M^e[:7u] + x^{8u} M^e[:5u] + x^{10u} M^e[:3u] \\ &\quad + x^{8u} M^e[:6u] + x^{12u} M^e[:2u] + (x^{7u} + x^{9u} + x^{13u}) R^e, \\ W^e[:14u] &= W^e[:7u] + x^{2u} W^e[:5u] + x^{4u} W^e[:3u] + x^{2u} W^e[:7u] + x^{4u} W^e[:5u] \\ &\quad + x^{6u} W^e[:3u] + x^{6u} W^e[:7u] + x^{8u} W^e[:5u] + x^{10u} W^e[:3u] \\ &\quad + x^{8u} W^e[:6u] + x^{12u} W^e[:2u] + (x^{7u} + x^{9u} + x^{13u}) R^e. \end{aligned}$$

For all integers $k \geq 2$ and $u = 2^{2k}$, the following identities hold.

$$\begin{aligned}
M^o[:14u] &= M^o[:7u] + x^{2u}M^o[:5u] + x^{4u}M^o[:3u] + x^{2u}M^o[:7u] + x^{4u}M^o[:5u] \\
&\quad + x^{6u}M^o[:3u] + x^{6u}M^o[:7u] + x^{8u}M^o[:5u] + x^{10u}M^o[:3u] \\
&\quad + x^{8u}M^o[:6u] + x^{12u}M^o[:2u] + (x^{7u} + x^{9u} + x^{13u})R^o, \\
W^o[:14u] &= W^o[:7u] + x^{2u}W^o[:5u] + x^{4u}W^o[:3u] + x^{2u}W^o[:7u] + x^{4u}W^o[:5u] \\
&\quad + x^{6u}W^o[:3u] + x^{6u}W^o[:7u] + x^{8u}W^o[:5u] + x^{10u}W^o[:3u] \\
&\quad + x^{8u}W^o[:6u] + x^{12u}W^o[:2u] + (x^{7u} + x^{9u} + x^{13u})R^o.
\end{aligned}$$

Proof. To prove Lemma 2.2 we first construct an automaton for each sequence concerned, and then transform the conditions on infinitely many k 's into finitely many conditions on the states of the automaton. In the following, we will prove that for $T = M_{0,0}^o$, for all integer $k \geq 2$ and $u = 2^{2k}$,

$$\begin{aligned}
T[:14u] &= T[:7u] + x^{2u}T[:5u] + x^{4u}T[:3u] + x^{2u}T[:7u] + x^{4u}T[:5u] + x^{6u}T[:3u] \\
&\quad + x^{6u}T[:7u] + x^{8u}T[:5u] + x^{10u}T[:3u] + x^{8u}T[:6u] + x^{12u}T[:2u].
\end{aligned}$$

The proof of the other 15 cases are similar. We break down the above identity into 7 parts:

$$\begin{aligned}
0 &= x^{4u}T[3u:4u] + x^{2u}T[5u:6u] + T[7u:8u], \\
0 &= x^{4u}T[4u:5u] + x^{2u}T[6u:7u] + T[8u:9u], \\
0 &= x^{6u}T[3u:4u] + T[9u:10u], \\
0 &= x^{10u}T[:u] + x^{6u}T[4u:5u] + T[10u:11u], \\
0 &= x^{10u}T[u:2u] + x^{6u}T[5u:6u] + T[11u:12u], \\
0 &= x^{12u}T[:u] + x^{10u}T[2u:3u] + x^{6u}T[6u:7u] + T[12u:13u], \\
0 &= x^{12u}T[u:2u] + x^{8u}T[5u:6u] + T[13u:14u],
\end{aligned}$$

which can be rewritten as

$$\begin{aligned}
(2.7) \quad 0 &= T[[11w]_2] + T[[101w]_2] + T[[111w]_2], \\
(2.8) \quad 0 &= T[[100w]_2] + T[[110w]_2] + T[[1000w]_2], \\
(2.9) \quad 0 &= T[[11w]_2] + T[[1001w]_2], \\
(2.10) \quad 0 &= T[[w]_2] + T[[100w]_2] + T[[1010w]_2], \\
(2.11) \quad 0 &= T[[1w]_2] + T[[101w]_2] + T[[1011w]_2], \\
(2.12) \quad 0 &= T[[w]_2] + T[[10w]_2] + T[[110w]_2] + T[[1100w]_2], \\
(2.13) \quad 0 &= T[[1w]_2] + T[[101w]_2] + T[[1101w]_2],
\end{aligned}$$

for all binary word w of length $2k$ and $w \neq 0^{2k}$, and

$$\begin{aligned}
(2.14) \quad 0 &= T[[11w]_2] + T[[101w]_2] + T[[111w]_2], \\
(2.15) \quad 0 &= T[[100w]_2] + T[[110w]_2] + T[[1000w]_2], \\
(2.16) \quad 0 &= T[[11w]_2] + T[[1001w]_2], \\
(2.17) \quad 0 &= T[[w]_2] + T[[100w]_2] + T[[1010w]_2], \\
(2.18) \quad 0 &= T[[1w]_2] + T[[101w]_2] + T[[1011w]_2], \\
(2.19) \quad 0 &= T[[w]_2] + T[[10w]_2] + T[[110w]_2] + T[[1100w]_2],
\end{aligned}$$

$$(2.20) \quad 0 = T[[1w]_2] + T[[101w]_2] + T[[1101w]_2],$$

for $w = 0^{2k}$.

First we calculate an 2-automaton that generates T from its minimal polynomial and its first terms. This automaton has 188 states; its transition function and output function can be found in the annex. Let $A(s, w)$ denote the state reached after reading w from right to left starting from the state s , and τ the output function. Define

$$E_{2k} = \{A(i, w) : |w| = 2k, w \neq 0^{2k}\}.$$

Identities (2.7) through (2.20) can be written as

$$(2.21) \quad 0 = \tau(A(s, 11)) + \tau(A(s, 101)) + \tau(A(s, 111)),$$

$$(2.22) \quad 0 = \tau(A(s, 100)) + \tau(A(s, 110)) + \tau(A(s, 1000)),$$

$$(2.23) \quad 0 = \tau(A(s, 11)) + \tau(A(s, 1001)),$$

$$(2.24) \quad 0 = \tau(A(s, \epsilon)) + \tau(A(s, 100)) + \tau(A(s, 1010)),$$

$$(2.25) \quad 0 = \tau(A(s, 1)) + \tau(A(s, 101)) + \tau(A(s, 1011)),$$

$$(2.26) \quad 0 = \tau(A(s, \epsilon)) + \tau(A(s, 10)) + \tau(A(s, 110)) + \tau(A(s, 1100)),$$

$$(2.27) \quad 0 = \tau(A(s, 1)) + \tau(A(s, 101)) + \tau(A(s, 1101)),$$

for all $s \in E_{2k}$, and

$$(2.28) \quad 0 = \tau(A(s, 11)) + \tau(A(s, 101)) + \tau(A(s, 111)),$$

$$(2.29) \quad 0 = \tau(A(s, 100)) + \tau(A(s, 110)) + \tau(A(s, 1000)),$$

$$(2.30) \quad 0 = \tau(A(s, 11)) + \tau(A(s, 1001)),$$

$$(2.31) \quad 0 = \tau(A(s, \epsilon)) + \tau(A(s, 100)) + \tau(A(s, 1010)),$$

$$(2.32) \quad 0 = \tau(A(s, 1)) + \tau(A(s, 101)) + \tau(A(s, 1011)),$$

$$(2.33) \quad 0 = \tau(A(s, \epsilon)) + \tau(A(s, 10)) + \tau(A(s, 110)) + \tau(A(s, 1100)),$$

$$(2.34) \quad 0 = \tau(A(s, 1)) + \tau(A(s, 101)) + \tau(A(s, 1101)),$$

for $s = A(i, 0^{2k})$. We find that $(A(i, 0^{32}), E_{32}) = (A(i, 0^{20}), E_{20})$, so that we only have to verify that identities (2.21) through (2.34) hold for $2 \leq k \leq 16$, which turns out to be true. $\square$

In the following lemma, we express M_{2k} , M_{2k+1} , W_{2k} , and W_{2k+1} in terms of M^e , M^o , W^e , and W^o .

Lemma 2.3. *For all integer $k \geq 2$, and $u = 2^{2k-1}$,*

$$M_{2k} = M^e[:7u] + x^{2u}M^e[:5u] + x^{4u}M^e[:3u] + x^{7u}R^e,$$

$$W_{2k} = W^e[:7u] + x^{2u}W^e[:5u] + x^{4u}W^e[:3u] + x^{7u}R^e.$$

For all integer $k \geq 2$, and $u = 2^{2k}$,

$$M_{2k+1} = M^o[:7u] + x^{2u}M^o[:5u] + x^{4u}M^o[:3u] + x^{7u}R^o,$$

$$W_{2k+1} = W^o[:7u] + x^{2u}W^o[:5u] + x^{4u}W^o[:3u] + x^{7u}R^o.$$

Proof. We call the four identities in Lemma 2.3 also by the name M_{2k} , W_{2k} , M_{2k+1} , and W_{2k+1} . For $n = 2$, the identities can be verified directly. For $n \geq 2$, we claim that

$$M_{2k} \wedge W_{2k} \Rightarrow M_{2k+1} \wedge W_{2k+1},$$

$$M_{2k+1} \wedge W_{2k+1} \Rightarrow M_{2k+2} \wedge W_{2k+2}.$$

We give the proof of

$$M_{2k} \wedge W_{2k} \Rightarrow M_{2k+1},$$

the other ones being similar. Set $u = 2^{2k}$ and $v = 2^{2k-1}$. By definition and induction hypothesis, the left side of identity M_{2k+1} is equal to

$$(2.35) \quad \begin{aligned} W_{2k} M_{2k} = & (W^e[:7v] + x^{2v} W^e[:5v] + x^{4v} W^e[:3v] + x^{7u} R^e) \times (M^e[:7v] \\ & + x^{2v} M^e[:5v] + x^{4v} M^e[:3v] + x^{7u} R^e); \end{aligned}$$

Call this expression *lhs*. Note that both sides of identity M_{2k+1} have the same term of highest degree $x^{14v} R^o$. Therefore we only need to prove that their difference is $O(x^{14v})$. Using Lemma 2.1 it can be seen that the right side of identity M_{2k+1} is congruent, modulo x^{14v} , to

$$W^e[:14v] M^e[:14v] + x^{4v} W^e[:10v] M^e[:10v] + x^{8v} W^e[:6v] M^e[:6v].$$

For all $n \leq 14$, replace the occurrences of $W^e[:n \cdot v]$ and $M^e[:n \cdot v]$ in the above expression by the reduction modulo $x^{n \cdot v}$ of the right side of the corresponding identity in Lemma 2.2 and get a new expression, which we call *rhs*. Define

$$\begin{aligned} X &:= x^v, \\ a_n &:= W^e[n \cdot v : (n+1) \cdot v] / X^n, \\ b_n &:= M^e[n \cdot v : (n+1) \cdot v] / X^n, \\ c &:= R^e. \end{aligned}$$

Using the notation introduced above, we can represent the expressions *lhs* (2.35) and *rhs* as polynomials in $\mathbb{F}_2[a_1, \dots, a_{14}, b_1, \dots, b_{14}, c][X]$. Note that it is not a problem that a_j commutes with b_k while W^e does not commute with M^e , because in the expressions concerned, the products of W^e -terms and M^e -terms are always in the same order. We let the computer do the simplification and check that the difference between these two polynomials is indeed $O(X^{14})$, which completes the proof. $\square$

Proof of Theorem 1.1. We prove the theorem for $\text{CF}(a, b)$; for $\text{CF}(b, a)$, the proof is similar. By Lemma 2.3, we have For all $0 \leq j, k \leq 1$,

$$\begin{aligned} \lim_{n \rightarrow \infty} M_{2n,j,k} &= M_{j,k}^e, \\ \lim_{n \rightarrow \infty} M_{2n+1,j,k} &= M_{j,k}^o, \\ \lim_{n \rightarrow \infty} W_{2n,j,k} &= W_{j,k}^e, \\ \lim_{n \rightarrow \infty} W_{2n+1,j,k} &= W_{j,k}^o. \end{aligned}$$

Let $z = 1/x$. By the convergence theorem and identity (2.2),

$$\text{CF}(a, b) = \frac{M_{0,1}^e(x)}{M_{0,0}^e(x)}.$$

By definition, $\phi(M^e, 0, 1)$ and $\phi(M^e, 0, 0)$ are minimal polynomials of $M_{0,1}^e$ and $M_{0,0}^e$. Therefore

$$P(x, y) = \text{Res}_t (\phi(M^e, 0, 1)(x, t), y^{12} \phi(M^e, 0, 1)(x, t/y))$$

is an annihilating polynomial of $f(x) = M_{0,1}^e / M_{0,0}^e$.

Define

$$Q(x, y) = q_4(x)y^4 + q_3(x)y^3 + q_2(x)y^2 + q_1(x)y + q_0(x),$$

where

$$\begin{aligned} q_0(x) &= x^{20} + x^{19} + x^{18} + x^{17} + x^{15} + x^{14} + x^{13} + x^{12} + x^9 + x^7, \\ q_1(x) &= x^{19} + x^{18} + x^{17} + x^{11} + x^{10} + x^8 + x^7 + x^5 + x^4 + x^3, \\ q_2(x) &= x^{18} + x^{14} + x^{10} + x^8 + x^6 + 1, \\ q_3(x) &= x^{19} + x^{18} + x^{17} + x^{11} + x^{10} + x^8 + x^7 + x^5 + x^4 + x^3, \\ q_4(x) &= x^{19} + x^{17} + x^{15} + x^{13} + x^{10} + x^9 + x^8 + x^7 + x^6. \end{aligned}$$

The polynomial $Q(x, y)$ is the candidate for the minimal polynomial of $f(x)$ found by Padé-Hermite approximation. To prove that it is indeed the minimal polynomial of $f(x)$, we only need to prove that it is an irreducible factor of $P(x, y)$ of multiplicity m and $R(x, y) := P(x, y)/Q(x, y)^m$ is not an annihilating polynomial of $f(x)$. We verify the first point directly. For the second point, we find that when we truncate $f(x)$ to order 216, and substitute it for y in $R(z, y)$, we get a series with valuation smaller than 216, which proves that $R(z, y)$ is not an annihilating polynomial of $f(x)$. Finally, $z^{12}Q(1/z, y)$ is the minimal polynomial of $\text{CF}(a, b) = f(1/z)$. $\square$

3. ANNEXE

All 16 polynomials are of the form

$$p_0(x) + p_3(x)y^3 + p_6(x)y^6 + p_9(x)y^9 + p_{12}(x)y^{12}.$$

The coefficients $p_j(x)$ and the 16 initial terms to determine the solutions uniquely are given below.

For $\phi(M^e, 0, 0)$:

$$\begin{aligned} p_0(x) &= x^{114} + x^{110} + x^{96} + x^{94} + x^{92} + x^{90} + x^{86} + x^{84} + x^{78} + x^{76} + x^{74} \\ &\quad + x^{68} + x^{66} + x^{64} + x^{60} + x^{52} + x^{50} + x^{48} + x^{44} + x^{38} + x^{36}, \\ p_3(x) &= x^{108} + x^{106} + x^{104} + x^{102} + x^{98} + x^{94} + x^{90} + x^{86} + x^{82} + x^{80} + x^{78} \\ &\quad + x^{74} + x^{72} + x^{70} + x^{68} + x^{64} + x^{52} + x^{50} + x^{46} + x^{44} + x^{40} + x^{36} \\ &\quad + x^{34} + x^{30} + x^{26} + x^{24} + x^{22} + x^{20}, \\ p_6(x) &= x^{110} + x^{108} + x^{106} + x^{104} + x^{102} + x^{98} + x^{96} + x^{94} + x^{92} + x^{90} + x^{86} \\ &\quad + x^{84} + x^{80} + x^{76} + x^{74} + x^{66} + x^{64} + x^{62} + x^{60} + x^{54} + x^{52} + x^{46} \\ &\quad + x^{44} + x^{42} + x^{40} + x^{26} + x^{24} + x^{12} + x^{10} + x^8 + x^6 + 1, \\ p_9(x) &= x^{112} + x^{110} + x^{106} + x^{100} + x^{90} + x^{86} + x^{82} + x^{80} + x^{78} + x^{74} + x^{72} \\ &\quad + x^{64} + x^{62} + x^{60} + x^{58} + x^{56} + x^{50} + x^{46} + x^{44} + x^{42} + x^{38} + x^{36} \\ &\quad + x^{32} + x^{30} + x^{28} + x^{24} + x^{22} + x^{14} + x^{12} + x^{10} + x^8 + x^6, \\ p_{12}(x) &= x^{112} + x^{48} + x^{32} + 1, \end{aligned}$$

and the initial terms are $[1, 0, 0, 0, 0, 0, 0, 0, 1, 0, 0, 0, 1, 0, 1, 0]$.

For $\phi(M^e, 0, 1)$:

$$\begin{aligned} p_0(x) &= x^{111} + x^{110} + x^{107} + x^{105} + x^{100} + x^{98} + x^{97} + x^{93} + x^{92} + x^{89} + x^{86} \\ &\quad + x^{84} + x^{83} + x^{81} + x^{80} + x^{78} + x^{76} + x^{73} + x^{72} + x^{71} + x^{70} + x^{66} \end{aligned}$$

$$\begin{aligned}
& + x^{65} + x^{63} + x^{62} + x^{61} + x^{59} + x^{57} + x^{54} + x^{52} + x^{51} + x^{49} + x^{48} \\
& + x^{47} + x^{45} + x^{44} + x^{43} + x^{40} + x^{39}, \\
p_3(x) &= x^{109} + x^{108} + x^{106} + x^{105} + x^{101} + x^{100} + x^{99} + x^{97} + x^{96} + x^{94} \\
& + x^{93} + x^{91} + x^{90} + x^{89} + x^{83} + x^{82} + x^{80} + x^{78} + x^{77} + x^{73} + x^{72} \\
& + x^{70} + x^{68} + x^{67} + x^{45} + x^{44} + x^{42} + x^{41} + x^{37} + x^{36} + x^{35} + x^{33} \\
& + x^{32} + x^{30} + x^{28} + x^{27}, \\
p_6(x) &= x^{108} + x^{104} + x^{102} + x^{98} + x^{96} + x^{92} + x^{90} + x^{88} + x^{86} + x^{84} + x^{82} \\
& + x^{80} + x^{78} + x^{76} + x^{74} + x^{72} + x^{70} + x^{68} + x^{66} + x^{62} + x^{60} + x^{54} \\
& + x^{52} + x^{48} + x^{46} + x^{44} + x^{42} + x^{38} + x^{36} + x^{32} + x^{30} + x^{28} + x^{26} \\
& + x^{24} + x^{22} + x^{18}, \\
p_9(x) &= x^{107} + x^{106} + x^{105} + x^{101} + x^{100} + x^{99} + x^{97} + x^{96} + x^{95} + x^{93} + x^{92} \\
& + x^{91} + x^{89} + x^{88} + x^{87} + x^{83} + x^{82} + x^{80} + x^{79} + x^{77} + x^{76} + x^{74} \\
& + x^{73} + x^{71} + x^{70} + x^{68} + x^{67} + x^{65} + x^{64} + x^{63} + x^{53} + x^{52} + x^{51} \\
& + x^{43} + x^{42} + x^{41} + x^{39} + x^{38} + x^{37} + x^{35} + x^{34} + x^{32} + x^{30} + x^{29} \\
& + x^{25} + x^{24} + x^{23} + x^{21} + x^{20} + x^{19} + x^{15} + x^{14} + x^{13} + x^{11} + x^{10} \\
& + x^9, \\
p_{12}(x) &= x^{106} + x^{104} + x^{98} + x^{96} + x^{90} + x^{88} + x^{82} + x^{80} + x^{74} + x^{72} + x^{66} \\
& + x^{64} + x^{58} + x^{56} + x^{50} + x^{48} + x^{26} + x^{24} + x^{18} + x^{16} + x^{10} + x^8 \\
& + x^2 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 1, 0, 1, 0, 1, 0, 1, 1, 0, 0, 0, 0, 1]$.

For $\phi(M^e, 1, 0)$:

$$\begin{aligned}
p_0(x) &= x^{111} + x^{110} + x^{107} + x^{105} + x^{100} + x^{98} + x^{97} + x^{93} + x^{92} + x^{89} + x^{86} \\
& + x^{84} + x^{83} + x^{81} + x^{80} + x^{78} + x^{76} + x^{73} + x^{72} + x^{71} + x^{70} + x^{66} \\
& + x^{65} + x^{63} + x^{62} + x^{61} + x^{59} + x^{57} + x^{54} + x^{52} + x^{51} + x^{49} + x^{48} \\
& + x^{47} + x^{45} + x^{44} + x^{43} + x^{40} + x^{39}, \\
p_3(x) &= x^{109} + x^{108} + x^{106} + x^{105} + x^{101} + x^{100} + x^{99} + x^{97} + x^{96} + x^{94} \\
& + x^{93} + x^{91} + x^{90} + x^{89} + x^{83} + x^{82} + x^{80} + x^{78} + x^{77} + x^{73} + x^{72} \\
& + x^{70} + x^{68} + x^{67} + x^{45} + x^{44} + x^{42} + x^{41} + x^{37} + x^{36} + x^{35} + x^{33} \\
& + x^{32} + x^{30} + x^{28} + x^{27}, \\
p_6(x) &= x^{108} + x^{104} + x^{102} + x^{98} + x^{96} + x^{92} + x^{90} + x^{88} + x^{86} + x^{84} + x^{82} \\
& + x^{80} + x^{78} + x^{76} + x^{74} + x^{72} + x^{70} + x^{68} + x^{66} + x^{62} + x^{60} + x^{54} \\
& + x^{52} + x^{48} + x^{46} + x^{44} + x^{42} + x^{38} + x^{36} + x^{32} + x^{30} + x^{28} + x^{26} \\
& + x^{24} + x^{22} + x^{18}, \\
p_9(x) &= x^{107} + x^{106} + x^{105} + x^{101} + x^{100} + x^{99} + x^{97} + x^{96} + x^{95} + x^{93} + x^{92} \\
& + x^{91} + x^{89} + x^{88} + x^{87} + x^{83} + x^{82} + x^{80} + x^{79} + x^{77} + x^{76} + x^{74}
\end{aligned}$$

$$\begin{aligned}
& + x^{73} + x^{71} + x^{70} + x^{68} + x^{67} + x^{65} + x^{64} + x^{63} + x^{53} + x^{52} + x^{51} \\
& + x^{43} + x^{42} + x^{41} + x^{39} + x^{38} + x^{37} + x^{35} + x^{34} + x^{32} + x^{30} + x^{29} \\
& + x^{25} + x^{24} + x^{23} + x^{21} + x^{20} + x^{19} + x^{15} + x^{14} + x^{13} + x^{11} + x^{10} \\
& + x^9, \\
p_{12}(x) = & x^{106} + x^{104} + x^{98} + x^{96} + x^{90} + x^{88} + x^{82} + x^{80} + x^{74} + x^{72} + x^{66} \\
& + x^{64} + x^{58} + x^{56} + x^{50} + x^{48} + x^{26} + x^{24} + x^{18} + x^{16} + x^{10} + x^8 \\
& + x^2 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 1, 0, 1, 0, 1, 0, 1, 1, 0, 0, 0, 0, 1]$.

For $\phi(M^e, 1, 1)$:

$$\begin{aligned}
p_0(x) = & x^{108} + x^{106} + x^{104} + x^{102} + x^{98} + x^{96} + x^{94} + x^{92} + x^{88} + x^{84} + x^{78} \\
& + x^{76} + x^{74} + x^{72} + x^{66} + x^{64} + x^{54} + x^{52} + x^{42}, \\
p_3(x) = & x^{102} + x^{100} + x^{98} + x^{96} + x^{94} + x^{90} + x^{86} + x^{82} + x^{74} + x^{70} + x^{68} \\
& + x^{66} + x^{64} + x^{60} + x^{56} + x^{44} + x^{42} + x^{40} + x^{38} + x^{30} + x^{26} + x^{18}, \\
p_6(x) = & x^{102} + x^{96} + x^{92} + x^{90} + x^{88} + x^{86} + x^{84} + x^{78} + x^{76} + x^{74} + x^{72} \\
& + x^{68} + x^{58} + x^{56} + x^{54} + x^{50} + x^{48} + x^{46} + x^{42} + x^{36} + x^{32} + x^{30} \\
& + x^{28} + x^{18} + x^{14} + x^{12} + x^8 + x^6 + x^4 + 1, \\
p_9(x) = & x^{100} + x^{98} + x^{96} + x^{90} + x^{88} + x^{82} + x^{80} + x^{78} + x^{74} + x^{72} + x^{68} \\
& + x^{66} + x^{62} + x^{60} + x^{46} + x^{44} + x^{42} + x^{40} + x^{38} + x^{32} + x^{30} + x^{26} \\
& + x^{18} + x^{16} + x^8 + x^6, \\
p_{12}(x) = & x^{100} + x^{96} + x^{84} + x^{80} + x^{68} + x^{64} + x^{52} + x^{48} + x^{20} + x^{16} + x^4 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 0, 1, 0, 0, 0, 1, 0, 0, 0, 0, 0]$.

For $\phi(W^e, 0, 0)$:

$$\begin{aligned}
p_0(x) = & x^{108} + x^{106} + x^{102} + x^{96} + x^{94} + x^{92} + x^{90} + x^{84} + x^{82} + x^{80} + x^{74} \\
& + x^{62} + x^{60} + x^{58} + x^{56} + x^{52} + x^{50} + x^{48} + x^{44} + x^{42} + x^{40} + x^{38} \\
& + x^{36}, \\
p_3(x) = & x^{102} + x^{100} + x^{88} + x^{86} + x^{84} + x^{78} + x^{76} + x^{70} + x^{68} + x^{62} + x^{60} \\
& + x^{56} + x^{54} + x^{52} + x^{48} + x^{46} + x^{44} + x^{38} + x^{36} + x^{32} + x^{30} + x^{28} \\
& + x^{22} + x^{20}, \\
p_6(x) = & x^{102} + x^{98} + x^{96} + x^{94} + x^{92} + x^{86} + x^{84} + x^{82} + x^{80} + x^{72} + x^{66} \\
& + x^{62} + x^{58} + x^{50} + x^{48} + x^{44} + x^{40} + x^{36} + x^{30} + x^{26} + x^{24} + x^{22} \\
& + x^{20} + x^{14} + x^8 + x^6 + x^4 + 1, \\
p_9(x) = & x^{100} + x^{98} + x^{96} + x^{90} + x^{88} + x^{82} + x^{80} + x^{78} + x^{74} + x^{72} + x^{68} \\
& + x^{66} + x^{62} + x^{60} + x^{46} + x^{44} + x^{42} + x^{40} + x^{38} + x^{32} + x^{30} + x^{26} \\
& + x^{18} + x^{16} + x^8 + x^6, \\
p_{12}(x) = & x^{100} + x^{96} + x^{84} + x^{80} + x^{68} + x^{64} + x^{52} + x^{48} + x^{20} + x^{16} + x^4 + 1,
\end{aligned}$$

and the initial terms are $[1, 0, 0, 0, 0, 0, 1, 0, 0, 0, 1, 0, 1, 0, 0, 0]$.

For $\phi(W^e, 0, 1)$:

$$\begin{aligned}
p_0(x) &= x^{111} + x^{109} + x^{108} + x^{102} + x^{101} + x^{99} + x^{98} + x^{97} + x^{95} + x^{89} + x^{88} \\
&\quad + x^{85} + x^{82} + x^{74} + x^{70} + x^{69} + x^{66} + x^{65} + x^{63} + x^{62} + x^{61} + x^{59} \\
&\quad + x^{57} + x^{56} + x^{54} + x^{53} + x^{50} + x^{47} + x^{46} + x^{45} + x^{43} + x^{40} + x^{39}, \\
p_3(x) &= x^{109} + x^{108} + x^{106} + x^{105} + x^{101} + x^{100} + x^{99} + x^{97} + x^{96} + x^{94} \\
&\quad + x^{93} + x^{91} + x^{90} + x^{89} + x^{83} + x^{82} + x^{80} + x^{78} + x^{77} + x^{73} + x^{72} \\
&\quad + x^{70} + x^{68} + x^{67} + x^{45} + x^{44} + x^{42} + x^{41} + x^{37} + x^{36} + x^{35} + x^{33} \\
&\quad + x^{32} + x^{30} + x^{28} + x^{27}, \\
p_6(x) &= x^{108} + x^{104} + x^{102} + x^{98} + x^{96} + x^{92} + x^{90} + x^{88} + x^{86} + x^{84} + x^{82} \\
&\quad + x^{80} + x^{78} + x^{76} + x^{74} + x^{72} + x^{70} + x^{68} + x^{66} + x^{62} + x^{60} + x^{54} \\
&\quad + x^{52} + x^{48} + x^{46} + x^{44} + x^{42} + x^{38} + x^{36} + x^{32} + x^{30} + x^{28} + x^{26} \\
&\quad + x^{24} + x^{22} + x^{18}, \\
p_9(x) &= x^{107} + x^{106} + x^{105} + x^{101} + x^{100} + x^{99} + x^{97} + x^{96} + x^{95} + x^{93} + x^{92} \\
&\quad + x^{91} + x^{89} + x^{88} + x^{87} + x^{83} + x^{82} + x^{80} + x^{79} + x^{77} + x^{76} + x^{74} \\
&\quad + x^{73} + x^{71} + x^{70} + x^{68} + x^{67} + x^{65} + x^{64} + x^{63} + x^{53} + x^{52} + x^{51} \\
&\quad + x^{43} + x^{42} + x^{41} + x^{39} + x^{38} + x^{37} + x^{35} + x^{34} + x^{32} + x^{30} + x^{29} \\
&\quad + x^{25} + x^{24} + x^{23} + x^{21} + x^{20} + x^{19} + x^{15} + x^{14} + x^{13} + x^{11} + x^{10} \\
&\quad + x^9, \\
p_{12}(x) &= x^{106} + x^{104} + x^{98} + x^{96} + x^{90} + x^{88} + x^{82} + x^{80} + x^{74} + x^{72} + x^{66} \\
&\quad + x^{64} + x^{58} + x^{56} + x^{50} + x^{48} + x^{26} + x^{24} + x^{18} + x^{16} + x^{10} + x^8 \\
&\quad + x^2 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 1, 0, 0, 0, 1, 0, 1, 1, 1, 1, 0, 1]$.

For $\phi(W^e, 1, 0)$:

$$\begin{aligned}
p_0(x) &= x^{111} + x^{109} + x^{108} + x^{102} + x^{101} + x^{99} + x^{98} + x^{97} + x^{95} + x^{89} + x^{88} \\
&\quad + x^{85} + x^{82} + x^{74} + x^{70} + x^{69} + x^{66} + x^{65} + x^{63} + x^{62} + x^{61} + x^{59} \\
&\quad + x^{57} + x^{56} + x^{54} + x^{53} + x^{50} + x^{47} + x^{46} + x^{45} + x^{43} + x^{40} + x^{39}, \\
p_3(x) &= x^{109} + x^{108} + x^{106} + x^{105} + x^{101} + x^{100} + x^{99} + x^{97} + x^{96} + x^{94} \\
&\quad + x^{93} + x^{91} + x^{90} + x^{89} + x^{83} + x^{82} + x^{80} + x^{78} + x^{77} + x^{73} + x^{72} \\
&\quad + x^{70} + x^{68} + x^{67} + x^{45} + x^{44} + x^{42} + x^{41} + x^{37} + x^{36} + x^{35} + x^{33} \\
&\quad + x^{32} + x^{30} + x^{28} + x^{27}, \\
p_6(x) &= x^{108} + x^{104} + x^{102} + x^{98} + x^{96} + x^{92} + x^{90} + x^{88} + x^{86} + x^{84} + x^{82} \\
&\quad + x^{80} + x^{78} + x^{76} + x^{74} + x^{72} + x^{70} + x^{68} + x^{66} + x^{62} + x^{60} + x^{54} \\
&\quad + x^{52} + x^{48} + x^{46} + x^{44} + x^{42} + x^{38} + x^{36} + x^{32} + x^{30} + x^{28} + x^{26} \\
&\quad + x^{24} + x^{22} + x^{18},
\end{aligned}$$

$$\begin{aligned}
p_9(x) &= x^{107} + x^{106} + x^{105} + x^{101} + x^{100} + x^{99} + x^{97} + x^{96} + x^{95} + x^{93} + x^{92} \\
&\quad + x^{91} + x^{89} + x^{88} + x^{87} + x^{83} + x^{82} + x^{80} + x^{79} + x^{77} + x^{76} + x^{74} \\
&\quad + x^{73} + x^{71} + x^{70} + x^{68} + x^{67} + x^{65} + x^{64} + x^{63} + x^{53} + x^{52} + x^{51} \\
&\quad + x^{43} + x^{42} + x^{41} + x^{39} + x^{38} + x^{37} + x^{35} + x^{34} + x^{32} + x^{30} + x^{29} \\
&\quad + x^{25} + x^{24} + x^{23} + x^{21} + x^{20} + x^{19} + x^{15} + x^{14} + x^{13} + x^{11} + x^{10} \\
&\quad + x^9, \\
p_{12}(x) &= x^{106} + x^{104} + x^{98} + x^{96} + x^{90} + x^{88} + x^{82} + x^{80} + x^{74} + x^{72} + x^{66} \\
&\quad + x^{64} + x^{58} + x^{56} + x^{50} + x^{48} + x^{26} + x^{24} + x^{18} + x^{16} + x^{10} + x^8 \\
&\quad + x^2 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 1, 0, 0, 0, 1, 0, 1, 1, 1, 1, 0, 1]$.

For $\phi(W^e, 1, 1)$:

$$\begin{aligned}
p_0(x) &= x^{114} + x^{112} + x^{106} + x^{104} + x^{100} + x^{96} + x^{94} + x^{92} + x^{88} + x^{86} + x^{80} \\
&\quad + x^{78} + x^{74} + x^{72} + x^{70} + x^{68} + x^{66} + x^{60} + x^{58} + x^{56} + x^{50} + x^{46} \\
&\quad + x^{42}, \\
p_3(x) &= x^{110} + x^{106} + x^{96} + x^{88} + x^{70} + x^{66} + x^{64} + x^{54} + x^{50} + x^{48} + x^{46} \\
&\quad + x^{42} + x^{40} + x^{38} + x^{34} + x^{32} + x^{22} + x^{18}, \\
p_6(x) &= x^{108} + x^{106} + x^{104} + x^{100} + x^{92} + x^{88} + x^{84} + x^{82} + x^{78} + x^{76} + x^{72} \\
&\quad + x^{62} + x^{52} + x^{46} + x^{38} + x^{36} + x^{30} + x^{28} + x^{18} + x^{16} + x^{10} + x^8 \\
&\quad + x^6 + 1, \\
p_9(x) &= x^{112} + x^{110} + x^{106} + x^{100} + x^{90} + x^{86} + x^{82} + x^{80} + x^{78} + x^{74} + x^{72} \\
&\quad + x^{64} + x^{62} + x^{60} + x^{58} + x^{56} + x^{50} + x^{46} + x^{44} + x^{42} + x^{38} + x^{36} \\
&\quad + x^{32} + x^{30} + x^{28} + x^{24} + x^{22} + x^{14} + x^{12} + x^{10} + x^8 + x^6, \\
p_{12}(x) &= x^{112} + x^{48} + x^{32} + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 0, 0, 0, 1, 0, 0, 0, 0, 0, 1, 0]$.

For $\phi(M^o, 0, 0)$:

$$\begin{aligned}
p_0(x) &= x^{111} + x^{110} + x^{109} + x^{106} + x^{105} + x^{104} + x^{102} + x^{101} + x^{100} + x^{98} \\
&\quad + x^{96} + x^{94} + x^{91} + x^{89} + x^{84} + x^{83} + x^{82} + x^{76} + x^{75} + x^{71} + x^{70} \\
&\quad + x^{65} + x^{62} + x^{61} + x^{59} + x^{54} + x^{52} + x^{50} + x^{49} + x^{47} + x^{46} + x^{43} \\
&\quad + x^{42} + x^{40} + x^{36}, \\
p_3(x) &= x^{110} + x^{109} + x^{104} + x^{102} + x^{101} + x^{99} + x^{96} + x^{95} + x^{94} + x^{93} + x^{92} \\
&\quad + x^{88} + x^{86} + x^{84} + x^{83} + x^{82} + x^{79} + x^{78} + x^{77} + x^{75} + x^{74} + x^{71} \\
&\quad + x^{66} + x^{64} + x^{60} + x^{58} + x^{56} + x^{54} + x^{48} + x^{45} + x^{44} + x^{42} + x^{40} \\
&\quad + x^{38} + x^{37} + x^{36} + x^{35} + x^{34} + x^{31} + x^{30} + x^{28} + x^{24}, \\
p_6(x) &= x^{110} + x^{109} + x^{105} + x^{100} + x^{99} + x^{97} + x^{96} + x^{94} + x^{93} + x^{89} + x^{88} \\
&\quad + x^{87} + x^{85} + x^{84} + x^{81} + x^{80} + x^{79} + x^{78} + x^{77} + x^{76} + x^{73} + x^{72}
\end{aligned}$$

$$\begin{aligned}
& + x^{71} + x^{70} + x^{69} + x^{64} + x^{63} + x^{62} + x^{61} + x^{60} + x^{53} + x^{52} + x^{51} \\
& + x^{50} + x^{49} + x^{48} + x^{47} + x^{45} + x^{44} + x^{42} + x^{39} + x^{38} + x^{37} + x^{34} \\
& + x^{33} + x^{32} + x^{29} + x^{27} + x^{26} + x^{25} + x^{20} + x^{19} + x^{16} + x^{12} + x^{10} \\
& + x^8 + x^2 + 1, \\
p_9(x) &= x^{110} + x^{108} + x^{106} + x^{104} + x^{98} + x^{96} + x^{94} + x^{92} + x^{78} + x^{76} + x^{74} \\
& + x^{72} + x^{66} + x^{64} + x^{62} + x^{60} + x^{30} + x^{28} + x^{26} + x^{24} + x^{18} + x^{16} \\
& + x^{14} + x^{12}, \\
p_{12}(x) &= x^{110} + x^{108} + x^{94} + x^{92} + x^{90} + x^{88} + x^{82} + x^{80} + x^{78} + x^{76} + x^{62} \\
& + x^{60} + x^{58} + x^{56} + x^{50} + x^{48} + x^{30} + x^{28} + x^{14} + x^{12} + x^{10} + x^8 \\
& + x^2 + 1,
\end{aligned}$$

and the initial terms are $[1, 0, 0, 0, 0, 0, 1, 1, 1, 1, 1, 0, 0, 1, 0, 1]$.

For $\phi(M^o, 0, 1)$:

$$\begin{aligned}
p_0(x) &= x^{108} + x^{105} + x^{98} + x^{97} + x^{96} + x^{95} + x^{94} + x^{92} + x^{91} + x^{86} + x^{85} \\
& + x^{83} + x^{82} + x^{77} + x^{75} + x^{74} + x^{72} + x^{70} + x^{63} + x^{61} + x^{60} + x^{55} \\
& + x^{53} + x^{52} + x^{51} + x^{50} + x^{47} + x^{46} + x^{45} + x^{44} + x^{43} + x^{42} + x^{41} \\
& + x^{40} + x^{39}, \\
p_3(x) &= x^{101} + x^{100} + x^{99} + x^{93} + x^{92} + x^{91} + x^{85} + x^{84} + x^{83} + x^{69} + x^{68} \\
& + x^{67} + x^{37} + x^{36} + x^{35} + x^{29} + x^{28} + x^{27}, \\
p_6(x) &= x^{102} + x^{100} + x^{98} + x^{90} + x^{88} + x^{86} + x^{82} + x^{80} + x^{78} + x^{74} + x^{72} \\
& + x^{70} + x^{66} + x^{64} + x^{62} + x^{54} + x^{52} + x^{48} + x^{46} + x^{42} + x^{40} + x^{38} \\
& + x^{30} + x^{28} + x^{26} + x^{22} + x^{20} + x^{18}, \\
p_9(x) &= x^{103} + x^{102} + x^{100} + x^{98} + x^{97} + x^{95} + x^{94} + x^{92} + x^{90} + x^{89} + x^{71} \\
& + x^{70} + x^{68} + x^{66} + x^{65} + x^{63} + x^{62} + x^{60} + x^{58} + x^{57} + x^{23} + x^{22} \\
& + x^{20} + x^{18} + x^{17} + x^{15} + x^{14} + x^{12} + x^{10} + x^9, \\
p_{12}(x) &= x^{104} + x^{100} + x^{96} + x^{92} + x^{84} + x^{80} + x^{72} + x^{68} + x^{64} + x^{60} + x^{52} \\
& + x^{48} + x^{24} + x^{20} + x^{16} + x^{12} + x^4 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 1, 0, 1, 0, 1, 0, 0, 0, 1, 0, 0, 0, 0]$.

For $\phi(M^o, 1, 0)$:

$$\begin{aligned}
p_0(x) &= x^{114} + x^{113} + x^{112} + x^{111} + x^{110} + x^{109} + x^{108} + x^{105} + x^{104} + x^{102} \\
& + x^{99} + x^{98} + x^{97} + x^{96} + x^{92} + x^{91} + x^{89} + x^{85} + x^{81} + x^{79} + x^{77} \\
& + x^{75} + x^{74} + x^{73} + x^{72} + x^{70} + x^{69} + x^{67} + x^{66} + x^{65} + x^{64} + x^{60} \\
& + x^{57} + x^{49} + x^{48} + x^{47} + x^{44} + x^{42} + x^{41} + x^{40} + x^{39}, \\
p_3(x) &= x^{113} + x^{112} + x^{111} + x^{109} + x^{108} + x^{107} + x^{89} + x^{88} + x^{87} + x^{85} \\
& + x^{84} + x^{83} + x^{81} + x^{80} + x^{79} + x^{77} + x^{76} + x^{75} + x^{73} + x^{72} + x^{71} \\
& + x^{69} + x^{68} + x^{67} + x^{49} + x^{48} + x^{47} + x^{45} + x^{44} + x^{43} + x^{33} + x^{32}
\end{aligned}$$

$$\begin{aligned}
& + x^{31} + x^{29} + x^{28} + x^{27}, \\
p_6(x) &= x^{114} + x^{112} + x^{108} + x^{104} + x^{102} + x^{98} + x^{96} + x^{94} + x^{70} + x^{68} + x^{66} \\
& + x^{54} + x^{52} + x^{50} + x^{46} + x^{44} + x^{42} + x^{38} + x^{36} + x^{34} + x^{26} + x^{24} \\
& + x^{20} + x^{18}, \\
p_9(x) &= x^{115} + x^{114} + x^{112} + x^{111} + x^{109} + x^{108} + x^{106} + x^{105} + x^{99} + x^{98} \\
& + x^{96} + x^{95} + x^{93} + x^{92} + x^{90} + x^{89} + x^{83} + x^{82} + x^{80} + x^{79} + x^{77} \\
& + x^{76} + x^{74} + x^{73} + x^{67} + x^{66} + x^{64} + x^{63} + x^{61} + x^{60} + x^{58} + x^{57} \\
& + x^{35} + x^{34} + x^{32} + x^{31} + x^{29} + x^{28} + x^{26} + x^{25} + x^{19} + x^{18} + x^{16} \\
& + x^{15} + x^{13} + x^{12} + x^{10} + x^9, \\
p_{12}(x) &= x^{116} + x^{108} + x^{100} + x^{96} + x^{92} + x^{84} + x^{80} + x^{76} + x^{68} + x^{64} + x^{60} \\
& + x^{48} + x^{36} + x^{28} + x^{20} + x^{16} + x^{12} + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 1, 0, 0, 0, 1, 0, 1, 0, 0, 0, 0, 0]$.

For $\phi(M^o, 1, 1)$:

$$\begin{aligned}
p_0(x) &= x^{111} + x^{106} + x^{104} + x^{99} + x^{96} + x^{94} + x^{93} + x^{92} + x^{89} + x^{88} + x^{87} \\
& + x^{86} + x^{85} + x^{84} + x^{83} + x^{82} + x^{81} + x^{74} + x^{73} + x^{71} + x^{69} + x^{66} \\
& + x^{65} + x^{63} + x^{61} + x^{59} + x^{58} + x^{54} + x^{53} + x^{52} + x^{48} + x^{47} + x^{46} \\
& + x^{44} + x^{42}, \\
p_3(x) &= x^{110} + x^{109} + x^{108} + x^{106} + x^{101} + x^{100} + x^{99} + x^{96} + x^{95} + x^{93} \\
& + x^{88} + x^{84} + x^{83} + x^{80} + x^{79} + x^{78} + x^{77} + x^{76} + x^{75} + x^{72} + x^{71} \\
& + x^{70} + x^{60} + x^{58} + x^{56} + x^{54} + x^{48} + x^{45} + x^{37} + x^{35} + x^{34} + x^{31} \\
& + x^{28} + x^{26}, \\
p_6(x) &= x^{110} + x^{109} + x^{108} + x^{106} + x^{105} + x^{99} + x^{97} + x^{96} + x^{94} + x^{93} + x^{92} \\
& + x^{89} + x^{88} + x^{87} + x^{86} + x^{85} + x^{82} + x^{81} + x^{79} + x^{77} + x^{74} + x^{73} \\
& + x^{71} + x^{69} + x^{68} + x^{66} + x^{64} + x^{63} + x^{62} + x^{61} + x^{60} + x^{54} + x^{53} \\
& + x^{51} + x^{50} + x^{49} + x^{47} + x^{45} + x^{44} + x^{39} + x^{38} + x^{37} + x^{34} + x^{33} \\
& + x^{29} + x^{27} + x^{25} + x^{24} + x^{22} + x^{20} + x^{19} + x^{14} + x^{10} + x^8 + x^2 + 1, \\
p_9(x) &= x^{110} + x^{108} + x^{106} + x^{104} + x^{98} + x^{96} + x^{94} + x^{92} + x^{78} + x^{76} + x^{74} \\
& + x^{72} + x^{66} + x^{64} + x^{62} + x^{60} + x^{30} + x^{28} + x^{26} + x^{24} + x^{18} + x^{16} \\
& + x^{14} + x^{12}, \\
p_{12}(x) &= x^{110} + x^{108} + x^{94} + x^{92} + x^{90} + x^{88} + x^{82} + x^{80} + x^{78} + x^{76} + x^{62} \\
& + x^{60} + x^{58} + x^{56} + x^{50} + x^{48} + x^{30} + x^{28} + x^{14} + x^{12} + x^{10} + x^8 \\
& + x^2 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 0, 0, 1, 0, 1, 0, 0, 0, 1, 1, 1]$.

For $\phi(W^o, 0, 0)$:

$$p_0(x) = x^{111} + x^{110} + x^{109} + x^{106} + x^{105} + x^{104} + x^{102} + x^{101} + x^{100} + x^{98}$$

$$\begin{aligned}
& + x^{96} + x^{94} + x^{91} + x^{89} + x^{84} + x^{83} + x^{82} + x^{76} + x^{75} + x^{71} + x^{70} \\
& + x^{65} + x^{62} + x^{61} + x^{59} + x^{54} + x^{52} + x^{50} + x^{49} + x^{47} + x^{46} + x^{43} \\
& + x^{42} + x^{40} + x^{36}, \\
p_3(x) &= x^{110} + x^{109} + x^{104} + x^{102} + x^{101} + x^{99} + x^{96} + x^{95} + x^{94} + x^{93} + x^{92} \\
& + x^{88} + x^{86} + x^{84} + x^{83} + x^{82} + x^{79} + x^{78} + x^{77} + x^{75} + x^{74} + x^{71} \\
& + x^{66} + x^{64} + x^{60} + x^{58} + x^{56} + x^{54} + x^{48} + x^{45} + x^{44} + x^{42} + x^{40} \\
& + x^{38} + x^{37} + x^{36} + x^{35} + x^{34} + x^{31} + x^{30} + x^{28} + x^{24}, \\
p_6(x) &= x^{110} + x^{109} + x^{105} + x^{100} + x^{99} + x^{97} + x^{96} + x^{94} + x^{93} + x^{89} + x^{88} \\
& + x^{87} + x^{85} + x^{84} + x^{81} + x^{80} + x^{79} + x^{78} + x^{77} + x^{76} + x^{73} + x^{72} \\
& + x^{71} + x^{70} + x^{69} + x^{64} + x^{63} + x^{62} + x^{61} + x^{60} + x^{53} + x^{52} + x^{51} \\
& + x^{50} + x^{49} + x^{48} + x^{47} + x^{45} + x^{44} + x^{42} + x^{39} + x^{38} + x^{37} + x^{34} \\
& + x^{33} + x^{32} + x^{29} + x^{27} + x^{26} + x^{25} + x^{20} + x^{19} + x^{16} + x^{12} + x^{10} \\
& + x^8 + x^2 + 1, \\
p_9(x) &= x^{110} + x^{108} + x^{106} + x^{104} + x^{98} + x^{96} + x^{94} + x^{92} + x^{78} + x^{76} + x^{74} \\
& + x^{72} + x^{66} + x^{64} + x^{62} + x^{60} + x^{30} + x^{28} + x^{26} + x^{24} + x^{18} + x^{16} \\
& + x^{14} + x^{12}, \\
p_{12}(x) &= x^{110} + x^{108} + x^{94} + x^{92} + x^{90} + x^{88} + x^{82} + x^{80} + x^{78} + x^{76} + x^{62} \\
& + x^{60} + x^{58} + x^{56} + x^{50} + x^{48} + x^{30} + x^{28} + x^{14} + x^{12} + x^{10} + x^8 \\
& + x^2 + 1,
\end{aligned}$$

and the initial terms are $[1, 0, 0, 0, 0, 0, 1, 1, 1, 1, 1, 0, 0, 1, 0, 1]$.

For $\phi(W^o, 0, 1)$:

$$\begin{aligned}
p_0(x) &= x^{114} + x^{113} + x^{112} + x^{111} + x^{110} + x^{109} + x^{108} + x^{105} + x^{104} + x^{102} \\
& + x^{99} + x^{98} + x^{97} + x^{96} + x^{92} + x^{91} + x^{89} + x^{85} + x^{81} + x^{79} + x^{77} \\
& + x^{75} + x^{74} + x^{73} + x^{72} + x^{70} + x^{69} + x^{67} + x^{66} + x^{65} + x^{64} + x^{60} \\
& + x^{57} + x^{49} + x^{48} + x^{47} + x^{44} + x^{42} + x^{41} + x^{40} + x^{39}, \\
p_3(x) &= x^{113} + x^{112} + x^{111} + x^{109} + x^{108} + x^{107} + x^{89} + x^{88} + x^{87} + x^{85} \\
& + x^{84} + x^{83} + x^{81} + x^{80} + x^{79} + x^{77} + x^{76} + x^{75} + x^{73} + x^{72} + x^{71} \\
& + x^{69} + x^{68} + x^{67} + x^{49} + x^{48} + x^{47} + x^{45} + x^{44} + x^{43} + x^{33} + x^{32} \\
& + x^{31} + x^{29} + x^{28} + x^{27}, \\
p_6(x) &= x^{114} + x^{112} + x^{108} + x^{104} + x^{102} + x^{98} + x^{96} + x^{94} + x^{70} + x^{68} + x^{66} \\
& + x^{54} + x^{52} + x^{50} + x^{46} + x^{44} + x^{42} + x^{38} + x^{36} + x^{34} + x^{26} + x^{24} \\
& + x^{20} + x^{18}, \\
p_9(x) &= x^{115} + x^{114} + x^{112} + x^{111} + x^{109} + x^{108} + x^{106} + x^{105} + x^{99} + x^{98} \\
& + x^{96} + x^{95} + x^{93} + x^{92} + x^{90} + x^{89} + x^{83} + x^{82} + x^{80} + x^{79} + x^{77} \\
& + x^{76} + x^{74} + x^{73} + x^{67} + x^{66} + x^{64} + x^{63} + x^{61} + x^{60} + x^{58} + x^{57}
\end{aligned}$$

$$\begin{aligned}
& + x^{35} + x^{34} + x^{32} + x^{31} + x^{29} + x^{28} + x^{26} + x^{25} + x^{19} + x^{18} + x^{16} \\
& + x^{15} + x^{13} + x^{12} + x^{10} + x^9, \\
p_{12}(x) = & x^{116} + x^{108} + x^{100} + x^{96} + x^{92} + x^{84} + x^{80} + x^{76} + x^{68} + x^{64} + x^{60} \\
& + x^{48} + x^{36} + x^{28} + x^{20} + x^{16} + x^{12} + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 1, 0, 0, 0, 1, 0, 1, 0, 0, 0, 0]$.

For $\phi(W^o, 1, 0)$:

$$\begin{aligned}
p_0(x) = & x^{108} + x^{105} + x^{98} + x^{97} + x^{96} + x^{95} + x^{94} + x^{92} + x^{91} + x^{86} + x^{85} \\
& + x^{83} + x^{82} + x^{77} + x^{75} + x^{74} + x^{72} + x^{70} + x^{63} + x^{61} + x^{60} + x^{55} \\
& + x^{53} + x^{52} + x^{51} + x^{50} + x^{47} + x^{46} + x^{45} + x^{44} + x^{43} + x^{42} + x^{41} \\
& + x^{40} + x^{39}, \\
p_3(x) = & x^{101} + x^{100} + x^{99} + x^{93} + x^{92} + x^{91} + x^{85} + x^{84} + x^{83} + x^{69} + x^{68} \\
& + x^{67} + x^{37} + x^{36} + x^{35} + x^{29} + x^{28} + x^{27}, \\
p_6(x) = & x^{102} + x^{100} + x^{98} + x^{90} + x^{88} + x^{86} + x^{82} + x^{80} + x^{78} + x^{74} + x^{72} \\
& + x^{70} + x^{66} + x^{64} + x^{62} + x^{54} + x^{52} + x^{48} + x^{46} + x^{42} + x^{40} + x^{38} \\
& + x^{30} + x^{28} + x^{26} + x^{22} + x^{20} + x^{18}, \\
p_9(x) = & x^{103} + x^{102} + x^{100} + x^{98} + x^{97} + x^{95} + x^{94} + x^{92} + x^{90} + x^{89} + x^{71} \\
& + x^{70} + x^{68} + x^{66} + x^{65} + x^{63} + x^{62} + x^{60} + x^{58} + x^{57} + x^{23} + x^{22} \\
& + x^{20} + x^{18} + x^{17} + x^{15} + x^{14} + x^{12} + x^{10} + x^9, \\
p_{12}(x) = & x^{104} + x^{100} + x^{96} + x^{92} + x^{84} + x^{80} + x^{72} + x^{68} + x^{64} + x^{60} + x^{52} \\
& + x^{48} + x^{24} + x^{20} + x^{16} + x^{12} + x^4 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 1, 0, 1, 0, 1, 0, 0, 0, 1, 0, 0, 0]$.

For $\phi(W^o, 1, 1)$:

$$\begin{aligned}
p_0(x) = & x^{111} + x^{106} + x^{104} + x^{99} + x^{96} + x^{94} + x^{93} + x^{92} + x^{89} + x^{88} + x^{87} \\
& + x^{86} + x^{85} + x^{84} + x^{83} + x^{82} + x^{81} + x^{74} + x^{73} + x^{71} + x^{69} + x^{66} \\
& + x^{65} + x^{63} + x^{61} + x^{59} + x^{58} + x^{54} + x^{53} + x^{52} + x^{48} + x^{47} + x^{46} \\
& + x^{44} + x^{42}, \\
p_3(x) = & x^{110} + x^{109} + x^{108} + x^{106} + x^{101} + x^{100} + x^{99} + x^{96} + x^{95} + x^{93} \\
& + x^{88} + x^{84} + x^{83} + x^{80} + x^{79} + x^{78} + x^{77} + x^{76} + x^{75} + x^{72} + x^{71} \\
& + x^{70} + x^{60} + x^{58} + x^{56} + x^{54} + x^{48} + x^{45} + x^{37} + x^{35} + x^{34} + x^{31} \\
& + x^{28} + x^{26}, \\
p_6(x) = & x^{110} + x^{109} + x^{108} + x^{106} + x^{105} + x^{99} + x^{97} + x^{96} + x^{94} + x^{93} + x^{92} \\
& + x^{89} + x^{88} + x^{87} + x^{86} + x^{85} + x^{82} + x^{81} + x^{79} + x^{77} + x^{74} + x^{73} \\
& + x^{71} + x^{69} + x^{68} + x^{66} + x^{64} + x^{63} + x^{62} + x^{61} + x^{60} + x^{54} + x^{53} \\
& + x^{51} + x^{50} + x^{49} + x^{47} + x^{45} + x^{44} + x^{39} + x^{38} + x^{37} + x^{34} + x^{33} \\
& + x^{29} + x^{27} + x^{25} + x^{24} + x^{22} + x^{20} + x^{19} + x^{14} + x^{10} + x^8 + x^2 + 1,
\end{aligned}$$

$$\begin{aligned}
p_9(x) &= x^{110} + x^{108} + x^{106} + x^{104} + x^{98} + x^{96} + x^{94} + x^{92} + x^{78} + x^{76} + x^{74} \\
&\quad + x^{72} + x^{66} + x^{64} + x^{62} + x^{60} + x^{30} + x^{28} + x^{26} + x^{24} + x^{18} + x^{16} \\
&\quad + x^{14} + x^{12}, \\
p_{12}(x) &= x^{110} + x^{108} + x^{94} + x^{92} + x^{90} + x^{88} + x^{82} + x^{80} + x^{78} + x^{76} + x^{62} \\
&\quad + x^{60} + x^{58} + x^{56} + x^{50} + x^{48} + x^{30} + x^{28} + x^{14} + x^{12} + x^{10} + x^8 \\
&\quad + x^2 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 0, 0, 1, 0, 1, 0, 0, 0, 1, 1, 1]$.

Below is the transition function and output function of an 2-automaton that generates $T = M_{0,0}^o$:

Transition function $(n, j) \mapsto \delta(n, j)$ ($\Lambda(n) := [\delta(n, 0), \delta(n, 1)]$):

n	$\Lambda(n)$	n	$\Lambda(n)$	n	$\Lambda(n)$	n	$\Lambda(n)$	n	$\Lambda(n)$
0	[1, 2]	38	[60, 61]	76	[99, 100]	114	[86, 32]	152	[163, 63]
1	[3, 4]	39	[62, 19]	77	[101, 98]	115	[39, 129]	153	[164, 58]
2	[5, 6]	40	[63, 64]	78	[102, 67]	116	[77, 125]	154	[121, 16]
3	[7, 8]	41	[59, 65]	79	[103, 104]	117	[133, 8]	155	[165, 157]
4	[9, 10]	42	[66, 67]	80	[105, 106]	118	[127, 93]	156	[8, 125]
5	[11, 12]	43	[68, 69]	81	[107, 50]	119	[124, 80]	157	[10, 37]
6	[13, 14]	44	[48, 70]	82	[108, 96]	120	[128, 86]	158	[12, 14]
7	[15, 16]	45	[71, 72]	83	[109, 19]	121	[39, 86]	159	[10, 9]
8	[17, 18]	46	[31, 44]	84	[110, 99]	122	[134, 14]	160	[166, 79]
9	[19, 20]	47	[11, 73]	85	[111, 99]	123	[37, 74]	161	[84, 74]
10	[10, 10]	48	[36, 74]	86	[61, 112]	124	[135, 123]	162	[167, 39]
11	[21, 22]	49	[75, 76]	87	[113, 63]	125	[16, 136]	163	[168, 86]
12	[23, 24]	50	[37, 9]	88	[114, 115]	126	[137, 138]	164	[133, 77]
13	[25, 26]	51	[77, 14]	89	[116, 53]	127	[139, 140]	165	[169, 79]
14	[27, 28]	52	[78, 79]	90	[117, 118]	128	[141, 20]	166	[170, 138]
15	[29, 30]	53	[36, 9]	91	[119, 120]	129	[142, 140]	167	[171, 22]
16	[31, 32]	54	[33, 80]	92	[46, 26]	130	[143, 27]	168	[172, 59]
17	[33, 34]	55	[81, 40]	93	[22, 121]	131	[144, 48]	169	[173, 161]
18	[35, 32]	56	[82, 44]	94	[122, 123]	132	[145, 114]	170	[174, 125]
19	[36, 10]	57	[83, 37]	95	[124, 34]	133	[146, 46]	171	[175, 40]
20	[37, 37]	58	[84, 37]	96	[31, 41]	134	[147, 112]	172	[176, 77]
21	[38, 39]	59	[10, 74]	97	[92, 76]	135	[148, 90]	173	[13, 125]
22	[40, 41]	60	[85, 86]	98	[31, 76]	136	[30, 92]	174	[177, 64]
23	[42, 43]	61	[86, 44]	99	[84, 9]	137	[149, 79]	175	[178, 10]
24	[35, 44]	62	[87, 77]	100	[73, 125]	138	[37, 10]	176	[179, 114]
25	[45, 34]	63	[40, 76]	101	[126, 43]	139	[150, 43]	177	[180, 43]
26	[40, 44]	64	[86, 41]	102	[73, 14]	140	[40, 32]	178	[181, 73]
27	[35, 41]	65	[77, 88]	103	[7, 12]	141	[151, 73]	179	[182, 39]
28	[30, 40]	66	[89, 90]	104	[127, 125]	142	[35, 76]	180	[183, 159]
29	[28, 46]	67	[36, 37]	105	[126, 91]	143	[152, 30]	181	[184, 61]
30	[47, 48]	68	[42, 91]	106	[128, 129]	144	[74, 10]	182	[55, 142]

31	[49, 24]	69	[81, 92]	107	[130, 77]	145	[153, 40]	183	[185, 90]
32	[50, 51]	70	[73, 93]	108	[129, 44]	146	[154, 39]	184	[186, 30]
33	[52, 53]	71	[94, 90]	109	[9, 37]	147	[155, 34]	185	[187, 72]
34	[54, 55]	72	[84, 10]	110	[131, 10]	148	[156, 157]	186	[120, 27]
35	[56, 18]	73	[95, 96]	111	[7, 73]	149	[158, 159]	187	[127, 14]
36	[57, 58]	74	[58, 50]	112	[86, 76]	150	[160, 161]		
37	[20, 59]	75	[97, 98]	113	[132, 30]	151	[162, 142]		

Output function $n \mapsto \tau(n)$:

n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$
0	0	27	0	54	1	81	0	108	0	135	1	162	1
1	0	28	0	55	0	82	0	109	0	136	0	163	0
2	0	29	0	56	0	83	0	110	1	137	0	164	1
3	0	30	0	57	0	84	1	111	0	138	1	165	0
4	0	31	1	58	1	85	0	112	1	139	1	166	1
5	0	32	1	59	0	86	1	113	1	140	0	167	1
6	0	33	1	60	0	87	1	114	1	141	1	168	0
7	0	34	1	61	1	88	1	115	1	142	0	169	0
8	1	35	0	62	1	89	0	116	0	143	0	170	1
9	0	36	0	63	0	90	1	117	1	144	1	171	1
10	0	37	1	64	1	91	1	118	1	145	1	172	0
11	0	38	0	65	0	92	1	119	1	146	1	173	0
12	0	39	1	66	0	93	0	120	1	147	0	174	1
13	0	40	0	67	0	94	0	121	1	148	1	175	1
14	0	41	0	68	0	95	1	122	0	149	0	176	0
15	0	42	0	69	0	96	1	123	1	150	1	177	1
16	1	43	0	70	1	97	1	124	1	151	1	178	1
17	1	44	0	71	0	98	1	125	1	152	0	179	0
18	0	45	0	72	1	99	1	126	0	153	1	180	1
19	0	46	1	73	1	100	1	127	1	154	1	181	1
20	1	47	0	74	1	101	0	128	1	155	0	182	0
21	0	48	0	75	1	102	1	129	0	156	1	183	1
22	0	49	1	76	1	103	0	130	0	157	0	184	1
23	0	50	1	77	0	104	1	131	1	158	0	185	1
24	0	51	0	78	1	105	0	132	1	159	0	186	1
25	0	52	1	79	0	106	1	133	1	160	1	187	1
26	0	53	0	80	0	107	0	134	0	161	1		

UNIVERSITÉ DE STRASBOURG, CNRS, IRMA UMR 7501, F-67000 STRASBOURG, FRANCE
Email address: `guoniu.han@unistra.fr`

SCHOOL OF MATHEMATICS AND STATISTICS, HUAZHONG UNIVERSITY OF SCIENCE AND TECHNOLOGY, WUHAN, PR CHINA
Email address: `huyining@hust.edu.cn`