

PROOF OF THE CONTINUED FRACTION CONJECTURE FOR

$$(a, b) = (z^2 + z, z^5 + z^4 + z^2 + z)$$

GUO-NIU HAN* AND YINING HU*

ABSTRACT. The continued fraction conjecture claims that the continued fraction defined by the (a, b) -Thue-Morse sequence is algebraic over $\mathbb{F}_2(x)$ for all pairs of distinct elements (a, b) in $\mathbb{F}_2[x] \setminus \mathbb{F}_2$. In this paper we prove the conjecture for $(a, b) = (z^2 + z, z^5 + z^4 + z^2 + z)$.

1. INTRODUCTION

Let $\mathbb{F}_2$ be the finite field of cardinality 2. Given a sequence of polynomials $\mathbf{a} = (a_0(z), a_1(z), a_2(z), \dots)$ of elements from $\mathbb{F}_2[z] \setminus \mathbb{F}_2$, we define the infinite continued fraction

$$(1.1) \quad \text{CF}(\mathbf{a}(z)) = \frac{1}{a_0(z) + \frac{1}{a_1(z) + \frac{1}{a_2(z) + \frac{1}{\ddots}}}}$$

as the limit of the finite continued fractions

$$(1.2) \quad \text{CF}_n(\mathbf{a}(z)) = \frac{1}{a_0(z) + \frac{1}{a_1(z) + \frac{1}{\ddots + \frac{1}{a_n(z)}}}} \in \mathbb{F}_2((1/z)).$$

Conversely, each power series in $\mathbb{F}_2[[1/z]]$ without constant term

$$(1.3) \quad c_{-1}z^{-1} + c_{-2}z^{-2} + c_{-3}z^{-3} + \dots$$

admits a continued fraction expansion of form (1.1).

Let a and b be two distinct elements from $\mathbb{F}_2[z] \setminus \mathbb{F}_2$. We consider the continued fraction (1.1) associated with the (a, b) -Thue-Morse sequence

$$\mathbf{t} = (t_0(z), t_1(z), t_2(z), \dots) = (a, b, b, a, b, a, a, b, \dots)$$

that we shall denote

$$(1.4) \quad \text{CF}(a, b) := \text{CF}(\mathbf{t}).$$

Date: 9/04/2020 22:09:54 .

2010 Mathematics Subject Classification. 11B85, 11J70, 11B50, 11Y65, 05A15.

Key words and phrases. automatic sequence, continued fraction, Thue-Morse sequence.

* This paper was automatically generated from a computer program developed by the authors, with 10.8 minutes CPU time used.

Theorem 1.1. *When $(a, b) = (z^2 + z, z^5 + z^4 + z^2 + z)$, the two power series $\text{CF}(a, b)$ and $\text{CF}(b, a)$ are algebraic over $\mathbb{F}_2(z)$, with minimal polynomials of the form*

$$p_4(z)y^4 + p_3(z)y^3 + p_2(z)y^2 + p_1(z)y + p_0(z) = 0$$

where, for $\text{CF}(a, b)$,

$$p_0(z) = z^{11} + z^{10} + z^9 + z^8 + z^7 + z^6 + z^5 + z^3 + z^2 + 1,$$

$$p_1(z) = z^{16} + z^{15} + z^{14} + z^6 + z^5 + z^4,$$

$$p_2(z) = z^{18} + z^{12} + z^8 + z^2,$$

$$p_3(z) = z^{16} + z^{15} + z^{14} + z^6 + z^5 + z^4,$$

$$p_4(z) = z^{14} + z^{12} + z^{11} + z^9 + z^7 + z^5 + z^3 + z^2 + 1,$$

and for $\text{CF}(b, a)$,

$$p_0(z) = z^{11} + z^9 + z^8 + z^7 + z^6 + z^5 + z^3 + z^2 + 1,$$

$$p_1(z) = z^{16} + z^{15} + z^{14} + z^6 + z^5 + z^4,$$

$$p_2(z) = z^{18} + z^{12} + z^8 + z^2,$$

$$p_3(z) = z^{16} + z^{15} + z^{14} + z^6 + z^5 + z^4,$$

$$p_4(z) = z^{14} + z^{12} + z^{11} + z^{10} + z^9 + z^7 + z^5 + z^3 + z^2 + 1.$$

2. PROOF

Define the sequences of polynomials $(P_n(z))_{n \geq 0}$ and $(Q_n(z))_{n \geq 0}$ by

$$\begin{pmatrix} P_n(z) & Q_n(z) \\ P_{n-1}(z) & Q_{n-1}(z) \end{pmatrix} := \begin{pmatrix} t_n(z) & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} t_{n-1}(z) & 1 \\ 1 & 0 \end{pmatrix} \cdots \begin{pmatrix} t_0(z) & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}.$$

By the basic properties of continued fractions, we have

$$(2.1) \quad \text{CF}_n(a, b) = \frac{P_n(z)}{Q_n(z)} \in \mathbb{F}_2((1/z)).$$

As formula (2.1) is not efficient for calculating $\text{CF}_n(a, b)$, a fast method is to be derived and described as follows. Define

$$\begin{aligned} M_n(x) &= x^{\deg(t_{2^n-1})} \begin{pmatrix} t_{2^n-1}(1/x) & 1 \\ 1 & 0 \end{pmatrix} \times x^{\deg(t_{2^n-2})} \begin{pmatrix} t_{2^n-2}(1/x) & 1 \\ 1 & 0 \end{pmatrix} \times \\ &\quad \cdots \times x^{\deg(t_0)} \begin{pmatrix} t_0(1/x) & 1 \\ 1 & 0 \end{pmatrix}, \\ W_n(x) &= x^{\deg(\bar{t}_{2^n-1})} \begin{pmatrix} \bar{t}_{2^n-1}(1/x) & 1 \\ 1 & 0 \end{pmatrix} \times x^{\deg(\bar{t}_{2^n-2})} \begin{pmatrix} \bar{t}_{2^n-2}(1/x) & 1 \\ 1 & 0 \end{pmatrix} \times \\ &\quad \cdots \times x^{\deg(\bar{t}_0)} \begin{pmatrix} \bar{t}_0(1/x) & 1 \\ 1 & 0 \end{pmatrix}, \end{aligned}$$

where $\bar{t}$ is the (b, a) -Thue-Morse sequence. By the properties of the Thue-Morse sequence, we have for $n \geq 0$

$$M_{n+1}(x) = W_n(x) \cdot M_n(x),$$

$$W_{n+1}(x) = M_n(x) \cdot W_n(x).$$

Let $x := 1/z$. For each non-zero polynomial $P(z)$, define $\tilde{P}(x)$ to be $P(1/x)$. Then,

$$\text{CF}_n(a, b) = \frac{P_n(z)}{Q_n(z)} = \frac{\tilde{P}_n(x)}{\tilde{Q}_n(x)} \in \mathbb{F}_2((x)) = \mathbb{F}_2((1/z)).$$

Comparing the definition of $M_n(x)$ with the definition of $P_n(x)$ and $Q_n(x)$, we see that

$$\begin{aligned} M_n(x)_{0,1} &= x^{d_n} \tilde{P}_{2^n-1}(x), \\ M_n(x)_{0,0} &= x^{d_n} \tilde{Q}_{2^n-1}(x), \end{aligned}$$

for some positive integer d_n . Hence,

$$(2.2) \quad \text{CF}_{2^{2^n}-1}(a, b) = \frac{\tilde{P}_{2^{2^n}-1}(x)}{\tilde{Q}_{2^{2^n}-1}(x)} = \frac{M_{2^n}(x)_{0,1}}{M_{2^n}(x)_{0,0}}.$$

Therefore, by the convergence theorem of continued fraction, the algebraicity of $\text{CF}(a, b)$ will be established if it is shown that both $M_{2^n}(x)_{0,1}$ and $M_{2^n}(x)_{0,0}$ converge to algebraic series in $\mathbb{F}_2[[x]]$.

Actually, we will prove that for all $0 \leq i, j \leq 1$ the four sequences $(M_{2^n}(x)_{i,j})_n$, $(M_{2^{n+1}}(x)_{i,j})_n$, $(W_{2^n}(x)_{i,j})_n$, and $(W_{2^{n+1}}(x)_{i,j})_n$ converge to algebraic series in $\mathbb{F}_2[[x]]$. For this purpose, we define four 2×2 matrices M^e, M^o, W^e, W^o as follows: For each $T \in \{M^e, M^o, W^e, W^o\}$ and $0 \leq i, j \leq 1$, $T_{i,j}$ is defined to be the unique solution in $\mathbb{F}_2[[x]]$ of the polynomial $\phi(T, i, j)$ under certain initial conditions; the polynomials $\phi(T, i, j)$ and initial conditions are given in Section 3. We will prove that these four matrices, whose components are algebraic by definition, are the limits of $(M_{2^n}(x))_n$, $(M_{2^{n+1}}(x))_n$, $(W_{2^n}(x))_n$, and $(W_{2^{n+1}}(x))_n$.

Let us explain how the polynomials $\phi(T, i, j)$ and initial conditions are found, and why the solutions exist and are unique. For $0 \leq i, j \leq 1$, the coefficients of the polynomial $\phi(M^e, i, j)$ (resp. $\phi(M^o, i, j)$, $\phi(W^e, i, j)$, and $\phi(W^o, i, j)$) are the Padé-Hermite approximants of type

$$(175, 175, 175, 175, 175)$$

of the vector

$$(1, T^3, T^6, T^9, T^{12}),$$

where $T = M_{12,i,j}$ (resp. $M_{11,i,j}$, $W_{12,i,j}$, and $W_{11,i,j}$), found by the Derksen algorithm. We take the first 8 terms of $M_{12,i,j}$ (resp. $M_{11,i,j}$, $W_{12,i,j}$, and $W_{11,i,j}$) as initial conditions for $\phi(M^e, i, j)$ (resp. $\phi(M^o, i, j)$, $\phi(W^e, i, j)$, and $\phi(W^o, i, j)$). The following fact will be used to ensure that the solution exists and is unique: let $P(x, y) \in \mathbb{F}_2[x, y]$ and for each series $f(x) = \sum_{n=0}^{\infty} a_n x^n \in \mathbb{F}_2[[x]]$ denote the partial sum $\sum_{j=0}^{n-1} a_j x^j$ by $f_n(x)$ for $n \geq 0$. If for some $n \geq 0$ and $a_0, a_1, \dots, a_{n-1} \in \mathbb{F}_2$ $P(x, \sum_{j=0}^{n-1} a_j x^j) = O(x^n)$ and $Q(x, y) := P(x, \sum_{j=0}^{n-1} a_j x^j + x^n y)$ can be written as $x^m \sum_{j=0}^{\infty} q_j(x) y^j$ where $q_j(x)$ are polynomials for $j \geq 0$, $q_1(0) = 1$, and $q_j(0) = 0$ for $j > 1$, then there exists a unique solution $f(x) \in \mathbb{F}_2[[x]]$ of $P(x, f(x)) = 0$ that satisfies the initial condition $f_n(x) = \sum_{j=0}^{n-1} a_j x^j$.

We state two lemmas concerning the four matrices M^e, M^o, W^e, W^o . The first one is about relations between them; the second, about the structure of the each matrix.

Lemma 2.1. *We have*

$$(2.3) \quad M^e = W^o \cdot M^o,$$

$$(2.4) \quad M^o = W^e \cdot M^e,$$

$$(2.5) \quad W^e = M^o \cdot W^o,$$

$$(2.6) \quad W^o = M^e \cdot W^e.$$

Proof. We give the proof of the identity

$$M_{0,0}^e = W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o,$$

the proofs of the others being similar.

First, we compute the minimal polynomials of $W_{0,0}^o M_{0,0}^o$ and $W_{0,1}^o M_{1,0}^o$. We know that

$$P(x, y) = \text{Res}_z (\phi(W^o, 0, 0)(x, z), z^{12} \cdot \phi(M^o, 0, 0)(x, y/z))$$

is an annihilating polynomial of $W_{0,0}^o M_{0,0}^o$. We use Padé-Hermite approximation to find a candidate for the minimal polynomial of $W_{0,0}^o M_{0,0}^o$, that will be called $\phi_0(x, y)$. To prove that $\phi_0(x, y)$ is indeed the minimal polynomial, it suffices to prove that it is an irreducible factor of $P(x, y)$ of multiplicity m and that $Q(x, y) := P(x, y)/\phi_0(x, y)^m$ is not an annihilating polynomial of $W_{0,0}^o M_{0,0}^o$. We verify the first point directly. For the second point, we truncate $W_{0,0}^o M_{0,0}^o$ to order 630 and substitute it for y in $Q(x, y)$. We get a series of valuation less than 630, which proves that $Q(x, y)$ is not an annihilating polynomial of $W_{0,0}^o M_{0,0}^o$. We find the minimal polynomial $\phi_1(x, y)$ of $W_{0,1}^o M_{1,0}^o$ in a similar way.

Now we prove that $\phi(M^e, 0, 0)$ is the minimal polynomial of $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$. We know that

$$S(x, y) = \text{Res}_z (\phi_0(x, z), \phi_1(x, y + z))$$

is an annihilating of $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$. We verify that $\phi(M^e, 0, 0)$ is an irreducible factor of $S(x, y)$ of multiplicity μ , and that $Q(x, y) := S(x, y)/\phi(M^e, 0, 0)^\mu$ is not an annihilating polynomial of $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$. To see the last point, we truncate $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$ to order 770 and substitute it for y in $Q(x, y)$. We get a series of valuation less than 770, and therefore $Q(x, y)$ is not an annihilating polynomial of $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$.

Finally, the first 16 terms of $M_{0,0}^e$ and $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$ coincide. As these first terms determine a unique solution of $\phi(M^e, 0, 0)$, we know that the two series are one and the same. $\square$

Define

$$R^e = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \quad \text{and} \quad R^o = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}.$$

Lemma 2.2. *For all integers $k \geq 2$ and $u = 2^{2k-1}$, the following identities hold.*

$$\begin{aligned} M^e[:14u] &= M^e[:7u] + x^{2u} M^e[:5u] + x^{3u} M^e[:4u] + x^{4u} M^e[:3u] + x^{5u} M^e[:2u] \\ &\quad + x^{2u} M^e[:7u] + x^{4u} M^e[:5u] + x^{5u} M^e[:4u] + x^{6u} M^e[:3u] \\ &\quad + x^{7u} M^e[:2u] + x^{3u} M^e[:7u] + x^{5u} M^e[:5u] + x^{6u} M^e[:4u] \\ &\quad + x^{7u} M^e[:3u] + x^{8u} M^e[:2u] + x^{5u} M^e[:7u] + x^{7u} M^e[:5u] \\ &\quad + x^{8u} M^e[:4u] + x^{9u} M^e[:3u] + x^{10u} M^e[:2u] + x^{8u} M^e[:6u] \\ &\quad + x^{10u} M^e[:4u] + x^{12u} M^e[:2u] + (x^{7u} + x^{9u} + x^{10u} + x^{12u}) R^e, \end{aligned}$$

$$\begin{aligned}
W^e[:14u] &= W^e[:7u] + x^{2u}W^e[:5u] + x^{3u}W^e[:4u] + x^{4u}W^e[:3u] + x^{5u}W^e[:2u] \\
&\quad + x^{2u}W^e[:7u] + x^{4u}W^e[:5u] + x^{5u}W^e[:4u] + x^{6u}W^e[:3u] \\
&\quad + x^{7u}W^e[:2u] + x^{3u}W^e[:7u] + x^{5u}W^e[:5u] + x^{6u}W^e[:4u] \\
&\quad + x^{7u}W^e[:3u] + x^{8u}W^e[:2u] + x^{5u}W^e[:7u] + x^{7u}W^e[:5u] \\
&\quad + x^{8u}W^e[:4u] + x^{9u}W^e[:3u] + x^{10u}W^e[:2u] + x^{8u}W^e[:6u] \\
&\quad + x^{10u}W^e[:4u] + x^{12u}W^e[:2u] + (x^{7u} + x^{9u} + x^{10u} + x^{12u})R^e.
\end{aligned}$$

For all integers $k \geq 2$ and $u = 2^{2k}$, the following identities hold.

$$\begin{aligned}
M^o[:14u] &= M^o[:7u] + x^{2u}M^o[:5u] + x^{3u}M^o[:4u] + x^{4u}M^o[:3u] + x^{5u}M^o[:2u] \\
&\quad + x^{2u}M^o[:7u] + x^{4u}M^o[:5u] + x^{5u}M^o[:4u] + x^{6u}M^o[:3u] \\
&\quad + x^{7u}M^o[:2u] + x^{3u}M^o[:7u] + x^{5u}M^o[:5u] + x^{6u}M^o[:4u] \\
&\quad + x^{7u}M^o[:3u] + x^{8u}M^o[:2u] + x^{5u}M^o[:7u] + x^{7u}M^o[:5u] \\
&\quad + x^{8u}M^o[:4u] + x^{9u}M^o[:3u] + x^{10u}M^o[:2u] + x^{8u}M^o[:6u] \\
&\quad + x^{10u}M^o[:4u] + x^{12u}M^o[:2u] + (x^{7u} + x^{9u} + x^{10u} + x^{12u})R^o, \\
W^o[:14u] &= W^o[:7u] + x^{2u}W^o[:5u] + x^{3u}W^o[:4u] + x^{4u}W^o[:3u] + x^{5u}W^o[:2u] \\
&\quad + x^{2u}W^o[:7u] + x^{4u}W^o[:5u] + x^{5u}W^o[:4u] + x^{6u}W^o[:3u] \\
&\quad + x^{7u}W^o[:2u] + x^{3u}W^o[:7u] + x^{5u}W^o[:5u] + x^{6u}W^o[:4u] \\
&\quad + x^{7u}W^o[:3u] + x^{8u}W^o[:2u] + x^{5u}W^o[:7u] + x^{7u}W^o[:5u] \\
&\quad + x^{8u}W^o[:4u] + x^{9u}W^o[:3u] + x^{10u}W^o[:2u] + x^{8u}W^o[:6u] \\
&\quad + x^{10u}W^o[:4u] + x^{12u}W^o[:2u] + (x^{7u} + x^{9u} + x^{10u} + x^{12u})R^o.
\end{aligned}$$

Proof. To prove Lemma 2.2 we first construct an automaton for each sequence concerned, and then transform the conditions on infinitely many k 's into finitely many conditions on the states of the automaton. In the following, we will prove that for $T = M_{0,0}^o$, for all integer $k \geq 2$ and $u = 2^{2k}$,

$$\begin{aligned}
T[:14u] &= T[:7u] + x^{2u}T[:5u] + x^{3u}T[:4u] + x^{4u}T[:3u] + x^{5u}T[:2u] + x^{2u}T[:7u] \\
&\quad + x^{4u}T[:5u] + x^{5u}T[:4u] + x^{6u}T[:3u] + x^{7u}T[:2u] + x^{3u}T[:7u] \\
&\quad + x^{5u}T[:5u] + x^{6u}T[:4u] + x^{7u}T[:3u] + x^{8u}T[:2u] + x^{5u}T[:7u] \\
&\quad + x^{7u}T[:5u] + x^{8u}T[:4u] + x^{9u}T[:3u] + x^{10u}T[:2u] + x^{8u}T[:6u] \\
&\quad + x^{10u}T[:4u] + x^{12u}T[:2u].
\end{aligned}$$

The proof of the other 15 cases are similar. We break down the above identity into 7 parts:

$$\begin{aligned}
0 &= x^{7u}T[:u] + x^{5u}T[2u:3u] + x^{4u}T[3u:4u] + x^{3u}T[4u:5u] \\
&\quad + x^{2u}T[5u:6u] + T[7u:8u], \\
0 &= x^{8u}T[:u] + x^{7u}T[u:2u] + x^{5u}T[3u:4u] + x^{4u}T[4u:5u] \\
&\quad + x^{3u}T[5u:6u] + x^{2u}T[6u:7u] + T[8u:9u], \\
0 &= x^{9u}T[:u] + x^{8u}T[u:2u] + x^{6u}T[3u:4u] + x^{3u}T[6u:7u] + T[9u:10u], \\
0 &= x^{9u}T[u:2u] + x^{7u}T[3u:4u] + x^{5u}T[5u:6u] + T[10u:11u],
\end{aligned}$$

$$0 = x^{9u}T[2u:3u] + x^{7u}T[4u:5u] + x^{5u}T[6u:7u] + T[11u:12u],$$

$$0 = x^{12u}T[:u] + x^{10u}T[2u:3u] + x^{8u}T[4u:5u] + T[12u:13u],$$

$$0 = x^{12u}T[u:2u] + x^{10u}T[3u:4u] + x^{8u}T[5u:6u] + T[13u:14u],$$

which can be rewritten as

$$(2.7) \quad \begin{aligned} 0 &= T[[w]_2] + T[[10w]_2] + T[[11w]_2] + T[[100w]_2] + T[[101w]_2] \\ &\quad + T[[111w]_2], \end{aligned}$$

$$(2.8) \quad \begin{aligned} 0 &= T[[w]_2] + T[[1w]_2] + T[[11w]_2] + T[[100w]_2] + T[[101w]_2] \\ &\quad + T[[110w]_2] + T[[1000w]_2], \end{aligned}$$

$$(2.9) \quad 0 = T[[w]_2] + T[[1w]_2] + T[[11w]_2] + T[[110w]_2] + T[[1001w]_2],$$

$$(2.10) \quad 0 = T[[1w]_2] + T[[11w]_2] + T[[101w]_2] + T[[1010w]_2],$$

$$(2.11) \quad 0 = T[[10w]_2] + T[[100w]_2] + T[[110w]_2] + T[[1011w]_2],$$

$$(2.12) \quad 0 = T[[w]_2] + T[[10w]_2] + T[[100w]_2] + T[[1100w]_2],$$

$$(2.13) \quad 0 = T[[1w]_2] + T[[11w]_2] + T[[101w]_2] + T[[1101w]_2],$$

for all binary word w of length $2k$ and $w \neq 0^{2k}$, and

$$(2.14) \quad \begin{aligned} 0 &= T[[w]_2] + T[[10w]_2] + T[[11w]_2] + T[[100w]_2] + T[[101w]_2] \\ &\quad + T[[111w]_2], \end{aligned}$$

$$(2.15) \quad \begin{aligned} 0 &= T[[w]_2] + T[[1w]_2] + T[[11w]_2] + T[[100w]_2] + T[[101w]_2] \\ &\quad + T[[110w]_2] + T[[1000w]_2], \end{aligned}$$

$$(2.16) \quad 0 = T[[w]_2] + T[[1w]_2] + T[[11w]_2] + T[[110w]_2] + T[[1001w]_2],$$

$$(2.17) \quad 0 = T[[1w]_2] + T[[11w]_2] + T[[101w]_2] + T[[1010w]_2],$$

$$(2.18) \quad 0 = T[[10w]_2] + T[[100w]_2] + T[[110w]_2] + T[[1011w]_2],$$

$$(2.19) \quad 0 = T[[w]_2] + T[[10w]_2] + T[[100w]_2] + T[[1100w]_2],$$

$$(2.20) \quad 0 = T[[1w]_2] + T[[11w]_2] + T[[101w]_2] + T[[1101w]_2],$$

for $w = 0^{2k}$.

First we calculate an 2-automaton that generates T from its minimal polynomial and its first terms. This automaton has 489 states; its transition function and output function can be found in the annex. Let $A(s, w)$ denote the state reached after reading w from right to left starting from the state s , and τ the output function. Define

$$E_{2k} = \{A(i, w) : |w| = 2k, w \neq 0^{2k}\}.$$

Identities (2.7) through (2.20) can be written as

$$(2.21) \quad \begin{aligned} 0 &= \tau(A(s, \epsilon)) + \tau(A(s, 10)) + \tau(A(s, 11)) + \tau(A(s, 100)) \\ &\quad + \tau(A(s, 101)) + \tau(A(s, 111)), \end{aligned}$$

$$(2.22) \quad \begin{aligned} 0 &= \tau(A(s, \epsilon)) + \tau(A(s, 1)) + \tau(A(s, 11)) + \tau(A(s, 100)) \\ &\quad + \tau(A(s, 101)) + \tau(A(s, 110)) + \tau(A(s, 1000)), \end{aligned}$$

$$(2.23) \quad \begin{aligned} 0 &= \tau(A(s, \epsilon)) + \tau(A(s, 1)) + \tau(A(s, 11)) + \tau(A(s, 110)) \\ &\quad + \tau(A(s, 1001)), \end{aligned}$$

$$(2.24) \quad 0 = \tau(A(s, 1)) + \tau(A(s, 11)) + \tau(A(s, 101)) + \tau(A(s, 1010)),$$

$$(2.25) \quad 0 = \tau(A(s, 10)) + \tau(A(s, 100)) + \tau(A(s, 110)) + \tau(A(s, 1011)),$$

$$(2.26) \quad 0 = \tau(A(s, \epsilon)) + \tau(A(s, 10)) + \tau(A(s, 100)) + \tau(A(s, 1100)),$$

$$(2.27) \quad 0 = \tau(A(s, 1)) + \tau(A(s, 11)) + \tau(A(s, 101)) + \tau(A(s, 1101)),$$

for all $s \in E_{2k}$, and

$$(2.28) \quad \begin{aligned} 0 &= \tau(A(s, \epsilon)) + \tau(A(s, 10)) + \tau(A(s, 11)) + \tau(A(s, 100)) \\ &\quad + \tau(A(s, 101)) + \tau(A(s, 111)), \end{aligned}$$

$$(2.29) \quad \begin{aligned} 0 &= \tau(A(s, \epsilon)) + \tau(A(s, 1)) + \tau(A(s, 11)) + \tau(A(s, 100)) \\ &\quad + \tau(A(s, 101)) + \tau(A(s, 110)) + \tau(A(s, 1000)), \end{aligned}$$

$$(2.30) \quad \begin{aligned} 0 &= \tau(A(s, \epsilon)) + \tau(A(s, 1)) + \tau(A(s, 11)) + \tau(A(s, 110)) \\ &\quad + \tau(A(s, 1001)), \end{aligned}$$

$$(2.31) \quad 0 = \tau(A(s, 1)) + \tau(A(s, 11)) + \tau(A(s, 101)) + \tau(A(s, 1010)),$$

$$(2.32) \quad 0 = \tau(A(s, 10)) + \tau(A(s, 100)) + \tau(A(s, 110)) + \tau(A(s, 1011)),$$

$$(2.33) \quad 0 = \tau(A(s, \epsilon)) + \tau(A(s, 10)) + \tau(A(s, 100)) + \tau(A(s, 1100)),$$

$$(2.34) \quad 0 = \tau(A(s, 1)) + \tau(A(s, 11)) + \tau(A(s, 101)) + \tau(A(s, 1101)),$$

for $s = A(i, 0^{2k})$. We find that $(A(i, 0^{22}), E_{22}) = (A(i, 0^{14}), E_{14})$, so that we only have to verify that identities (2.21) through (2.34) hold for $2 \leq k \leq 11$, which turns out to be true. $\square$

In the following lemma, we express M_{2k} , M_{2k+1} , W_{2k} , and W_{2k+1} in terms of M^e , M^o , W^e , and W^o .

Lemma 2.3. *For all integer $k \geq 2$, and $u = 2^{2k-1}$,*

$$\begin{aligned} M_{2k} &= M^e[:7u] + x^{2u}M^e[:5u] + x^{3u}M^e[:4u] + x^{4u}M^e[:3u] + x^{5u}M^e[:2u] \\ &\quad + x^{7u}R^e, \\ W_{2k} &= W^e[:7u] + x^{2u}W^e[:5u] + x^{3u}W^e[:4u] + x^{4u}W^e[:3u] + x^{5u}W^e[:2u] \\ &\quad + x^{7u}R^e. \end{aligned}$$

For all integer $k \geq 2$, and $u = 2^{2k}$,

$$\begin{aligned} M_{2k+1} &= M^o[:7u] + x^{2u}M^o[:5u] + x^{3u}M^o[:4u] + x^{4u}M^o[:3u] + x^{5u}M^o[:2u] \\ &\quad + x^{7u}R^o, \\ W_{2k+1} &= W^o[:7u] + x^{2u}W^o[:5u] + x^{3u}W^o[:4u] + x^{4u}W^o[:3u] + x^{5u}W^o[:2u] \\ &\quad + x^{7u}R^o. \end{aligned}$$

Proof. We call the four identities in Lemma 2.3 also by the name M_{2k} , W_{2k} , M_{2k+1} , and W_{2k+1} . For $n = 2$, the identities can be verified directly. For $n \geq 2$, we claim that

$$\begin{aligned} M_{2k} \wedge W_{2k} &\Rightarrow M_{2k+1} \wedge W_{2k+1}, \\ M_{2k+1} \wedge W_{2k+1} &\Rightarrow M_{2k+2} \wedge W_{2k+2}. \end{aligned}$$

We give the proof of

$$M_{2k} \wedge W_{2k} \Rightarrow M_{2k+1},$$

the other ones being similar. Set $u = 2^{2k}$ and $v = 2^{2k-1}$. By definition and induction hypothesis, the left side of identity M_{2k+1} is equal to

$$(2.35) \quad \begin{aligned} W_{2k}M_{2k} = & (W^e[:7v] + x^{2v}W^e[:5v] + x^{3v}W^e[:4v] + x^{4v}W^e[:3v] \\ & + x^{5v}W^e[:2v] + x^{7u}R^e) \times (M^e[:7v] + x^{2v}M^e[:5v] + x^{3v}M^e[:4v] \\ & + x^{4v}M^e[:3v] + x^{5v}M^e[:2v] + x^{7u}R^e); \end{aligned}$$

Call this expression *lhs*. Note that both sides of identity M_{2k+1} have the same term of highest degree $x^{14v}R^o$. Therefore we only need to prove that their difference is $O(x^{14v})$. Using Lemma 2.1 it can be seen that the right side of identity M_{2k+1} is congruent, modulo x^{14v} , to

$$\begin{aligned} & W^e[:14v]M^e[:14v] + x^{4v}W^e[:10v]M^e[:10v] + x^{6v}W^e[:8v]M^e[:8v] \\ & + x^{8v}W^e[:6v]M^e[:6v] + x^{10v}W^e[:4v]M^e[:4v]. \end{aligned}$$

For all $n \leq 14$, replace the occurrences of $W^e[:n \cdot v]$ and $M^e[:n \cdot v]$ in the above expression by the reduction modulo $x^{n \cdot v}$ of the right side of the corresponding identity in Lemma 2.2 and get a new expression, which we call *rhs*. Define

$$\begin{aligned} X &:= x^v, \\ a_n &:= W^e[n \cdot v : (n+1) \cdot v] / X^n, \\ b_n &:= M^e[n \cdot v : (n+1) \cdot v] / X^n, \\ c &:= R^e. \end{aligned}$$

Using the notation introduced above, we can represent the expressions *lhs* (2.35) and *rhs* as polynomials in $\mathbb{F}_2[a_1, \dots, a_{14}, b_1, \dots, b_{14}, c][X]$. Note that it is not a problem that a_j commutes with b_k while W^e does not commute with M^e , because in the expressions concerned, the products of W^e -terms and M^e -terms are always in the same order. We let the computer do the simplification and check that the difference between these two polynomials is indeed $O(X^{14})$, which completes the proof. $\square$

Proof of Theorem 1.1. We prove the theorem for $\text{CF}(a, b)$; for $\text{CF}(b, a)$, the proof is similar. By Lemma 2.3, we have For all $0 \leq j, k \leq 1$,

$$\begin{aligned} \lim_{n \rightarrow \infty} M_{2n,j,k} &= M_{j,k}^e, \\ \lim_{n \rightarrow \infty} M_{2n+1,j,k} &= M_{j,k}^o, \\ \lim_{n \rightarrow \infty} W_{2n,j,k} &= W_{j,k}^e, \\ \lim_{n \rightarrow \infty} W_{2n+1,j,k} &= W_{j,k}^o. \end{aligned}$$

Let $z = 1/x$. By the convergence theorem and identity (2.2),

$$\text{CF}(a, b) = \frac{M_{0,1}^e(x)}{M_{0,0}^e(x)}.$$

By definition, $\phi(M^e, 0, 1)$ and $\phi(M^e, 0, 0)$ are minimal polynomials of $M_{0,1}^e$ and $M_{0,0}^e$. Therefore

$$P(x, y) = \text{Res}_t (\phi(M^e, 0, 1)(x, t), y^{12}\phi(M^e, 0, 1)(x, t/y))$$

is an annihilating polynomial of $f(x) = M_{0,1}^e/M_{0,0}^e$.

Define

$$Q(x, y) = q_4(x)y^4 + q_3(x)y^3 + q_2(x)y^2 + q_1(x)y + q_0(x),$$

where

$$\begin{aligned}
q_0(x) &= x^{18} + x^{16} + x^{15} + x^{13} + x^{12} + x^{11} + x^{10} + x^9 + x^8 + x^7, \\
q_1(x) &= x^{14} + x^{13} + x^{12} + x^4 + x^3 + x^2, \\
q_2(x) &= x^{16} + x^{10} + x^6 + 1, \\
q_3(x) &= x^{14} + x^{13} + x^{12} + x^4 + x^3 + x^2, \\
q_4(x) &= x^{18} + x^{16} + x^{15} + x^{13} + x^{11} + x^9 + x^7 + x^6 + x^4.
\end{aligned}$$

The polynomial $Q(x, y)$ is the candidate for the minimal polynomial of $f(x)$ found by Padé-Hermite approximation. To prove that it is indeed the minimal polynomial of $f(x)$, we only need to prove that it is an irreducible factor of $P(x, y)$ of multiplicity m and $R(x, y) := P(x, y)/Q(x, y)^m$ is not an annihilating polynomial of $f(x)$. We verify the first point directly. For the second point, we find that when we truncate $f(x)$ to order 216, and substitute it for y in $R(z, y)$, we get a series with valuation smaller than 216, which proves that $R(z, y)$ is not an annihilating polynomial of $f(x)$. Finally, $z^{12}Q(1/z, y)$ is the minimal polynomial of $\text{CF}(a, b) = f(1/z)$. $\square$

3. ANNEXE

All 16 polynomials are of the form

$$p_0(x) + p_3(x)y^3 + p_6(x)y^6 + p_9(x)y^9 + p_{12}(x)y^{12}.$$

The coefficients $p_j(x)$ and the 16 initial terms to determine the solutions uniquely are given below.

For $\phi(M^e, 0, 0)$:

$$\begin{aligned}
p_0(x) &= x^{108} + x^{104} + x^{102} + x^{100} + x^{98} + x^{92} + x^{90} + x^{88} + x^{86} + x^{82} + x^{78} \\
&\quad + x^{74} + x^{72} + x^{66} + x^{60} + x^{58} + x^{54} + x^{50} + x^{46} + x^{44} + x^{42} + x^{40} \\
&\quad + x^{34} + x^{32} + x^{30} + x^{28} + x^{24}, \\
p_3(x) &= x^{96} + x^{94} + x^{84} + x^{78} + x^{72} + x^{68} + x^{66} + x^{60} + x^{58} + x^{54} + x^{44} \\
&\quad + x^{38} + x^{32} + x^{28} + x^{26} + x^{20} + x^{18} + x^{16}, \\
p_6(x) &= x^{100} + x^{96} + x^{92} + x^{86} + x^{78} + x^{76} + x^{68} + x^{66} + x^{64} + x^{62} + x^{60} \\
&\quad + x^{58} + x^{52} + x^{48} + x^{44} + x^{40} + x^{38} + x^{36} + x^{28} + x^{26} + x^{24} + x^{22} \\
&\quad + x^{18} + x^{16} + x^8 + x^6 + x^4 + 1, \\
p_9(x) &= x^{92} + x^{90} + x^{84} + x^{80} + x^{76} + x^{70} + x^{66} + x^{60} + x^{56} + x^{50} + x^{46} \\
&\quad + x^{40} + x^{36} + x^{30} + x^{26} + x^{20} + x^{16} + x^{12} + x^6 + x^4, \\
p_{12}(x) &= x^{96} + x^{88} + x^{80} + x^{16} + x^8 + 1,
\end{aligned}$$

and the initial terms are $[1, 0, 0, 0, 0, 0, 0, 0, 1, 0, 1, 0, 0, 0, 0, 0]$.

For $\phi(M^e, 0, 1)$:

$$\begin{aligned}
p_0(x) &= x^{99} + x^{96} + x^{95} + x^{92} + x^{91} + x^{90} + x^{89} + x^{88} + x^{87} + x^{85} + x^{83} \\
&\quad + x^{82} + x^{78} + x^{74} + x^{71} + x^{68} + x^{63} + x^{60} + x^{59} + x^{58} + x^{55} + x^{54} \\
&\quad + x^{53} + x^{51} + x^{48} + x^{44} + x^{43} + x^{42} + x^{41} + x^{36} + x^{35} + x^{34} + x^{33}, \\
p_3(x) &= x^{69} + x^{68} + x^{67} + x^{66} + x^{63} + x^{62} + x^{57} + x^{56} + x^{55} + x^{54} + x^{53}
\end{aligned}$$

$$\begin{aligned}
& + x^{52} + x^{51} + x^{50} + x^{49} + x^{48} + x^{47} + x^{46} + x^{45} + x^{44} + x^{43} + x^{42} \\
& + x^{41} + x^{40} + x^{39} + x^{38} + x^{37} + x^{36} + x^{35} + x^{34} + x^{33} + x^{32} + x^{31} \\
& + x^{30} + x^{25} + x^{24} + x^{21} + x^{20} + x^{19} + x^{18}, \\
p_6(x) &= x^{75} + x^{73} + x^{72} + x^{70} + x^{69} + x^{66} + x^{65} + x^{62} + x^{61} + x^{58} + x^{57} \\
& + x^{55} + x^{54} + x^{52} + x^{35} + x^{33} + x^{32} + x^{30} + x^{29} + x^{26} + x^{25} + x^{22} \\
& + x^{21} + x^{18} + x^{17} + x^{15} + x^{14} + x^{12}, \\
p_9(x) &= x^{81} + x^{80} + x^{73} + x^{72} + x^{71} + x^{70} + x^{69} + x^{68} + x^{65} + x^{64} + x^{63} \\
& + x^{62} + x^{59} + x^{58} + x^{57} + x^{56} + x^{55} + x^{54} + x^{47} + x^{46} + x^{41} + x^{40} \\
& + x^{33} + x^{32} + x^{31} + x^{30} + x^{29} + x^{28} + x^{25} + x^{24} + x^{23} + x^{22} + x^{19} \\
& + x^{18} + x^{17} + x^{16} + x^{15} + x^{14} + x^7 + x^6, \\
p_{12}(x) &= x^{87} + x^{84} + x^{79} + x^{76} + x^{75} + x^{72} + x^{71} + x^{68} + x^{67} + x^{64} + x^{63} \\
& + x^{60} + x^{59} + x^{56} + x^{55} + x^{52} + x^{51} + x^{48} + x^{47} + x^{44} + x^{43} + x^{40} \\
& + x^{39} + x^{36} + x^{35} + x^{32} + x^{31} + x^{28} + x^{27} + x^{24} + x^{23} + x^{20} + x^{19} \\
& + x^{16} + x^{15} + x^{12} + x^{11} + x^8 + x^3 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 1, 1, 1, 1, 1, 1, 0, 1, 0, 0, 1, 0, 0]$.

For $\phi(M^e, 1, 0)$:

$$\begin{aligned}
p_0(x) &= x^{99} + x^{96} + x^{95} + x^{92} + x^{91} + x^{90} + x^{89} + x^{88} + x^{87} + x^{85} + x^{83} \\
& + x^{82} + x^{78} + x^{74} + x^{71} + x^{68} + x^{63} + x^{60} + x^{59} + x^{58} + x^{55} + x^{54} \\
& + x^{53} + x^{51} + x^{48} + x^{44} + x^{43} + x^{42} + x^{41} + x^{36} + x^{35} + x^{34} + x^{33}, \\
p_3(x) &= x^{69} + x^{68} + x^{67} + x^{66} + x^{63} + x^{62} + x^{57} + x^{56} + x^{55} + x^{54} + x^{53} \\
& + x^{52} + x^{51} + x^{50} + x^{49} + x^{48} + x^{47} + x^{46} + x^{45} + x^{44} + x^{43} + x^{42} \\
& + x^{41} + x^{40} + x^{39} + x^{38} + x^{37} + x^{36} + x^{35} + x^{34} + x^{33} + x^{32} + x^{31} \\
& + x^{30} + x^{25} + x^{24} + x^{21} + x^{20} + x^{19} + x^{18}, \\
p_6(x) &= x^{75} + x^{73} + x^{72} + x^{70} + x^{69} + x^{66} + x^{65} + x^{62} + x^{61} + x^{58} + x^{57} \\
& + x^{55} + x^{54} + x^{52} + x^{35} + x^{33} + x^{32} + x^{30} + x^{29} + x^{26} + x^{25} + x^{22} \\
& + x^{21} + x^{18} + x^{17} + x^{15} + x^{14} + x^{12}, \\
p_9(x) &= x^{81} + x^{80} + x^{73} + x^{72} + x^{71} + x^{70} + x^{69} + x^{68} + x^{65} + x^{64} + x^{63} \\
& + x^{62} + x^{59} + x^{58} + x^{57} + x^{56} + x^{55} + x^{54} + x^{47} + x^{46} + x^{41} + x^{40} \\
& + x^{33} + x^{32} + x^{31} + x^{30} + x^{29} + x^{28} + x^{25} + x^{24} + x^{23} + x^{22} + x^{19} \\
& + x^{18} + x^{17} + x^{16} + x^{15} + x^{14} + x^7 + x^6, \\
p_{12}(x) &= x^{87} + x^{84} + x^{79} + x^{76} + x^{75} + x^{72} + x^{71} + x^{68} + x^{67} + x^{64} + x^{63} \\
& + x^{60} + x^{59} + x^{56} + x^{55} + x^{52} + x^{51} + x^{48} + x^{47} + x^{44} + x^{43} + x^{40} \\
& + x^{39} + x^{36} + x^{35} + x^{32} + x^{31} + x^{28} + x^{27} + x^{24} + x^{23} + x^{20} + x^{19} \\
& + x^{16} + x^{15} + x^{12} + x^{11} + x^8 + x^3 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 1, 1, 1, 1, 1, 1, 0, 1, 0, 0, 1, 0, 0]$.

For $\phi(M^e, 1, 1)$:

$$\begin{aligned}
p_0(x) &= x^{90} + x^{86} + x^{82} + x^{76} + x^{74} + x^{70} + x^{68} + x^{52} + x^{48} + x^{44} + x^{42}, \\
p_3(x) &= x^{78} + x^{76} + x^{72} + x^{70} + x^{64} + x^{58} + x^{56} + x^{54} + x^{52} + x^{38} + x^{36} \\
&\quad + x^{32} + x^{30} + x^{24} + x^{18} + x^{16} + x^{14} + x^{12}, \\
p_6(x) &= x^{82} + x^{78} + x^{76} + x^{74} + x^{72} + x^{70} + x^{54} + x^{52} + x^{50} + x^{44} + x^{42} \\
&\quad + x^{40} + x^{38} + x^{36} + x^{34} + x^{32} + x^{30} + x^{14} + x^{12} + x^{10} + x^4 + 1, \\
p_9(x) &= x^{74} + x^{72} + x^{68} + x^{62} + x^{60} + x^{58} + x^{56} + x^{50} + x^{46} + x^{44} + x^{34} \\
&\quad + x^{32} + x^{28} + x^{22} + x^{20} + x^{18} + x^{16} + x^{10} + x^6 + x^4, \\
p_{12}(x) &= x^{78} + x^{72} + x^{70} + x^{64} + x^{62} + x^{56} + x^{54} + x^{48} + x^{46} + x^{40} + x^{38} \\
&\quad + x^{32} + x^{30} + x^{24} + x^{22} + x^{16} + x^{14} + x^8 + x^6 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 1, 0, 1, 0, 1, 0, 0, 0, 1, 0]$.

For $\phi(W^e, 0, 0)$:

$$\begin{aligned}
p_0(x) &= x^{90} + x^{86} + x^{82} + x^{78} + x^{76} + x^{72} + x^{64} + x^{62} + x^{60} + x^{58} + x^{54} \\
&\quad + x^{50} + x^{48} + x^{46} + x^{44} + x^{42} + x^{36} + x^{28} + x^{24}, \\
p_3(x) &= x^{78} + x^{76} + x^{72} + x^{70} + x^{66} + x^{58} + x^{56} + x^{38} + x^{36} + x^{32} + x^{30} \\
&\quad + x^{26} + x^{18} + x^{16}, \\
p_6(x) &= x^{82} + x^{78} + x^{76} + x^{74} + x^{72} + x^{60} + x^{58} + x^{54} + x^{52} + x^{50} + x^{48} \\
&\quad + x^{44} + x^{42} + x^{40} + x^{38} + x^{36} + x^{34} + x^{32} + x^{20} + x^{18} + x^{14} + x^{12} \\
&\quad + x^{10} + x^8 + x^4 + 1, \\
p_9(x) &= x^{74} + x^{72} + x^{68} + x^{62} + x^{60} + x^{58} + x^{56} + x^{50} + x^{46} + x^{44} + x^{34} \\
&\quad + x^{32} + x^{28} + x^{22} + x^{20} + x^{18} + x^{16} + x^{10} + x^6 + x^4, \\
p_{12}(x) &= x^{78} + x^{72} + x^{70} + x^{64} + x^{62} + x^{56} + x^{54} + x^{48} + x^{46} + x^{40} + x^{38} \\
&\quad + x^{32} + x^{30} + x^{24} + x^{22} + x^{16} + x^{14} + x^8 + x^6 + 1,
\end{aligned}$$

and the initial terms are $[1, 0, 0, 0, 1, 0, 1, 0, 0, 0, 1, 0, 1, 0, 1, 0]$.

For $\phi(W^e, 0, 1)$:

$$\begin{aligned}
p_0(x) &= x^{99} + x^{96} + x^{95} + x^{92} + x^{91} + x^{90} + x^{89} + x^{88} + x^{87} + x^{85} + x^{82} \\
&\quad + x^{81} + x^{80} + x^{77} + x^{73} + x^{70} + x^{63} + x^{60} + x^{56} + x^{54} + x^{51} + x^{50} \\
&\quad + x^{48} + x^{45} + x^{44} + x^{43} + x^{37} + x^{36} + x^{33}, \\
p_3(x) &= x^{69} + x^{68} + x^{67} + x^{66} + x^{63} + x^{62} + x^{57} + x^{56} + x^{55} + x^{54} + x^{53} \\
&\quad + x^{52} + x^{51} + x^{50} + x^{49} + x^{48} + x^{47} + x^{46} + x^{45} + x^{44} + x^{43} + x^{42} \\
&\quad + x^{41} + x^{40} + x^{39} + x^{38} + x^{37} + x^{36} + x^{35} + x^{34} + x^{33} + x^{32} + x^{31} \\
&\quad + x^{30} + x^{25} + x^{24} + x^{21} + x^{20} + x^{19} + x^{18}, \\
p_6(x) &= x^{75} + x^{73} + x^{72} + x^{70} + x^{69} + x^{66} + x^{65} + x^{62} + x^{61} + x^{58} + x^{57} \\
&\quad + x^{55} + x^{54} + x^{52} + x^{35} + x^{33} + x^{32} + x^{30} + x^{29} + x^{26} + x^{25} + x^{22} \\
&\quad + x^{21} + x^{18} + x^{17} + x^{15} + x^{14} + x^{12},
\end{aligned}$$

$$\begin{aligned}
p_9(x) &= x^{81} + x^{80} + x^{73} + x^{72} + x^{71} + x^{70} + x^{69} + x^{68} + x^{65} + x^{64} + x^{63} \\
&\quad + x^{62} + x^{59} + x^{58} + x^{57} + x^{56} + x^{55} + x^{54} + x^{47} + x^{46} + x^{41} + x^{40} \\
&\quad + x^{33} + x^{32} + x^{31} + x^{30} + x^{29} + x^{28} + x^{25} + x^{24} + x^{23} + x^{22} + x^{19} \\
&\quad + x^{18} + x^{17} + x^{16} + x^{15} + x^{14} + x^7 + x^6, \\
p_{12}(x) &= x^{87} + x^{84} + x^{79} + x^{76} + x^{75} + x^{72} + x^{71} + x^{68} + x^{67} + x^{64} + x^{63} \\
&\quad + x^{60} + x^{59} + x^{56} + x^{55} + x^{52} + x^{51} + x^{48} + x^{47} + x^{44} + x^{43} + x^{40} \\
&\quad + x^{39} + x^{36} + x^{35} + x^{32} + x^{31} + x^{28} + x^{27} + x^{24} + x^{23} + x^{20} + x^{19} \\
&\quad + x^{16} + x^{15} + x^{12} + x^{11} + x^8 + x^3 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 1, 1, 1, 0, 1, 1, 1, 1, 1, 0, 0]$.

For $\phi(W^e, 1, 0)$:

$$\begin{aligned}
p_0(x) &= x^{99} + x^{96} + x^{95} + x^{92} + x^{91} + x^{90} + x^{89} + x^{88} + x^{87} + x^{85} + x^{82} \\
&\quad + x^{81} + x^{80} + x^{77} + x^{73} + x^{70} + x^{63} + x^{60} + x^{56} + x^{54} + x^{51} + x^{50} \\
&\quad + x^{48} + x^{45} + x^{44} + x^{43} + x^{37} + x^{36} + x^{33}, \\
p_3(x) &= x^{69} + x^{68} + x^{67} + x^{66} + x^{63} + x^{62} + x^{57} + x^{56} + x^{55} + x^{54} + x^{53} \\
&\quad + x^{52} + x^{51} + x^{50} + x^{49} + x^{48} + x^{47} + x^{46} + x^{45} + x^{44} + x^{43} + x^{42} \\
&\quad + x^{41} + x^{40} + x^{39} + x^{38} + x^{37} + x^{36} + x^{35} + x^{34} + x^{33} + x^{32} + x^{31} \\
&\quad + x^{30} + x^{25} + x^{24} + x^{21} + x^{20} + x^{19} + x^{18}, \\
p_6(x) &= x^{75} + x^{73} + x^{72} + x^{70} + x^{69} + x^{66} + x^{65} + x^{62} + x^{61} + x^{58} + x^{57} \\
&\quad + x^{55} + x^{54} + x^{52} + x^{35} + x^{33} + x^{32} + x^{30} + x^{29} + x^{26} + x^{25} + x^{22} \\
&\quad + x^{21} + x^{18} + x^{17} + x^{15} + x^{14} + x^{12}, \\
p_9(x) &= x^{81} + x^{80} + x^{73} + x^{72} + x^{71} + x^{70} + x^{69} + x^{68} + x^{65} + x^{64} + x^{63} \\
&\quad + x^{62} + x^{59} + x^{58} + x^{57} + x^{56} + x^{55} + x^{54} + x^{47} + x^{46} + x^{41} + x^{40} \\
&\quad + x^{33} + x^{32} + x^{31} + x^{30} + x^{29} + x^{28} + x^{25} + x^{24} + x^{23} + x^{22} + x^{19} \\
&\quad + x^{18} + x^{17} + x^{16} + x^{15} + x^{14} + x^7 + x^6, \\
p_{12}(x) &= x^{87} + x^{84} + x^{79} + x^{76} + x^{75} + x^{72} + x^{71} + x^{68} + x^{67} + x^{64} + x^{63} \\
&\quad + x^{60} + x^{59} + x^{56} + x^{55} + x^{52} + x^{51} + x^{48} + x^{47} + x^{44} + x^{43} + x^{40} \\
&\quad + x^{39} + x^{36} + x^{35} + x^{32} + x^{31} + x^{28} + x^{27} + x^{24} + x^{23} + x^{20} + x^{19} \\
&\quad + x^{16} + x^{15} + x^{12} + x^{11} + x^8 + x^3 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 1, 1, 1, 0, 1, 1, 1, 1, 1, 0, 0]$.

For $\phi(W^e, 1, 1)$:

$$\begin{aligned}
p_0(x) &= x^{108} + x^{104} + x^{102} + x^{100} + x^{98} + x^{96} + x^{90} + x^{86} + x^{82} + x^{80} + x^{76} \\
&\quad + x^{68} + x^{66} + x^{64} + x^{60} + x^{56} + x^{54} + x^{52} + x^{48} + x^{46} + x^{42}, \\
p_3(x) &= x^{96} + x^{94} + x^{82} + x^{76} + x^{72} + x^{68} + x^{66} + x^{52} + x^{42} + x^{36} + x^{32} \\
&\quad + x^{28} + x^{26} + x^{16} + x^{14} + x^{12}, \\
p_6(x) &= x^{100} + x^{96} + x^{92} + x^{88} + x^{86} + x^{82} + x^{76} + x^{72} + x^{68} + x^{66} + x^{62}
\end{aligned}$$

$$\begin{aligned}
& + x^{60} + x^{54} + x^{52} + x^{48} + x^{44} + x^{42} + x^{40} + x^{36} + x^{32} + x^{28} + x^{26} \\
& + x^{22} + x^{16} + x^{14} + x^6 + x^4 + 1, \\
p_9(x) &= x^{92} + x^{90} + x^{84} + x^{80} + x^{76} + x^{70} + x^{66} + x^{60} + x^{56} + x^{50} + x^{46} \\
& + x^{40} + x^{36} + x^{30} + x^{26} + x^{20} + x^{16} + x^{12} + x^6 + x^4, \\
p_{12}(x) &= x^{96} + x^{88} + x^{80} + x^{16} + x^8 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 1, 0, 1, 0, 0, 0]$.

For $\phi(M^o, 0, 0)$:

$$\begin{aligned}
p_0(x) &= x^{99} + x^{96} + x^{95} + x^{93} + x^{92} + x^{91} + x^{90} + x^{88} + x^{86} + x^{84} + x^{83} \\
& + x^{82} + x^{80} + x^{79} + x^{78} + x^{76} + x^{72} + x^{68} + x^{67} + x^{64} + x^{63} + x^{60} \\
& + x^{58} + x^{55} + x^{53} + x^{51} + x^{50} + x^{48} + x^{47} + x^{45} + x^{44} + x^{38} + x^{35} \\
& + x^{33} + x^{31} + x^{27} + x^{24}, \\
p_3(x) &= x^{83} + x^{80} + x^{77} + x^{74} + x^{73} + x^{71} + x^{64} + x^{61} + x^{59} + x^{57} + x^{53} \\
& + x^{52} + x^{49} + x^{48} + x^{45} + x^{44} + x^{43} + x^{41} + x^{36} + x^{34} + x^{32} + x^{31} \\
& + x^{29} + x^{28} + x^{25} + x^{20} + x^{19} + x^{16}, \\
p_6(x) &= x^{91} + x^{88} + x^{85} + x^{83} + x^{82} + x^{80} + x^{79} + x^{78} + x^{75} + x^{74} + x^{72} \\
& + x^{70} + x^{69} + x^{67} + x^{66} + x^{65} + x^{63} + x^{61} + x^{60} + x^{53} + x^{51} + x^{49} \\
& + x^{44} + x^{42} + x^{41} + x^{40} + x^{39} + x^{38} + x^{37} + x^{36} + x^{35} + x^{34} + x^{33} \\
& + x^{30} + x^{28} + x^{27} + x^{26} + x^{24} + x^{23} + x^{17} + x^{16} + x^{12} + x^3 + 1, \\
p_9(x) &= x^{79} + x^{76} + x^{75} + x^{72} + x^{63} + x^{60} + x^{59} + x^{56} + x^{47} + x^{44} + x^{43} \\
& + x^{40} + x^{31} + x^{28} + x^{27} + x^{24} + x^{15} + x^{12} + x^{11} + x^8, \\
p_{12}(x) &= x^{87} + x^{84} + x^{71} + x^{68} + x^{67} + x^{64} + x^{55} + x^{52} + x^{51} + x^{48} + x^{39} \\
& + x^{36} + x^{35} + x^{32} + x^{23} + x^{20} + x^{19} + x^{16} + x^3 + 1,
\end{aligned}$$

and the initial terms are $[1, 0, 0, 0, 1, 0, 1, 1, 1, 1, 1, 0, 1, 0, 1, 1]$.

For $\phi(M^o, 0, 1)$:

$$\begin{aligned}
p_0(x) &= x^{94} + x^{92} + x^{86} + x^{85} + x^{84} + x^{83} + x^{81} + x^{80} + x^{77} + x^{74} + x^{73} \\
& + x^{70} + x^{69} + x^{68} + x^{66} + x^{65} + x^{59} + x^{56} + x^{55} + x^{54} + x^{52} + x^{50} \\
& + x^{48} + x^{47} + x^{46} + x^{45} + x^{44} + x^{43} + x^{41} + x^{38} + x^{36} + x^{34} + x^{33}, \\
p_3(x) &= x^{64} + x^{63} + x^{61} + x^{59} + x^{58} + x^{24} + x^{23} + x^{21} + x^{19} + x^{18}, \\
p_6(x) &= x^{70} + x^{64} + x^{62} + x^{56} + x^{54} + x^{50} + x^{48} + x^{46} + x^{44} + x^{42} + x^{40} \\
& + x^{38} + x^{36} + x^{34} + x^{32} + x^{28} + x^{26} + x^{20} + x^{18} + x^{12}, \\
p_9(x) &= x^{76} + x^{75} + x^{74} + x^{72} + x^{71} + x^{70} + x^{60} + x^{59} + x^{58} + x^{56} + x^{55} \\
& + x^{54} + x^{44} + x^{43} + x^{42} + x^{40} + x^{39} + x^{38} + x^{28} + x^{27} + x^{26} + x^{24} \\
& + x^{23} + x^{22} + x^{12} + x^{11} + x^{10} + x^8 + x^7 + x^6, \\
p_{12}(x) &= x^{82} + x^{80} + x^{78} + x^{76} + x^{74} + x^{72} + x^{70} + x^{68} + x^{62} + x^{60} + x^{58} \\
& + x^{56} + x^{54} + x^{52} + x^{46} + x^{44} + x^{42} + x^{40} + x^{38} + x^{36} + x^{30} + x^{28}
\end{aligned}$$

$$+ x^{26} + x^{24} + x^{22} + x^{20} + x^{14} + x^{12} + x^{10} + x^8 + x^6 + x^4 + x^2 + 1,$$

and the initial terms are $[0, 0, 1, 1, 1, 1, 0, 0, 1, 1, 0, 0, 0, 0, 0]$.

For $\phi(M^o, 1, 0)$:

$$\begin{aligned} p_0(x) &= x^{108} + x^{104} + x^{100} + x^{99} + x^{94} + x^{93} + x^{92} + x^{91} + x^{90} + x^{89} + x^{88} \\ &\quad + x^{85} + x^{83} + x^{76} + x^{75} + x^{67} + x^{66} + x^{65} + x^{64} + x^{63} + x^{61} + x^{60} \\ &\quad + x^{57} + x^{56} + x^{53} + x^{49} + x^{46} + x^{45} + x^{44} + x^{43} + x^{42} + x^{39} + x^{33}, \\ p_3(x) &= x^{78} + x^{77} + x^{76} + x^{72} + x^{71} + x^{70} + x^{66} + x^{65} + x^{64} + x^{60} + x^{59} \\ &\quad + x^{58} + x^{38} + x^{37} + x^{36} + x^{32} + x^{31} + x^{30} + x^{26} + x^{25} + x^{24} + x^{20} \\ &\quad + x^{19} + x^{18}, \\ p_6(x) &= x^{84} + x^{82} + x^{78} + x^{74} + x^{72} + x^{64} + x^{62} + x^{58} + x^{54} + x^{52} + x^{44} \\ &\quad + x^{42} + x^{38} + x^{34} + x^{32} + x^{24} + x^{22} + x^{18} + x^{14} + x^{12}, \\ p_9(x) &= x^{90} + x^{89} + x^{87} + x^{85} + x^{83} + x^{82} + x^{78} + x^{77} + x^{75} + x^{74} + x^{70} \\ &\quad + x^{69} + x^{67} + x^{66} + x^{62} + x^{61} + x^{59} + x^{58} + x^{54} + x^{53} + x^{51} + x^{50} \\ &\quad + x^{46} + x^{45} + x^{43} + x^{42} + x^{38} + x^{37} + x^{35} + x^{34} + x^{30} + x^{29} + x^{27} \\ &\quad + x^{26} + x^{22} + x^{21} + x^{19} + x^{18} + x^{14} + x^{13} + x^{11} + x^9 + x^7 + x^6, \\ p_{12}(x) &= x^{96} + x^{84} + x^{80} + x^{76} + x^{68} + x^{60} + x^{52} + x^{44} + x^{36} + x^{28} + x^{20} \\ &\quad + x^{16} + x^{12} + 1, \end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 1, 1, 1, 0, 1, 1, 1, 0, 1, 1, 0]$.

For $\phi(M^o, 1, 1)$:

$$\begin{aligned} p_0(x) &= x^{99} + x^{96} + x^{95} + x^{93} + x^{92} + x^{91} + x^{90} + x^{88} + x^{87} + x^{86} + x^{83} \\ &\quad + x^{82} + x^{81} + x^{80} + x^{77} + x^{75} + x^{74} + x^{73} + x^{70} + x^{68} + x^{63} + x^{62} \\ &\quad + x^{61} + x^{58} + x^{55} + x^{52} + x^{50} + x^{49} + x^{48} + x^{47} + x^{46} + x^{45} + x^{44} \\ &\quad + x^{43} + x^{42}, \\ p_3(x) &= x^{83} + x^{80} + x^{77} + x^{74} + x^{73} + x^{68} + x^{67} + x^{65} + x^{63} + x^{62} + x^{61} \\ &\quad + x^{60} + x^{57} + x^{56} + x^{53} + x^{52} + x^{49} + x^{48} + x^{45} + x^{44} + x^{43} + x^{41} \\ &\quad + x^{36} + x^{34} + x^{32} + x^{29} + x^{27} + x^{24} + x^{23} + x^{22}, \\ p_6(x) &= x^{91} + x^{88} + x^{85} + x^{83} + x^{82} + x^{80} + x^{78} + x^{76} + x^{75} + x^{74} + x^{73} \\ &\quad + x^{72} + x^{71} + x^{68} + x^{67} + x^{63} + x^{62} + x^{60} + x^{59} + x^{58} + x^{57} + x^{56} \\ &\quad + x^{54} + x^{53} + x^{49} + x^{48} + x^{44} + x^{42} + x^{41} + x^{40} + x^{38} + x^{37} + x^{35} \\ &\quad + x^{34} + x^{31} + x^{29} + x^{27} + x^{25} + x^{24} + x^{23} + x^{22} + x^{21} + x^{19} + x^{18} \\ &\quad + x^{14} + x^{12} + x^{11} + x^8 + x^3 + 1, \\ p_9(x) &= x^{79} + x^{76} + x^{75} + x^{72} + x^{63} + x^{60} + x^{59} + x^{56} + x^{47} + x^{44} + x^{43} \\ &\quad + x^{40} + x^{31} + x^{28} + x^{27} + x^{24} + x^{15} + x^{12} + x^{11} + x^8, \\ p_{12}(x) &= x^{87} + x^{84} + x^{71} + x^{68} + x^{67} + x^{64} + x^{55} + x^{52} + x^{51} + x^{48} + x^{39} \\ &\quad + x^{36} + x^{35} + x^{32} + x^{23} + x^{20} + x^{19} + x^{16} + x^3 + 1, \end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 0, 0, 1, 0, 1, 1, 0, 1, 0, 1, 1]$.

For $\phi(W^o, 0, 0)$:

$$\begin{aligned}
p_0(x) &= x^{99} + x^{96} + x^{95} + x^{93} + x^{92} + x^{91} + x^{90} + x^{88} + x^{86} + x^{84} + x^{83} \\
&\quad + x^{82} + x^{80} + x^{79} + x^{78} + x^{76} + x^{72} + x^{68} + x^{67} + x^{64} + x^{63} + x^{60} \\
&\quad + x^{58} + x^{55} + x^{53} + x^{51} + x^{50} + x^{48} + x^{47} + x^{45} + x^{44} + x^{38} + x^{35} \\
&\quad + x^{33} + x^{31} + x^{27} + x^{24}, \\
p_3(x) &= x^{83} + x^{80} + x^{77} + x^{74} + x^{73} + x^{71} + x^{64} + x^{61} + x^{59} + x^{57} + x^{53} \\
&\quad + x^{52} + x^{49} + x^{48} + x^{45} + x^{44} + x^{43} + x^{41} + x^{36} + x^{34} + x^{32} + x^{31} \\
&\quad + x^{29} + x^{28} + x^{25} + x^{20} + x^{19} + x^{16}, \\
p_6(x) &= x^{91} + x^{88} + x^{85} + x^{83} + x^{82} + x^{80} + x^{79} + x^{78} + x^{75} + x^{74} + x^{72} \\
&\quad + x^{70} + x^{69} + x^{67} + x^{66} + x^{65} + x^{63} + x^{61} + x^{60} + x^{53} + x^{51} + x^{49} \\
&\quad + x^{44} + x^{42} + x^{41} + x^{40} + x^{39} + x^{38} + x^{37} + x^{36} + x^{35} + x^{34} + x^{33} \\
&\quad + x^{30} + x^{28} + x^{27} + x^{26} + x^{24} + x^{23} + x^{17} + x^{16} + x^{12} + x^3 + 1, \\
p_9(x) &= x^{79} + x^{76} + x^{75} + x^{72} + x^{63} + x^{60} + x^{59} + x^{56} + x^{47} + x^{44} + x^{43} \\
&\quad + x^{40} + x^{31} + x^{28} + x^{27} + x^{24} + x^{15} + x^{12} + x^{11} + x^8, \\
p_{12}(x) &= x^{87} + x^{84} + x^{71} + x^{68} + x^{67} + x^{64} + x^{55} + x^{52} + x^{51} + x^{48} + x^{39} \\
&\quad + x^{36} + x^{35} + x^{32} + x^{23} + x^{20} + x^{19} + x^{16} + x^3 + 1,
\end{aligned}$$

and the initial terms are $[1, 0, 0, 0, 1, 0, 1, 1, 1, 1, 1, 0, 1, 0, 1, 1]$.

For $\phi(W^o, 0, 1)$:

$$\begin{aligned}
p_0(x) &= x^{108} + x^{104} + x^{100} + x^{99} + x^{94} + x^{93} + x^{92} + x^{91} + x^{90} + x^{89} + x^{88} \\
&\quad + x^{85} + x^{83} + x^{76} + x^{75} + x^{67} + x^{66} + x^{65} + x^{64} + x^{63} + x^{61} + x^{60} \\
&\quad + x^{57} + x^{56} + x^{53} + x^{49} + x^{46} + x^{45} + x^{44} + x^{43} + x^{42} + x^{39} + x^{33}, \\
p_3(x) &= x^{78} + x^{77} + x^{76} + x^{72} + x^{71} + x^{70} + x^{66} + x^{65} + x^{64} + x^{60} + x^{59} \\
&\quad + x^{58} + x^{38} + x^{37} + x^{36} + x^{32} + x^{31} + x^{30} + x^{26} + x^{25} + x^{24} + x^{20} \\
&\quad + x^{19} + x^{18}, \\
p_6(x) &= x^{84} + x^{82} + x^{78} + x^{74} + x^{72} + x^{64} + x^{62} + x^{58} + x^{54} + x^{52} + x^{44} \\
&\quad + x^{42} + x^{38} + x^{34} + x^{32} + x^{24} + x^{22} + x^{18} + x^{14} + x^{12}, \\
p_9(x) &= x^{90} + x^{89} + x^{87} + x^{85} + x^{83} + x^{82} + x^{78} + x^{77} + x^{75} + x^{74} + x^{70} \\
&\quad + x^{69} + x^{67} + x^{66} + x^{62} + x^{61} + x^{59} + x^{58} + x^{54} + x^{53} + x^{51} + x^{50} \\
&\quad + x^{46} + x^{45} + x^{43} + x^{42} + x^{38} + x^{37} + x^{35} + x^{34} + x^{30} + x^{29} + x^{27} \\
&\quad + x^{26} + x^{22} + x^{21} + x^{19} + x^{18} + x^{14} + x^{13} + x^{11} + x^9 + x^7 + x^6, \\
p_{12}(x) &= x^{96} + x^{84} + x^{80} + x^{76} + x^{68} + x^{60} + x^{52} + x^{44} + x^{36} + x^{28} + x^{20} \\
&\quad + x^{16} + x^{12} + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 1, 1, 1, 0, 1, 1, 1, 0, 1, 1, 0]$.

For $\phi(W^o, 1, 0)$:

$$p_0(x) = x^{94} + x^{92} + x^{86} + x^{85} + x^{84} + x^{83} + x^{81} + x^{80} + x^{77} + x^{74} + x^{73}$$

$$\begin{aligned}
& + x^{70} + x^{69} + x^{68} + x^{66} + x^{65} + x^{59} + x^{56} + x^{55} + x^{54} + x^{52} + x^{50} \\
& + x^{48} + x^{47} + x^{46} + x^{45} + x^{44} + x^{43} + x^{41} + x^{38} + x^{36} + x^{34} + x^{33}, \\
p_3(x) &= x^{64} + x^{63} + x^{61} + x^{59} + x^{58} + x^{24} + x^{23} + x^{21} + x^{19} + x^{18}, \\
p_6(x) &= x^{70} + x^{64} + x^{62} + x^{56} + x^{54} + x^{50} + x^{48} + x^{46} + x^{44} + x^{42} + x^{40} \\
& + x^{38} + x^{36} + x^{34} + x^{32} + x^{28} + x^{26} + x^{20} + x^{18} + x^{12}, \\
p_9(x) &= x^{76} + x^{75} + x^{74} + x^{72} + x^{71} + x^{70} + x^{60} + x^{59} + x^{58} + x^{56} + x^{55} \\
& + x^{54} + x^{44} + x^{43} + x^{42} + x^{40} + x^{39} + x^{38} + x^{28} + x^{27} + x^{26} + x^{24} \\
& + x^{23} + x^{22} + x^{12} + x^{11} + x^{10} + x^8 + x^7 + x^6, \\
p_{12}(x) &= x^{82} + x^{80} + x^{78} + x^{76} + x^{74} + x^{72} + x^{70} + x^{68} + x^{62} + x^{60} + x^{58} \\
& + x^{56} + x^{54} + x^{52} + x^{46} + x^{44} + x^{42} + x^{40} + x^{38} + x^{36} + x^{30} + x^{28} \\
& + x^{26} + x^{24} + x^{22} + x^{20} + x^{14} + x^{12} + x^{10} + x^8 + x^6 + x^4 + x^2 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 1, 1, 1, 1, 0, 0, 1, 1, 0, 0, 0, 0, 0, 0]$.

For $\phi(W^o, 1, 1)$:

$$\begin{aligned}
p_0(x) &= x^{99} + x^{96} + x^{95} + x^{93} + x^{92} + x^{91} + x^{90} + x^{88} + x^{87} + x^{86} + x^{83} \\
& + x^{82} + x^{81} + x^{80} + x^{77} + x^{75} + x^{74} + x^{73} + x^{70} + x^{68} + x^{63} + x^{62} \\
& + x^{61} + x^{58} + x^{55} + x^{52} + x^{50} + x^{49} + x^{48} + x^{47} + x^{46} + x^{45} + x^{44} \\
& + x^{43} + x^{42}, \\
p_3(x) &= x^{83} + x^{80} + x^{77} + x^{74} + x^{73} + x^{68} + x^{67} + x^{65} + x^{63} + x^{62} + x^{61} \\
& + x^{60} + x^{57} + x^{56} + x^{53} + x^{52} + x^{49} + x^{48} + x^{45} + x^{44} + x^{43} + x^{41} \\
& + x^{36} + x^{34} + x^{32} + x^{29} + x^{27} + x^{24} + x^{23} + x^{22}, \\
p_6(x) &= x^{91} + x^{88} + x^{85} + x^{83} + x^{82} + x^{80} + x^{78} + x^{76} + x^{75} + x^{74} + x^{73} \\
& + x^{72} + x^{71} + x^{68} + x^{67} + x^{63} + x^{62} + x^{60} + x^{59} + x^{58} + x^{57} + x^{56} \\
& + x^{54} + x^{53} + x^{49} + x^{48} + x^{44} + x^{42} + x^{41} + x^{40} + x^{38} + x^{37} + x^{35} \\
& + x^{34} + x^{31} + x^{29} + x^{27} + x^{25} + x^{24} + x^{23} + x^{22} + x^{21} + x^{19} + x^{18} \\
& + x^{14} + x^{12} + x^{11} + x^8 + x^3 + 1, \\
p_9(x) &= x^{79} + x^{76} + x^{75} + x^{72} + x^{63} + x^{60} + x^{59} + x^{56} + x^{47} + x^{44} + x^{43} \\
& + x^{40} + x^{31} + x^{28} + x^{27} + x^{24} + x^{15} + x^{12} + x^{11} + x^8, \\
p_{12}(x) &= x^{87} + x^{84} + x^{71} + x^{68} + x^{67} + x^{64} + x^{55} + x^{52} + x^{51} + x^{48} + x^{39} \\
& + x^{36} + x^{35} + x^{32} + x^{23} + x^{20} + x^{19} + x^{16} + x^3 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 0, 0, 1, 0, 1, 1, 0, 1, 0, 1, 1]$.

Below is the transition function and output function of an 2-automaton that generates $T = M_{0,0}^o$:

Transition function $(n, j) \mapsto \delta(n, j)$ ($\Lambda(n) := [\delta(n, 0), \delta(n, 1)]$):

n	$\Lambda(n)$	n	$\Lambda(n)$	n	$\Lambda(n)$	n	$\Lambda(n)$	n	$\Lambda(n)$
0	[1, 2]	98	[170, 128]	196	[304, 305]	294	[82, 366]	392	[33, 330]
1	[3, 4]	99	[52, 150]	197	[306, 49]	295	[367, 111]	393	[8, 445]

2	[5, 6]	100	[171, 172]	198	[180, 307]	296	[28, 368]	394	[7, 326]
3	[7, 8]	101	[173, 174]	199	[148, 180]	297	[369, 276]	395	[446, 156]
4	[9, 10]	102	[175, 176]	200	[186, 145]	298	[254, 370]	396	[447, 12]
5	[11, 12]	103	[172, 177]	201	[162, 160]	299	[24, 371]	397	[32, 335]
6	[13, 14]	104	[162, 178]	202	[180, 308]	300	[372, 373]	398	[448, 172]
7	[15, 16]	105	[57, 179]	203	[309, 60]	301	[72, 374]	399	[321, 449]
8	[17, 18]	106	[145, 180]	204	[44, 40]	302	[375, 279]	400	[310, 323]
9	[19, 20]	107	[181, 182]	205	[12, 13]	303	[205, 376]	401	[450, 59]
10	[21, 22]	108	[183, 176]	206	[43, 123]	304	[377, 280]	402	[451, 326]
11	[23, 24]	109	[37, 184]	207	[310, 311]	305	[279, 378]	403	[439, 14]
12	[25, 26]	110	[185, 38]	208	[312, 47]	306	[379, 191]	404	[115, 358]
13	[27, 28]	111	[186, 148]	209	[144, 54]	307	[80, 380]	405	[452, 333]
14	[29, 30]	112	[187, 159]	210	[50, 313]	308	[238, 381]	406	[326, 445]
15	[31, 32]	113	[60, 153]	211	[135, 60]	309	[382, 84]	407	[8, 327]
16	[33, 34]	114	[188, 189]	212	[120, 314]	310	[383, 91]	408	[311, 316]
17	[35, 36]	115	[190, 191]	213	[301, 42]	311	[384, 385]	409	[317, 349]
18	[37, 38]	116	[192, 21]	214	[311, 315]	312	[386, 26]	410	[361, 449]
19	[39, 40]	117	[193, 194]	215	[124, 43]	313	[264, 225]	411	[315, 314]
20	[41, 42]	118	[195, 83]	216	[316, 121]	314	[385, 387]	412	[41, 361]
21	[43, 44]	119	[76, 196]	217	[122, 44]	315	[388, 389]	413	[300, 41]
22	[45, 41]	120	[197, 198]	218	[317, 12]	316	[271, 73]	414	[443, 136]
23	[46, 47]	121	[99, 199]	219	[318, 180]	317	[390, 27]	415	[360, 449]
24	[48, 49]	122	[200, 77]	220	[179, 156]	318	[391, 80]	416	[357, 122]
25	[50, 51]	123	[201, 202]	221	[154, 135]	319	[392, 393]	417	[453, 151]
26	[52, 48]	124	[203, 85]	222	[149, 179]	320	[394, 231]	418	[51, 55]
27	[53, 54]	125	[204, 205]	223	[58, 319]	321	[395, 92]	419	[135, 152]
28	[55, 56]	126	[206, 207]	224	[136, 56]	322	[243, 23]	420	[454, 51]
29	[57, 58]	127	[208, 209]	225	[320, 144]	323	[275, 105]	421	[364, 326]
30	[59, 60]	128	[210, 29]	226	[51, 136]	324	[70, 200]	422	[42, 322]
31	[61, 62]	129	[26, 211]	227	[321, 322]	325	[288, 396]	423	[447, 349]
32	[63, 17]	130	[212, 213]	228	[323, 315]	326	[94, 260]	424	[356, 121]
33	[64, 65]	131	[214, 215]	229	[324, 119]	327	[16, 397]	425	[455, 54]
34	[65, 66]	132	[216, 214]	230	[118, 165]	328	[398, 259]	426	[331, 136]
35	[67, 68]	133	[217, 218]	231	[116, 325]	329	[399, 205]	427	[167, 339]
36	[69, 70]	134	[219, 220]	232	[326, 327]	330	[228, 400]	428	[328, 32]
37	[71, 72]	135	[221, 222]	233	[46, 141]	331	[401, 385]	429	[456, 160]
38	[73, 74]	136	[223, 224]	234	[328, 115]	332	[402, 108]	430	[146, 307]
39	[75, 76]	137	[225, 226]	235	[329, 330]	333	[273, 230]	431	[297, 14]
40	[77, 78]	138	[227, 228]	236	[173, 296]	334	[403, 214]	432	[299, 14]
41	[79, 80]	139	[229, 230]	237	[134, 59]	335	[62, 404]	433	[457, 8]
42	[81, 82]	140	[231, 232]	238	[143, 53]	336	[405, 235]	434	[458, 388]
43	[83, 84]	141	[209, 69]	239	[53, 140]	337	[108, 406]	435	[459, 408]
44	[85, 86]	142	[90, 233]	240	[152, 157]	338	[380, 85]	436	[460, 288]
45	[87, 88]	143	[234, 235]	241	[331, 55]	339	[4, 407]	437	[461, 206]

46	[89, 90]	144	[110, 236]	242	[147, 145]	340	[408, 409]	438	[462, 101]
47	[91, 92]	145	[237, 238]	243	[155, 48]	341	[410, 20]	439	[463, 240]
48	[93, 94]	146	[239, 240]	244	[332, 333]	342	[411, 21]	440	[464, 384]
49	[95, 96]	147	[241, 201]	245	[334, 10]	343	[412, 413]	441	[233, 104]
50	[97, 24]	148	[242, 243]	246	[183, 293]	344	[414, 223]	442	[30, 88]
51	[98, 99]	149	[244, 245]	247	[115, 335]	345	[415, 217]	443	[465, 105]
52	[100, 16]	150	[230, 109]	248	[148, 146]	346	[213, 416]	444	[252, 91]
53	[101, 93]	151	[246, 247]	249	[158, 141]	347	[417, 224]	445	[466, 467]
54	[102, 103]	152	[220, 198]	250	[336, 167]	348	[222, 418]	446	[468, 190]
55	[104, 76]	153	[112, 248]	251	[333, 337]	349	[68, 254]	447	[469, 104]
56	[105, 106]	154	[249, 81]	252	[313, 55]	350	[419, 388]	448	[470, 392]
57	[107, 108]	155	[250, 95]	253	[338, 148]	351	[420, 374]	449	[374, 419]
58	[109, 110]	156	[245, 251]	254	[320, 53]	352	[421, 256]	450	[226, 373]
59	[111, 112]	157	[84, 252]	255	[35, 294]	353	[422, 267]	451	[471, 466]
60	[88, 113]	158	[253, 254]	256	[329, 34]	354	[267, 423]	452	[472, 282]
61	[114, 115]	159	[255, 255]	257	[182, 339]	355	[424, 193]	453	[473, 17]
62	[116, 117]	160	[256, 257]	258	[114, 32]	356	[425, 113]	454	[199, 27]
63	[118, 119]	161	[258, 259]	259	[9, 340]	357	[426, 285]	455	[474, 273]
64	[120, 121]	162	[260, 261]	260	[164, 119]	358	[189, 427]	456	[475, 260]
65	[122, 123]	163	[262, 242]	261	[295, 174]	359	[428, 102]	457	[476, 246]
66	[124, 122]	164	[263, 264]	262	[341, 305]	360	[429, 69]	458	[312, 141]
67	[125, 126]	165	[86, 265]	263	[342, 343]	361	[373, 430]	459	[347, 40]
68	[127, 128]	166	[266, 267]	264	[344, 51]	362	[431, 412]	460	[123, 348]
69	[47, 129]	167	[18, 190]	265	[345, 346]	363	[432, 65]	461	[39, 348]
70	[130, 131]	168	[235, 268]	266	[347, 348]	364	[433, 95]	462	[477, 36]
71	[132, 133]	169	[269, 77]	267	[349, 299]	365	[434, 311]	463	[478, 178]
72	[134, 135]	170	[270, 271]	268	[333, 177]	366	[435, 343]	464	[437, 305]
73	[136, 137]	171	[272, 245]	269	[350, 46]	367	[436, 346]	465	[479, 145]
74	[138, 126]	172	[236, 273]	270	[351, 146]	368	[437, 133]	466	[334, 340]
75	[139, 140]	173	[274, 275]	271	[159, 178]	369	[438, 319]	467	[182, 168]
76	[141, 142]	174	[276, 277]	272	[352, 115]	370	[128, 141]	468	[480, 165]
77	[143, 144]	175	[278, 279]	273	[185, 184]	371	[47, 142]	469	[169, 152]
78	[145, 146]	176	[280, 281]	274	[353, 354]	372	[318, 146]	470	[181, 167]
79	[147, 148]	177	[282, 283]	275	[170, 46]	373	[144, 140]	471	[336, 182]
80	[149, 58]	178	[259, 284]	276	[146, 308]	374	[187, 162]	472	[332, 172]
81	[150, 151]	179	[191, 63]	277	[355, 303]	375	[13, 298]	473	[481, 184]
82	[152, 153]	180	[285, 276]	278	[356, 314]	376	[434, 323]	474	[482, 294]
83	[154, 59]	181	[286, 231]	279	[323, 316]	377	[439, 298]	475	[483, 38]
84	[155, 150]	182	[261, 101]	280	[349, 13]	378	[45, 301]	476	[171, 333]
85	[58, 156]	183	[287, 288]	281	[357, 43]	379	[440, 296]	477	[265, 25]
86	[60, 157]	184	[198, 289]	282	[166, 325]	380	[128, 47]	478	[484, 18]
87	[158, 47]	185	[290, 210]	283	[172, 337]	381	[59, 152]	479	[78, 285]
88	[159, 160]	186	[291, 239]	284	[32, 358]	382	[441, 313]	480	[485, 98]
89	[127, 46]	187	[292, 189]	285	[48, 151]	383	[351, 180]	481	[277, 89]

90	[161, 162]	188	[1, 167]	286	[359, 182]	384	[344, 313]	482	[486, 237]
91	[150, 49]	189	[175, 293]	287	[360, 322]	385	[161, 159]	483	[487, 79]
92	[55, 137]	190	[163, 294]	288	[301, 361]	386	[442, 135]	484	[488, 174]
93	[163, 36]	191	[295, 296]	289	[362, 354]	387	[313, 136]	485	[304, 133]
94	[164, 165]	192	[297, 298]	290	[363, 131]	388	[179, 319]	486	[435, 354]
95	[166, 117]	193	[12, 299]	291	[309, 152]	389	[141, 129]	487	[362, 343]
96	[167, 168]	194	[300, 301]	292	[364, 8]	390	[443, 55]	488	[74, 221]
97	[169, 60]	195	[302, 303]	293	[20, 365]	391	[444, 128]		

Output function $n \mapsto \tau(n)$:

n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$
0	0	70	0	140	0	210	1	280	0	350	1	420	1
1	0	71	0	141	1	211	1	281	1	351	1	421	1
2	0	72	0	142	0	212	0	282	1	352	1	422	0
3	0	73	0	143	1	213	0	283	0	353	0	423	1
4	0	74	1	144	1	214	0	284	0	354	0	424	1
5	0	75	0	145	0	215	0	285	1	355	1	425	1
6	0	76	1	146	0	216	0	286	1	356	1	426	1
7	0	77	1	147	1	217	0	287	0	357	1	427	0
8	0	78	0	148	1	218	0	288	0	358	1	428	1
9	0	79	1	149	0	219	0	289	0	359	1	429	0
10	1	80	0	150	0	220	1	290	1	360	0	430	0
11	0	81	0	151	0	221	1	291	0	361	1	431	0
12	1	82	1	152	1	222	0	292	1	362	0	432	1
13	0	83	1	153	1	223	0	293	1	363	1	433	1
14	1	84	0	154	1	224	0	294	1	364	1	434	0
15	0	85	0	155	0	225	0	295	1	365	0	435	1
16	0	86	0	156	1	226	1	296	1	366	1	436	1
17	0	87	1	157	0	227	1	297	0	367	1	437	0
18	0	88	0	158	1	228	1	298	0	368	0	438	0
19	0	89	0	159	0	229	0	299	1	369	0	439	1
20	1	90	0	160	1	230	0	300	0	370	1	440	0
21	1	91	0	161	0	231	0	301	0	371	0	441	0
22	1	92	1	162	1	232	1	302	0	372	0	442	0
23	0	93	1	163	1	233	0	303	1	373	1	443	0
24	1	94	1	164	1	234	1	304	1	374	1	444	0
25	1	95	1	165	0	235	1	305	1	375	0	445	1
26	1	96	0	166	1	236	0	306	0	376	0	446	1
27	0	97	1	167	0	237	0	307	0	377	1	447	1
28	1	98	1	168	1	238	1	308	1	378	1	448	1
29	1	99	1	169	1	239	0	309	0	379	0	449	1
30	0	100	1	170	1	240	1	310	1	380	1	450	1
31	0	101	0	171	1	241	1	311	0	381	0	451	0
32	0	102	1	172	0	242	1	312	0	382	0	452	0

33	0	103	0	173	0	243	0	313	0	383	1	453	1
34	0	104	1	174	0	244	0	314	0	384	0	454	1
35	0	105	1	175	1	245	1	315	1	385	0	455	1
36	0	106	0	176	0	246	0	316	0	386	0	456	0
37	0	107	1	177	1	247	1	317	0	387	0	457	1
38	0	108	0	178	0	248	1	318	0	388	1	458	0
39	0	109	0	179	1	249	1	319	0	389	1	459	1
40	1	110	1	180	1	250	0	320	0	390	0	460	1
41	1	111	0	181	1	251	1	321	1	391	0	461	0
42	0	112	1	182	1	252	0	322	0	392	0	462	0
43	1	113	0	183	0	253	1	323	1	393	0	463	1
44	0	114	0	184	1	254	0	324	0	394	0	464	0
45	1	115	1	185	1	255	0	325	0	395	1	465	0
46	0	116	0	186	0	256	1	326	1	396	1	466	1
47	0	117	1	187	1	257	1	327	0	397	0	467	1
48	1	118	0	188	0	258	0	328	1	398	1	468	1
49	1	119	1	189	1	259	0	329	1	399	1	469	1
50	1	120	0	190	1	260	1	330	1	400	1	470	1
51	1	121	1	191	1	261	1	331	1	401	1	471	0
52	1	122	0	192	0	262	1	332	0	402	0	472	0
53	0	123	1	193	1	263	1	333	1	403	1	473	1
54	1	124	0	194	0	264	0	334	1	404	1	474	1
55	1	125	0	195	0	265	0	335	0	405	0	475	0
56	1	126	1	196	1	266	1	336	0	406	1	476	1
57	1	127	0	197	0	267	0	337	0	407	0	477	0
58	0	128	1	198	1	268	1	338	1	408	0	478	1
59	0	129	1	199	1	269	1	339	0	409	0	479	0
60	0	130	0	200	0	270	1	340	0	410	1	480	1
61	0	131	0	201	1	271	0	341	1	411	1	481	1
62	0	132	0	202	1	272	1	342	1	412	1	482	1
63	0	133	0	203	0	273	1	343	1	413	0	483	0
64	0	134	0	204	0	274	0	344	0	414	0	484	1
65	0	135	1	205	1	275	1	345	0	415	0	485	1
66	0	136	0	206	1	276	0	346	0	416	1	486	1
67	0	137	0	207	1	277	1	347	1	417	1	487	0
68	0	138	1	208	0	278	1	348	0	418	1	488	1
69	0	139	0	209	1	279	1	349	0	419	1		

UNIVERSITÉ DE STRASBOURG, CNRS, IRMA UMR 7501, F-67000 STRASBOURG, FRANCE
Email address: `guoniu.han@unistra.fr`

SCHOOL OF MATHEMATICS AND STATISTICS, HUAZHONG UNIVERSITY OF SCIENCE AND TECHNOLOGY, WUHAN, PR CHINA
Email address: `huyining@hust.edu.cn`