

PROOF OF THE CONTINUED FRACTION CONJECTURE FOR

$$(a, b) = (z^2 + z, z^5 + z^4 + z + 1)$$

GUO-NIU HAN* AND YINING HU*

ABSTRACT. The continued fraction conjecture claims that the continued fraction defined by the (a, b) -Thue-Morse sequence is algebraic over $\mathbb{F}_2(x)$ for all pairs of distinct elements (a, b) in $\mathbb{F}_2[x] \setminus \mathbb{F}_2$. In this paper we prove the conjecture for $(a, b) = (z^2 + z, z^5 + z^4 + z + 1)$.

1. INTRODUCTION

Let $\mathbb{F}_2$ be the finite field of cardinality 2. Given a sequence of polynomials $\mathbf{a} = (a_0(z), a_1(z), a_2(z), \dots)$ of elements from $\mathbb{F}_2[z] \setminus \mathbb{F}_2$, we define the infinite continued fraction

$$(1.1) \quad \text{CF}(\mathbf{a}(z)) = \frac{1}{a_0(z) + \frac{1}{a_1(z) + \frac{1}{a_2(z) + \frac{1}{\ddots}}}}$$

as the limit of the finite continued fractions

$$(1.2) \quad \text{CF}_n(\mathbf{a}(z)) = \frac{1}{a_0(z) + \frac{1}{a_1(z) + \frac{1}{\ddots + \frac{1}{a_n(z)}}}} \in \mathbb{F}_2((1/z)).$$

Conversely, each power series in $\mathbb{F}_2[[1/z]]$ without constant term

$$(1.3) \quad c_{-1}z^{-1} + c_{-2}z^{-2} + c_{-3}z^{-3} + \dots$$

admits a continued fraction expansion of form (1.1).

Let a and b be two distinct elements from $\mathbb{F}_2[z] \setminus \mathbb{F}_2$. We consider the continued fraction (1.1) associated with the (a, b) -Thue-Morse sequence

$$\mathbf{t} = (t_0(z), t_1(z), t_2(z), \dots) = (a, b, b, a, b, a, a, b, \dots)$$

that we shall denote

$$(1.4) \quad \text{CF}(a, b) := \text{CF}(\mathbf{t}).$$

Date: 9/04/2020 22:45:36 .

2010 Mathematics Subject Classification. 11B85, 11J70, 11B50, 11Y65, 05A15.

Key words and phrases. automatic sequence, continued fraction, Thue-Morse sequence.

* This paper was automatically generated from a computer program developed by the authors, with 9.2 minutes CPU time used.

Theorem 1.1. *When $(a, b) = (z^2 + z, z^5 + z^4 + z + 1)$, the two power series $\text{CF}(a, b)$ and $\text{CF}(b, a)$ are algebraic over $\mathbb{F}_2(z)$, with minimal polynomials of the form*

$$p_4(z)y^4 + p_3(z)y^3 + p_2(z)y^2 + p_1(z)y + p_0(z) = 0$$

where, for $\text{CF}(a, b)$,

$$\begin{aligned} p_0(z) &= z^{17} + z^{16} + z^{15} + z^{11} + z^{10} + z^2 + z + 1, \\ p_1(z) &= z^{22} + z^{21} + z^{20} + z^{16} + z^{13} + z^{11} + z^6 + z^5 + z^3 + z, \\ p_2(z) &= z^{24} + z^{18} + z^{14} + z^{12} + z^8 + z^6 + z^4 + z^2, \\ p_3(z) &= z^{22} + z^{21} + z^{20} + z^{16} + z^{13} + z^{11} + z^6 + z^5 + z^3 + z, \\ p_4(z) &= z^{20} + z^{18} + z^{17} + z^{15} + z^{14} + z^{12} + z^{11} + z^{10} + z^6 + z^4 + z, \end{aligned}$$

and for $\text{CF}(b, a)$,

$$\begin{aligned} p_0(z) &= z^{17} + z^{15} + z^{11} + z^{10} + z^4 + z^2 + z, \\ p_1(z) &= z^{22} + z^{21} + z^{20} + z^{16} + z^{13} + z^{11} + z^6 + z^5 + z^3 + z, \\ p_2(z) &= z^{24} + z^{18} + z^{14} + z^{12} + z^8 + z^6 + z^4 + z^2, \\ p_3(z) &= z^{22} + z^{21} + z^{20} + z^{16} + z^{13} + z^{11} + z^6 + z^5 + z^3 + z, \\ p_4(z) &= z^{20} + z^{18} + z^{17} + z^{16} + z^{15} + z^{14} + z^{12} + z^{11} + z^{10} + z^6 + z + 1. \end{aligned}$$

2. PROOF

Define the sequences of polynomials $(P_n(z))_{n \geq 0}$ and $(Q_n(z))_{n \geq 0}$ by

$$\begin{pmatrix} P_n(z) & Q_n(z) \\ P_{n-1}(z) & Q_{n-1}(z) \end{pmatrix} := \begin{pmatrix} t_n(z) & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} t_{n-1}(z) & 1 \\ 1 & 0 \end{pmatrix} \cdots \begin{pmatrix} t_0(z) & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}.$$

By the basic properties of continued fractions, we have

$$(2.1) \quad \text{CF}_n(a, b) = \frac{P_n(z)}{Q_n(z)} \in \mathbb{F}_2((1/z)).$$

As formula (2.1) is not efficient for calculating $\text{CF}_n(a, b)$, a fast method is to be derived and described as follows. Define

$$\begin{aligned} M_n(x) &= x^{\deg(t_{2^n-1})} \begin{pmatrix} t_{2^n-1}(1/x) & 1 \\ 1 & 0 \end{pmatrix} \times x^{\deg(t_{2^n-2})} \begin{pmatrix} t_{2^n-2}(1/x) & 1 \\ 1 & 0 \end{pmatrix} \times \\ &\quad \cdots \times x^{\deg(t_0)} \begin{pmatrix} t_0(1/x) & 1 \\ 1 & 0 \end{pmatrix}, \\ W_n(x) &= x^{\deg(\bar{t}_{2^n-1})} \begin{pmatrix} \bar{t}_{2^n-1}(1/x) & 1 \\ 1 & 0 \end{pmatrix} \times x^{\deg(\bar{t}_{2^n-2})} \begin{pmatrix} \bar{t}_{2^n-2}(1/x) & 1 \\ 1 & 0 \end{pmatrix} \times \\ &\quad \cdots \times x^{\deg(\bar{t}_0)} \begin{pmatrix} \bar{t}_0(1/x) & 1 \\ 1 & 0 \end{pmatrix}, \end{aligned}$$

where $\bar{t}$ is the (b, a) -Thue-Morse sequence. By the properties of the Thue-Morse sequence, we have for $n \geq 0$

$$\begin{aligned} M_{n+1}(x) &= W_n(x) \cdot M_n(x), \\ W_{n+1}(x) &= M_n(x) \cdot W_n(x). \end{aligned}$$

Let $x := 1/z$. For each non-zero polynomial $P(z)$, define $\tilde{P}(x)$ to be $P(1/x)$. Then,

$$\text{CF}_n(a, b) = \frac{P_n(z)}{Q_n(z)} = \frac{\tilde{P}_n(x)}{\tilde{Q}_n(x)} \in \mathbb{F}_2((x)) = \mathbb{F}_2((1/z)).$$

Comparing the definition of $M_n(x)$ with the definition of $P_n(x)$ and $Q_n(x)$, we see that

$$\begin{aligned} M_n(x)_{0,1} &= x^{d_n} \tilde{P}_{2^n-1}(x), \\ M_n(x)_{0,0} &= x^{d_n} \tilde{Q}_{2^n-1}(x), \end{aligned}$$

for some positive integer d_n . Hence,

$$(2.2) \quad \text{CF}_{2^{2^n}-1}(a, b) = \frac{\tilde{P}_{2^{2^n}-1}(x)}{\tilde{Q}_{2^{2^n}-1}(x)} = \frac{M_{2^n}(x)_{0,1}}{M_{2^n}(x)_{0,0}}.$$

Therefore, by the convergence theorem of continued fraction, the algebraicity of $\text{CF}(a, b)$ will be established if it is shown that both $M_{2^n}(x)_{0,1}$ and $M_{2^n}(x)_{0,0}$ converge to algebraic series in $\mathbb{F}_2[[x]]$.

Actually, we will prove that for all $0 \leq i, j \leq 1$ the four sequences $(M_{2^n}(x)_{i,j})_n$, $(M_{2^{n+1}}(x)_{i,j})_n$, $(W_{2^n}(x)_{i,j})_n$, and $(W_{2^{n+1}}(x)_{i,j})_n$ converge to algebraic series in $\mathbb{F}_2[[x]]$. For this purpose, we define four 2×2 matrices M^e, M^o, W^e, W^o as follows: For each $T \in \{M^e, M^o, W^e, W^o\}$ and $0 \leq i, j \leq 1$, $T_{i,j}$ is defined to be the unique solution in $\mathbb{F}_2[[x]]$ of the polynomial $\phi(T, i, j)$ under certain initial conditions; the polynomials $\phi(T, i, j)$ and initial conditions are given in Section 3. We will prove that these four matrices, whose components are algebraic by definition, are the limits of $(M_{2^n}(x))_n$, $(M_{2^{n+1}}(x))_n$, $(W_{2^n}(x))_n$, and $(W_{2^{n+1}}(x))_n$.

Let us explain how the polynomials $\phi(T, i, j)$ and initial conditions are found, and why the solutions exist and are unique. For $0 \leq i, j \leq 1$, the coefficients of the polynomial $\phi(M^e, i, j)$ (resp. $\phi(M^o, i, j)$, $\phi(W^e, i, j)$, and $\phi(W^o, i, j)$) are the Padé-Hermite approximants of type

$$(175, 175, 175, 175, 175)$$

of the vector

$$(1, T^3, T^6, T^9, T^{12}),$$

where $T = M_{12,i,j}$ (resp. $M_{11,i,j}$, $W_{12,i,j}$, and $W_{11,i,j}$), found by the Derksen algorithm. We take the first 8 terms of $M_{12,i,j}$ (resp. $M_{11,i,j}$, $W_{12,i,j}$, and $W_{11,i,j}$) as initial conditions for $\phi(M^e, i, j)$ (resp. $\phi(M^o, i, j)$, $\phi(W^e, i, j)$, and $\phi(W^o, i, j)$). The following fact will be used to ensure that the solution exists and is unique: let $P(x, y) \in \mathbb{F}_2[x, y]$ and for each series $f(x) = \sum_{n=0}^{\infty} a_n x^n \in \mathbb{F}_2[[x]]$ denote the partial sum $\sum_{j=0}^{n-1} a_j x^j$ by $f_n(x)$ for $n \geq 0$. If for some $n \geq 0$ and $a_0, a_1, \dots, a_{n-1} \in \mathbb{F}_2$ $P(x, \sum_{j=0}^{n-1} a_j x^j) = O(x^n)$ and $Q(x, y) := P(x, \sum_{j=0}^{n-1} a_j x^j + x^n y)$ can be written as $x^m \sum_{j=0}^{\infty} q_j(x) y^j$ where $q_j(x)$ are polynomials for $j \geq 0$, $q_1(0) = 1$, and $q_j(0) = 0$ for $j > 1$, then there exists a unique solution $f(x) \in \mathbb{F}_2[[x]]$ of $P(x, f(x)) = 0$ that satisfies the initial condition $f_n(x) = \sum_{j=0}^{n-1} a_j x^j$.

We state two lemmas concerning the four matrices M^e, M^o, W^e, W^o . The first one is about relations between them; the second, about the structure of the each matrix.

Lemma 2.1. *We have*

$$(2.3) \quad M^e = W^o \cdot M^o,$$

$$(2.4) \quad M^o = W^e \cdot M^e,$$

$$(2.5) \quad W^e = M^o \cdot W^o,$$

$$(2.6) \quad W^o = M^e \cdot W^e.$$

Proof. We give the proof of the identity

$$M_{0,0}^e = W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o,$$

the proofs of the others being similar.

First, we compute the minimal polynomials of $W_{0,0}^o M_{0,0}^o$ and $W_{0,1}^o M_{1,0}^o$. We know that

$$P(x, y) = \text{Res}_z (\phi(W^o, 0, 0)(x, z), z^{12} \cdot \phi(M^o, 0, 0)(x, y/z))$$

is an annihilating polynomial of $W_{0,0}^o M_{0,0}^o$. We use Padé-Hermite approximation to find a candidate for the minimal polynomial of $W_{0,0}^o M_{0,0}^o$, that will be called $\phi_0(x, y)$. To prove that $\phi_0(x, y)$ is indeed the minimal polynomial, it suffices to prove that it is an irreducible factor of $P(x, y)$ of multiplicity m and that $Q(x, y) := P(x, y)/\phi_0(x, y)^m$ is not an annihilating polynomial of $W_{0,0}^o M_{0,0}^o$. We verify the first point directly. For the second point, we truncate $W_{0,0}^o M_{0,0}^o$ to order 630 and substitute it for y in $Q(x, y)$. We get a series of valuation less than 630, which proves that $Q(x, y)$ is not an annihilating polynomial of $W_{0,0}^o M_{0,0}^o$. We find the minimal polynomial $\phi_1(x, y)$ of $W_{0,1}^o M_{1,0}^o$ in a similar way.

Now we prove that $\phi(M^e, 0, 0)$ is the minimal polynomial of $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$. We know that

$$S(x, y) = \text{Res}_z (\phi_0(x, z), \phi_1(x, y + z))$$

is an annihilating of $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$. We verify that $\phi(M^e, 0, 0)$ is an irreducible factor of $S(x, y)$ of multiplicity μ , and that $Q(x, y) := S(x, y)/\phi(M^e, 0, 0)^\mu$ is not an annihilating polynomial of $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$. To see the last point, we truncate $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$ to order 770 and substitute it for y in $Q(x, y)$. We get a series of valuation less than 770, and therefore $Q(x, y)$ is not an annihilating polynomial of $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$.

Finally, the first 16 terms of $M_{0,0}^e$ and $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$ coincide. As these first terms determine a unique solution of $\phi(M^e, 0, 0)$, we know that the two series are one and the same. $\square$

Define

$$R^e = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \quad \text{and} \quad R^o = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}.$$

Lemma 2.2. *For all integers $k \geq 2$ and $u = 2^{2k-1}$, the following identities hold.*

$$\begin{aligned} M^e[:14u] &= M^e[:7u] + x^{2u} M^e[:5u] + x^{6u} M^e[:u] + x^{2u} M^e[:7u] + x^{4u} M^e[:5u] \\ &\quad + x^{8u} M^e[:u] + x^{4u} M^e[:7u] + x^{6u} M^e[:5u] + x^{10u} M^e[:u] \\ &\quad + x^{8u} M^e[:6u] + x^{10u} M^e[:4u] + x^{12u} M^e[:2u] \\ &\quad + (x^{7u} + x^{9u} + x^{11u}) R^e, \\ W^e[:14u] &= W^e[:7u] + x^{2u} W^e[:5u] + x^{6u} W^e[:u] + x^{2u} W^e[:7u] + x^{4u} W^e[:5u] \\ &\quad + x^{8u} W^e[:u] + x^{4u} W^e[:7u] + x^{6u} W^e[:5u] + x^{10u} W^e[:u] \end{aligned}$$

$$\begin{aligned}
& + x^{8u}W^e[:6u] + x^{10u}W^e[:4u] + x^{12u}W^e[:2u] \\
& + (x^{7u} + x^{9u} + x^{11u})R^e.
\end{aligned}$$

For all integers $k \geq 2$ and $u = 2^{2k}$, the following identities hold.

$$\begin{aligned}
M^o[:14u] &= M^o[:7u] + x^{2u}M^o[:5u] + x^{6u}M^o[:u] + x^{2u}M^o[:7u] + x^{4u}M^o[:5u] \\
& + x^{8u}M^o[:u] + x^{4u}M^o[:7u] + x^{6u}M^o[:5u] + x^{10u}M^o[:u] \\
& + x^{8u}M^o[:6u] + x^{10u}M^o[:4u] + x^{12u}M^o[:2u] \\
& + (x^{7u} + x^{9u} + x^{11u})R^o, \\
W^o[:14u] &= W^o[:7u] + x^{2u}W^o[:5u] + x^{6u}W^o[:u] + x^{2u}W^o[:7u] + x^{4u}W^o[:5u] \\
& + x^{8u}W^o[:u] + x^{4u}W^o[:7u] + x^{6u}W^o[:5u] + x^{10u}W^o[:u] \\
& + x^{8u}W^o[:6u] + x^{10u}W^o[:4u] + x^{12u}W^o[:2u] \\
& + (x^{7u} + x^{9u} + x^{11u})R^o.
\end{aligned}$$

Proof. To prove Lemma 2.2 we first construct an automaton for each sequence concerned, and then transform the conditions on infinitely many k 's into finitely many conditions on the states of the automaton. In the following, we will prove that for $T = M_{0,0}^o$, for all integer $k \geq 2$ and $u = 2^{2k}$,

$$\begin{aligned}
T[:14u] &= T[:7u] + x^{2u}T[:5u] + x^{6u}T[:u] + x^{2u}T[:7u] + x^{4u}T[:5u] + x^{8u}T[:u] \\
& + x^{4u}T[:7u] + x^{6u}T[:5u] + x^{10u}T[:u] + x^{8u}T[:6u] + x^{10u}T[:4u] \\
& + x^{12u}T[:2u].
\end{aligned}$$

The proof of the other 15 cases are similar. We break down the above identity into 7 parts:

$$\begin{aligned}
0 &= x^{6u}T[u:2u] + x^{2u}T[5u:6u] + T[7u:8u], \\
0 &= x^{6u}T[2u:3u] + x^{2u}T[6u:7u] + T[8u:9u], \\
0 &= x^{8u}T[u:2u] + x^{6u}T[3u:4u] + x^{4u}T[5u:6u] + T[9u:10u], \\
0 &= x^{8u}T[2u:3u] + x^{6u}T[4u:5u] + x^{4u}T[6u:7u] + T[10u:11u], \\
0 &= x^{10u}T[u:2u] + x^{8u}T[3u:4u] + T[11u:12u], \\
0 &= x^{12u}T[:u] + x^{10u}T[2u:3u] + x^{8u}T[4u:5u] + T[12u:13u], \\
0 &= x^{12u}T[u:2u] + x^{10u}T[3u:4u] + x^{8u}T[5u:6u] + T[13u:14u],
\end{aligned}$$

which can be rewritten as

$$\begin{aligned}
(2.7) \quad 0 &= T[[1w]_2] + T[[101w]_2] + T[[111w]_2], \\
(2.8) \quad 0 &= T[[10w]_2] + T[[110w]_2] + T[[1000w]_2], \\
(2.9) \quad 0 &= T[[1w]_2] + T[[11w]_2] + T[[101w]_2] + T[[1001w]_2], \\
(2.10) \quad 0 &= T[[10w]_2] + T[[100w]_2] + T[[110w]_2] + T[[1010w]_2], \\
(2.11) \quad 0 &= T[[1w]_2] + T[[11w]_2] + T[[1011w]_2], \\
(2.12) \quad 0 &= T[[w]_2] + T[[10w]_2] + T[[100w]_2] + T[[1100w]_2], \\
(2.13) \quad 0 &= T[[1w]_2] + T[[11w]_2] + T[[101w]_2] + T[[1101w]_2],
\end{aligned}$$

for all binary word w of length $2k$ and $w \neq 0^{2k}$, and

$$(2.14) \quad 0 = T[[1w]_2] + T[[101w]_2] + T[[111w]_2],$$

$$(2.15) \quad 0 = T[[10w]_2] + T[[110w]_2] + T[[1000w]_2],$$

$$(2.16) \quad 0 = T[[1w]_2] + T[[11w]_2] + T[[101w]_2] + T[[1001w]_2],$$

$$(2.17) \quad 0 = T[[10w]_2] + T[[100w]_2] + T[[110w]_2] + T[[1010w]_2],$$

$$(2.18) \quad 0 = T[[1w]_2] + T[[11w]_2] + T[[1011w]_2],$$

$$(2.19) \quad 0 = T[[w]_2] + T[[10w]_2] + T[[100w]_2] + T[[1100w]_2],$$

$$(2.20) \quad 0 = T[[1w]_2] + T[[11w]_2] + T[[101w]_2] + T[[1101w]_2],$$

for $w = 0^{2k}$.

First we calculate an 2-automaton that generates T from its minimal polynomial and its first terms. This automaton has 283 states; its transition function and output function can be found in the annex. Let $A(s, w)$ denote the state reached after reading w from right to left starting from the state s , and τ the output function. Define

$$E_{2k} = \{A(i, w) : |w| = 2k, w \neq 0^{2k}\}.$$

Identities (2.7) through (2.20) can be written as

$$(2.21) \quad 0 = \tau(A(s, 1)) + \tau(A(s, 101)) + \tau(A(s, 111)),$$

$$(2.22) \quad 0 = \tau(A(s, 10)) + \tau(A(s, 110)) + \tau(A(s, 1000)),$$

$$(2.23) \quad 0 = \tau(A(s, 1)) + \tau(A(s, 11)) + \tau(A(s, 101)) + \tau(A(s, 1001)),$$

$$(2.24) \quad 0 = \tau(A(s, 10)) + \tau(A(s, 100)) + \tau(A(s, 110)) + \tau(A(s, 1010)),$$

$$(2.25) \quad 0 = \tau(A(s, 1)) + \tau(A(s, 11)) + \tau(A(s, 1011)),$$

$$(2.26) \quad 0 = \tau(A(s, \epsilon)) + \tau(A(s, 10)) + \tau(A(s, 100)) + \tau(A(s, 1100)),$$

$$(2.27) \quad 0 = \tau(A(s, 1)) + \tau(A(s, 11)) + \tau(A(s, 101)) + \tau(A(s, 1101)),$$

for all $s \in E_{2k}$, and

$$(2.28) \quad 0 = \tau(A(s, 1)) + \tau(A(s, 101)) + \tau(A(s, 111)),$$

$$(2.29) \quad 0 = \tau(A(s, 10)) + \tau(A(s, 110)) + \tau(A(s, 1000)),$$

$$(2.30) \quad 0 = \tau(A(s, 1)) + \tau(A(s, 11)) + \tau(A(s, 101)) + \tau(A(s, 1001)),$$

$$(2.31) \quad 0 = \tau(A(s, 10)) + \tau(A(s, 100)) + \tau(A(s, 110)) + \tau(A(s, 1010)),$$

$$(2.32) \quad 0 = \tau(A(s, 1)) + \tau(A(s, 11)) + \tau(A(s, 1011)),$$

$$(2.33) \quad 0 = \tau(A(s, \epsilon)) + \tau(A(s, 10)) + \tau(A(s, 100)) + \tau(A(s, 1100)),$$

$$(2.34) \quad 0 = \tau(A(s, 1)) + \tau(A(s, 11)) + \tau(A(s, 101)) + \tau(A(s, 1101)),$$

for $s = A(i, 0^{2k})$. We find that $(A(i, 0^{22}), E_{22}) = (A(i, 0^{18}), E_{18})$, so that we only have to verify that identities (2.21) through (2.34) hold for $2 \leq k \leq 11$, which turns out to be true. $\square$

In the following lemma, we express M_{2k} , M_{2k+1} , W_{2k} , and W_{2k+1} in terms of M^e , M^o , W^e , and W^o .

Lemma 2.3. *For all integer $k \geq 2$, and $u = 2^{2k-1}$,*

$$M_{2k} = M^e[:7u] + x^{2u}M^e[:5u] + x^{6u}M^e[:u] + x^{7u}R^e,$$

$$W_{2k} = W^e[:7u] + x^{2u}W^e[:5u] + x^{6u}W^e[:u] + x^{7u}R^e.$$

For all integer $k \geq 2$, and $u = 2^{2k}$,

$$\begin{aligned} M_{2k+1} &= M^o[:7u] + x^{2u}M^o[:5u] + x^{6u}M^o[:u] + x^{7u}R^o, \\ W_{2k+1} &= W^o[:7u] + x^{2u}W^o[:5u] + x^{6u}W^o[:u] + x^{7u}R^o. \end{aligned}$$

Proof. We call the four identities in Lemma 2.3 also by the name M_{2k} , W_{2k} , M_{2k+1} , and W_{2k+1} . For $n = 2$, the identities can be verified directly. For $n \geq 2$, we claim that

$$\begin{aligned} M_{2k} \wedge W_{2k} &\Rightarrow M_{2k+1} \wedge W_{2k+1}, \\ M_{2k+1} \wedge W_{2k+1} &\Rightarrow M_{2k+2} \wedge W_{2k+2}. \end{aligned}$$

We give the proof of

$$M_{2k} \wedge W_{2k} \Rightarrow M_{2k+1},$$

the other ones being similar. Set $u = 2^{2k}$ and $v = 2^{2k-1}$. By definition and induction hypothesis, the left side of identity M_{2k+1} is equal to

$$\begin{aligned} (2.35) \quad W_{2k}M_{2k} &= (W^e[:7v] + x^{2v}W^e[:5v] + x^{6v}W^e[:v] + x^{7u}R^e) \times (M^e[:7v] \\ &\quad + x^{2v}M^e[:5v] + x^{6v}M^e[:v] + x^{7u}R^e); \end{aligned}$$

Call this expression *lhs*. Note that both sides of identity M_{2k+1} have the same term of highest degree $x^{14v}R^o$. Therefore we only need to prove that their difference is $O(x^{14v})$. Using Lemma 2.1 it can be seen that the right side of identity M_{2k+1} is congruent, modulo x^{14v} , to

$$W^e[:14v]M^e[:14v] + x^{4v}W^e[:10v]M^e[:10v] + x^{12v}W^e[:2v]M^e[:2v].$$

For all $n \leq 14$, replace the occurrences of $W^e[:n \cdot v]$ and $M^e[:n \cdot v]$ in the above expression by the reduction modulo $x^{n \cdot v}$ of the right side of the corresponding identity in Lemma 2.2 and get a new expression, which we call *rhs*. Define

$$\begin{aligned} X &:= x^v, \\ a_n &:= W^e[n \cdot v : (n+1) \cdot v] / X^n, \\ b_n &:= M^e[n \cdot v : (n+1) \cdot v] / X^n, \\ c &:= R^e. \end{aligned}$$

Using the notation introduced above, we can represent the expressions *lhs* (2.35) and *rhs* as polynomials in $\mathbb{F}_2[a_1, \dots, a_{14}, b_1, \dots, b_{14}, c][X]$. Note that it is not a problem that a_j commutes with b_k while W^e does not commute with M^e , because in the expressions concerned, the products of W^e -terms and M^e -terms are always in the same order. We let the computer do the simplification and check that the difference between these two polynomials is indeed $O(X^{14})$, which completes the proof. $\square$

Proof of Theorem 1.1. We prove the theorem for $\text{CF}(a, b)$; for $\text{CF}(b, a)$, the proof is similar. By Lemma 2.3, we have For all $0 \leq j, k \leq 1$,

$$\begin{aligned} \lim_{n \rightarrow \infty} M_{2n,j,k} &= M_{j,k}^e, \\ \lim_{n \rightarrow \infty} M_{2n+1,j,k} &= M_{j,k}^o, \\ \lim_{n \rightarrow \infty} W_{2n,j,k} &= W_{j,k}^e, \\ \lim_{n \rightarrow \infty} W_{2n+1,j,k} &= W_{j,k}^o. \end{aligned}$$

Let $z = 1/x$. By the convergence theorem and identity (2.2),

$$\text{CF}(a, b) = \frac{M_{0,1}^e(x)}{M_{0,0}^e(x)}.$$

By definition, $\phi(M^e, 0, 1)$ and $\phi(M^e, 0, 0)$ are minimal polynomials of $M_{0,1}^e$ and $M_{0,0}^e$. Therefore

$$P(x, y) = \text{Res}_t (\phi(M^e, 0, 1)(x, t), y^{12}\phi(M^e, 0, 1)(x, t/y))$$

is an annihilating polynomial of $f(x) = M_{0,1}^e/M_{0,0}^e$.

Define

$$Q(x, y) = q_4(x)y^4 + q_3(x)y^3 + q_2(x)y^2 + q_1(x)y + q_0(x),$$

where

$$\begin{aligned} q_0(x) &= x^{24} + x^{23} + x^{22} + x^{14} + x^{13} + x^9 + x^8 + x^7, \\ q_1(x) &= x^{23} + x^{21} + x^{19} + x^{18} + x^{13} + x^{11} + x^8 + x^4 + x^3 + x^2, \\ q_2(x) &= x^{22} + x^{20} + x^{18} + x^{16} + x^{12} + x^{10} + x^6 + 1, \\ q_3(x) &= x^{23} + x^{21} + x^{19} + x^{18} + x^{13} + x^{11} + x^8 + x^4 + x^3 + x^2, \\ q_4(x) &= x^{23} + x^{20} + x^{18} + x^{14} + x^{13} + x^{12} + x^{10} + x^9 + x^7 + x^6 + x^4. \end{aligned}$$

The polynomial $Q(x, y)$ is the candidate for the minimal polynomial of $f(x)$ found by Padé-Hermite approximation. To prove that it is indeed the minimal polynomial of $f(x)$, we only need to prove that it is an irreducible factor of $P(x, y)$ of multiplicity m and $R(x, y) := P(x, y)/Q(x, y)^m$ is not an annihilating polynomial of $f(x)$. We verify the first point directly. For the second point, we find that when we truncate $f(x)$ to order 216, and substitute it for y in $R(z, y)$, we get a series with valuation smaller than 216, which proves that $R(z, y)$ is not an annihilating polynomial of $f(x)$. Finally, $z^{12}Q(1/z, y)$ is the minimal polynomial of $\text{CF}(a, b) = f(1/z)$. $\square$

3. ANNEXE

All 16 polynomials are of the form

$$p_0(x) + p_3(x)y^3 + p_6(x)y^6 + p_9(x)y^9 + p_{12}(x)y^{12}.$$

The coefficients $p_j(x)$ and the 16 initial terms to determine the solutions uniquely are given below.

For $\phi(M^e, 0, 0)$:

$$\begin{aligned} p_0(x) &= x^{138} + x^{132} + x^{128} + x^{126} + x^{110} + x^{102} + x^{100} + x^{98} + x^{90} + x^{88} \\ &\quad + x^{84} + x^{78} + x^{76} + x^{74} + x^{72} + x^{70} + x^{66} + x^{62} + x^{60} + x^{58} + x^{54} \\ &\quad + x^{46} + x^{44} + x^{42} + x^{38} + x^{36} + x^{34} + x^{32} + x^{30} + x^{28} + x^{24}, \\ p_3(x) &= x^{134} + x^{130} + x^{128} + x^{122} + x^{118} + x^{104} + x^{102} + x^{98} + x^{96} + x^{92} \\ &\quad + x^{90} + x^{88} + x^{84} + x^{80} + x^{74} + x^{72} + x^{70} + x^{56} + x^{52} + x^{48} + x^{46} \\ &\quad + x^{44} + x^{40} + x^{36} + x^{34} + x^{32} + x^{30} + x^{26} + x^{24} + x^{20} + x^{18} + x^{16}, \\ p_6(x) &= x^{128} + x^{126} + x^{124} + x^{122} + x^{118} + x^{116} + x^{110} + x^{108} + x^{102} + x^{100} \\ &\quad + x^{96} + x^{86} + x^{84} + x^{72} + x^{70} + x^{68} + x^{66} + x^{62} + x^{60} + x^{58} + x^{56} \\ &\quad + x^{48} + x^{42} + x^{40} + x^{30} + x^{28} + x^{26} + x^{22} + x^{20} + x^{18} + x^8 + x^6 + x^4 \end{aligned}$$

$$\begin{aligned}
& + 1, \\
p_9(x) &= x^{136} + x^{134} + x^{132} + x^{130} + x^{124} + x^{122} + x^{116} + x^{112} + x^{110} + x^{98} \\
& + x^{94} + x^{88} + x^{82} + x^{80} + x^{74} + x^{70} + x^{68} + x^{54} + x^{50} + x^{46} + x^{44} \\
& + x^{42} + x^{40} + x^{38} + x^{36} + x^{30} + x^{28} + x^{26} + x^{24} + x^{22} + x^{20} + x^{12} \\
& + x^6 + x^4, \\
p_{12}(x) &= x^{136} + x^{128} + x^{56} + x^{48} + x^{24} + x^{16} + x^8 + 1,
\end{aligned}$$

and the initial terms are $[1, 0, 0, 0, 0, 0, 0, 0, 1, 0, 1, 0, 0, 0, 0, 0]$.

For $\phi(M^e, 0, 1)$:

$$\begin{aligned}
p_0(x) &= x^{129} + x^{128} + x^{124} + x^{122} + x^{121} + x^{115} + x^{113} + x^{112} + x^{111} + x^{110} \\
& + x^{108} + x^{106} + x^{105} + x^{104} + x^{101} + x^{92} + x^{88} + x^{87} + x^{84} + x^{83} \\
& + x^{82} + x^{81} + x^{75} + x^{74} + x^{72} + x^{67} + x^{63} + x^{59} + x^{58} + x^{57} + x^{55} \\
& + x^{52} + x^{50} + x^{49} + x^{48} + x^{47} + x^{42} + x^{38} + x^{37} + x^{36} + x^{35} + x^{34} \\
& + x^{33}, \\
p_3(x) &= x^{127} + x^{125} + x^{123} + x^{122} + x^{117} + x^{115} + x^{112} + x^{108} + x^{107} + x^{106} \\
& + x^{95} + x^{93} + x^{91} + x^{90} + x^{85} + x^{83} + x^{80} + x^{79} + x^{77} + x^{76} + x^{69} \\
& + x^{67} + x^{64} + x^{60} + x^{59} + x^{58} + x^{39} + x^{37} + x^{35} + x^{34} + x^{29} + x^{27} \\
& + x^{24} + x^{20} + x^{19} + x^{18}, \\
p_6(x) &= x^{126} + x^{118} + x^{116} + x^{114} + x^{112} + x^{108} + x^{106} + x^{104} + x^{102} + x^{94} \\
& + x^{92} + x^{90} + x^{88} + x^{84} + x^{80} + x^{74} + x^{72} + x^{70} + x^{68} + x^{66} + x^{64} \\
& + x^{62} + x^{60} + x^{58} + x^{56} + x^{52} + x^{50} + x^{48} + x^{42} + x^{38} + x^{34} + x^{32} \\
& + x^{26} + x^{16} + x^{14} + x^{12}, \\
p_9(x) &= x^{125} + x^{123} + x^{120} + x^{119} + x^{117} + x^{116} + x^{113} + x^{111} + x^{108} + x^{107} \\
& + x^{105} + x^{104} + x^{103} + x^{101} + x^{98} + x^{97} + x^{94} + x^{93} + x^{90} + x^{87} \\
& + x^{86} + x^{84} + x^{83} + x^{82} + x^{81} + x^{79} + x^{77} + x^{76} + x^{75} + x^{70} + x^{69} \\
& + x^{68} + x^{65} + x^{61} + x^{60} + x^{59} + x^{58} + x^{56} + x^{54} + x^{51} + x^{50} + x^{48} \\
& + x^{47} + x^{46} + x^{43} + x^{41} + x^{39} + x^{38} + x^{35} + x^{30} + x^{28} + x^{25} + x^{24} \\
& + x^{23} + x^{22} + x^{21} + x^{20} + x^{19} + x^{18} + x^{16} + x^{15} + x^{14} + x^{11} + x^9 \\
& + x^7 + x^6, \\
p_{12}(x) &= x^{124} + x^{120} + x^{116} + x^{112} + x^{108} + x^{104} + x^{100} + x^{96} + x^{92} + x^{88} \\
& + x^{84} + x^{80} + x^{76} + x^{72} + x^{68} + x^{64} + x^{60} + x^{56} + x^{52} + x^{48} + x^{12} \\
& + x^8 + x^4 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 1, 1, 1, 1, 1, 1, 0, 1, 0, 1, 1, 1, 1]$.

For $\phi(M^e, 1, 0)$:

$$\begin{aligned}
p_0(x) &= x^{129} + x^{128} + x^{124} + x^{122} + x^{121} + x^{115} + x^{113} + x^{112} + x^{111} + x^{110} \\
& + x^{108} + x^{106} + x^{105} + x^{104} + x^{101} + x^{92} + x^{88} + x^{87} + x^{84} + x^{83}
\end{aligned}$$

$$\begin{aligned}
& + x^{82} + x^{81} + x^{75} + x^{74} + x^{72} + x^{67} + x^{63} + x^{59} + x^{58} + x^{57} + x^{55} \\
& + x^{52} + x^{50} + x^{49} + x^{48} + x^{47} + x^{42} + x^{38} + x^{37} + x^{36} + x^{35} + x^{34} \\
& + x^{33}, \\
p_3(x) &= x^{127} + x^{125} + x^{123} + x^{122} + x^{117} + x^{115} + x^{112} + x^{108} + x^{107} + x^{106} \\
& + x^{95} + x^{93} + x^{91} + x^{90} + x^{85} + x^{83} + x^{80} + x^{79} + x^{77} + x^{76} + x^{69} \\
& + x^{67} + x^{64} + x^{60} + x^{59} + x^{58} + x^{39} + x^{37} + x^{35} + x^{34} + x^{29} + x^{27} \\
& + x^{24} + x^{20} + x^{19} + x^{18}, \\
p_6(x) &= x^{126} + x^{118} + x^{116} + x^{114} + x^{112} + x^{108} + x^{106} + x^{104} + x^{102} + x^{94} \\
& + x^{92} + x^{90} + x^{88} + x^{84} + x^{80} + x^{74} + x^{72} + x^{70} + x^{68} + x^{66} + x^{64} \\
& + x^{62} + x^{60} + x^{58} + x^{56} + x^{52} + x^{50} + x^{48} + x^{42} + x^{38} + x^{34} + x^{32} \\
& + x^{26} + x^{16} + x^{14} + x^{12}, \\
p_9(x) &= x^{125} + x^{123} + x^{120} + x^{119} + x^{117} + x^{116} + x^{113} + x^{111} + x^{108} + x^{107} \\
& + x^{105} + x^{104} + x^{103} + x^{101} + x^{98} + x^{97} + x^{94} + x^{93} + x^{90} + x^{87} \\
& + x^{86} + x^{84} + x^{83} + x^{82} + x^{81} + x^{79} + x^{77} + x^{76} + x^{75} + x^{70} + x^{69} \\
& + x^{68} + x^{65} + x^{61} + x^{60} + x^{59} + x^{58} + x^{56} + x^{54} + x^{51} + x^{50} + x^{48} \\
& + x^{47} + x^{46} + x^{43} + x^{41} + x^{39} + x^{38} + x^{35} + x^{30} + x^{28} + x^{25} + x^{24} \\
& + x^{23} + x^{22} + x^{21} + x^{20} + x^{19} + x^{18} + x^{16} + x^{15} + x^{14} + x^{11} + x^9 \\
& + x^7 + x^6, \\
p_{12}(x) &= x^{124} + x^{120} + x^{116} + x^{112} + x^{108} + x^{104} + x^{100} + x^{96} + x^{92} + x^{88} \\
& + x^{84} + x^{80} + x^{76} + x^{72} + x^{68} + x^{64} + x^{60} + x^{56} + x^{52} + x^{48} + x^{12} \\
& + x^8 + x^4 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 1, 1, 1, 1, 1, 1, 1, 0, 1, 0, 1, 1, 1, 1]$.

For $\phi(M^e, 1, 1)$:

$$\begin{aligned}
p_0(x) &= x^{120} + x^{118} + x^{114} + x^{112} + x^{108} + x^{106} + x^{102} + x^{98} + x^{96} + x^{94} \\
& + x^{90} + x^{88} + x^{82} + x^{72} + x^{70} + x^{62} + x^{58} + x^{50} + x^{48} + x^{44} + x^{42}, \\
p_3(x) &= x^{114} + x^{102} + x^{96} + x^{94} + x^{82} + x^{72} + x^{70} + x^{68} + x^{66} + x^{64} + x^{60} \\
& + x^{52} + x^{48} + x^{44} + x^{38} + x^{36} + x^{32} + x^{30} + x^{26} + x^{16} + x^{14} + x^{12}, \\
p_6(x) &= x^{114} + x^{112} + x^{110} + x^{108} + x^{106} + x^{104} + x^{102} + x^{98} + x^{96} + x^{88} \\
& + x^{86} + x^{84} + x^{82} + x^{76} + x^{70} + x^{68} + x^{64} + x^{60} + x^{58} + x^{56} + x^{50} \\
& + x^{44} + x^{40} + x^{34} + x^{26} + x^{22} + x^{16} + x^{12} + x^8 + x^6 + x^4 + 1, \\
p_9(x) &= x^{112} + x^{110} + x^{108} + x^{106} + x^{104} + x^{102} + x^{90} + x^{88} + x^{86} + x^{84} \\
& + x^{76} + x^{72} + x^{66} + x^{64} + x^{62} + x^{56} + x^{54} + x^{52} + x^{48} + x^{46} + x^{42} \\
& + x^{40} + x^{38} + x^{36} + x^{34} + x^{32} + x^{26} + x^{24} + x^{22} + x^{14} + x^6 + x^4, \\
p_{12}(x) &= x^{112} + x^{96} + x^{80} + x^{64} + x^{48} + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 1, 0, 1, 0, 1, 0, 1, 0, 0, 0, 1, 0]$.

For $\phi(W^e, 0, 0)$:

$$\begin{aligned}
p_0(x) &= x^{120} + x^{118} + x^{116} + x^{114} + x^{112} + x^{110} + x^{106} + x^{104} + x^{98} + x^{92} \\
&\quad + x^{86} + x^{82} + x^{68} + x^{64} + x^{62} + x^{60} + x^{58} + x^{56} + x^{54} + x^{52} + x^{48} \\
&\quad + x^{46} + x^{34} + x^{32} + x^{30} + x^{28} + x^{24}, \\
p_3(x) &= x^{114} + x^{110} + x^{100} + x^{98} + x^{92} + x^{88} + x^{84} + x^{78} + x^{76} + x^{72} + x^{68} \\
&\quad + x^{66} + x^{64} + x^{62} + x^{58} + x^{56} + x^{44} + x^{40} + x^{34} + x^{32} + x^{24} + x^{22} \\
&\quad + x^{18} + x^{16}, \\
p_6(x) &= x^{114} + x^{112} + x^{106} + x^{104} + x^{94} + x^{92} + x^{88} + x^{86} + x^{84} + x^{78} + x^{76} \\
&\quad + x^{72} + x^{64} + x^{62} + x^{60} + x^{50} + x^{38} + x^{36} + x^{30} + x^{28} + x^{18} + x^{14} \\
&\quad + x^{12} + x^6 + x^4 + 1, \\
p_9(x) &= x^{112} + x^{110} + x^{108} + x^{106} + x^{104} + x^{102} + x^{90} + x^{88} + x^{86} + x^{84} \\
&\quad + x^{76} + x^{72} + x^{66} + x^{64} + x^{62} + x^{56} + x^{54} + x^{52} + x^{48} + x^{46} + x^{42} \\
&\quad + x^{40} + x^{38} + x^{36} + x^{34} + x^{32} + x^{26} + x^{24} + x^{22} + x^{14} + x^6 + x^4, \\
p_{12}(x) &= x^{112} + x^{96} + x^{80} + x^{64} + x^{48} + 1,
\end{aligned}$$

and the initial terms are $[1, 0, 0, 0, 1, 0, 1, 0, 0, 0, 1, 0, 1, 0, 1, 0]$.

For $\phi(W^e, 0, 1)$:

$$\begin{aligned}
p_0(x) &= x^{129} + x^{127} + x^{126} + x^{125} + x^{122} + x^{121} + x^{120} + x^{119} + x^{118} + x^{115} \\
&\quad + x^{114} + x^{113} + x^{112} + x^{111} + x^{110} + x^{106} + x^{105} + x^{104} + x^{103} \\
&\quad + x^{102} + x^{100} + x^{95} + x^{94} + x^{87} + x^{82} + x^{80} + x^{79} + x^{77} + x^{76} + x^{74} \\
&\quad + x^{71} + x^{70} + x^{65} + x^{64} + x^{61} + x^{60} + x^{57} + x^{54} + x^{51} + x^{48} + x^{47} \\
&\quad + x^{44} + x^{41} + x^{40} + x^{37} + x^{36} + x^{33}, \\
p_3(x) &= x^{127} + x^{125} + x^{123} + x^{122} + x^{117} + x^{115} + x^{112} + x^{108} + x^{107} + x^{106} \\
&\quad + x^{95} + x^{93} + x^{91} + x^{90} + x^{85} + x^{83} + x^{80} + x^{79} + x^{77} + x^{76} + x^{69} \\
&\quad + x^{67} + x^{64} + x^{60} + x^{59} + x^{58} + x^{39} + x^{37} + x^{35} + x^{34} + x^{29} + x^{27} \\
&\quad + x^{24} + x^{20} + x^{19} + x^{18}, \\
p_6(x) &= x^{126} + x^{118} + x^{116} + x^{114} + x^{112} + x^{108} + x^{106} + x^{104} + x^{102} + x^{94} \\
&\quad + x^{92} + x^{90} + x^{88} + x^{84} + x^{80} + x^{74} + x^{72} + x^{70} + x^{68} + x^{66} + x^{64} \\
&\quad + x^{62} + x^{60} + x^{58} + x^{56} + x^{52} + x^{50} + x^{48} + x^{42} + x^{38} + x^{34} + x^{32} \\
&\quad + x^{26} + x^{16} + x^{14} + x^{12}, \\
p_9(x) &= x^{125} + x^{123} + x^{120} + x^{119} + x^{117} + x^{116} + x^{113} + x^{111} + x^{108} + x^{107} \\
&\quad + x^{105} + x^{104} + x^{103} + x^{101} + x^{98} + x^{97} + x^{94} + x^{93} + x^{90} + x^{87} \\
&\quad + x^{86} + x^{84} + x^{83} + x^{82} + x^{81} + x^{79} + x^{77} + x^{76} + x^{75} + x^{70} + x^{69} \\
&\quad + x^{68} + x^{65} + x^{61} + x^{60} + x^{59} + x^{58} + x^{56} + x^{54} + x^{51} + x^{50} + x^{48} \\
&\quad + x^{47} + x^{46} + x^{43} + x^{41} + x^{39} + x^{38} + x^{35} + x^{30} + x^{28} + x^{25} + x^{24} \\
&\quad + x^{23} + x^{22} + x^{21} + x^{20} + x^{19} + x^{18} + x^{16} + x^{15} + x^{14} + x^{11} + x^9
\end{aligned}$$

$$\begin{aligned}
& + x^7 + x^6, \\
p_{12}(x) = & x^{124} + x^{120} + x^{116} + x^{112} + x^{108} + x^{104} + x^{100} + x^{96} + x^{92} + x^{88} \\
& + x^{84} + x^{80} + x^{76} + x^{72} + x^{68} + x^{64} + x^{60} + x^{56} + x^{52} + x^{48} + x^{12} \\
& + x^8 + x^4 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 1, 1, 1, 1, 1, 1, 0, 1, 1, 0, 0]$.

For $\phi(W^e, 1, 0)$:

$$\begin{aligned}
p_0(x) = & x^{129} + x^{127} + x^{126} + x^{125} + x^{122} + x^{121} + x^{120} + x^{119} + x^{118} + x^{115} \\
& + x^{114} + x^{113} + x^{112} + x^{111} + x^{110} + x^{106} + x^{105} + x^{104} + x^{103} \\
& + x^{102} + x^{100} + x^{95} + x^{94} + x^{87} + x^{82} + x^{80} + x^{79} + x^{77} + x^{76} + x^{74} \\
& + x^{71} + x^{70} + x^{65} + x^{64} + x^{61} + x^{60} + x^{57} + x^{54} + x^{51} + x^{48} + x^{47} \\
& + x^{44} + x^{41} + x^{40} + x^{37} + x^{36} + x^{33}, \\
p_3(x) = & x^{127} + x^{125} + x^{123} + x^{122} + x^{117} + x^{115} + x^{112} + x^{108} + x^{107} + x^{106} \\
& + x^{95} + x^{93} + x^{91} + x^{90} + x^{85} + x^{83} + x^{80} + x^{79} + x^{77} + x^{76} + x^{69} \\
& + x^{67} + x^{64} + x^{60} + x^{59} + x^{58} + x^{39} + x^{37} + x^{35} + x^{34} + x^{29} + x^{27} \\
& + x^{24} + x^{20} + x^{19} + x^{18}, \\
p_6(x) = & x^{126} + x^{118} + x^{116} + x^{114} + x^{112} + x^{108} + x^{106} + x^{104} + x^{102} + x^{94} \\
& + x^{92} + x^{90} + x^{88} + x^{84} + x^{80} + x^{74} + x^{72} + x^{70} + x^{68} + x^{66} + x^{64} \\
& + x^{62} + x^{60} + x^{58} + x^{56} + x^{52} + x^{50} + x^{48} + x^{42} + x^{38} + x^{34} + x^{32} \\
& + x^{26} + x^{16} + x^{14} + x^{12}, \\
p_9(x) = & x^{125} + x^{123} + x^{120} + x^{119} + x^{117} + x^{116} + x^{113} + x^{111} + x^{108} + x^{107} \\
& + x^{105} + x^{104} + x^{103} + x^{101} + x^{98} + x^{97} + x^{94} + x^{93} + x^{90} + x^{87} \\
& + x^{86} + x^{84} + x^{83} + x^{82} + x^{81} + x^{79} + x^{77} + x^{76} + x^{75} + x^{70} + x^{69} \\
& + x^{68} + x^{65} + x^{61} + x^{60} + x^{59} + x^{58} + x^{56} + x^{54} + x^{51} + x^{50} + x^{48} \\
& + x^{47} + x^{46} + x^{43} + x^{41} + x^{39} + x^{38} + x^{35} + x^{30} + x^{28} + x^{25} + x^{24} \\
& + x^{23} + x^{22} + x^{21} + x^{20} + x^{19} + x^{18} + x^{16} + x^{15} + x^{14} + x^{11} + x^9 \\
& + x^7 + x^6, \\
p_{12}(x) = & x^{124} + x^{120} + x^{116} + x^{112} + x^{108} + x^{104} + x^{100} + x^{96} + x^{92} + x^{88} \\
& + x^{84} + x^{80} + x^{76} + x^{72} + x^{68} + x^{64} + x^{60} + x^{56} + x^{52} + x^{48} + x^{12} \\
& + x^8 + x^4 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 1, 1, 1, 1, 1, 1, 0, 1, 1, 0, 0]$.

For $\phi(W^e, 1, 1)$:

$$\begin{aligned}
p_0(x) = & x^{138} + x^{136} + x^{134} + x^{128} + x^{126} + x^{124} + x^{118} + x^{116} + x^{114} + x^{110} \\
& + x^{108} + x^{106} + x^{100} + x^{94} + x^{92} + x^{84} + x^{78} + x^{76} + x^{72} + x^{68} + x^{66} \\
& + x^{64} + x^{62} + x^{56} + x^{50} + x^{46} + x^{42}, \\
p_3(x) = & x^{130} + x^{128} + x^{124} + x^{120} + x^{114} + x^{110} + x^{108} + x^{104} + x^{98} + x^{96}
\end{aligned}$$

$$\begin{aligned}
& + x^{94} + x^{86} + x^{78} + x^{76} + x^{72} + x^{70} + x^{66} + x^{64} + x^{60} + x^{52} + x^{42} \\
& + x^{40} + x^{36} + x^{28} + x^{26} + x^{16} + x^{14} + x^{12}, \\
p_6(x) &= x^{134} + x^{132} + x^{128} + x^{126} + x^{120} + x^{116} + x^{114} + x^{112} + x^{108} + x^{104} \\
& + x^{96} + x^{94} + x^{90} + x^{88} + x^{78} + x^{76} + x^{74} + x^{72} + x^{70} + x^{68} + x^{62} \\
& + x^{60} + x^{58} + x^{56} + x^{52} + x^{46} + x^{44} + x^{40} + x^{34} + x^{26} + x^{22} + x^{20} \\
& + x^{14} + x^6 + x^4 + 1, \\
p_9(x) &= x^{136} + x^{134} + x^{132} + x^{130} + x^{124} + x^{122} + x^{116} + x^{112} + x^{110} + x^{98} \\
& + x^{94} + x^{88} + x^{82} + x^{80} + x^{74} + x^{70} + x^{68} + x^{54} + x^{50} + x^{46} + x^{44} \\
& + x^{42} + x^{40} + x^{38} + x^{36} + x^{30} + x^{28} + x^{26} + x^{24} + x^{22} + x^{20} + x^{12} \\
& + x^6 + x^4, \\
p_{12}(x) &= x^{136} + x^{128} + x^{56} + x^{48} + x^{24} + x^{16} + x^8 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 1, 0, 1, 0, 0, 0]$.

For $\phi(M^\circ, 0, 0)$:

$$\begin{aligned}
p_0(x) &= x^{129} + x^{128} + x^{127} + x^{124} + x^{123} + x^{119} + x^{118} + x^{117} + x^{114} + x^{110} \\
& + x^{109} + x^{103} + x^{102} + x^{99} + x^{98} + x^{95} + x^{94} + x^{93} + x^{92} + x^{91} + x^{90} \\
& + x^{89} + x^{85} + x^{83} + x^{81} + x^{80} + x^{75} + x^{74} + x^{72} + x^{67} + x^{65} + x^{64} \\
& + x^{60} + x^{56} + x^{55} + x^{54} + x^{52} + x^{50} + x^{47} + x^{43} + x^{41} + x^{40} + x^{35} \\
& + x^{33} + x^{31} + x^{28} + x^{24}, \\
p_3(x) &= x^{128} + x^{127} + x^{126} + x^{120} + x^{117} + x^{116} + x^{113} + x^{112} + x^{111} + x^{110} \\
& + x^{104} + x^{102} + x^{98} + x^{96} + x^{95} + x^{94} + x^{88} + x^{86} + x^{85} + x^{84} + x^{81} \\
& + x^{69} + x^{68} + x^{65} + x^{64} + x^{63} + x^{58} + x^{56} + x^{42} + x^{40} + x^{39} + x^{34} \\
& + x^{32} + x^{30} + x^{29} + x^{28} + x^{25} + x^{24} + x^{23} + x^{16}, \\
p_6(x) &= x^{128} + x^{127} + x^{126} + x^{123} + x^{122} + x^{117} + x^{116} + x^{112} + x^{111} + x^{110} \\
& + x^{109} + x^{107} + x^{106} + x^{104} + x^{103} + x^{100} + x^{99} + x^{98} + x^{94} + x^{93} \\
& + x^{92} + x^{88} + x^{87} + x^{85} + x^{84} + x^{82} + x^{80} + x^{79} + x^{76} + x^{73} + x^{72} \\
& + x^{65} + x^{64} + x^{62} + x^{60} + x^{57} + x^{56} + x^{55} + x^{52} + x^{51} + x^{50} + x^{49} \\
& + x^{45} + x^{44} + x^{43} + x^{41} + x^{38} + x^{37} + x^{36} + x^{35} + x^{34} + x^{32} + x^{28} \\
& + x^{27} + x^{23} + x^{22} + x^{20} + x^{19} + x^{17} + x^{15} + x^4 + 1, \\
p_9(x) &= x^{128} + x^{124} + x^{120} + x^{116} + x^{112} + x^{104} + x^{96} + x^{92} + x^{88} + x^{84} \\
& + x^{80} + x^{72} + x^{64} + x^{60} + x^{56} + x^{52} + x^{44} + x^{36} + x^{28} + x^{20} + x^{16} \\
& + x^8, \\
p_{12}(x) &= x^{128} + x^{108} + x^{100} + x^{76} + x^{68} + x^{48} + x^{44} + x^{36} + x^{28} + x^{20} + x^{16} \\
& + x^{12} + x^4 + 1,
\end{aligned}$$

and the initial terms are $[1, 0, 0, 0, 1, 0, 1, 1, 1, 1, 0, 1, 0, 0, 1, 1]$.

For $\phi(M^o, 0, 1)$:

$$\begin{aligned}
p_0(x) &= x^{124} + x^{121} + x^{120} + x^{115} + x^{107} + x^{106} + x^{105} + x^{104} + x^{103} + x^{102} \\
&\quad + x^{101} + x^{99} + x^{97} + x^{94} + x^{93} + x^{90} + x^{89} + x^{87} + x^{84} + x^{82} + x^{80} \\
&\quad + x^{78} + x^{76} + x^{71} + x^{68} + x^{67} + x^{66} + x^{65} + x^{64} + x^{63} + x^{62} + x^{56} \\
&\quad + x^{54} + x^{53} + x^{52} + x^{45} + x^{43} + x^{40} + x^{36} + x^{35} + x^{34} + x^{33}, \\
p_3(x) &= x^{117} + x^{113} + x^{112} + x^{111} + x^{110} + x^{108} + x^{107} + x^{106} + x^{85} + x^{81} \\
&\quad + x^{80} + x^{79} + x^{78} + x^{76} + x^{75} + x^{74} + x^{69} + x^{65} + x^{64} + x^{63} + x^{62} \\
&\quad + x^{60} + x^{59} + x^{58} + x^{29} + x^{25} + x^{24} + x^{23} + x^{22} + x^{20} + x^{19} + x^{18}, \\
p_6(x) &= x^{118} + x^{108} + x^{106} + x^{104} + x^{94} + x^{84} + x^{82} + x^{80} + x^{74} + x^{64} + x^{62} \\
&\quad + x^{60} + x^{58} + x^{48} + x^{44} + x^{42} + x^{36} + x^{30} + x^{28} + x^{26} + x^{24} + x^{22} \\
&\quad + x^{20} + x^{16} + x^{14} + x^{12}, \\
p_9(x) &= x^{119} + x^{115} + x^{114} + x^{113} + x^{112} + x^{111} + x^{110} + x^{108} + x^{106} + x^{105} \\
&\quad + x^{103} + x^{102} + x^{87} + x^{83} + x^{82} + x^{81} + x^{80} + x^{79} + x^{78} + x^{76} + x^{74} \\
&\quad + x^{73} + x^{71} + x^{70} + x^{55} + x^{51} + x^{50} + x^{49} + x^{48} + x^{47} + x^{46} + x^{44} \\
&\quad + x^{42} + x^{41} + x^{38} + x^{35} + x^{34} + x^{33} + x^{32} + x^{31} + x^{30} + x^{28} + x^{26} \\
&\quad + x^{25} + x^{22} + x^{19} + x^{18} + x^{17} + x^{16} + x^{15} + x^{14} + x^{12} + x^{10} + x^9 \\
&\quad + x^7 + x^6, \\
p_{12}(x) &= x^{120} + x^{112} + x^{104} + x^{100} + x^{96} + x^{88} + x^{80} + x^{72} + x^{68} + x^{64} + x^{56} \\
&\quad + x^{48} + x^{36} + x^{20} + x^8 + x^4 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 1, 1, 1, 1, 0, 0, 1, 1, 0, 0, 0, 0, 0, 0]$.

For $\phi(M^o, 1, 0)$:

$$\begin{aligned}
p_0(x) &= x^{138} + x^{137} + x^{136} + x^{135} + x^{134} + x^{133} + x^{132} + x^{131} + x^{129} + x^{128} \\
&\quad + x^{127} + x^{126} + x^{124} + x^{121} + x^{120} + x^{117} + x^{114} + x^{111} + x^{110} \\
&\quad + x^{108} + x^{107} + x^{104} + x^{103} + x^{100} + x^{99} + x^{98} + x^{97} + x^{96} + x^{95} \\
&\quad + x^{93} + x^{92} + x^{88} + x^{87} + x^{86} + x^{85} + x^{82} + x^{81} + x^{80} + x^{79} + x^{77} \\
&\quad + x^{76} + x^{74} + x^{71} + x^{70} + x^{68} + x^{66} + x^{64} + x^{62} + x^{61} + x^{60} + x^{58} \\
&\quad + x^{57} + x^{53} + x^{51} + x^{50} + x^{49} + x^{48} + x^{47} + x^{44} + x^{39} + x^{37} + x^{36} \\
&\quad + x^{33}, \\
p_3(x) &= x^{137} + x^{132} + x^{131} + x^{130} + x^{129} + x^{124} + x^{123} + x^{122} + x^{121} + x^{116} \\
&\quad + x^{115} + x^{114} + x^{113} + x^{108} + x^{107} + x^{106} + x^{105} + x^{100} + x^{99} + x^{98} \\
&\quad + x^{97} + x^{92} + x^{91} + x^{90} + x^{73} + x^{68} + x^{67} + x^{66} + x^{65} + x^{60} + x^{59} \\
&\quad + x^{58} + x^{49} + x^{44} + x^{43} + x^{42} + x^{41} + x^{36} + x^{35} + x^{34} + x^{33} + x^{28} \\
&\quad + x^{27} + x^{26} + x^{25} + x^{20} + x^{19} + x^{18}, \\
p_6(x) &= x^{138} + x^{134} + x^{128} + x^{126} + x^{120} + x^{118} + x^{114} + x^{112} + x^{106} + x^{102} \\
&\quad + x^{96} + x^{90} + x^{88} + x^{86} + x^{84} + x^{80} + x^{78} + x^{76} + x^{66} + x^{62} + x^{56}
\end{aligned}$$

$$\begin{aligned}
& + x^{50} + x^{48} + x^{46} + x^{44} + x^{42} + x^{40} + x^{38} + x^{36} + x^{32} + x^{30} + x^{28} \\
& + x^{22} + x^{18} + x^{14} + x^{12}, \\
p_9(x) = & x^{139} + x^{134} + x^{133} + x^{132} + x^{129} + x^{127} + x^{125} + x^{124} + x^{121} + x^{119} \\
& + x^{117} + x^{116} + x^{113} + x^{111} + x^{109} + x^{108} + x^{105} + x^{103} + x^{101} \\
& + x^{100} + x^{97} + x^{95} + x^{93} + x^{92} + x^{89} + x^{87} + x^{85} + x^{84} + x^{81} + x^{79} \\
& + x^{77} + x^{76} + x^{73} + x^{71} + x^{69} + x^{68} + x^{65} + x^{63} + x^{61} + x^{60} + x^{59} \\
& + x^{57} + x^{55} + x^{54} + x^{27} + x^{22} + x^{21} + x^{20} + x^{17} + x^{15} + x^{13} + x^{12} \\
& + x^{11} + x^9 + x^7 + x^6, \\
p_{12}(x) = & x^{140} + x^{136} + x^{132} + x^{128} + x^{120} + x^{116} + x^{104} + x^{100} + x^{88} + x^{84} \\
& + x^{72} + x^{68} + x^{60} + x^{48} + x^{28} + x^{24} + x^{20} + x^{16} + x^{12} + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 1, 1, 1, 1, 1, 1, 0, 0, 1, 1, 0]$.

For $\phi(M^o, 1, 1)$:

$$\begin{aligned}
p_0(x) = & x^{129} + x^{124} + x^{116} + x^{115} + x^{114} + x^{109} + x^{108} + x^{106} + x^{104} + x^{100} \\
& + x^{89} + x^{85} + x^{84} + x^{81} + x^{75} + x^{73} + x^{69} + x^{68} + x^{67} + x^{65} + x^{64} \\
& + x^{60} + x^{57} + x^{56} + x^{55} + x^{53} + x^{50} + x^{48} + x^{44} + x^{43} + x^{42}, \\
p_3(x) = & x^{128} + x^{127} + x^{124} + x^{122} + x^{117} + x^{114} + x^{113} + x^{112} + x^{111} + x^{102} \\
& + x^{98} + x^{96} + x^{95} + x^{92} + x^{90} + x^{86} + x^{85} + x^{82} + x^{81} + x^{76} + x^{74} \\
& + x^{69} + x^{66} + x^{65} + x^{64} + x^{63} + x^{62} + x^{58} + x^{42} + x^{40} + x^{39} + x^{38} \\
& + x^{36} + x^{30} + x^{29} + x^{26} + x^{25} + x^{24} + x^{23} + x^{22}, \\
p_6(x) = & x^{128} + x^{127} + x^{124} + x^{123} + x^{122} + x^{118} + x^{117} + x^{116} + x^{114} + x^{111} \\
& + x^{110} + x^{109} + x^{107} + x^{103} + x^{102} + x^{100} + x^{99} + x^{98} + x^{93} + x^{92} \\
& + x^{90} + x^{87} + x^{85} + x^{84} + x^{82} + x^{80} + x^{79} + x^{74} + x^{73} + x^{72} + x^{70} \\
& + x^{68} + x^{66} + x^{65} + x^{58} + x^{57} + x^{55} + x^{52} + x^{51} + x^{49} + x^{48} + x^{45} \\
& + x^{43} + x^{42} + x^{41} + x^{40} + x^{37} + x^{35} + x^{32} + x^{27} + x^{26} + x^{24} + x^{23} \\
& + x^{22} + x^{20} + x^{19} + x^{17} + x^{16} + x^{15} + x^{14} + x^{12} + x^8 + x^4 + 1, \\
p_9(x) = & x^{128} + x^{124} + x^{120} + x^{116} + x^{112} + x^{104} + x^{96} + x^{92} + x^{88} + x^{84} \\
& + x^{80} + x^{72} + x^{64} + x^{60} + x^{56} + x^{52} + x^{44} + x^{36} + x^{28} + x^{20} + x^{16} \\
& + x^8, \\
p_{12}(x) = & x^{128} + x^{108} + x^{100} + x^{76} + x^{68} + x^{48} + x^{44} + x^{36} + x^{28} + x^{20} + x^{16} \\
& + x^{12} + x^4 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 0, 0, 1, 0, 1, 0, 1, 0, 0, 1, 1]$.

For $\phi(W^o, 0, 0)$:

$$\begin{aligned}
p_0(x) = & x^{129} + x^{128} + x^{127} + x^{124} + x^{123} + x^{119} + x^{118} + x^{117} + x^{114} + x^{110} \\
& + x^{109} + x^{103} + x^{102} + x^{99} + x^{98} + x^{95} + x^{94} + x^{93} + x^{92} + x^{91} + x^{90} \\
& + x^{89} + x^{85} + x^{83} + x^{81} + x^{80} + x^{75} + x^{74} + x^{72} + x^{67} + x^{65} + x^{64}
\end{aligned}$$

$$\begin{aligned}
& + x^{60} + x^{56} + x^{55} + x^{54} + x^{52} + x^{50} + x^{47} + x^{43} + x^{41} + x^{40} + x^{35} \\
& + x^{33} + x^{31} + x^{28} + x^{24}, \\
p_3(x) &= x^{128} + x^{127} + x^{126} + x^{120} + x^{117} + x^{116} + x^{113} + x^{112} + x^{111} + x^{110} \\
& + x^{104} + x^{102} + x^{98} + x^{96} + x^{95} + x^{94} + x^{88} + x^{86} + x^{85} + x^{84} + x^{81} \\
& + x^{69} + x^{68} + x^{65} + x^{64} + x^{63} + x^{58} + x^{56} + x^{42} + x^{40} + x^{39} + x^{34} \\
& + x^{32} + x^{30} + x^{29} + x^{28} + x^{25} + x^{24} + x^{23} + x^{16}, \\
p_6(x) &= x^{128} + x^{127} + x^{126} + x^{123} + x^{122} + x^{117} + x^{116} + x^{112} + x^{111} + x^{110} \\
& + x^{109} + x^{107} + x^{106} + x^{104} + x^{103} + x^{100} + x^{99} + x^{98} + x^{94} + x^{93} \\
& + x^{92} + x^{88} + x^{87} + x^{85} + x^{84} + x^{82} + x^{80} + x^{79} + x^{76} + x^{73} + x^{72} \\
& + x^{65} + x^{64} + x^{62} + x^{60} + x^{57} + x^{56} + x^{55} + x^{52} + x^{51} + x^{50} + x^{49} \\
& + x^{45} + x^{44} + x^{43} + x^{41} + x^{38} + x^{37} + x^{36} + x^{35} + x^{34} + x^{32} + x^{28} \\
& + x^{27} + x^{23} + x^{22} + x^{20} + x^{19} + x^{17} + x^{15} + x^4 + 1, \\
p_9(x) &= x^{128} + x^{124} + x^{120} + x^{116} + x^{112} + x^{104} + x^{96} + x^{92} + x^{88} + x^{84} \\
& + x^{80} + x^{72} + x^{64} + x^{60} + x^{56} + x^{52} + x^{44} + x^{36} + x^{28} + x^{20} + x^{16} \\
& + x^8, \\
p_{12}(x) &= x^{128} + x^{108} + x^{100} + x^{76} + x^{68} + x^{48} + x^{44} + x^{36} + x^{28} + x^{20} + x^{16} \\
& + x^{12} + x^4 + 1,
\end{aligned}$$

and the initial terms are $[1, 0, 0, 0, 1, 0, 1, 1, 1, 1, 0, 1, 0, 0, 1, 1]$.

For $\phi(W^o, 0, 1)$:

$$\begin{aligned}
p_0(x) &= x^{138} + x^{137} + x^{136} + x^{135} + x^{134} + x^{133} + x^{132} + x^{131} + x^{129} + x^{128} \\
& + x^{127} + x^{126} + x^{124} + x^{121} + x^{120} + x^{117} + x^{114} + x^{111} + x^{110} \\
& + x^{108} + x^{107} + x^{104} + x^{103} + x^{100} + x^{99} + x^{98} + x^{97} + x^{96} + x^{95} \\
& + x^{93} + x^{92} + x^{88} + x^{87} + x^{86} + x^{85} + x^{82} + x^{81} + x^{80} + x^{79} + x^{77} \\
& + x^{76} + x^{74} + x^{71} + x^{70} + x^{68} + x^{66} + x^{64} + x^{62} + x^{61} + x^{60} + x^{58} \\
& + x^{57} + x^{53} + x^{51} + x^{50} + x^{49} + x^{48} + x^{47} + x^{44} + x^{39} + x^{37} + x^{36} \\
& + x^{33}, \\
p_3(x) &= x^{137} + x^{132} + x^{131} + x^{130} + x^{129} + x^{124} + x^{123} + x^{122} + x^{121} + x^{116} \\
& + x^{115} + x^{114} + x^{113} + x^{108} + x^{107} + x^{106} + x^{105} + x^{100} + x^{99} + x^{98} \\
& + x^{97} + x^{92} + x^{91} + x^{90} + x^{73} + x^{68} + x^{67} + x^{66} + x^{65} + x^{60} + x^{59} \\
& + x^{58} + x^{49} + x^{44} + x^{43} + x^{42} + x^{41} + x^{36} + x^{35} + x^{34} + x^{33} + x^{28} \\
& + x^{27} + x^{26} + x^{25} + x^{20} + x^{19} + x^{18}, \\
p_6(x) &= x^{138} + x^{134} + x^{128} + x^{126} + x^{120} + x^{118} + x^{114} + x^{112} + x^{106} + x^{102} \\
& + x^{96} + x^{90} + x^{88} + x^{86} + x^{84} + x^{80} + x^{78} + x^{76} + x^{66} + x^{62} + x^{56} \\
& + x^{50} + x^{48} + x^{46} + x^{44} + x^{42} + x^{40} + x^{38} + x^{36} + x^{32} + x^{30} + x^{28} \\
& + x^{22} + x^{18} + x^{14} + x^{12},
\end{aligned}$$

$$\begin{aligned}
p_9(x) &= x^{139} + x^{134} + x^{133} + x^{132} + x^{129} + x^{127} + x^{125} + x^{124} + x^{121} + x^{119} \\
&\quad + x^{117} + x^{116} + x^{113} + x^{111} + x^{109} + x^{108} + x^{105} + x^{103} + x^{101} \\
&\quad + x^{100} + x^{97} + x^{95} + x^{93} + x^{92} + x^{89} + x^{87} + x^{85} + x^{84} + x^{81} + x^{79} \\
&\quad + x^{77} + x^{76} + x^{73} + x^{71} + x^{69} + x^{68} + x^{65} + x^{63} + x^{61} + x^{60} + x^{59} \\
&\quad + x^{57} + x^{55} + x^{54} + x^{27} + x^{22} + x^{21} + x^{20} + x^{17} + x^{15} + x^{13} + x^{12} \\
&\quad + x^{11} + x^9 + x^7 + x^6, \\
p_{12}(x) &= x^{140} + x^{136} + x^{132} + x^{128} + x^{120} + x^{116} + x^{104} + x^{100} + x^{88} + x^{84} \\
&\quad + x^{72} + x^{68} + x^{60} + x^{48} + x^{28} + x^{24} + x^{20} + x^{16} + x^{12} + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 1, 1, 1, 1, 1, 1, 0, 0, 1, 1, 0]$.

For $\phi(W^o, 1, 0)$:

$$\begin{aligned}
p_0(x) &= x^{124} + x^{121} + x^{120} + x^{115} + x^{107} + x^{106} + x^{105} + x^{104} + x^{103} + x^{102} \\
&\quad + x^{101} + x^{99} + x^{97} + x^{94} + x^{93} + x^{90} + x^{89} + x^{87} + x^{84} + x^{82} + x^{80} \\
&\quad + x^{78} + x^{76} + x^{71} + x^{68} + x^{67} + x^{66} + x^{65} + x^{64} + x^{63} + x^{62} + x^{56} \\
&\quad + x^{54} + x^{53} + x^{52} + x^{45} + x^{43} + x^{40} + x^{36} + x^{35} + x^{34} + x^{33}, \\
p_3(x) &= x^{117} + x^{113} + x^{112} + x^{111} + x^{110} + x^{108} + x^{107} + x^{106} + x^{85} + x^{81} \\
&\quad + x^{80} + x^{79} + x^{78} + x^{76} + x^{75} + x^{74} + x^{69} + x^{65} + x^{64} + x^{63} + x^{62} \\
&\quad + x^{60} + x^{59} + x^{58} + x^{29} + x^{25} + x^{24} + x^{23} + x^{22} + x^{20} + x^{19} + x^{18}, \\
p_6(x) &= x^{118} + x^{108} + x^{106} + x^{104} + x^{94} + x^{84} + x^{82} + x^{80} + x^{74} + x^{64} + x^{62} \\
&\quad + x^{60} + x^{58} + x^{48} + x^{44} + x^{42} + x^{36} + x^{30} + x^{28} + x^{26} + x^{24} + x^{22} \\
&\quad + x^{20} + x^{16} + x^{14} + x^{12}, \\
p_9(x) &= x^{119} + x^{115} + x^{114} + x^{113} + x^{112} + x^{111} + x^{110} + x^{108} + x^{106} + x^{105} \\
&\quad + x^{103} + x^{102} + x^{87} + x^{83} + x^{82} + x^{81} + x^{80} + x^{79} + x^{78} + x^{76} + x^{74} \\
&\quad + x^{73} + x^{71} + x^{70} + x^{55} + x^{51} + x^{50} + x^{49} + x^{48} + x^{47} + x^{46} + x^{44} \\
&\quad + x^{42} + x^{41} + x^{38} + x^{35} + x^{34} + x^{33} + x^{32} + x^{31} + x^{30} + x^{28} + x^{26} \\
&\quad + x^{25} + x^{22} + x^{19} + x^{18} + x^{17} + x^{16} + x^{15} + x^{14} + x^{12} + x^{10} + x^9 \\
&\quad + x^7 + x^6, \\
p_{12}(x) &= x^{120} + x^{112} + x^{104} + x^{100} + x^{96} + x^{88} + x^{80} + x^{72} + x^{68} + x^{64} + x^{56} \\
&\quad + x^{48} + x^{36} + x^{20} + x^8 + x^4 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 1, 1, 1, 1, 0, 0, 1, 1, 0, 0, 0, 0, 0, 0]$.

For $\phi(W^o, 1, 1)$:

$$\begin{aligned}
p_0(x) &= x^{129} + x^{124} + x^{116} + x^{115} + x^{114} + x^{109} + x^{108} + x^{106} + x^{104} + x^{100} \\
&\quad + x^{89} + x^{85} + x^{84} + x^{81} + x^{75} + x^{73} + x^{69} + x^{68} + x^{67} + x^{65} + x^{64} \\
&\quad + x^{60} + x^{57} + x^{56} + x^{55} + x^{53} + x^{50} + x^{48} + x^{44} + x^{43} + x^{42}, \\
p_3(x) &= x^{128} + x^{127} + x^{124} + x^{122} + x^{117} + x^{114} + x^{113} + x^{112} + x^{111} + x^{102} \\
&\quad + x^{98} + x^{96} + x^{95} + x^{92} + x^{90} + x^{86} + x^{85} + x^{82} + x^{81} + x^{76} + x^{74}
\end{aligned}$$

$$\begin{aligned}
& + x^{69} + x^{66} + x^{65} + x^{64} + x^{63} + x^{62} + x^{58} + x^{42} + x^{40} + x^{39} + x^{38} \\
& + x^{36} + x^{30} + x^{29} + x^{26} + x^{25} + x^{24} + x^{23} + x^{22}, \\
p_6(x) = & x^{128} + x^{127} + x^{124} + x^{123} + x^{122} + x^{118} + x^{117} + x^{116} + x^{114} + x^{111} \\
& + x^{110} + x^{109} + x^{107} + x^{103} + x^{102} + x^{100} + x^{99} + x^{98} + x^{93} + x^{92} \\
& + x^{90} + x^{87} + x^{85} + x^{84} + x^{82} + x^{80} + x^{79} + x^{74} + x^{73} + x^{72} + x^{70} \\
& + x^{68} + x^{66} + x^{65} + x^{58} + x^{57} + x^{55} + x^{52} + x^{51} + x^{49} + x^{48} + x^{45} \\
& + x^{43} + x^{42} + x^{41} + x^{40} + x^{37} + x^{35} + x^{32} + x^{27} + x^{26} + x^{24} + x^{23} \\
& + x^{22} + x^{20} + x^{19} + x^{17} + x^{16} + x^{15} + x^{14} + x^{12} + x^8 + x^4 + 1, \\
p_9(x) = & x^{128} + x^{124} + x^{120} + x^{116} + x^{112} + x^{104} + x^{96} + x^{92} + x^{88} + x^{84} \\
& + x^{80} + x^{72} + x^{64} + x^{60} + x^{56} + x^{52} + x^{44} + x^{36} + x^{28} + x^{20} + x^{16} \\
& + x^8, \\
p_{12}(x) = & x^{128} + x^{108} + x^{100} + x^{76} + x^{68} + x^{48} + x^{44} + x^{36} + x^{28} + x^{20} + x^{16} \\
& + x^{12} + x^4 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 0, 0, 1, 0, 1, 0, 1, 0, 0, 1, 1]$.

Below is the transition function and output function of an 2-automaton that generates $T = M_{0,0}^o$:

Transition function $(n, j) \mapsto \delta(n, j)$ ($\Lambda(n) := [\delta(n, 0), \delta(n, 1)]$):

n	$\Lambda(n)$	n	$\Lambda(n)$	n	$\Lambda(n)$	n	$\Lambda(n)$	n	$\Lambda(n)$
0	[1, 2]	57	[96, 10]	114	[161, 162]	171	[191, 192]	228	[198, 250]
1	[3, 4]	58	[97, 98]	115	[56, 163]	172	[193, 194]	229	[71, 17]
2	[5, 6]	59	[1, 36]	116	[143, 132]	173	[195, 196]	230	[251, 156]
3	[7, 8]	60	[99, 45]	117	[164, 73]	174	[197, 198]	231	[252, 248]
4	[9, 10]	61	[100, 101]	118	[165, 75]	175	[199, 200]	232	[253, 254]
5	[11, 12]	62	[102, 103]	119	[166, 72]	176	[201, 63]	233	[255, 153]
6	[13, 14]	63	[8, 14]	120	[167, 168]	177	[202, 203]	234	[256, 257]
7	[15, 16]	64	[96, 14]	121	[169, 170]	178	[204, 205]	235	[203, 258]
8	[17, 18]	65	[104, 30]	122	[171, 7]	179	[206, 137]	236	[140, 259]
9	[19, 20]	66	[35, 48]	123	[116, 95]	180	[207, 208]	237	[215, 260]
10	[16, 21]	67	[8, 10]	124	[115, 90]	181	[209, 126]	238	[261, 254]
11	[22, 23]	68	[105, 35]	125	[172, 110]	182	[210, 211]	239	[262, 263]
12	[24, 25]	69	[106, 39]	126	[13, 111]	183	[212, 213]	240	[264, 236]
13	[26, 27]	70	[107, 33]	127	[173, 36]	184	[214, 215]	241	[33, 42]
14	[28, 29]	71	[31, 108]	128	[38, 48]	185	[216, 151]	242	[238, 265]
15	[1, 30]	72	[109, 14]	129	[174, 175]	186	[217, 155]	243	[230, 92]
16	[31, 32]	73	[110, 111]	130	[13, 98]	187	[218, 64]	244	[91, 33]
17	[33, 34]	74	[112, 8]	131	[94, 103]	188	[32, 219]	245	[35, 45]
18	[35, 32]	75	[113, 98]	132	[103, 116]	189	[120, 39]	246	[108, 236]
19	[36, 37]	76	[114, 101]	133	[116, 103]	190	[220, 221]	247	[220, 266]
20	[38, 32]	77	[115, 116]	134	[120, 34]	191	[222, 92]	248	[182, 34]
21	[30, 39]	78	[35, 108]	135	[49, 108]	192	[182, 39]	249	[176, 33]

22	[40, 41]	79	[117, 118]	136	[41, 45]	193	[173, 30]	250	[47, 45]
23	[30, 42]	80	[95, 103]	137	[99, 48]	194	[47, 108]	251	[267, 49]
24	[43, 44]	81	[119, 30]	138	[44, 34]	195	[223, 97]	252	[45, 236]
25	[38, 45]	82	[49, 45]	139	[176, 44]	196	[110, 98]	253	[268, 221]
26	[46, 39]	83	[120, 37]	140	[41, 108]	197	[108, 224]	254	[120, 42]
27	[47, 48]	84	[41, 48]	141	[177, 170]	198	[36, 34]	255	[269, 92]
28	[49, 48]	85	[43, 33]	142	[178, 179]	199	[225, 226]	256	[270, 232]
29	[33, 37]	86	[47, 32]	143	[89, 116]	200	[106, 34]	257	[9, 14]
30	[50, 51]	87	[121, 67]	144	[95, 95]	201	[227, 96]	258	[99, 32]
31	[52, 53]	88	[122, 4]	145	[12, 96]	202	[228, 229]	259	[44, 37]
32	[21, 54]	89	[123, 124]	146	[97, 111]	203	[44, 42]	260	[38, 108]
33	[55, 56]	90	[90, 90]	147	[40, 92]	204	[230, 41]	261	[235, 224]
34	[57, 58]	91	[125, 126]	148	[180, 181]	205	[36, 39]	262	[48, 229]
35	[59, 60]	92	[127, 128]	149	[179, 97]	206	[107, 44]	263	[33, 39]
36	[61, 62]	93	[129, 130]	150	[92, 45]	207	[231, 232]	264	[248, 271]
37	[63, 64]	94	[80, 131]	151	[182, 37]	208	[113, 111]	265	[272, 189]
38	[65, 66]	95	[132, 133]	152	[99, 108]	209	[233, 179]	266	[273, 213]
39	[4, 67]	96	[134, 135]	153	[106, 42]	210	[234, 118]	267	[274, 275]
40	[68, 69]	97	[23, 136]	154	[92, 48]	211	[90, 116]	268	[276, 26]
41	[70, 71]	98	[137, 138]	155	[41, 32]	212	[235, 236]	269	[277, 83]
42	[72, 73]	99	[139, 140]	156	[36, 42]	213	[106, 37]	270	[278, 279]
43	[74, 75]	100	[141, 58]	157	[104, 36]	214	[237, 219]	271	[31, 48]
44	[76, 77]	101	[142, 72]	158	[92, 108]	215	[46, 42]	272	[225, 265]
45	[69, 78]	102	[62, 143]	159	[44, 39]	216	[238, 226]	273	[268, 266]
46	[79, 80]	103	[124, 144]	160	[31, 45]	217	[119, 36]	274	[280, 35]
47	[81, 82]	104	[145, 146]	161	[183, 175]	218	[239, 185]	275	[182, 42]
48	[83, 84]	105	[147, 29]	162	[109, 10]	219	[60, 205]	276	[45, 224]
49	[85, 86]	106	[148, 123]	163	[102, 95]	220	[240, 241]	277	[269, 41]
50	[87, 88]	107	[149, 57]	164	[184, 185]	221	[242, 215]	278	[264, 224]
51	[89, 90]	108	[29, 150]	165	[11, 7]	222	[243, 138]	279	[30, 37]
52	[91, 44]	109	[151, 152]	166	[186, 110]	223	[244, 245]	280	[281, 282]
53	[92, 32]	110	[153, 154]	167	[187, 181]	224	[82, 23]	281	[222, 41]
54	[49, 32]	111	[155, 156]	168	[103, 90]	225	[246, 159]	282	[46, 34]
55	[93, 88]	112	[157, 158]	169	[188, 189]	226	[247, 248]		
56	[94, 95]	113	[159, 160]	170	[190, 26]	227	[249, 28]		

Output function $n \mapsto \tau(n)$:

n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$
0	0	41	0	82	1	123	0	164	0	205	0	246	0
1	0	42	1	83	1	124	1	165	0	206	0	247	1
2	0	43	1	84	0	125	1	166	0	207	0	248	1
3	0	44	1	85	1	126	0	167	1	208	1	249	0
4	0	45	0	86	0	127	1	168	1	209	0	250	0
5	0	46	0	87	1	128	1	169	1	210	1	251	1

6	0	47	0	88	1	129	0	170	1	211	0	252	0
7	0	48	1	89	0	130	0	171	1	212	1	253	0
8	0	49	1	90	0	131	1	172	1	213	0	254	1
9	0	50	1	91	1	132	1	173	1	214	0	255	0
10	1	51	0	92	1	133	0	174	0	215	0	256	1
11	0	52	1	93	0	134	1	175	0	216	1	257	0
12	1	53	1	94	1	135	1	176	0	217	0	258	0
13	0	54	1	95	1	136	0	177	0	218	1	259	1
14	1	55	0	96	1	137	0	178	1	219	0	260	1
15	0	56	1	97	1	138	1	179	0	220	1	261	1
16	1	57	1	98	0	139	0	180	0	221	1	262	1
17	0	58	1	99	0	140	0	181	0	222	1	263	0
18	0	59	0	100	0	141	0	182	1	223	1	264	1
19	0	60	0	101	1	142	1	183	1	224	1	265	0
20	1	61	0	102	0	143	0	184	0	225	0	266	0
21	1	62	0	103	1	144	1	185	1	226	1	267	1
22	0	63	0	104	1	145	1	186	0	227	0	268	0
23	1	64	1	105	0	146	1	187	1	228	0	269	0
24	1	65	1	106	0	147	0	188	1	229	1	270	1
25	1	66	0	107	0	148	0	189	1	230	1	271	1
26	0	67	0	108	0	149	0	190	1	231	0	272	0
27	0	68	0	109	1	150	1	191	1	232	0	273	0
28	1	69	0	110	0	151	1	192	1	233	0	274	1
29	0	70	0	111	0	152	0	193	1	234	1	275	1
30	1	71	1	112	1	153	0	194	0	235	1	276	0
31	1	72	1	113	1	154	1	195	1	236	0	277	0
32	1	73	0	114	1	155	0	196	0	237	0	278	1
33	0	74	1	115	1	156	0	197	0	238	1	279	1
34	1	75	1	116	0	157	1	198	0	239	1	280	1
35	0	76	1	117	0	158	1	199	0	240	1	281	1
36	0	77	1	118	0	159	1	200	0	241	0	282	0
37	0	78	0	119	0	160	1	201	0	242	1		
38	1	79	0	120	1	161	1	202	0	243	1		
39	0	80	1	121	1	162	1	203	1	244	1		
40	0	81	0	122	1	163	0	204	1	245	0		

UNIVERSITÉ DE STRASBOURG, CNRS, IRMA UMR 7501, F-67000 STRASBOURG, FRANCE
Email address: `guoniu.han@unistra.fr`

SCHOOL OF MATHEMATICS AND STATISTICS, HUAZHONG UNIVERSITY OF SCIENCE AND TECHNOLOGY, WUHAN, PR CHINA
Email address: `huyining@hust.edu.cn`