

PROOF OF THE CONTINUED FRACTION CONJECTURE FOR

$$(a, b) = (z^3 + z^2 + z, z^4 + z^3 + z^2 + z)$$

GUO-NIU HAN* AND YINING HU*

ABSTRACT. The continued fraction conjecture claims that the continued fraction defined by the (a, b) -Thue-Morse sequence is algebraic over $\mathbb{F}_2(x)$ for all pairs of distinct elements (a, b) in $\mathbb{F}_2[x] \setminus \mathbb{F}_2$. In this paper we prove the conjecture for $(a, b) = (z^3 + z^2 + z, z^4 + z^3 + z^2 + z)$.

1. INTRODUCTION

Let $\mathbb{F}_2$ be the finite field of cardinality 2. Given a sequence of polynomials $\mathbf{a} = (a_0(z), a_1(z), a_2(z), \dots)$ of elements from $\mathbb{F}_2[z] \setminus \mathbb{F}_2$, we define the infinite continued fraction

$$(1.1) \quad \text{CF}(\mathbf{a}(z)) = \frac{1}{a_0(z) + \frac{1}{a_1(z) + \frac{1}{a_2(z) + \frac{1}{\ddots}}}}$$

as the limit of the finite continued fractions

$$(1.2) \quad \text{CF}_n(\mathbf{a}(z)) = \frac{1}{a_0(z) + \frac{1}{a_1(z) + \frac{1}{\ddots + \frac{1}{a_n(z)}}}} \in \mathbb{F}_2((1/z)).$$

Conversely, each power series in $\mathbb{F}_2[[1/z]]$ without constant term

$$(1.3) \quad c_{-1}z^{-1} + c_{-2}z^{-2} + c_{-3}z^{-3} + \dots$$

admits a continued fraction expansion of form (1.1).

Let a and b be two distinct elements from $\mathbb{F}_2[z] \setminus \mathbb{F}_2$. We consider the continued fraction (1.1) associated with the (a, b) -Thue-Morse sequence

$$\mathbf{t} = (t_0(z), t_1(z), t_2(z), \dots) = (a, b, b, a, b, a, a, b, \dots)$$

that we shall denote

$$(1.4) \quad \text{CF}(a, b) := \text{CF}(\mathbf{t}).$$

Date: 9/04/2020 23:32:17 .

2010 Mathematics Subject Classification. 11B85, 11J70, 11B50, 11Y65, 05A15.

Key words and phrases. automatic sequence, continued fraction, Thue-Morse sequence.

* This paper was automatically generated from a computer program developed by the authors, with 14.1 minutes CPU time used.

Theorem 1.1. *When $(a, b) = (z^3 + z^2 + z, z^4 + z^3 + z^2 + z)$, the two power series $\text{CF}(a, b)$ and $\text{CF}(b, a)$ are algebraic over $\mathbb{F}_2(z)$, with minimal polynomials of the form*

$$p_4(z)y^4 + p_3(z)y^3 + p_2(z)y^2 + p_1(z)y + p_0(z) = 0$$

where, for $\text{CF}(a, b)$,

$$\begin{aligned} p_0(z) &= z^{15} + z^{13} + z^{12} + z^{11} + z^9 + z^5 + z^4 + z^3 + z^2 + 1, \\ p_1(z) &= z^{19} + z^{17} + z^{16} + z^{15} + z^{14} + z^{13} + z^{12} + z^{10} + z^9 + z^7 + z^6 + z^4, \\ p_2(z) &= z^{22} + z^{16} + z^{14} + z^{12} + z^6 + z^2, \\ p_3(z) &= z^{19} + z^{17} + z^{16} + z^{15} + z^{14} + z^{13} + z^{12} + z^{10} + z^9 + z^7 + z^6 + z^4, \\ p_4(z) &= z^{16} + z^{15} + z^{13} + z^{11} + z^9 + z^6 + z^5 + z^4 + z^3 + z^2 + 1, \end{aligned}$$

and for $\text{CF}(b, a)$,

$$\begin{aligned} p_0(z) &= z^{15} + z^{13} + z^{12} + z^{11} + z^{10} + z^9 + z^5 + z^4 + z^3 + z^2 + 1, \\ p_1(z) &= z^{19} + z^{17} + z^{16} + z^{15} + z^{14} + z^{13} + z^{12} + z^{10} + z^9 + z^7 + z^6 + z^4, \\ p_2(z) &= z^{22} + z^{16} + z^{14} + z^{12} + z^6 + z^2, \\ p_3(z) &= z^{19} + z^{17} + z^{16} + z^{15} + z^{14} + z^{13} + z^{12} + z^{10} + z^9 + z^7 + z^6 + z^4, \\ p_4(z) &= z^{16} + z^{15} + z^{13} + z^{11} + z^{10} + z^9 + z^6 + z^5 + z^4 + z^3 + z^2 + 1. \end{aligned}$$

2. PROOF

Define the sequences of polynomials $(P_n(z))_{n \geq 0}$ and $(Q_n(z))_{n \geq 0}$ by

$$\begin{pmatrix} P_n(z) & Q_n(z) \\ P_{n-1}(z) & Q_{n-1}(z) \end{pmatrix} := \begin{pmatrix} t_n(z) & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} t_{n-1}(z) & 1 \\ 1 & 0 \end{pmatrix} \cdots \begin{pmatrix} t_0(z) & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}.$$

By the basic properties of continued fractions, we have

$$(2.1) \quad \text{CF}_n(a, b) = \frac{P_n(z)}{Q_n(z)} \in \mathbb{F}_2((1/z)).$$

As formula (2.1) is not efficient for calculating $\text{CF}_n(a, b)$, a fast method is to be derived and described as follows. Define

$$\begin{aligned} M_n(x) &= x^{\deg(t_{2^n-1})} \begin{pmatrix} t_{2^n-1}(1/x) & 1 \\ 1 & 0 \end{pmatrix} \times x^{\deg(t_{2^n-2})} \begin{pmatrix} t_{2^n-2}(1/x) & 1 \\ 1 & 0 \end{pmatrix} \times \\ &\quad \cdots \times x^{\deg(t_0)} \begin{pmatrix} t_0(1/x) & 1 \\ 1 & 0 \end{pmatrix}, \\ W_n(x) &= x^{\deg(\bar{t}_{2^n-1})} \begin{pmatrix} \bar{t}_{2^n-1}(1/x) & 1 \\ 1 & 0 \end{pmatrix} \times x^{\deg(\bar{t}_{2^n-2})} \begin{pmatrix} \bar{t}_{2^n-2}(1/x) & 1 \\ 1 & 0 \end{pmatrix} \times \\ &\quad \cdots \times x^{\deg(\bar{t}_0)} \begin{pmatrix} \bar{t}_0(1/x) & 1 \\ 1 & 0 \end{pmatrix}, \end{aligned}$$

where $\bar{t}$ is the (b, a) -Thue-Morse sequence. By the properties of the Thue-Morse sequence, we have for $n \geq 0$

$$\begin{aligned} M_{n+1}(x) &= W_n(x) \cdot M_n(x), \\ W_{n+1}(x) &= M_n(x) \cdot W_n(x). \end{aligned}$$

Let $x := 1/z$. For each non-zero polynomial $P(z)$, define $\tilde{P}(x)$ to be $P(1/x)$. Then,

$$\text{CF}_n(a, b) = \frac{P_n(z)}{Q_n(z)} = \frac{\tilde{P}_n(x)}{\tilde{Q}_n(x)} \in \mathbb{F}_2((x)) = \mathbb{F}_2((1/z)).$$

Comparing the definition of $M_n(x)$ with the definition of $P_n(x)$ and $Q_n(x)$, we see that

$$\begin{aligned} M_n(x)_{0,1} &= x^{d_n} \tilde{P}_{2^n-1}(x), \\ M_n(x)_{0,0} &= x^{d_n} \tilde{Q}_{2^n-1}(x), \end{aligned}$$

for some positive integer d_n . Hence,

$$(2.2) \quad \text{CF}_{2^{2^n}-1}(a, b) = \frac{\tilde{P}_{2^{2^n}-1}(x)}{\tilde{Q}_{2^{2^n}-1}(x)} = \frac{M_{2^n}(x)_{0,1}}{M_{2^n}(x)_{0,0}}.$$

Therefore, by the convergence theorem of continued fraction, the algebraicity of $\text{CF}(a, b)$ will be established if it is shown that both $M_{2^n}(x)_{0,1}$ and $M_{2^n}(x)_{0,0}$ converge to algebraic series in $\mathbb{F}_2[[x]]$.

Actually, we will prove that for all $0 \leq i, j \leq 1$ the four sequences $(M_{2^n}(x)_{i,j})_n$, $(M_{2^{n+1}}(x)_{i,j})_n$, $(W_{2^n}(x)_{i,j})_n$, and $(W_{2^{n+1}}(x)_{i,j})_n$ converge to algebraic series in $\mathbb{F}_2[[x]]$. For this purpose, we define four 2×2 matrices M^e, M^o, W^e, W^o as follows: For each $T \in \{M^e, M^o, W^e, W^o\}$ and $0 \leq i, j \leq 1$, $T_{i,j}$ is defined to be the unique solution in $\mathbb{F}_2[[x]]$ of the polynomial $\phi(T, i, j)$ under certain initial conditions; the polynomials $\phi(T, i, j)$ and initial conditions are given in Section 3. We will prove that these four matrices, whose components are algebraic by definition, are the limits of $(M_{2^n}(x))_n$, $(M_{2^{n+1}}(x))_n$, $(W_{2^n}(x))_n$, and $(W_{2^{n+1}}(x))_n$.

Let us explain how the polynomials $\phi(T, i, j)$ and initial conditions are found, and why the solutions exist and are unique. For $0 \leq i, j \leq 1$, the coefficients of the polynomial $\phi(M^e, i, j)$ (resp. $\phi(M^o, i, j)$, $\phi(W^e, i, j)$, and $\phi(W^o, i, j)$) are the Padé-Hermite approximants of type

$$(175, 175, 175, 175, 175)$$

of the vector

$$(1, T^3, T^6, T^9, T^{12}),$$

where $T = M_{12,i,j}$ (resp. $M_{11,i,j}$, $W_{12,i,j}$, and $W_{11,i,j}$), found by the Derksen algorithm. We take the first 8 terms of $M_{12,i,j}$ (resp. $M_{11,i,j}$, $W_{12,i,j}$, and $W_{11,i,j}$) as initial conditions for $\phi(M^e, i, j)$ (resp. $\phi(M^o, i, j)$, $\phi(W^e, i, j)$, and $\phi(W^o, i, j)$). The following fact will be used to ensure that the solution exists and is unique: let $P(x, y) \in \mathbb{F}_2[x, y]$ and for each series $f(x) = \sum_{n=0}^{\infty} a_n x^n \in \mathbb{F}_2[[x]]$ denote the partial sum $\sum_{j=0}^{n-1} a_j x^j$ by $f_n(x)$ for $n \geq 0$. If for some $n \geq 0$ and $a_0, a_1, \dots, a_{n-1} \in \mathbb{F}_2$ $P(x, \sum_{j=0}^{n-1} a_j x^j) = O(x^n)$ and $Q(x, y) := P(x, \sum_{j=0}^{n-1} a_j x^j + x^n y)$ can be written as $x^m \sum_{j=0}^{\infty} q_j(x) y^j$ where $q_j(x)$ are polynomials for $j \geq 0$, $q_1(0) = 1$, and $q_j(0) = 0$ for $j > 1$, then there exists a unique solution $f(x) \in \mathbb{F}_2[[x]]$ of $P(x, f(x)) = 0$ that satisfies the initial condition $f_n(x) = \sum_{j=0}^{n-1} a_j x^j$.

We state two lemmas concerning the four matrices M^e, M^o, W^e, W^o . The first one is about relations between them; the second, about the structure of the each matrix.

Lemma 2.1. *We have*

$$(2.3) \quad M^e = W^o \cdot M^o,$$

$$(2.4) \quad M^o = W^e \cdot M^e,$$

$$(2.5) \quad W^e = M^o \cdot W^o,$$

$$(2.6) \quad W^o = M^e \cdot W^e.$$

Proof. We give the proof of the identity

$$M_{0,0}^e = W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o,$$

the proofs of the others being similar.

First, we compute the minimal polynomials of $W_{0,0}^o M_{0,0}^o$ and $W_{0,1}^o M_{1,0}^o$. We know that

$$P(x, y) = \text{Res}_z (\phi(W^o, 0, 0)(x, z), z^{12} \cdot \phi(M^o, 0, 0)(x, y/z))$$

is an annihilating polynomial of $W_{0,0}^o M_{0,0}^o$. We use Padé-Hermite approximation to find a candidate for the minimal polynomial of $W_{0,0}^o M_{0,0}^o$, that will be called $\phi_0(x, y)$. To prove that $\phi_0(x, y)$ is indeed the minimal polynomial, it suffices to prove that it is an irreducible factor of $P(x, y)$ of multiplicity m and that $Q(x, y) := P(x, y)/\phi_0(x, y)^m$ is not an annihilating polynomial of $W_{0,0}^o M_{0,0}^o$. We verify the first point directly. For the second point, we truncate $W_{0,0}^o M_{0,0}^o$ to order 630 and substitute it for y in $Q(x, y)$. We get a series of valuation less than 630, which proves that $Q(x, y)$ is not an annihilating polynomial of $W_{0,0}^o M_{0,0}^o$. We find the minimal polynomial $\phi_1(x, y)$ of $W_{0,1}^o M_{1,0}^o$ in a similar way.

Now we prove that $\phi(M^e, 0, 0)$ is the minimal polynomial of $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$. We know that

$$S(x, y) = \text{Res}_z (\phi_0(x, z), \phi_1(x, y + z))$$

is an annihilating of $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$. We verify that $\phi(M^e, 0, 0)$ is an irreducible factor of $S(x, y)$ of multiplicity μ , and that $Q(x, y) := S(x, y)/\phi(M^e, 0, 0)^\mu$ is not an annihilating polynomial of $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$. To see the last point, we truncate $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$ to order 770 and substitute it for y in $Q(x, y)$. We get a series of valuation less than 770, and therefore $Q(x, y)$ is not an annihilating polynomial of $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$.

Finally, the first 16 terms of $M_{0,0}^e$ and $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$ coincide. As these first terms determine a unique solution of $\phi(M^e, 0, 0)$, we know that the two series are one and the same. $\square$

Define

$$R^e = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \quad \text{and} \quad R^o = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}.$$

Lemma 2.2. *For all integers $k \geq 2$ and $u = 2^{2k-1}$, the following identities hold.*

$$\begin{aligned} M^e[:14u] &= M^e[:7u] + x^{2u} M^e[:5u] + x^{3u} M^e[:4u] + x^{5u} M^e[:2u] + x^{6u} M^e[:u] \\ &\quad + x^{2u} M^e[:7u] + x^{4u} M^e[:5u] + x^{5u} M^e[:4u] + x^{7u} M^e[:2u] \\ &\quad + x^{8u} M^e[:u] + x^{3u} M^e[:7u] + x^{5u} M^e[:5u] + x^{6u} M^e[:4u] \\ &\quad + x^{8u} M^e[:2u] + x^{9u} M^e[:u] + x^{4u} M^e[:7u] + x^{6u} M^e[:5u] \\ &\quad + x^{7u} M^e[:4u] + x^{9u} M^e[:2u] + x^{10u} M^e[:u] + x^{5u} M^e[:7u] \\ &\quad + x^{7u} M^e[:5u] + x^{8u} M^e[:4u] + x^{10u} M^e[:2u] + x^{11u} M^e[:u] \end{aligned}$$

$$\begin{aligned}
& + x^{6u} M^e[:7u] + x^{8u} M^e[:5u] + x^{9u} M^e[:4u] + x^{11u} M^e[:2u] \\
& + x^{12u} M^e[:u] + x^{7u} M^e[:7u] + x^{8u} M^e[:6u] + x^{11u} M^e[:3u] \\
& + x^{13u} M^e[:u] + x^{12u} M^e[:2u] \\
& + (x^{7u} + x^{9u} + x^{10u} + x^{11u} + x^{12u} + x^{13u}) R^e, \\
W^e[:14u] = & W^e[:7u] + x^{2u} W^e[:5u] + x^{3u} W^e[:4u] + x^{5u} W^e[:2u] + x^{6u} W^e[:u] \\
& + x^{2u} W^e[:7u] + x^{4u} W^e[:5u] + x^{5u} W^e[:4u] + x^{7u} W^e[:2u] \\
& + x^{8u} W^e[:u] + x^{3u} W^e[:7u] + x^{5u} W^e[:5u] + x^{6u} W^e[:4u] \\
& + x^{8u} W^e[:2u] + x^{9u} W^e[:u] + x^{4u} W^e[:7u] + x^{6u} W^e[:5u] \\
& + x^{7u} W^e[:4u] + x^{9u} W^e[:2u] + x^{10u} W^e[:u] + x^{5u} W^e[:7u] \\
& + x^{7u} W^e[:5u] + x^{8u} W^e[:4u] + x^{10u} W^e[:2u] + x^{11u} W^e[:u] \\
& + x^{6u} W^e[:7u] + x^{8u} W^e[:5u] + x^{9u} W^e[:4u] + x^{11u} W^e[:2u] \\
& + x^{12u} W^e[:u] + x^{7u} W^e[:7u] + x^{8u} W^e[:6u] + x^{11u} W^e[:3u] \\
& + x^{13u} W^e[:u] + x^{12u} W^e[:2u] \\
& + (x^{7u} + x^{9u} + x^{10u} + x^{11u} + x^{12u} + x^{13u}) R^e.
\end{aligned}$$

For all integers $k \geq 2$ and $u = 2^{2k}$, the following identities hold.

$$\begin{aligned}
M^o[:14u] = & M^o[:7u] + x^{2u} M^o[:5u] + x^{3u} M^o[:4u] + x^{5u} M^o[:2u] + x^{6u} M^o[:u] \\
& + x^{2u} M^o[:7u] + x^{4u} M^o[:5u] + x^{5u} M^o[:4u] + x^{7u} M^o[:2u] \\
& + x^{8u} M^o[:u] + x^{3u} M^o[:7u] + x^{5u} M^o[:5u] + x^{6u} M^o[:4u] \\
& + x^{8u} M^o[:2u] + x^{9u} M^o[:u] + x^{4u} M^o[:7u] + x^{6u} M^o[:5u] \\
& + x^{7u} M^o[:4u] + x^{9u} M^o[:2u] + x^{10u} M^o[:u] + x^{5u} M^o[:7u] \\
& + x^{7u} M^o[:5u] + x^{8u} M^o[:4u] + x^{10u} M^o[:2u] + x^{11u} M^o[:u] \\
& + x^{6u} M^o[:7u] + x^{8u} M^o[:5u] + x^{9u} M^o[:4u] + x^{11u} M^o[:2u] \\
& + x^{12u} M^o[:u] + x^{7u} M^o[:7u] + x^{8u} M^o[:6u] + x^{11u} M^o[:3u] \\
& + x^{13u} M^o[:u] + x^{12u} M^o[:2u] \\
& + (x^{7u} + x^{9u} + x^{10u} + x^{11u} + x^{12u} + x^{13u}) R^o, \\
W^o[:14u] = & W^o[:7u] + x^{2u} W^o[:5u] + x^{3u} W^o[:4u] + x^{5u} W^o[:2u] + x^{6u} W^o[:u] \\
& + x^{2u} W^o[:7u] + x^{4u} W^o[:5u] + x^{5u} W^o[:4u] + x^{7u} W^o[:2u] \\
& + x^{8u} W^o[:u] + x^{3u} W^o[:7u] + x^{5u} W^o[:5u] + x^{6u} W^o[:4u] \\
& + x^{8u} W^o[:2u] + x^{9u} W^o[:u] + x^{4u} W^o[:7u] + x^{6u} W^o[:5u] \\
& + x^{7u} W^o[:4u] + x^{9u} W^o[:2u] + x^{10u} W^o[:u] + x^{5u} W^o[:7u] \\
& + x^{7u} W^o[:5u] + x^{8u} W^o[:4u] + x^{10u} W^o[:2u] + x^{11u} W^o[:u] \\
& + x^{6u} W^o[:7u] + x^{8u} W^o[:5u] + x^{9u} W^o[:4u] + x^{11u} W^o[:2u] \\
& + x^{12u} W^o[:u] + x^{7u} W^o[:7u] + x^{8u} W^o[:6u] + x^{11u} W^o[:3u] \\
& + x^{13u} W^o[:u] + x^{12u} W^o[:2u] \\
& + (x^{7u} + x^{9u} + x^{10u} + x^{11u} + x^{12u} + x^{13u}) R^o.
\end{aligned}$$

Proof. To prove Lemma 2.2 we first construct an automaton for each sequence concerned, and then transform the conditions on infinitely many k 's into finitely many conditions on the states of the automaton. In the following, we will prove that for $T = M_{0,0}^0$, for all integer $k \geq 2$ and $u = 2^{2k}$,

$$\begin{aligned}
T[:14u] = & T[:7u] + x^{2u}T[:5u] + x^{3u}T[:4u] + x^{5u}T[:2u] + x^{6u}T[:u] + x^{2u}T[:7u] \\
& + x^{4u}T[:5u] + x^{5u}T[:4u] + x^{7u}T[:2u] + x^{8u}T[:u] + x^{3u}T[:7u] \\
& + x^{5u}T[:5u] + x^{6u}T[:4u] + x^{8u}T[:2u] + x^{9u}T[:u] + x^{4u}T[:7u] \\
& + x^{6u}T[:5u] + x^{7u}T[:4u] + x^{9u}T[:2u] + x^{10u}T[:u] + x^{5u}T[:7u] \\
& + x^{7u}T[:5u] + x^{8u}T[:4u] + x^{10u}T[:2u] + x^{11u}T[:u] + x^{6u}T[:7u] \\
& + x^{8u}T[:5u] + x^{9u}T[:4u] + x^{11u}T[:2u] + x^{12u}T[:u] + x^{7u}T[:7u] \\
& + x^{8u}T[:6u] + x^{11u}T[:3u] + x^{13u}T[:u] + x^{12u}T[:2u].
\end{aligned}$$

The proof of the other 15 cases are similar. We break down the above identity into 7 parts:

$$\begin{aligned}
0 = & x^{6u}T[u:2u] + x^{5u}T[2u:3u] + x^{3u}T[4u:5u] + x^{2u}T[5u:6u] \\
& + T[7u:8u], \\
0 = & x^{8u}T[:u] + x^{6u}T[2u:3u] + x^{5u}T[3u:4u] + x^{3u}T[5u:6u] \\
& + x^{2u}T[6u:7u] + T[8u:9u], \\
0 = & x^{9u}T[:u] + x^{7u}T[2u:3u] + x^{6u}T[3u:4u] + x^{4u}T[5u:6u] \\
& + x^{3u}T[6u:7u] + T[9u:10u], \\
0 = & x^{8u}T[2u:3u] + x^{7u}T[3u:4u] + x^{5u}T[5u:6u] + x^{4u}T[6u:7u] \\
& + T[10u:11u], \\
0 = & x^{11u}T[:u] + x^{10u}T[u:2u] + x^{9u}T[2u:3u] + x^{8u}T[3u:4u] \\
& + x^{6u}T[5u:6u] + x^{5u}T[6u:7u] + T[11u:12u], \\
0 = & x^{9u}T[3u:4u] + x^{7u}T[5u:6u] + x^{6u}T[6u:7u] + T[12u:13u], \\
0 = & x^{13u}T[:u] + x^{12u}T[u:2u] + x^{11u}T[2u:3u] + x^{8u}T[5u:6u] \\
& + x^{7u}T[6u:7u] + T[13u:14u],
\end{aligned}$$

which can be rewritten as

$$(2.7) \quad 0 = T[[1w]_2] + T[[10w]_2] + T[[100w]_2] + T[[101w]_2] + T[[111w]_2],$$

$$0 = T[[w]_2] + T[[10w]_2] + T[[11w]_2] + T[[101w]_2] + T[[110w]_2]$$

$$(2.8) \quad + T[[1000w]_2],$$

$$0 = T[[w]_2] + T[[10w]_2] + T[[11w]_2] + T[[101w]_2] + T[[110w]_2]$$

$$(2.9) \quad + T[[1001w]_2],$$

$$(2.10) \quad 0 = T[[10w]_2] + T[[11w]_2] + T[[101w]_2] + T[[110w]_2] + T[[1010w]_2],$$

$$0 = T[[w]_2] + T[[1w]_2] + T[[10w]_2] + T[[11w]_2] + T[[101w]_2]$$

$$(2.11) \quad + T[[110w]_2] + T[[1011w]_2],$$

$$(2.12) \quad 0 = T[[11w]_2] + T[[101w]_2] + T[[110w]_2] + T[[1100w]_2],$$

$$0 = T[[w]_2] + T[[1w]_2] + T[[10w]_2] + T[[101w]_2] + T[[110w]_2]$$

$$(2.13) \quad + T[[1101w]_2],$$

for all binary word w of length $2k$ and $w \neq 0^{2k}$, and

$$(2.14) \quad 0 = T[[1w]_2] + T[[10w]_2] + T[[100w]_2] + T[[101w]_2] + T[[111w]_2],$$

$$0 = T[[w]_2] + T[[10w]_2] + T[[11w]_2] + T[[101w]_2] + T[[110w]_2]$$

$$(2.15) \quad + T[[1000w]_2],$$

$$0 = T[[w]_2] + T[[10w]_2] + T[[11w]_2] + T[[101w]_2] + T[[110w]_2]$$

$$(2.16) \quad + T[[1001w]_2],$$

$$(2.17) \quad 0 = T[[10w]_2] + T[[11w]_2] + T[[101w]_2] + T[[110w]_2] + T[[1010w]_2],$$

$$0 = T[[w]_2] + T[[1w]_2] + T[[10w]_2] + T[[11w]_2] + T[[101w]_2]$$

$$(2.18) \quad + T[[110w]_2] + T[[1011w]_2],$$

$$(2.19) \quad 0 = T[[11w]_2] + T[[101w]_2] + T[[110w]_2] + T[[1100w]_2],$$

$$0 = T[[w]_2] + T[[1w]_2] + T[[10w]_2] + T[[101w]_2] + T[[110w]_2]$$

$$(2.20) \quad + T[[1101w]_2],$$

for $w = 0^{2k}$.

First we calculate an 2-automaton that generates T from its minimal polynomial and its first terms. This automaton has 580 states; its transition function and output function can be found in the annex. Let $A(s, w)$ denote the state reached after reading w from right to left starting from the state s , and τ the output function. Define

$$E_{2k} = \{A(i, w) : |w| = 2k, w \neq 0^{2k}\}.$$

Identities (2.7) through (2.20) can be written as

$$(2.21) \quad \begin{aligned} 0 &= \tau(A(s, 1)) + \tau(A(s, 10)) + \tau(A(s, 100)) + \tau(A(s, 101)) \\ &\quad + \tau(A(s, 111)), \end{aligned}$$

$$(2.22) \quad \begin{aligned} 0 &= \tau(A(s, \epsilon)) + \tau(A(s, 10)) + \tau(A(s, 11)) + \tau(A(s, 101)) \\ &\quad + \tau(A(s, 110)) + \tau(A(s, 1000)), \end{aligned}$$

$$(2.23) \quad \begin{aligned} 0 &= \tau(A(s, \epsilon)) + \tau(A(s, 10)) + \tau(A(s, 11)) + \tau(A(s, 101)) \\ &\quad + \tau(A(s, 110)) + \tau(A(s, 1001)), \end{aligned}$$

$$(2.24) \quad \begin{aligned} 0 &= \tau(A(s, 10)) + \tau(A(s, 11)) + \tau(A(s, 101)) + \tau(A(s, 110)) \\ &\quad + \tau(A(s, 1010)), \end{aligned}$$

$$(2.25) \quad \begin{aligned} 0 &= \tau(A(s, \epsilon)) + \tau(A(s, 1)) + \tau(A(s, 10)) + \tau(A(s, 11)) + \tau(A(s, 101)) \\ &\quad + \tau(A(s, 110)) + \tau(A(s, 1011)), \end{aligned}$$

$$(2.26) \quad \begin{aligned} 0 &= \tau(A(s, 11)) + \tau(A(s, 101)) + \tau(A(s, 110)) + \tau(A(s, 1100)), \\ 0 &= \tau(A(s, \epsilon)) + \tau(A(s, 1)) + \tau(A(s, 10)) + \tau(A(s, 101)) \end{aligned}$$

$$(2.27) \quad + \tau(A(s, 110)) + \tau(A(s, 1101)),$$

for all $s \in E_{2k}$, and

$$(2.28) \quad \begin{aligned} 0 &= \tau(A(s, 1)) + \tau(A(s, 10)) + \tau(A(s, 100)) + \tau(A(s, 101)) \\ &\quad + \tau(A(s, 111)), \end{aligned}$$

$$0 = \tau(A(s, \epsilon)) + \tau(A(s, 10)) + \tau(A(s, 11)) + \tau(A(s, 101))$$

$$\begin{aligned}
(2.29) \quad & + \tau(A(s, 110)) + \tau(A(s, 1000)), \\
& 0 = \tau(A(s, \epsilon)) + \tau(A(s, 10)) + \tau(A(s, 11)) + \tau(A(s, 101)) \\
(2.30) \quad & + \tau(A(s, 110)) + \tau(A(s, 1001)), \\
& 0 = \tau(A(s, 10)) + \tau(A(s, 11)) + \tau(A(s, 101)) + \tau(A(s, 110)) \\
(2.31) \quad & + \tau(A(s, 1010)), \\
& 0 = \tau(A(s, \epsilon)) + \tau(A(s, 1)) + \tau(A(s, 10)) + \tau(A(s, 11)) + \tau(A(s, 101)) \\
(2.32) \quad & + \tau(A(s, 110)) + \tau(A(s, 1011)), \\
(2.33) \quad & 0 = \tau(A(s, 11)) + \tau(A(s, 101)) + \tau(A(s, 110)) + \tau(A(s, 1100)), \\
& 0 = \tau(A(s, \epsilon)) + \tau(A(s, 1)) + \tau(A(s, 10)) + \tau(A(s, 101)) \\
(2.34) \quad & + \tau(A(s, 110)) + \tau(A(s, 1101)),
\end{aligned}$$

for $s = A(i, 0^{2k})$. We find that $(A(i, 0^{24}), E_{24}) = (A(i, 0^{16}), E_{16})$, so that we only have to verify that identities (2.21) through (2.34) hold for $2 \leq k \leq 12$, which turns out to be true. $\square$

In the following lemma, we express M_{2k} , M_{2k+1} , W_{2k} , and W_{2k+1} in terms of M^e , M^o , W^e , and W^o .

Lemma 2.3. *For all integer $k \geq 2$, and $u = 2^{2k-1}$,*

$$\begin{aligned}
M_{2k} &= M^e[:7u] + x^{2u} M^e[:5u] + x^{3u} M^e[:4u] + x^{5u} M^e[:2u] + x^{6u} M^e[:u] \\
&\quad + x^{7u} R^e, \\
W_{2k} &= W^e[:7u] + x^{2u} W^e[:5u] + x^{3u} W^e[:4u] + x^{5u} W^e[:2u] + x^{6u} W^e[:u] \\
&\quad + x^{7u} R^e.
\end{aligned}$$

For all integer $k \geq 2$, and $u = 2^{2k}$,

$$\begin{aligned}
M_{2k+1} &= M^o[:7u] + x^{2u} M^o[:5u] + x^{3u} M^o[:4u] + x^{5u} M^o[:2u] + x^{6u} M^o[:u] \\
&\quad + x^{7u} R^o, \\
W_{2k+1} &= W^o[:7u] + x^{2u} W^o[:5u] + x^{3u} W^o[:4u] + x^{5u} W^o[:2u] + x^{6u} W^o[:u] \\
&\quad + x^{7u} R^o.
\end{aligned}$$

Proof. We call the four identities in Lemma 2.3 also by the name M_{2k} , W_{2k} , M_{2k+1} , and W_{2k+1} . For $n = 2$, the identities can be verified directly. For $n \geq 2$, we claim that

$$\begin{aligned}
M_{2k} \wedge W_{2k} &\Rightarrow M_{2k+1} \wedge W_{2k+1}, \\
M_{2k+1} \wedge W_{2k+1} &\Rightarrow M_{2k+2} \wedge W_{2k+2}.
\end{aligned}$$

We give the proof of

$$M_{2k} \wedge W_{2k} \Rightarrow M_{2k+1},$$

the other ones being similar. Set $u = 2^{2k}$ and $v = 2^{2k-1}$. By definition and induction hypothesis, the left side of identity M_{2k+1} is equal to

$$\begin{aligned}
(2.35) \quad & W_{2k} M_{2k} = (W^e[:7v] + x^{2v} W^e[:5v] + x^{3v} W^e[:4v] + x^{5v} W^e[:2v] + x^{6v} W^e[:v] \\
&\quad + x^{7v} R^e) \times (M^e[:7v] + x^{2v} M^e[:5v] + x^{3v} M^e[:4v] + x^{5v} M^e[:2v] \\
&\quad + x^{6v} M^e[:v] + x^{7v} R^e);
\end{aligned}$$

Call this expression *lhs*. Note that both sides of identity M_{2k+1} have the same term of highest degree $x^{14v}R^o$. Therefore we only need to prove that their difference is $O(x^{14v})$. Using Lemma 2.1 it can be seen that the right side of identity M_{2k+1} is congruent, modulo x^{14v} , to

$$\begin{aligned} W^e[:14v]M^e[:14v] + x^{4v}W^e[:10v]M^e[:10v] + x^{6v}W^e[:8v]M^e[:8v] \\ + x^{10v}W^e[:4v]M^e[:4v] + x^{12v}W^e[:2v]M^e[:2v]. \end{aligned}$$

For all $n \leq 14$, replace the occurrences of $W^e[:n \cdot v]$ and $M^e[:n \cdot v]$ in the above expression by the reduction modulo $x^{n \cdot v}$ of the right side of the corresponding identity in Lemma 2.2 and get a new expression, which we call *rhs*. Define

$$\begin{aligned} X &:= x^v, \\ a_n &:= W^e[n \cdot v : (n+1) \cdot v] / X^n, \\ b_n &:= M^e[n \cdot v : (n+1) \cdot v] / X^n, \\ c &:= R^e. \end{aligned}$$

Using the notation introduced above, we can represent the expressions *lhs* (2.35) and *rhs* as polynomials in $\mathbb{F}_2[a_1, \dots, a_{14}, b_1, \dots, b_{14}, c][X]$. Note that it is not a problem that a_j commutes with b_k while W^e does not commute with M^e , because in the expressions concerned, the products of W^e -terms and M^e -terms are always in the same order. We let the computer do the simplification and check that the difference between these two polynomials is indeed $O(X^{14})$, which completes the proof. $\square$

Proof of Theorem 1.1. We prove the theorem for $\text{CF}(a, b)$; for $\text{CF}(b, a)$, the proof is similar. By Lemma 2.3, we have For all $0 \leq j, k \leq 1$,

$$\begin{aligned} \lim_{n \rightarrow \infty} M_{2n,j,k} &= M_{j,k}^e, \\ \lim_{n \rightarrow \infty} M_{2n+1,j,k} &= M_{j,k}^o, \\ \lim_{n \rightarrow \infty} W_{2n,j,k} &= W_{j,k}^e, \\ \lim_{n \rightarrow \infty} W_{2n+1,j,k} &= W_{j,k}^o. \end{aligned}$$

Let $z = 1/x$. By the convergence theorem and identity (2.2),

$$\text{CF}(a, b) = \frac{M_{0,1}^e(x)}{M_{0,0}^e(x)}.$$

By definition, $\phi(M^e, 0, 1)$ and $\phi(M^e, 0, 0)$ are minimal polynomials of $M_{0,1}^e$ and $M_{0,0}^e$. Therefore

$$P(x, y) = \text{Res}_t (\phi(M^e, 0, 1)(x, t), y^{12}\phi(M^e, 0, 1)(x, t/y))$$

is an annihilating polynomial of $f(x) = M_{0,1}^e/M_{0,0}^e$.

Define

$$Q(x, y) = q_4(x)y^4 + q_3(x)y^3 + q_2(x)y^2 + q_1(x)y + q_0(x),$$

where

$$\begin{aligned} q_0(x) &= x^{22} + x^{20} + x^{19} + x^{18} + x^{17} + x^{13} + x^{11} + x^{10} + x^9 + x^7, \\ q_1(x) &= x^{18} + x^{16} + x^{15} + x^{13} + x^{12} + x^{10} + x^9 + x^8 + x^7 + x^6 + x^5 + x^3, \\ q_2(x) &= x^{20} + x^{16} + x^{10} + x^8 + x^6 + 1, \end{aligned}$$

$$\begin{aligned}
q_3(x) &= x^{18} + x^{16} + x^{15} + x^{13} + x^{12} + x^{10} + x^9 + x^8 + x^7 + x^6 + x^5 + x^3, \\
q_4(x) &= x^{22} + x^{20} + x^{19} + x^{18} + x^{17} + x^{16} + x^{13} + x^{11} + x^9 + x^7 + x^6.
\end{aligned}$$

The polynomial $Q(x, y)$ is the candidate for the minimal polynomial of $f(x)$ found by Padé-Hermite approximation. To prove that it is indeed the minimal polynomial of $f(x)$, we only need to prove that it is an irreducible factor of $P(x, y)$ of multiplicity m and $R(x, y) := P(x, y)/Q(x, y)^m$ is not an annihilating polynomial of $f(x)$. We verify the first point directly. For the second point, we find that when we truncate $f(x)$ to order 216, and substitute it for y in $R(z, y)$, we get a series with valuation smaller than 216, which proves that $R(z, y)$ is not an annihilating polynomial of $f(x)$. Finally, $z^{12}Q(1/z, y)$ is the minimal polynomial of $\text{CF}(a, b) = f(1/z)$. $\square$

3. ANNEXE

All 16 polynomials are of the form

$$p_0(x) + p_3(x)y^3 + p_6(x)y^6 + p_9(x)y^9 + p_{12}(x)y^{12}.$$

The coefficients $p_j(x)$ and the 16 initial terms to determine the solutions uniquely are given below.

For $\phi(M^e, 0, 0)$:

$$\begin{aligned}
p_0(x) &= x^{120} + x^{118} + x^{112} + x^{110} + x^{106} + x^{104} + x^{102} + x^{98} + x^{96} + x^{94} \\
&\quad + x^{92} + x^{88} + x^{82} + x^{76} + x^{74} + x^{72} + x^{66} + x^{58} + x^{56} + x^{54} + x^{52} \\
&\quad + x^{40} + x^{36}, \\
p_3(x) &= x^{108} + x^{106} + x^{104} + x^{96} + x^{90} + x^{78} + x^{66} + x^{64} + x^{62} + x^{58} + x^{54} \\
&\quad + x^{50} + x^{48} + x^{40} + x^{36} + x^{32} + x^{28} + x^{26} + x^{22} + x^{20}, \\
p_6(x) &= x^{112} + x^{110} + x^{104} + x^{102} + x^{98} + x^{96} + x^{94} + x^{90} + x^{88} + x^{84} + x^{80} \\
&\quad + x^{78} + x^{74} + x^{68} + x^{66} + x^{64} + x^{62} + x^{58} + x^{56} + x^{54} + x^{44} + x^{42} \\
&\quad + x^{40} + x^{38} + x^{36} + x^{28} + x^{26} + x^{18} + x^{16} + x^{10} + x^6 + x^4 + x^2 + 1, \\
p_9(x) &= x^{104} + x^{102} + x^{100} + x^{98} + x^{94} + x^{92} + x^{90} + x^{88} + x^{84} + x^{78} + x^{76} \\
&\quad + x^{74} + x^{72} + x^{70} + x^{62} + x^{60} + x^{50} + x^{48} + x^{46} + x^{42} + x^{40} + x^{36} \\
&\quad + x^{34} + x^{30} + x^{26} + x^{24} + x^{22} + x^{20} + x^{18} + x^{16} + x^{14} + x^{12} + x^8 \\
&\quad + x^6, \\
p_{12}(x) &= x^{108} + x^{106} + x^{104} + x^{100} + x^{98} + x^{96} + x^{92} + x^{90} + x^{88} + x^{84} + x^{82} \\
&\quad + x^{80} + x^{60} + x^{58} + x^{56} + x^{52} + x^{50} + x^{48} + x^{44} + x^{42} + x^{40} + x^{36} \\
&\quad + x^{34} + x^{32} + x^{28} + x^{26} + x^{24} + x^{20} + x^{18} + x^{16} + x^{12} + x^{10} + x^8 \\
&\quad + x^4 + x^2 + 1,
\end{aligned}$$

and the initial terms are $[1, 0, 0, 0, 0, 0, 0, 0, 1, 0, 1, 0, 1, 0, 1, 0]$.

For $\phi(M^e, 0, 1)$:

$$\begin{aligned}
p_0(x) &= x^{117} + x^{115} + x^{114} + x^{113} + x^{112} + x^{111} + x^{110} + x^{105} + x^{104} + x^{103} \\
&\quad + x^{101} + x^{100} + x^{99} + x^{97} + x^{96} + x^{94} + x^{91} + x^{90} + x^{87} + x^{85} + x^{81} \\
&\quad + x^{78} + x^{77} + x^{76} + x^{75} + x^{74} + x^{73} + x^{72} + x^{71} + x^{70} + x^{68} + x^{66}
\end{aligned}$$

$$\begin{aligned}
& + x^{61} + x^{60} + x^{58} + x^{55} + x^{50} + x^{48} + x^{45} + x^{43} + x^{41} + x^{40} + x^{39}, \\
p_3(x) &= x^{87} + x^{83} + x^{77} + x^{75} + x^{73} + x^{67} + x^{63} + x^{59} + x^{53} + x^{51} + x^{49} \\
& + x^{47} + x^{37} + x^{35} + x^{33} + x^{27}, \\
p_6(x) &= x^{93} + x^{91} + x^{90} + x^{89} + x^{88} + x^{86} + x^{85} + x^{83} + x^{82} + x^{80} + x^{79} \\
& + x^{77} + x^{76} + x^{74} + x^{69} + x^{67} + x^{66} + x^{64} + x^{63} + x^{60} + x^{59} + x^{56} \\
& + x^{55} + x^{52} + x^{51} + x^{49} + x^{48} + x^{46} + x^{45} + x^{43} + x^{42} + x^{41} + x^{40} \\
& + x^{38} + x^{25} + x^{23} + x^{22} + x^{21} + x^{20} + x^{18}, \\
p_9(x) &= x^{99} + x^{93} + x^{91} + x^{89} + x^{85} + x^{81} + x^{79} + x^{73} + x^{71} + x^{69} + x^{67} \\
& + x^{65} + x^{57} + x^{55} + x^{49} + x^{47} + x^{45} + x^{41} + x^{39} + x^{37} + x^{33} + x^{31} \\
& + x^{29} + x^{25} + x^{21} + x^{19} + x^{13} + x^9, \\
p_{12}(x) &= x^{105} + x^{103} + x^{102} + x^{100} + x^{97} + x^{95} + x^{94} + x^{93} + x^{92} + x^{91} + x^{90} \\
& + x^{88} + x^{85} + x^{83} + x^{82} + x^{80} + x^{57} + x^{55} + x^{54} + x^{52} + x^{49} + x^{47} \\
& + x^{46} + x^{45} + x^{44} + x^{43} + x^{42} + x^{40} + x^{37} + x^{35} + x^{34} + x^{32} + x^{25} \\
& + x^{23} + x^{22} + x^{20} + x^{17} + x^{15} + x^{14} + x^{13} + x^{12} + x^{11} + x^{10} + x^8 \\
& + x^5 + x^3 + x^2 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 1, 1, 0, 1, 1, 0, 1, 0, 0, 1, 1, 1, 0]$.

For $\phi(M^e, 1, 0)$:

$$\begin{aligned}
p_0(x) &= x^{117} + x^{115} + x^{114} + x^{113} + x^{112} + x^{111} + x^{110} + x^{105} + x^{104} + x^{103} \\
& + x^{101} + x^{100} + x^{99} + x^{97} + x^{96} + x^{94} + x^{91} + x^{90} + x^{87} + x^{85} + x^{81} \\
& + x^{78} + x^{77} + x^{76} + x^{75} + x^{74} + x^{73} + x^{72} + x^{71} + x^{70} + x^{68} + x^{66} \\
& + x^{61} + x^{60} + x^{58} + x^{55} + x^{50} + x^{48} + x^{45} + x^{43} + x^{41} + x^{40} + x^{39}, \\
p_3(x) &= x^{87} + x^{83} + x^{77} + x^{75} + x^{73} + x^{67} + x^{63} + x^{59} + x^{53} + x^{51} + x^{49} \\
& + x^{47} + x^{37} + x^{35} + x^{33} + x^{27}, \\
p_6(x) &= x^{93} + x^{91} + x^{90} + x^{89} + x^{88} + x^{86} + x^{85} + x^{83} + x^{82} + x^{80} + x^{79} \\
& + x^{77} + x^{76} + x^{74} + x^{69} + x^{67} + x^{66} + x^{64} + x^{63} + x^{60} + x^{59} + x^{56} \\
& + x^{55} + x^{52} + x^{51} + x^{49} + x^{48} + x^{46} + x^{45} + x^{43} + x^{42} + x^{41} + x^{40} \\
& + x^{38} + x^{25} + x^{23} + x^{22} + x^{21} + x^{20} + x^{18}, \\
p_9(x) &= x^{99} + x^{93} + x^{91} + x^{89} + x^{85} + x^{81} + x^{79} + x^{73} + x^{71} + x^{69} + x^{67} \\
& + x^{65} + x^{57} + x^{55} + x^{49} + x^{47} + x^{45} + x^{41} + x^{39} + x^{37} + x^{33} + x^{31} \\
& + x^{29} + x^{25} + x^{21} + x^{19} + x^{13} + x^9, \\
p_{12}(x) &= x^{105} + x^{103} + x^{102} + x^{100} + x^{97} + x^{95} + x^{94} + x^{93} + x^{92} + x^{91} + x^{90} \\
& + x^{88} + x^{85} + x^{83} + x^{82} + x^{80} + x^{57} + x^{55} + x^{54} + x^{52} + x^{49} + x^{47} \\
& + x^{46} + x^{45} + x^{44} + x^{43} + x^{42} + x^{40} + x^{37} + x^{35} + x^{34} + x^{32} + x^{25} \\
& + x^{23} + x^{22} + x^{20} + x^{17} + x^{15} + x^{14} + x^{13} + x^{12} + x^{11} + x^{10} + x^8 \\
& + x^5 + x^3 + x^2 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 1, 1, 0, 1, 1, 0, 1, 0, 0, 1, 1, 1, 0]$.

For $\phi(M^e, 1, 1)$:

$$\begin{aligned} p_0(x) = & x^{114} + x^{112} + x^{108} + x^{106} + x^{100} + x^{98} + x^{90} + x^{86} + x^{82} + x^{80} + x^{78} \\ & + x^{76} + x^{72} + x^{70} + x^{68} + x^{62} + x^{58} + x^{56} + x^{52} + x^{50} + x^{48} + x^{44} \\ & + x^{42}, \end{aligned}$$

$$\begin{aligned} p_3(x) = & x^{102} + x^{100} + x^{98} + x^{96} + x^{88} + x^{84} + x^{82} + x^{80} + x^{78} + x^{74} + x^{72} \\ & + x^{68} + x^{66} + x^{58} + x^{52} + x^{48} + x^{38} + x^{36} + x^{34} + x^{28} + x^{26} + x^{24} \\ & + x^{22} + x^{18}, \end{aligned}$$

$$\begin{aligned} p_6(x) = & x^{106} + x^{104} + x^{100} + x^{98} + x^{92} + x^{88} + x^{86} + x^{80} + x^{78} + x^{72} + x^{66} \\ & + x^{62} + x^{54} + x^{46} + x^{40} + x^{38} + x^{34} + x^{32} + x^{28} + x^{26} + x^{18} + x^{16} \\ & + x^{10} + x^4 + x^2 + 1, \end{aligned}$$

$$\begin{aligned} p_9(x) = & x^{98} + x^{96} + x^{94} + x^{88} + x^{80} + x^{76} + x^{74} + x^{68} + x^{66} + x^{64} + x^{62} \\ & + x^{60} + x^{52} + x^{48} + x^{44} + x^{42} + x^{40} + x^{38} + x^{34} + x^{32} + x^{28} + x^{26} \\ & + x^{24} + x^{20} + x^{18} + x^{14} + x^8 + x^6, \end{aligned}$$

$$\begin{aligned} p_{12}(x) = & x^{102} + x^{100} + x^{98} + x^{96} + x^{94} + x^{92} + x^{90} + x^{88} + x^{86} + x^{84} + x^{82} \\ & + x^{80} + x^{54} + x^{52} + x^{50} + x^{48} + x^{46} + x^{44} + x^{42} + x^{40} + x^{38} + x^{36} \\ & + x^{34} + x^{32} + x^{22} + x^{20} + x^{18} + x^{16} + x^{14} + x^{12} + x^{10} + x^8 + x^6 + x^4 \\ & + x^2 + 1, \end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 0, 1, 0, 1, 0, 0, 0, 1, 0, 0, 0]$.

For $\phi(W^e, 0, 0)$:

$$\begin{aligned} p_0(x) = & x^{114} + x^{112} + x^{108} + x^{106} + x^{102} + x^{96} + x^{94} + x^{92} + x^{90} + x^{86} + x^{84} \\ & + x^{82} + x^{78} + x^{76} + x^{74} + x^{72} + x^{70} + x^{64} + x^{60} + x^{58} + x^{54} + x^{48} \\ & + x^{46} + x^{44} + x^{42} + x^{40} + x^{36}, \end{aligned}$$

$$\begin{aligned} p_3(x) = & x^{102} + x^{100} + x^{98} + x^{96} + x^{90} + x^{80} + x^{78} + x^{74} + x^{72} + x^{68} + x^{66} \\ & + x^{58} + x^{52} + x^{50} + x^{44} + x^{40} + x^{38} + x^{28} + x^{22} + x^{20}, \end{aligned}$$

$$\begin{aligned} p_6(x) = & x^{106} + x^{104} + x^{100} + x^{98} + x^{94} + x^{88} + x^{86} + x^{84} + x^{76} + x^{72} + x^{66} \\ & + x^{62} + x^{52} + x^{34} + x^{28} + x^{26} + x^{20} + x^{18} + x^{14} + x^{12} + x^{10} + x^4 \\ & + x^2 + 1, \end{aligned}$$

$$\begin{aligned} p_9(x) = & x^{98} + x^{96} + x^{94} + x^{88} + x^{80} + x^{76} + x^{74} + x^{68} + x^{66} + x^{64} + x^{62} \\ & + x^{60} + x^{52} + x^{48} + x^{44} + x^{42} + x^{40} + x^{38} + x^{34} + x^{32} + x^{28} + x^{26} \\ & + x^{24} + x^{20} + x^{18} + x^{14} + x^8 + x^6, \end{aligned}$$

$$\begin{aligned} p_{12}(x) = & x^{102} + x^{100} + x^{98} + x^{96} + x^{94} + x^{92} + x^{90} + x^{88} + x^{86} + x^{84} + x^{82} \\ & + x^{80} + x^{54} + x^{52} + x^{50} + x^{48} + x^{46} + x^{44} + x^{42} + x^{40} + x^{38} + x^{36} \\ & + x^{34} + x^{32} + x^{22} + x^{20} + x^{18} + x^{16} + x^{14} + x^{12} + x^{10} + x^8 + x^6 + x^4 \\ & + x^2 + 1, \end{aligned}$$

and the initial terms are $[1, 0, 0, 0, 0, 0, 1, 0, 1, 0, 0, 0, 0, 0, 0]$.

For $\phi(W^e, 0, 1)$:

$$\begin{aligned}
p_0(x) &= x^{117} + x^{115} + x^{114} + x^{113} + x^{112} + x^{111} + x^{110} + x^{105} + x^{104} + x^{103} \\
&\quad + x^{100} + x^{98} + x^{93} + x^{88} + x^{85} + x^{84} + x^{77} + x^{74} + x^{72} + x^{71} + x^{70} \\
&\quad + x^{68} + x^{67} + x^{61} + x^{56} + x^{55} + x^{54} + x^{53} + x^{52} + x^{51} + x^{50} + x^{49} \\
&\quad + x^{47} + x^{45} + x^{44} + x^{43} + x^{41} + x^{40} + x^{39}, \\
p_3(x) &= x^{87} + x^{83} + x^{77} + x^{75} + x^{73} + x^{67} + x^{63} + x^{59} + x^{53} + x^{51} + x^{49} \\
&\quad + x^{47} + x^{37} + x^{35} + x^{33} + x^{27}, \\
p_6(x) &= x^{93} + x^{91} + x^{90} + x^{89} + x^{88} + x^{86} + x^{85} + x^{83} + x^{82} + x^{80} + x^{79} \\
&\quad + x^{77} + x^{76} + x^{74} + x^{69} + x^{67} + x^{66} + x^{64} + x^{63} + x^{60} + x^{59} + x^{56} \\
&\quad + x^{55} + x^{52} + x^{51} + x^{49} + x^{48} + x^{46} + x^{45} + x^{43} + x^{42} + x^{41} + x^{40} \\
&\quad + x^{38} + x^{25} + x^{23} + x^{22} + x^{21} + x^{20} + x^{18}, \\
p_9(x) &= x^{99} + x^{93} + x^{91} + x^{89} + x^{85} + x^{81} + x^{79} + x^{73} + x^{71} + x^{69} + x^{67} \\
&\quad + x^{65} + x^{57} + x^{55} + x^{49} + x^{47} + x^{45} + x^{41} + x^{39} + x^{37} + x^{33} + x^{31} \\
&\quad + x^{29} + x^{25} + x^{21} + x^{19} + x^{13} + x^9, \\
p_{12}(x) &= x^{105} + x^{103} + x^{102} + x^{100} + x^{97} + x^{95} + x^{94} + x^{93} + x^{92} + x^{91} + x^{90} \\
&\quad + x^{88} + x^{85} + x^{83} + x^{82} + x^{80} + x^{57} + x^{55} + x^{54} + x^{52} + x^{49} + x^{47} \\
&\quad + x^{46} + x^{45} + x^{44} + x^{43} + x^{42} + x^{40} + x^{37} + x^{35} + x^{34} + x^{32} + x^{25} \\
&\quad + x^{23} + x^{22} + x^{20} + x^{17} + x^{15} + x^{14} + x^{13} + x^{12} + x^{11} + x^{10} + x^8 \\
&\quad + x^5 + x^3 + x^2 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 1, 1, 0, 0, 1, 1, 1, 0, 1, 1, 1, 0]$.

For $\phi(W^e, 1, 0)$:

$$\begin{aligned}
p_0(x) &= x^{117} + x^{115} + x^{114} + x^{113} + x^{112} + x^{111} + x^{110} + x^{105} + x^{104} + x^{103} \\
&\quad + x^{100} + x^{98} + x^{93} + x^{88} + x^{85} + x^{84} + x^{77} + x^{74} + x^{72} + x^{71} + x^{70} \\
&\quad + x^{68} + x^{67} + x^{61} + x^{56} + x^{55} + x^{54} + x^{53} + x^{52} + x^{51} + x^{50} + x^{49} \\
&\quad + x^{47} + x^{45} + x^{44} + x^{43} + x^{41} + x^{40} + x^{39}, \\
p_3(x) &= x^{87} + x^{83} + x^{77} + x^{75} + x^{73} + x^{67} + x^{63} + x^{59} + x^{53} + x^{51} + x^{49} \\
&\quad + x^{47} + x^{37} + x^{35} + x^{33} + x^{27}, \\
p_6(x) &= x^{93} + x^{91} + x^{90} + x^{89} + x^{88} + x^{86} + x^{85} + x^{83} + x^{82} + x^{80} + x^{79} \\
&\quad + x^{77} + x^{76} + x^{74} + x^{69} + x^{67} + x^{66} + x^{64} + x^{63} + x^{60} + x^{59} + x^{56} \\
&\quad + x^{55} + x^{52} + x^{51} + x^{49} + x^{48} + x^{46} + x^{45} + x^{43} + x^{42} + x^{41} + x^{40} \\
&\quad + x^{38} + x^{25} + x^{23} + x^{22} + x^{21} + x^{20} + x^{18}, \\
p_9(x) &= x^{99} + x^{93} + x^{91} + x^{89} + x^{85} + x^{81} + x^{79} + x^{73} + x^{71} + x^{69} + x^{67} \\
&\quad + x^{65} + x^{57} + x^{55} + x^{49} + x^{47} + x^{45} + x^{41} + x^{39} + x^{37} + x^{33} + x^{31} \\
&\quad + x^{29} + x^{25} + x^{21} + x^{19} + x^{13} + x^9,
\end{aligned}$$

$$\begin{aligned}
p_{12}(x) = & x^{105} + x^{103} + x^{102} + x^{100} + x^{97} + x^{95} + x^{94} + x^{93} + x^{92} + x^{91} + x^{90} \\
& + x^{88} + x^{85} + x^{83} + x^{82} + x^{80} + x^{57} + x^{55} + x^{54} + x^{52} + x^{49} + x^{47} \\
& + x^{46} + x^{45} + x^{44} + x^{43} + x^{42} + x^{40} + x^{37} + x^{35} + x^{34} + x^{32} + x^{25} \\
& + x^{23} + x^{22} + x^{20} + x^{17} + x^{15} + x^{14} + x^{13} + x^{12} + x^{11} + x^{10} + x^8 \\
& + x^5 + x^3 + x^2 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 1, 1, 0, 0, 1, 1, 1, 0, 1, 1, 1, 0]$.

For $\phi(W^e, 1, 1)$:

$$\begin{aligned}
p_0(x) = & x^{120} + x^{118} + x^{112} + x^{110} + x^{108} + x^{102} + x^{100} + x^{96} + x^{92} + x^{88} \\
& + x^{80} + x^{76} + x^{72} + x^{66} + x^{64} + x^{58} + x^{56} + x^{52} + x^{50} + x^{44} + x^{42}, \\
p_3(x) = & x^{108} + x^{106} + x^{104} + x^{94} + x^{90} + x^{88} + x^{86} + x^{84} + x^{82} + x^{80} + x^{76} \\
& + x^{74} + x^{68} + x^{58} + x^{54} + x^{52} + x^{48} + x^{38} + x^{36} + x^{30} + x^{22} + x^{18}, \\
p_6(x) = & x^{112} + x^{110} + x^{104} + x^{102} + x^{100} + x^{96} + x^{90} + x^{88} + x^{84} + x^{82} + x^{74} \\
& + x^{72} + x^{68} + x^{66} + x^{62} + x^{56} + x^{50} + x^{48} + x^{46} + x^{44} + x^{40} + x^{36} \\
& + x^{26} + x^{24} + x^{20} + x^{14} + x^{12} + x^{10} + x^6 + x^4 + x^2 + 1, \\
p_9(x) = & x^{104} + x^{102} + x^{100} + x^{98} + x^{94} + x^{92} + x^{90} + x^{88} + x^{84} + x^{78} + x^{76} \\
& + x^{74} + x^{72} + x^{70} + x^{62} + x^{60} + x^{50} + x^{48} + x^{46} + x^{42} + x^{40} + x^{36} \\
& + x^{34} + x^{30} + x^{26} + x^{24} + x^{22} + x^{20} + x^{18} + x^{16} + x^{14} + x^{12} + x^8 \\
& + x^6, \\
p_{12}(x) = & x^{108} + x^{106} + x^{104} + x^{100} + x^{98} + x^{96} + x^{92} + x^{90} + x^{88} + x^{84} + x^{82} \\
& + x^{80} + x^{60} + x^{58} + x^{56} + x^{52} + x^{50} + x^{48} + x^{44} + x^{42} + x^{40} + x^{36} \\
& + x^{34} + x^{32} + x^{28} + x^{26} + x^{24} + x^{20} + x^{18} + x^{16} + x^{12} + x^{10} + x^8 \\
& + x^4 + x^2 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 0, 0, 0, 1, 0, 1, 0, 0, 0, 1, 0]$.

For $\phi(M^o, 0, 0)$:

$$\begin{aligned}
p_0(x) = & x^{117} + x^{115} + x^{114} + x^{113} + x^{112} + x^{110} + x^{107} + x^{103} + x^{99} + x^{96} \\
& + x^{95} + x^{94} + x^{93} + x^{91} + x^{89} + x^{87} + x^{86} + x^{76} + x^{73} + x^{72} + x^{70} \\
& + x^{69} + x^{64} + x^{62} + x^{61} + x^{58} + x^{57} + x^{52} + x^{49} + x^{47} + x^{43} + x^{39} \\
& + x^{36}, \\
p_3(x) = & x^{101} + x^{99} + x^{98} + x^{97} + x^{96} + x^{94} + x^{91} + x^{87} + x^{85} + x^{82} + x^{81} \\
& + x^{79} + x^{75} + x^{74} + x^{73} + x^{72} + x^{70} + x^{68} + x^{64} + x^{63} + x^{59} + x^{56} \\
& + x^{55} + x^{54} + x^{53} + x^{51} + x^{47} + x^{45} + x^{44} + x^{43} + x^{42} + x^{41} + x^{40} \\
& + x^{39} + x^{37} + x^{28} + x^{27} + x^{24}, \\
p_6(x) = & x^{109} + x^{107} + x^{106} + x^{104} + x^{103} + x^{100} + x^{99} + x^{95} + x^{93} + x^{91} \\
& + x^{90} + x^{88} + x^{87} + x^{86} + x^{84} + x^{83} + x^{82} + x^{79} + x^{77} + x^{76} + x^{73} \\
& + x^{72} + x^{68} + x^{66} + x^{58} + x^{57} + x^{56} + x^{55} + x^{54} + x^{53} + x^{51} + x^{49}
\end{aligned}$$

$$\begin{aligned}
& + x^{48} + x^{46} + x^{43} + x^{38} + x^{34} + x^{32} + x^{31} + x^{30} + x^{29} + x^{28} + x^{27} \\
& + x^{24} + x^{22} + x^{17} + x^{16} + x^{14} + x^{13} + x^{11} + x^{10} + x^8 + x^5 + x^3 + x^2 \\
& + 1, \\
p_9(x) &= x^{97} + x^{95} + x^{94} + x^{92} + x^{49} + x^{47} + x^{46} + x^{44} + x^{17} + x^{15} + x^{14} \\
& + x^{12}, \\
p_{12}(x) &= x^{105} + x^{103} + x^{102} + x^{100} + x^{93} + x^{91} + x^{90} + x^{88} + x^{85} + x^{83} + x^{82} \\
& + x^{80} + x^{57} + x^{55} + x^{54} + x^{52} + x^{45} + x^{43} + x^{42} + x^{40} + x^{37} + x^{35} \\
& + x^{34} + x^{32} + x^{25} + x^{23} + x^{22} + x^{20} + x^{13} + x^{11} + x^{10} + x^8 + x^5 + x^3 \\
& + x^2 + 1,
\end{aligned}$$

and the initial terms are $[1, 0, 0, 0, 0, 0, 1, 1, 0, 1, 0, 1, 0, 1, 1, 1]$.

For $\phi(M^o, 0, 1)$:

$$\begin{aligned}
p_0(x) &= x^{120} + x^{111} + x^{110} + x^{109} + x^{106} + x^{102} + x^{101} + x^{98} + x^{97} + x^{96} \\
& + x^{95} + x^{94} + x^{90} + x^{87} + x^{86} + x^{84} + x^{82} + x^{78} + x^{75} + x^{74} + x^{71} \\
& + x^{68} + x^{67} + x^{66} + x^{64} + x^{63} + x^{61} + x^{60} + x^{59} + x^{58} + x^{54} + x^{49} \\
& + x^{48} + x^{46} + x^{45} + x^{43} + x^{40} + x^{39}, \\
p_3(x) &= x^{90} + x^{88} + x^{87} + x^{85} + x^{84} + x^{82} + x^{81} + x^{79} + x^{78} + x^{76} + x^{75} \\
& + x^{73} + x^{72} + x^{70} + x^{69} + x^{67} + x^{66} + x^{64} + x^{63} + x^{61} + x^{60} + x^{58} \\
& + x^{57} + x^{55} + x^{54} + x^{52} + x^{51} + x^{50} + x^{49} + x^{47} + x^{46} + x^{44} + x^{43} \\
& + x^{42} + x^{41} + x^{39} + x^{38} + x^{36} + x^{35} + x^{33} + x^{32} + x^{30} + x^{29} + x^{27}, \\
p_6(x) &= x^{96} + x^{90} + x^{76} + x^{70} + x^{64} + x^{58} + x^{56} + x^{52} + x^{50} + x^{48} + x^{46} \\
& + x^{42} + x^{40} + x^{36} + x^{34} + x^{32} + x^{30} + x^{26} + x^{24} + x^{18}, \\
p_9(x) &= x^{102} + x^{100} + x^{99} + x^{98} + x^{97} + x^{96} + x^{95} + x^{94} + x^{93} + x^{92} + x^{91} \\
& + x^{89} + x^{54} + x^{52} + x^{51} + x^{50} + x^{49} + x^{48} + x^{47} + x^{46} + x^{45} + x^{44} \\
& + x^{43} + x^{41} + x^{22} + x^{20} + x^{19} + x^{18} + x^{17} + x^{16} + x^{15} + x^{14} + x^{13} \\
& + x^{12} + x^{11} + x^9, \\
p_{12}(x) &= x^{108} + x^{104} + x^{100} + x^{96} + x^{92} + x^{84} + x^{80} + x^{60} + x^{56} + x^{52} + x^{48} \\
& + x^{44} + x^{36} + x^{32} + x^{28} + x^{24} + x^{20} + x^{16} + x^{12} + x^4 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 1, 1, 0, 1, 1, 0, 0, 0, 0, 0, 1, 1, 1]$.

For $\phi(M^o, 1, 0)$:

$$\begin{aligned}
p_0(x) &= x^{124} + x^{115} + x^{114} + x^{113} + x^{112} + x^{110} + x^{106} + x^{105} + x^{104} + x^{103} \\
& + x^{102} + x^{100} + x^{98} + x^{97} + x^{96} + x^{94} + x^{93} + x^{89} + x^{88} + x^{84} + x^{82} \\
& + x^{79} + x^{78} + x^{77} + x^{75} + x^{70} + x^{68} + x^{67} + x^{64} + x^{63} + x^{60} + x^{54} \\
& + x^{53} + x^{52} + x^{45} + x^{44} + x^{43} + x^{40} + x^{39}, \\
p_3(x) &= x^{94} + x^{92} + x^{91} + x^{89} + x^{88} + x^{86} + x^{85} + x^{83} + x^{78} + x^{76} + x^{75} \\
& + x^{73} + x^{72} + x^{69} + x^{68} + x^{65} + x^{64} + x^{62} + x^{61} + x^{59} + x^{38} + x^{36}
\end{aligned}$$

$$\begin{aligned}
& + x^{35} + x^{33} + x^{32} + x^{30} + x^{29} + x^{27}, \\
p_6(x) &= x^{100} + x^{94} + x^{88} + x^{84} + x^{82} + x^{80} + x^{78} + x^{76} + x^{74} + x^{72} + x^{70} \\
& + x^{66} + x^{56} + x^{50} + x^{44} + x^{38} + x^{32} + x^{26} + x^{24} + x^{18}, \\
p_9(x) &= x^{106} + x^{104} + x^{103} + x^{102} + x^{101} + x^{100} + x^{99} + x^{98} + x^{97} + x^{96} \\
& + x^{95} + x^{94} + x^{93} + x^{92} + x^{91} + x^{89} + x^{58} + x^{56} + x^{55} + x^{54} + x^{53} \\
& + x^{52} + x^{51} + x^{50} + x^{49} + x^{48} + x^{47} + x^{46} + x^{45} + x^{44} + x^{43} + x^{41} \\
& + x^{26} + x^{24} + x^{23} + x^{22} + x^{21} + x^{20} + x^{19} + x^{18} + x^{17} + x^{16} + x^{15} \\
& + x^{14} + x^{13} + x^{12} + x^{11} + x^9, \\
p_{12}(x) &= x^{112} + x^{108} + x^{104} + x^{96} + x^{84} + x^{80} + x^{64} + x^{60} + x^{56} + x^{48} + x^{36} \\
& + x^{28} + x^{24} + x^{16} + x^4 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 1, 1, 0, 0, 1, 1, 1, 1, 1, 0, 0]$.

For $\phi(M^o, 1, 1)$:

$$\begin{aligned}
p_0(x) &= x^{117} + x^{115} + x^{114} + x^{113} + x^{112} + x^{110} + x^{107} + x^{105} + x^{102} + x^{101} \\
& + x^{100} + x^{98} + x^{94} + x^{92} + x^{90} + x^{88} + x^{85} + x^{84} + x^{82} + x^{80} + x^{79} \\
& + x^{76} + x^{73} + x^{72} + x^{67} + x^{65} + x^{64} + x^{59} + x^{58} + x^{56} + x^{53} + x^{52} \\
& + x^{49} + x^{46} + x^{45} + x^{44} + x^{42}, \\
p_3(x) &= x^{101} + x^{99} + x^{98} + x^{97} + x^{96} + x^{94} + x^{91} + x^{89} + x^{86} + x^{84} + x^{83} \\
& + x^{81} + x^{80} + x^{76} + x^{75} + x^{74} + x^{72} + x^{69} + x^{68} + x^{67} + x^{66} + x^{65} \\
& + x^{62} + x^{61} + x^{60} + x^{58} + x^{54} + x^{53} + x^{52} + x^{51} + x^{45} + x^{43} + x^{42} \\
& + x^{41} + x^{40} + x^{37} + x^{36} + x^{33} + x^{30} + x^{29} + x^{28} + x^{26}, \\
p_6(x) &= x^{109} + x^{107} + x^{106} + x^{104} + x^{103} + x^{100} + x^{99} + x^{97} + x^{94} + x^{92} \\
& + x^{86} + x^{85} + x^{80} + x^{77} + x^{75} + x^{73} + x^{68} + x^{66} + x^{65} + x^{62} + x^{59} \\
& + x^{58} + x^{57} + x^{54} + x^{52} + x^{51} + x^{50} + x^{48} + x^{43} + x^{41} + x^{37} + x^{35} \\
& + x^{31} + x^{30} + x^{28} + x^{26} + x^{25} + x^{23} + x^{21} + x^{20} + x^{18} + x^{16} + x^{15} \\
& + x^{13} + x^{12} + x^{11} + x^{10} + x^8 + x^5 + x^3 + x^2 + 1, \\
p_9(x) &= x^{97} + x^{95} + x^{94} + x^{92} + x^{49} + x^{47} + x^{46} + x^{44} + x^{17} + x^{15} + x^{14} \\
& + x^{12}, \\
p_{12}(x) &= x^{105} + x^{103} + x^{102} + x^{100} + x^{93} + x^{91} + x^{90} + x^{88} + x^{85} + x^{83} + x^{82} \\
& + x^{80} + x^{57} + x^{55} + x^{54} + x^{52} + x^{45} + x^{43} + x^{42} + x^{40} + x^{37} + x^{35} \\
& + x^{34} + x^{32} + x^{25} + x^{23} + x^{22} + x^{20} + x^{13} + x^{11} + x^{10} + x^8 + x^5 + x^3 \\
& + x^2 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 0, 1, 0, 1, 1, 1, 1, 1, 0, 1]$.

For $\phi(W^o, 0, 0)$:

$$\begin{aligned}
p_0(x) &= x^{117} + x^{115} + x^{114} + x^{113} + x^{112} + x^{110} + x^{107} + x^{103} + x^{99} + x^{96} \\
& + x^{95} + x^{94} + x^{93} + x^{91} + x^{89} + x^{87} + x^{86} + x^{76} + x^{73} + x^{72} + x^{70}
\end{aligned}$$

$$\begin{aligned}
& + x^{69} + x^{64} + x^{62} + x^{61} + x^{58} + x^{57} + x^{52} + x^{49} + x^{47} + x^{43} + x^{39} \\
& + x^{36}, \\
p_3(x) &= x^{101} + x^{99} + x^{98} + x^{97} + x^{96} + x^{94} + x^{91} + x^{87} + x^{85} + x^{82} + x^{81} \\
& + x^{79} + x^{75} + x^{74} + x^{73} + x^{72} + x^{70} + x^{68} + x^{64} + x^{63} + x^{59} + x^{56} \\
& + x^{55} + x^{54} + x^{53} + x^{51} + x^{47} + x^{45} + x^{44} + x^{43} + x^{42} + x^{41} + x^{40} \\
& + x^{39} + x^{37} + x^{28} + x^{27} + x^{24}, \\
p_6(x) &= x^{109} + x^{107} + x^{106} + x^{104} + x^{103} + x^{100} + x^{99} + x^{95} + x^{93} + x^{91} \\
& + x^{90} + x^{88} + x^{87} + x^{86} + x^{84} + x^{83} + x^{82} + x^{79} + x^{77} + x^{76} + x^{73} \\
& + x^{72} + x^{68} + x^{66} + x^{58} + x^{57} + x^{56} + x^{55} + x^{54} + x^{53} + x^{51} + x^{49} \\
& + x^{48} + x^{46} + x^{43} + x^{38} + x^{34} + x^{32} + x^{31} + x^{30} + x^{29} + x^{28} + x^{27} \\
& + x^{24} + x^{22} + x^{17} + x^{16} + x^{14} + x^{13} + x^{11} + x^{10} + x^8 + x^5 + x^3 + x^2 \\
& + 1, \\
p_9(x) &= x^{97} + x^{95} + x^{94} + x^{92} + x^{49} + x^{47} + x^{46} + x^{44} + x^{17} + x^{15} + x^{14} \\
& + x^{12}, \\
p_{12}(x) &= x^{105} + x^{103} + x^{102} + x^{100} + x^{93} + x^{91} + x^{90} + x^{88} + x^{85} + x^{83} + x^{82} \\
& + x^{80} + x^{57} + x^{55} + x^{54} + x^{52} + x^{45} + x^{43} + x^{42} + x^{40} + x^{37} + x^{35} \\
& + x^{34} + x^{32} + x^{25} + x^{23} + x^{22} + x^{20} + x^{13} + x^{11} + x^{10} + x^8 + x^5 + x^3 \\
& + x^2 + 1,
\end{aligned}$$

and the initial terms are $[1, 0, 0, 0, 0, 0, 1, 1, 0, 1, 0, 1, 0, 1, 1, 1]$.

For $\phi(W^o, 0, 1)$:

$$\begin{aligned}
p_0(x) &= x^{124} + x^{115} + x^{114} + x^{113} + x^{112} + x^{110} + x^{106} + x^{105} + x^{104} + x^{103} \\
& + x^{102} + x^{100} + x^{98} + x^{97} + x^{96} + x^{94} + x^{93} + x^{89} + x^{88} + x^{84} + x^{82} \\
& + x^{79} + x^{78} + x^{77} + x^{75} + x^{70} + x^{68} + x^{67} + x^{64} + x^{63} + x^{60} + x^{54} \\
& + x^{53} + x^{52} + x^{45} + x^{44} + x^{43} + x^{40} + x^{39}, \\
p_3(x) &= x^{94} + x^{92} + x^{91} + x^{89} + x^{88} + x^{86} + x^{85} + x^{83} + x^{78} + x^{76} + x^{75} \\
& + x^{73} + x^{72} + x^{69} + x^{68} + x^{65} + x^{64} + x^{62} + x^{61} + x^{59} + x^{38} + x^{36} \\
& + x^{35} + x^{33} + x^{32} + x^{30} + x^{29} + x^{27}, \\
p_6(x) &= x^{100} + x^{94} + x^{88} + x^{84} + x^{82} + x^{80} + x^{78} + x^{76} + x^{74} + x^{72} + x^{70} \\
& + x^{66} + x^{56} + x^{50} + x^{44} + x^{38} + x^{32} + x^{26} + x^{24} + x^{18}, \\
p_9(x) &= x^{106} + x^{104} + x^{103} + x^{102} + x^{101} + x^{100} + x^{99} + x^{98} + x^{97} + x^{96} \\
& + x^{95} + x^{94} + x^{93} + x^{92} + x^{91} + x^{89} + x^{58} + x^{56} + x^{55} + x^{54} + x^{53} \\
& + x^{52} + x^{51} + x^{50} + x^{49} + x^{48} + x^{47} + x^{46} + x^{45} + x^{44} + x^{43} + x^{41} \\
& + x^{26} + x^{24} + x^{23} + x^{22} + x^{21} + x^{20} + x^{19} + x^{18} + x^{17} + x^{16} + x^{15} \\
& + x^{14} + x^{13} + x^{12} + x^{11} + x^9, \\
p_{12}(x) &= x^{112} + x^{108} + x^{104} + x^{96} + x^{84} + x^{80} + x^{64} + x^{60} + x^{56} + x^{48} + x^{36}
\end{aligned}$$

$$+ x^{28} + x^{24} + x^{16} + x^4 + 1,$$

and the initial terms are $[0, 0, 0, 0, 1, 1, 0, 0, 1, 1, 1, 1, 1, 0, 0]$.

For $\phi(W^o, 1, 0)$:

$$\begin{aligned} p_0(x) = & x^{120} + x^{111} + x^{110} + x^{109} + x^{106} + x^{102} + x^{101} + x^{98} + x^{97} + x^{96} \\ & + x^{95} + x^{94} + x^{90} + x^{87} + x^{86} + x^{84} + x^{82} + x^{78} + x^{75} + x^{74} + x^{71} \\ & + x^{68} + x^{67} + x^{66} + x^{64} + x^{63} + x^{61} + x^{60} + x^{59} + x^{58} + x^{54} + x^{49} \\ & + x^{48} + x^{46} + x^{45} + x^{43} + x^{40} + x^{39}, \end{aligned}$$

$$\begin{aligned} p_3(x) = & x^{90} + x^{88} + x^{87} + x^{85} + x^{84} + x^{82} + x^{81} + x^{79} + x^{78} + x^{76} + x^{75} \\ & + x^{73} + x^{72} + x^{70} + x^{69} + x^{67} + x^{66} + x^{64} + x^{63} + x^{61} + x^{60} + x^{58} \\ & + x^{57} + x^{55} + x^{54} + x^{52} + x^{51} + x^{50} + x^{49} + x^{47} + x^{46} + x^{44} + x^{43} \\ & + x^{42} + x^{41} + x^{39} + x^{38} + x^{36} + x^{35} + x^{33} + x^{32} + x^{30} + x^{29} + x^{27}, \end{aligned}$$

$$\begin{aligned} p_6(x) = & x^{96} + x^{90} + x^{76} + x^{70} + x^{64} + x^{58} + x^{56} + x^{52} + x^{50} + x^{48} + x^{46} \\ & + x^{42} + x^{40} + x^{36} + x^{34} + x^{32} + x^{30} + x^{26} + x^{24} + x^{18}, \end{aligned}$$

$$\begin{aligned} p_9(x) = & x^{102} + x^{100} + x^{99} + x^{98} + x^{97} + x^{96} + x^{95} + x^{94} + x^{93} + x^{92} + x^{91} \\ & + x^{89} + x^{54} + x^{52} + x^{51} + x^{50} + x^{49} + x^{48} + x^{47} + x^{46} + x^{45} + x^{44} \\ & + x^{43} + x^{41} + x^{22} + x^{20} + x^{19} + x^{18} + x^{17} + x^{16} + x^{15} + x^{14} + x^{13} \\ & + x^{12} + x^{11} + x^9, \end{aligned}$$

$$\begin{aligned} p_{12}(x) = & x^{108} + x^{104} + x^{100} + x^{96} + x^{92} + x^{84} + x^{80} + x^{60} + x^{56} + x^{52} + x^{48} \\ & + x^{44} + x^{36} + x^{32} + x^{28} + x^{24} + x^{20} + x^{16} + x^{12} + x^4 + 1, \end{aligned}$$

and the initial terms are $[0, 0, 0, 1, 1, 0, 1, 1, 0, 0, 0, 0, 1, 1, 1]$.

For $\phi(W^o, 1, 1)$:

$$\begin{aligned} p_0(x) = & x^{117} + x^{115} + x^{114} + x^{113} + x^{112} + x^{110} + x^{107} + x^{105} + x^{102} + x^{101} \\ & + x^{100} + x^{98} + x^{94} + x^{92} + x^{90} + x^{88} + x^{85} + x^{84} + x^{82} + x^{80} + x^{79} \\ & + x^{76} + x^{73} + x^{72} + x^{67} + x^{65} + x^{64} + x^{59} + x^{58} + x^{56} + x^{53} + x^{52} \\ & + x^{49} + x^{46} + x^{45} + x^{44} + x^{42}, \end{aligned}$$

$$\begin{aligned} p_3(x) = & x^{101} + x^{99} + x^{98} + x^{97} + x^{96} + x^{94} + x^{91} + x^{89} + x^{86} + x^{84} + x^{83} \\ & + x^{81} + x^{80} + x^{76} + x^{75} + x^{74} + x^{72} + x^{69} + x^{68} + x^{67} + x^{66} + x^{65} \\ & + x^{62} + x^{61} + x^{60} + x^{58} + x^{54} + x^{53} + x^{52} + x^{51} + x^{45} + x^{43} + x^{42} \\ & + x^{41} + x^{40} + x^{37} + x^{36} + x^{33} + x^{30} + x^{29} + x^{28} + x^{26}, \end{aligned}$$

$$\begin{aligned} p_6(x) = & x^{109} + x^{107} + x^{106} + x^{104} + x^{103} + x^{100} + x^{99} + x^{97} + x^{94} + x^{92} \\ & + x^{86} + x^{85} + x^{80} + x^{77} + x^{75} + x^{73} + x^{68} + x^{66} + x^{65} + x^{62} + x^{59} \\ & + x^{58} + x^{57} + x^{54} + x^{52} + x^{51} + x^{50} + x^{48} + x^{43} + x^{41} + x^{37} + x^{35} \\ & + x^{31} + x^{30} + x^{28} + x^{26} + x^{25} + x^{23} + x^{21} + x^{20} + x^{18} + x^{16} + x^{15} \\ & + x^{13} + x^{12} + x^{11} + x^{10} + x^8 + x^5 + x^3 + x^2 + 1, \end{aligned}$$

$$p_9(x) = x^{97} + x^{95} + x^{94} + x^{92} + x^{49} + x^{47} + x^{46} + x^{44} + x^{17} + x^{15} + x^{14}$$

$$\begin{aligned}
& + x^{12}, \\
p_{12}(x) = & x^{105} + x^{103} + x^{102} + x^{100} + x^{93} + x^{91} + x^{90} + x^{88} + x^{85} + x^{83} + x^{82} \\
& + x^{80} + x^{57} + x^{55} + x^{54} + x^{52} + x^{45} + x^{43} + x^{42} + x^{40} + x^{37} + x^{35} \\
& + x^{34} + x^{32} + x^{25} + x^{23} + x^{22} + x^{20} + x^{13} + x^{11} + x^{10} + x^8 + x^5 + x^3 \\
& + x^2 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 0, 0, 1, 0, 1, 1, 1, 1, 1, 0, 1]$.

Below is the transition function and output function of an 2-automaton that generates $T = M_{0,0}^o$:

Transition function $(n, j) \mapsto \delta(n, j)$ ($\Lambda(n) := [\delta(n, 0), \delta(n, 1)]$):

n	$\Lambda(n)$	n	$\Lambda(n)$	n	$\Lambda(n)$	n	$\Lambda(n)$	n	$\Lambda(n)$
0	[1, 2]	117	[195, 196]	234	[341, 174]	351	[439, 418]	468	[383, 327]
1	[3, 4]	118	[197, 91]	235	[35, 154]	352	[301, 440]	469	[156, 13]
2	[5, 6]	119	[194, 198]	236	[338, 316]	353	[271, 416]	470	[358, 380]
3	[7, 8]	120	[86, 199]	237	[10, 342]	354	[441, 442]	471	[512, 356]
4	[9, 10]	121	[200, 201]	238	[343, 126]	355	[269, 404]	472	[349, 363]
5	[11, 12]	122	[202, 203]	239	[344, 143]	356	[249, 224]	473	[513, 50]
6	[13, 14]	123	[204, 205]	240	[343, 340]	357	[443, 444]	474	[142, 388]
7	[15, 16]	124	[206, 207]	241	[345, 346]	358	[262, 445]	475	[512, 364]
8	[17, 18]	125	[208, 209]	242	[347, 334]	359	[446, 29]	476	[514, 178]
9	[19, 20]	126	[210, 211]	243	[348, 347]	360	[447, 448]	477	[238, 515]
10	[21, 22]	127	[212, 213]	244	[349, 350]	361	[438, 24]	478	[516, 303]
11	[23, 24]	128	[214, 215]	245	[144, 351]	362	[449, 450]	479	[517, 304]
12	[25, 26]	129	[216, 217]	246	[315, 352]	363	[199, 99]	480	[518, 98]
13	[27, 17]	130	[218, 219]	247	[353, 129]	364	[451, 452]	481	[519, 229]
14	[28, 29]	131	[220, 221]	248	[125, 340]	365	[284, 453]	482	[520, 197]
15	[30, 31]	132	[222, 223]	249	[354, 318]	366	[454, 455]	483	[440, 278]
16	[32, 33]	133	[224, 225]	250	[319, 124]	367	[456, 423]	484	[521, 424]
17	[34, 35]	134	[226, 25]	251	[119, 321]	368	[457, 95]	485	[245, 97]
18	[36, 37]	135	[227, 75]	252	[355, 119]	369	[205, 200]	486	[80, 522]
19	[38, 39]	136	[111, 228]	253	[150, 356]	370	[458, 192]	487	[523, 524]
20	[40, 41]	137	[229, 230]	254	[357, 127]	371	[255, 210]	488	[525, 215]
21	[42, 43]	138	[228, 231]	255	[47, 358]	372	[273, 189]	489	[526, 242]
22	[44, 45]	139	[232, 233]	256	[126, 326]	373	[459, 229]	490	[527, 64]
23	[46, 47]	140	[234, 207]	257	[359, 139]	374	[460, 461]	491	[528, 529]
24	[48, 49]	141	[109, 235]	258	[360, 361]	375	[462, 295]	492	[530, 274]
25	[32, 50]	142	[236, 237]	259	[362, 363]	376	[463, 464]	493	[398, 531]
26	[51, 52]	143	[102, 238]	260	[151, 149]	377	[425, 465]	494	[104, 532]
27	[53, 54]	144	[239, 240]	261	[152, 309]	378	[466, 202]	495	[533, 534]
28	[55, 56]	145	[241, 242]	262	[150, 364]	379	[467, 468]	496	[535, 452]
29	[57, 58]	146	[243, 241]	263	[113, 12]	380	[469, 299]	497	[448, 6]
30	[59, 60]	147	[244, 245]	264	[365, 175]	381	[400, 470]	498	[464, 427]
31	[22, 61]	148	[246, 247]	265	[366, 115]	382	[471, 472]	499	[536, 270]

32	[16, 62]	149	[248, 249]	266	[367, 138]	383	[410, 466]	500	[537, 469]
33	[63, 64]	150	[250, 251]	267	[368, 41]	384	[473, 278]	501	[538, 539]
34	[65, 66]	151	[252, 250]	268	[329, 369]	385	[474, 220]	502	[540, 239]
35	[67, 68]	152	[84, 253]	269	[158, 35]	386	[409, 74]	503	[541, 542]
36	[69, 70]	153	[225, 254]	270	[163, 333]	387	[475, 476]	504	[543, 544]
37	[71, 72]	154	[90, 255]	271	[163, 347]	388	[299, 87]	505	[476, 545]
38	[73, 74]	155	[68, 256]	272	[370, 159]	389	[355, 45]	506	[546, 531]
39	[75, 76]	156	[257, 258]	273	[53, 371]	390	[477, 312]	507	[547, 224]
40	[77, 78]	157	[259, 260]	274	[372, 373]	391	[12, 182]	508	[548, 293]
41	[79, 80]	158	[261, 262]	275	[317, 372]	392	[478, 328]	509	[549, 550]
42	[81, 82]	159	[88, 263]	276	[374, 167]	393	[479, 129]	510	[551, 552]
43	[83, 84]	160	[264, 265]	277	[127, 342]	394	[333, 334]	511	[553, 542]
44	[85, 86]	161	[266, 267]	278	[175, 32]	395	[129, 347]	512	[554, 290]
45	[87, 88]	162	[268, 269]	279	[325, 340]	396	[480, 380]	513	[555, 263]
46	[89, 90]	163	[270, 271]	280	[52, 182]	397	[481, 187]	514	[556, 557]
47	[91, 92]	164	[272, 273]	281	[375, 334]	398	[482, 361]	515	[176, 33]
48	[93, 94]	165	[274, 275]	282	[376, 377]	399	[362, 350]	516	[313, 332]
49	[95, 96]	166	[276, 277]	283	[166, 378]	400	[180, 45]	517	[499, 334]
50	[97, 98]	167	[278, 15]	284	[379, 51]	401	[122, 150]	518	[558, 124]
51	[99, 100]	168	[279, 280]	285	[154, 380]	402	[381, 47]	519	[559, 138]
52	[101, 102]	169	[281, 282]	286	[381, 35]	403	[358, 155]	520	[507, 486]
53	[103, 104]	170	[283, 284]	287	[171, 364]	404	[319, 483]	521	[560, 50]
54	[98, 105]	171	[285, 83]	288	[375, 353]	405	[35, 358]	522	[122, 171]
55	[106, 107]	172	[286, 287]	289	[382, 346]	406	[38, 57]	523	[561, 54]
56	[108, 109]	173	[288, 289]	290	[309, 123]	407	[8, 32]	524	[186, 562]
57	[110, 111]	174	[190, 290]	291	[44, 119]	408	[481, 137]	525	[345, 563]
58	[74, 110]	175	[291, 63]	292	[340, 383]	409	[484, 485]	526	[331, 314]
59	[112, 113]	176	[280, 292]	293	[47, 154]	410	[114, 486]	527	[490, 380]
60	[114, 115]	177	[219, 108]	294	[34, 47]	411	[126, 383]	528	[116, 564]
61	[48, 116]	178	[293, 294]	295	[314, 314]	412	[153, 123]	529	[478, 170]
62	[31, 114]	179	[196, 295]	296	[121, 149]	413	[487, 350]	530	[385, 310]
63	[117, 118]	180	[296, 297]	297	[321, 323]	414	[328, 488]	531	[187, 137]
64	[119, 120]	181	[298, 299]	298	[384, 327]	415	[489, 347]	532	[495, 49]
65	[121, 122]	182	[24, 293]	299	[312, 366]	416	[333, 353]	533	[348, 333]
66	[123, 124]	183	[300, 301]	300	[385, 184]	417	[347, 353]	534	[565, 377]
67	[125, 126]	184	[263, 302]	301	[162, 386]	418	[334, 129]	535	[341, 182]
68	[9, 127]	185	[303, 304]	302	[379, 113]	419	[353, 163]	536	[479, 163]
69	[128, 129]	186	[305, 306]	303	[387, 165]	420	[490, 155]	537	[157, 14]
70	[130, 131]	187	[230, 307]	304	[332, 314]	421	[147, 371]	538	[560, 33]
71	[132, 133]	188	[168, 175]	305	[344, 388]	422	[482, 491]	539	[510, 316]
72	[134, 51]	189	[308, 309]	306	[30, 312]	423	[58, 136]	540	[181, 174]
73	[135, 136]	190	[146, 37]	307	[57, 138]	424	[492, 167]	541	[503, 39]
74	[137, 137]	191	[113, 52]	308	[389, 204]	425	[372, 493]	542	[139, 57]
75	[138, 136]	192	[45, 120]	309	[72, 206]	426	[484, 494]	543	[374, 378]

76	[139, 39]	193	[183, 310]	310	[292, 67]	427	[172, 149]	544	[13, 339]
77	[140, 141]	194	[311, 312]	311	[390, 391]	428	[495, 116]	545	[49, 564]
78	[142, 143]	195	[313, 314]	312	[294, 392]	429	[496, 486]	546	[506, 137]
79	[144, 145]	196	[315, 131]	313	[393, 394]	430	[365, 8]	547	[513, 33]
80	[146, 144]	197	[132, 316]	314	[394, 395]	431	[176, 50]	548	[508, 364]
81	[147, 54]	198	[317, 318]	315	[396, 259]	432	[497, 388]	549	[118, 562]
82	[116, 148]	199	[153, 319]	316	[191, 188]	433	[318, 373]	550	[173, 186]
83	[149, 150]	200	[320, 153]	317	[397, 398]	434	[370, 323]	551	[497, 143]
84	[151, 122]	201	[321, 159]	318	[399, 400]	435	[161, 498]	552	[498, 386]
85	[152, 153]	202	[160, 175]	319	[253, 401]	436	[499, 353]	553	[55, 566]
86	[154, 155]	203	[322, 157]	320	[402, 403]	437	[177, 43]	554	[567, 159]
87	[134, 113]	204	[120, 323]	321	[404, 405]	438	[170, 488]	555	[384, 141]
88	[156, 157]	205	[45, 321]	322	[406, 28]	439	[387, 500]	556	[487, 363]
89	[158, 47]	206	[324, 162]	323	[4, 407]	440	[354, 372]	557	[186, 179]
90	[120, 159]	207	[51, 12]	324	[408, 409]	441	[367, 58]	558	[568, 412]
91	[160, 8]	208	[325, 126]	325	[306, 410]	442	[501, 56]	559	[569, 232]
92	[161, 162]	209	[326, 327]	326	[209, 411]	443	[359, 136]	560	[570, 407]
93	[128, 163]	210	[308, 153]	327	[211, 412]	444	[502, 485]	561	[571, 572]
94	[164, 165]	211	[169, 328]	328	[107, 248]	445	[309, 319]	562	[573, 270]
95	[166, 167]	212	[329, 330]	329	[413, 414]	446	[503, 57]	563	[574, 4]
96	[168, 8]	213	[320, 309]	330	[412, 291]	447	[140, 327]	564	[534, 394]
97	[169, 170]	214	[331, 332]	331	[415, 247]	448	[504, 184]	565	[575, 576]
98	[149, 171]	215	[129, 333]	332	[332, 332]	449	[505, 330]	566	[82, 577]
99	[172, 122]	216	[334, 163]	333	[416, 417]	450	[37, 145]	567	[578, 522]
100	[173, 118]	217	[314, 332]	334	[418, 419]	451	[324, 498]	568	[558, 483]
101	[52, 174]	218	[335, 124]	335	[420, 235]	452	[31, 366]	569	[559, 58]
102	[175, 176]	219	[336, 330]	336	[421, 79]	453	[383, 141]	570	[496, 115]
103	[177, 178]	220	[10, 337]	337	[422, 423]	454	[326, 141]	571	[505, 369]
104	[118, 179]	221	[322, 13]	338	[424, 425]	455	[8, 176]	572	[49, 148]
105	[180, 119]	222	[338, 133]	339	[426, 75]	456	[506, 187]	573	[565, 579]
106	[181, 182]	223	[157, 339]	340	[427, 428]	457	[507, 115]	574	[162, 511]
107	[183, 184]	224	[340, 326]	341	[429, 256]	458	[508, 356]	575	[335, 483]
108	[170, 185]	225	[311, 31]	342	[20, 232]	459	[40, 509]	576	[561, 371]
109	[117, 186]	226	[112, 51]	343	[430, 431]	460	[510, 133]	577	[36, 144]
110	[136, 39]	227	[135, 139]	344	[432, 433]	461	[498, 511]	578	[567, 323]
111	[137, 187]	228	[39, 138]	345	[434, 212]	462	[489, 333]	579	[223, 451]
112	[188, 101]	229	[136, 57]	346	[237, 435]	463	[480, 155]		
113	[189, 190]	230	[138, 139]	347	[304, 216]	464	[42, 178]		
114	[60, 191]	231	[187, 187]	348	[436, 418]	465	[357, 10]		
115	[61, 192]	232	[39, 58]	349	[437, 438]	466	[312, 114]		
116	[193, 194]	233	[58, 139]	350	[192, 89]	467	[477, 31]		

Output function $n \mapsto \tau(n)$:

n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$
0	0	83	0	166	1	249	1	332	0	415	1	498	0
1	0	84	0	167	0	250	1	333	1	416	1	499	0
2	0	85	0	168	0	251	1	334	1	417	0	500	1
3	0	86	0	169	1	252	0	335	1	418	1	501	1
4	0	87	0	170	1	253	1	336	0	419	0	502	0
5	0	88	1	171	0	254	1	337	0	420	1	503	1
6	0	89	0	172	1	255	1	338	0	421	0	504	1
7	0	90	0	173	1	256	0	339	1	422	0	505	1
8	1	91	1	174	0	257	1	340	1	423	0	506	1
9	0	92	1	175	0	258	1	341	1	424	0	507	0
10	0	93	1	176	0	259	1	342	1	425	0	508	0
11	0	94	0	177	0	260	0	343	1	426	1	509	0
12	1	95	1	178	1	261	0	344	1	427	1	510	1
13	0	96	0	179	0	262	1	345	0	428	0	511	0
14	0	97	1	180	1	263	0	346	0	429	1	512	1
15	0	98	0	181	0	264	1	347	0	430	1	513	0
16	1	99	1	182	1	265	1	348	0	431	0	514	1
17	1	100	1	183	0	266	1	349	0	432	1	515	0
18	1	101	0	184	0	267	0	350	0	433	1	516	0
19	0	102	0	185	1	268	1	351	1	434	0	517	0
20	1	103	0	186	1	269	0	352	1	435	1	518	0
21	0	104	0	187	1	270	0	353	0	436	0	519	0
22	0	105	1	188	0	271	0	354	1	437	0	520	0
23	0	106	0	189	0	272	0	355	0	438	1	521	1
24	1	107	0	190	0	273	0	356	1	439	1	522	1
25	1	108	1	191	0	274	0	357	1	440	1	523	1
26	1	109	0	192	0	275	0	358	1	441	1	524	1
27	0	110	0	193	0	276	1	359	1	442	1	525	0
28	0	111	0	194	1	277	1	360	1	443	1	526	1
29	0	112	0	195	0	278	0	361	1	444	0	527	1
30	0	113	0	196	0	279	0	362	1	445	0	528	0
31	0	114	0	197	0	280	0	363	1	446	1	529	0
32	1	115	1	198	0	281	1	364	0	447	1	530	0
33	0	116	0	199	1	282	0	365	1	448	1	531	1
34	1	117	0	200	1	283	1	366	1	449	1	532	0
35	0	118	0	201	1	284	1	367	1	450	0	533	0
36	1	119	1	202	1	285	0	368	0	451	0	534	1
37	0	120	0	203	0	286	1	369	0	452	0	535	1
38	0	121	1	204	0	287	0	370	0	453	0	536	0
39	1	122	1	205	0	288	1	371	1	454	1	537	1
40	1	123	0	206	0	289	1	372	0	455	1	538	1
41	1	124	0	207	1	290	0	373	1	456	1	539	1
42	0	125	0	208	0	291	0	374	1	457	0	540	0

43	0	126	0	209	1	292	1	375	1	458	0	541	1
44	0	127	1	210	0	293	1	376	0	459	1	542	1
45	0	128	1	211	1	294	1	377	0	460	1	543	1
46	0	129	1	212	1	295	1	378	1	461	0	544	0
47	1	130	1	213	1	296	1	379	1	462	1	545	1
48	1	131	0	214	1	297	1	380	1	463	0	546	1
49	1	132	0	215	1	298	0	381	1	464	0	547	0
50	1	133	1	216	1	299	1	382	1	465	1	548	0
51	1	134	0	217	1	300	0	383	0	466	1	549	0
52	0	135	0	218	1	301	1	384	0	467	1	550	1
53	0	136	0	219	0	302	1	385	0	468	0	551	1
54	0	137	0	220	0	303	1	386	1	469	1	552	0
55	0	138	1	221	0	304	0	387	1	470	1	553	0
56	1	139	1	222	0	305	1	388	1	471	1	554	1
57	0	140	1	223	1	306	0	389	0	472	0	555	0
58	0	141	0	224	1	307	0	390	1	473	0	556	1
59	0	142	0	225	1	308	0	391	1	474	0	557	1
60	0	143	0	226	0	309	0	392	0	475	1	558	0
61	1	144	1	227	0	310	1	393	0	476	1	559	0
62	0	145	0	228	1	311	1	394	1	477	1	560	1
63	0	146	0	229	0	312	1	395	1	478	0	561	1
64	1	147	0	230	1	313	0	396	0	479	0	562	1
65	1	148	0	231	1	314	1	397	0	480	0	563	1
66	0	149	0	232	1	315	0	398	0	481	0	564	1
67	0	150	1	233	0	316	0	399	1	482	0	565	1
68	0	151	0	234	1	317	0	400	1	483	1	566	0
69	1	152	0	235	0	318	1	401	1	484	1	567	1
70	1	153	1	236	0	319	1	402	1	485	1	568	0
71	0	154	0	237	0	320	1	403	1	486	0	569	0
72	0	155	0	238	1	321	1	404	1	487	1	570	1
73	0	156	1	239	1	322	0	405	0	488	0	571	1
74	0	157	1	240	1	323	0	406	0	489	1	572	1
75	1	158	0	241	0	324	0	407	1	490	1	573	1
76	1	159	1	242	0	325	0	408	0	491	0	574	1
77	1	160	1	243	0	326	1	409	1	492	0	575	1
78	0	161	1	244	0	327	1	410	0	493	0	576	1
79	1	162	1	245	1	328	0	411	0	494	0	577	1
80	0	163	0	246	0	329	1	412	1	495	0	578	1
81	0	164	0	247	0	330	1	413	1	496	1	579	1
82	0	165	0	248	0	331	1	414	0	497	1		

UNIVERSITÉ DE STRASBOURG, CNRS, IRMA UMR 7501, F-67000 STRASBOURG, FRANCE
Email address: `guoniu.han@unistra.fr`

SCHOOL OF MATHEMATICS AND STATISTICS, HUAZHONG UNIVERSITY OF SCIENCE AND TECHNOLOGY, WUHAN, PR CHINA
Email address: `huyining@hust.edu.cn`