

PROOF OF THE CONTINUED FRACTION CONJECTURE FOR

$$(a, b) = (z^3 + z^2 + z, z^4 + z^2 + 1)$$

GUO-NIU HAN* AND YINING HU*

ABSTRACT. The continued fraction conjecture claims that the continued fraction defined by the (a, b) -Thue-Morse sequence is algebraic over $\mathbb{F}_2(x)$ for all pairs of distinct elements (a, b) in $\mathbb{F}_2[x] \setminus \mathbb{F}_2$. In this paper we prove the conjecture for $(a, b) = (z^3 + z^2 + z, z^4 + z^2 + 1)$.

1. INTRODUCTION

Let $\mathbb{F}_2$ be the finite field of cardinality 2. Given a sequence of polynomials $\mathbf{a} = (a_0(z), a_1(z), a_2(z), \dots)$ of elements from $\mathbb{F}_2[z] \setminus \mathbb{F}_2$, we define the infinite continued fraction

$$(1.1) \quad \text{CF}(\mathbf{a}(z)) = \frac{1}{a_0(z) + \frac{1}{a_1(z) + \frac{1}{a_2(z) + \frac{1}{\ddots}}}}$$

as the limit of the finite continued fractions

$$(1.2) \quad \text{CF}_n(\mathbf{a}(z)) = \frac{1}{a_0(z) + \frac{1}{a_1(z) + \frac{1}{\ddots + \frac{1}{a_n(z)}}}} \in \mathbb{F}_2((1/z)).$$

Conversely, each power series in $\mathbb{F}_2[[1/z]]$ without constant term

$$(1.3) \quad c_{-1}z^{-1} + c_{-2}z^{-2} + c_{-3}z^{-3} + \dots$$

admits a continued fraction expansion of form (1.1).

Let a and b be two distinct elements from $\mathbb{F}_2[z] \setminus \mathbb{F}_2$. We consider the continued fraction (1.1) associated with the (a, b) -Thue-Morse sequence

$$\mathbf{t} = (t_0(z), t_1(z), t_2(z), \dots) = (a, b, b, a, b, a, a, b, \dots)$$

that we shall denote

$$(1.4) \quad \text{CF}(a, b) := \text{CF}(\mathbf{t}).$$

Date: 9/04/2020 23:46:31 .

2010 Mathematics Subject Classification. 11B85, 11J70, 11B50, 11Y65, 05A15.

Key words and phrases. automatic sequence, continued fraction, Thue-Morse sequence.

* This paper was automatically generated from a computer program developed by the authors, with 6.9 minutes CPU time used.

Theorem 1.1. *When $(a, b) = (z^3 + z^2 + z, z^4 + z^2 + 1)$, the two power series $\text{CF}(a, b)$ and $\text{CF}(b, a)$ are algebraic over $\mathbb{F}_2(z)$, with minimal polynomials of the form*

$$p_4(z)y^4 + p_3(z)y^3 + p_2(z)y^2 + p_1(z)y + p_0(z) = 0$$

where, for $\text{CF}(a, b)$,

$$\begin{aligned} p_0(z) &= z^{13} + z^{12} + z^{11} + z^9 + z^4 + z + 1, \\ p_1(z) &= z^{17} + z^{13} + z^9 + z^5 + z^3 + z, \\ p_2(z) &= z^{20} + z^{14} + z^{12} + z^4 + z^2, \\ p_3(z) &= z^{17} + z^{13} + z^9 + z^5 + z^3 + z, \\ p_4(z) &= z^{14} + z^{13} + z^{12} + z^{11} + z^9 + z^8 + z^4 + z^2 + z, \end{aligned}$$

and for $\text{CF}(b, a)$,

$$\begin{aligned} p_0(z) &= z^{13} + z^{12} + z^{11} + z^9 + z^8 + z^4 + z, \\ p_1(z) &= z^{17} + z^{13} + z^9 + z^5 + z^3 + z, \\ p_2(z) &= z^{20} + z^{14} + z^{12} + z^4 + z^2, \\ p_3(z) &= z^{17} + z^{13} + z^9 + z^5 + z^3 + z, \\ p_4(z) &= z^{14} + z^{13} + z^{12} + z^{11} + z^9 + z^4 + z^2 + z + 1. \end{aligned}$$

2. PROOF

Define the sequences of polynomials $(P_n(z))_{n \geq 0}$ and $(Q_n(z))_{n \geq 0}$ by

$$\begin{pmatrix} P_n(z) & Q_n(z) \\ P_{n-1}(z) & Q_{n-1}(z) \end{pmatrix} := \begin{pmatrix} t_n(z) & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} t_{n-1}(z) & 1 \\ 1 & 0 \end{pmatrix} \cdots \begin{pmatrix} t_0(z) & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}.$$

By the basic properties of continued fractions, we have

$$(2.1) \quad \text{CF}_n(a, b) = \frac{P_n(z)}{Q_n(z)} \in \mathbb{F}_2((1/z)).$$

As formula (2.1) is not efficient for calculating $\text{CF}_n(a, b)$, a fast method is to be derived and described as follows. Define

$$\begin{aligned} M_n(x) &= x^{\deg(t_{2^n-1})} \begin{pmatrix} t_{2^n-1}(1/x) & 1 \\ 1 & 0 \end{pmatrix} \times x^{\deg(t_{2^n-2})} \begin{pmatrix} t_{2^n-2}(1/x) & 1 \\ 1 & 0 \end{pmatrix} \times \\ &\quad \cdots \times x^{\deg(t_0)} \begin{pmatrix} t_0(1/x) & 1 \\ 1 & 0 \end{pmatrix}, \\ W_n(x) &= x^{\deg(\bar{t}_{2^n-1})} \begin{pmatrix} \bar{t}_{2^n-1}(1/x) & 1 \\ 1 & 0 \end{pmatrix} \times x^{\deg(\bar{t}_{2^n-2})} \begin{pmatrix} \bar{t}_{2^n-2}(1/x) & 1 \\ 1 & 0 \end{pmatrix} \times \\ &\quad \cdots \times x^{\deg(\bar{t}_0)} \begin{pmatrix} \bar{t}_0(1/x) & 1 \\ 1 & 0 \end{pmatrix}, \end{aligned}$$

where $\bar{t}$ is the (b, a) -Thue-Morse sequence. By the properties of the Thue-Morse sequence, we have for $n \geq 0$

$$\begin{aligned} M_{n+1}(x) &= W_n(x) \cdot M_n(x), \\ W_{n+1}(x) &= M_n(x) \cdot W_n(x). \end{aligned}$$

Let $x := 1/z$. For each non-zero polynomial $P(z)$, define $\tilde{P}(x)$ to be $P(1/x)$. Then,

$$\text{CF}_n(a, b) = \frac{P_n(z)}{Q_n(z)} = \frac{\tilde{P}_n(x)}{\tilde{Q}_n(x)} \in \mathbb{F}_2((x)) = \mathbb{F}_2((1/z)).$$

Comparing the definition of $M_n(x)$ with the definition of $P_n(x)$ and $Q_n(x)$, we see that

$$\begin{aligned} M_n(x)_{0,1} &= x^{d_n} \tilde{P}_{2^n-1}(x), \\ M_n(x)_{0,0} &= x^{d_n} \tilde{Q}_{2^n-1}(x), \end{aligned}$$

for some positive integer d_n . Hence,

$$(2.2) \quad \text{CF}_{2^{2^n}-1}(a, b) = \frac{\tilde{P}_{2^{2^n}-1}(x)}{\tilde{Q}_{2^{2^n}-1}(x)} = \frac{M_{2^n}(x)_{0,1}}{M_{2^n}(x)_{0,0}}.$$

Therefore, by the convergence theorem of continued fraction, the algebraicity of $\text{CF}(a, b)$ will be established if it is shown that both $M_{2^n}(x)_{0,1}$ and $M_{2^n}(x)_{0,0}$ converge to algebraic series in $\mathbb{F}_2[[x]]$.

Actually, we will prove that for all $0 \leq i, j \leq 1$ the four sequences $(M_{2^n}(x)_{i,j})_n$, $(M_{2^{n+1}}(x)_{i,j})_n$, $(W_{2^n}(x)_{i,j})_n$, and $(W_{2^{n+1}}(x)_{i,j})_n$ converge to algebraic series in $\mathbb{F}_2[[x]]$. For this purpose, we define four 2×2 matrices M^e, M^o, W^e, W^o as follows: For each $T \in \{M^e, M^o, W^e, W^o\}$ and $0 \leq i, j \leq 1$, $T_{i,j}$ is defined to be the unique solution in $\mathbb{F}_2[[x]]$ of the polynomial $\phi(T, i, j)$ under certain initial conditions; the polynomials $\phi(T, i, j)$ and initial conditions are given in Section 3. We will prove that these four matrices, whose components are algebraic by definition, are the limits of $(M_{2^n}(x))_n$, $(M_{2^{n+1}}(x))_n$, $(W_{2^n}(x))_n$, and $(W_{2^{n+1}}(x))_n$.

Let us explain how the polynomials $\phi(T, i, j)$ and initial conditions are found, and why the solutions exist and are unique. For $0 \leq i, j \leq 1$, the coefficients of the polynomial $\phi(M^e, i, j)$ (resp. $\phi(M^o, i, j)$, $\phi(W^e, i, j)$, and $\phi(W^o, i, j)$) are the Padé-Hermite approximants of type

$$(175, 175, 175, 175, 175)$$

of the vector

$$(1, T^3, T^6, T^9, T^{12}),$$

where $T = M_{12,i,j}$ (resp. $M_{11,i,j}$, $W_{12,i,j}$, and $W_{11,i,j}$), found by the Derksen algorithm. We take the first 8 terms of $M_{12,i,j}$ (resp. $M_{11,i,j}$, $W_{12,i,j}$, and $W_{11,i,j}$) as initial conditions for $\phi(M^e, i, j)$ (resp. $\phi(M^o, i, j)$, $\phi(W^e, i, j)$, and $\phi(W^o, i, j)$). The following fact will be used to ensure that the solution exists and is unique: let $P(x, y) \in \mathbb{F}_2[x, y]$ and for each series $f(x) = \sum_{n=0}^{\infty} a_n x^n \in \mathbb{F}_2[[x]]$ denote the partial sum $\sum_{j=0}^{n-1} a_j x^j$ by $f_n(x)$ for $n \geq 0$. If for some $n \geq 0$ and $a_0, a_1, \dots, a_{n-1} \in \mathbb{F}_2$ $P(x, \sum_{j=0}^{n-1} a_j x^j) = O(x^n)$ and $Q(x, y) := P(x, \sum_{j=0}^{n-1} a_j x^j + x^n y)$ can be written as $x^m \sum_{j=0}^{\infty} q_j(x) y^j$ where $q_j(x)$ are polynomials for $j \geq 0$, $q_1(0) = 1$, and $q_j(0) = 0$ for $j > 1$, then there exists a unique solution $f(x) \in \mathbb{F}_2[[x]]$ of $P(x, f(x)) = 0$ that satisfies the initial condition $f_n(x) = \sum_{j=0}^{n-1} a_j x^j$.

We state two lemmas concerning the four matrices M^e, M^o, W^e, W^o . The first one is about relations between them; the second, about the structure of the each matrix.

Lemma 2.1. *We have*

$$(2.3) \quad M^e = W^o \cdot M^o,$$

$$(2.4) \quad M^o = W^e \cdot M^e,$$

$$(2.5) \quad W^e = M^o \cdot W^o,$$

$$(2.6) \quad W^o = M^e \cdot W^e.$$

Proof. We give the proof of the identity

$$M_{0,0}^e = W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o,$$

the proofs of the others being similar.

First, we compute the minimal polynomials of $W_{0,0}^o M_{0,0}^o$ and $W_{0,1}^o M_{1,0}^o$. We know that

$$P(x, y) = \text{Res}_z (\phi(W^o, 0, 0)(x, z), z^{12} \cdot \phi(M^o, 0, 0)(x, y/z))$$

is an annihilating polynomial of $W_{0,0}^o M_{0,0}^o$. We use Padé-Hermite approximation to find a candidate for the minimal polynomial of $W_{0,0}^o M_{0,0}^o$, that will be called $\phi_0(x, y)$. To prove that $\phi_0(x, y)$ is indeed the minimal polynomial, it suffices to prove that it is an irreducible factor of $P(x, y)$ of multiplicity m and that $Q(x, y) := P(x, y)/\phi_0(x, y)^m$ is not an annihilating polynomial of $W_{0,0}^o M_{0,0}^o$. We verify the first point directly. For the second point, we truncate $W_{0,0}^o M_{0,0}^o$ to order 630 and substitute it for y in $Q(x, y)$. We get a series of valuation less than 630, which proves that $Q(x, y)$ is not an annihilating polynomial of $W_{0,0}^o M_{0,0}^o$. We find the minimal polynomial $\phi_1(x, y)$ of $W_{0,1}^o M_{1,0}^o$ in a similar way.

Now we prove that $\phi(M^e, 0, 0)$ is the minimal polynomial of $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$. We know that

$$S(x, y) = \text{Res}_z (\phi_0(x, z), \phi_1(x, y + z))$$

is an annihilating of $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$. We verify that $\phi(M^e, 0, 0)$ is an irreducible factor of $S(x, y)$ of multiplicity μ , and that $Q(x, y) := S(x, y)/\phi(M^e, 0, 0)^\mu$ is not an annihilating polynomial of $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$. To see the last point, we truncate $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$ to order 770 and substitute it for y in $Q(x, y)$. We get a series of valuation less than 770, and therefore $Q(x, y)$ is not an annihilating polynomial of $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$.

Finally, the first 16 terms of $M_{0,0}^e$ and $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$ coincide. As these first terms determine a unique solution of $\phi(M^e, 0, 0)$, we know that the two series are one and the same. $\square$

Define

$$R^e = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \quad \text{and} \quad R^o = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}.$$

Lemma 2.2. *For all integers $k \geq 2$ and $u = 2^{2k-1}$, the following identities hold.*

$$\begin{aligned} M^e[:14u] &= M^e[:7u] + x^u M^e[:6u] + x^{3u} M^e[:4u] + x^{5u} M^e[:2u] + x^u M^e[:7u] \\ &\quad + x^{2u} M^e[:6u] + x^{4u} M^e[:4u] + x^{6u} M^e[:2u] + x^{2u} M^e[:7u] \\ &\quad + x^{3u} M^e[:6u] + x^{5u} M^e[:4u] + x^{7u} M^e[:2u] + x^{4u} M^e[:7u] \\ &\quad + x^{5u} M^e[:6u] + x^{7u} M^e[:4u] + x^{9u} M^e[:2u] + x^{5u} M^e[:7u] \\ &\quad + x^{6u} M^e[:6u] + x^{8u} M^e[:4u] + x^{10u} M^e[:2u] + x^{7u} M^e[:7u] \\ &\quad + x^{9u} M^e[:5u] + x^{11u} M^e[:3u] + x^{13u} M^e[:u] \end{aligned}$$

$$\begin{aligned}
& + (x^{7u} + x^{8u} + x^{9u} + x^{11u} + x^{12u})R^e, \\
W^e[:14u] &= W^e[:7u] + x^u W^e[:6u] + x^{3u} W^e[:4u] + x^{5u} W^e[:2u] + x^u W^e[:7u] \\
& + x^{2u} W^e[:6u] + x^{4u} W^e[:4u] + x^{6u} W^e[:2u] + x^{2u} W^e[:7u] \\
& + x^{3u} W^e[:6u] + x^{5u} W^e[:4u] + x^{7u} W^e[:2u] + x^{4u} W^e[:7u] \\
& + x^{5u} W^e[:6u] + x^{7u} W^e[:4u] + x^{9u} W^e[:2u] + x^{5u} W^e[:7u] \\
& + x^{6u} W^e[:6u] + x^{8u} W^e[:4u] + x^{10u} W^e[:2u] + x^{7u} W^e[:7u] \\
& + x^{9u} W^e[:5u] + x^{11u} W^e[:3u] + x^{13u} W^e[:u] \\
& + (x^{7u} + x^{8u} + x^{9u} + x^{11u} + x^{12u})R^e.
\end{aligned}$$

For all integers $k \geq 2$ and $u = 2^{2k}$, the following identities hold.

$$\begin{aligned}
M^o[:14u] &= M^o[:7u] + x^u M^o[:6u] + x^{3u} M^o[:4u] + x^{5u} M^o[:2u] + x^u M^o[:7u] \\
& + x^{2u} M^o[:6u] + x^{4u} M^o[:4u] + x^{6u} M^o[:2u] + x^{2u} M^o[:7u] \\
& + x^{3u} M^o[:6u] + x^{5u} M^o[:4u] + x^{7u} M^o[:2u] + x^{4u} M^o[:7u] \\
& + x^{5u} M^o[:6u] + x^{7u} M^o[:4u] + x^{9u} M^o[:2u] + x^{5u} M^o[:7u] \\
& + x^{6u} M^o[:6u] + x^{8u} M^o[:4u] + x^{10u} M^o[:2u] + x^{7u} M^o[:7u] \\
& + x^{9u} M^o[:5u] + x^{11u} M^o[:3u] + x^{13u} M^o[:u] \\
& + (x^{7u} + x^{8u} + x^{9u} + x^{11u} + x^{12u})R^o, \\
W^o[:14u] &= W^o[:7u] + x^u W^o[:6u] + x^{3u} W^o[:4u] + x^{5u} W^o[:2u] + x^u W^o[:7u] \\
& + x^{2u} W^o[:6u] + x^{4u} W^o[:4u] + x^{6u} W^o[:2u] + x^{2u} W^o[:7u] \\
& + x^{3u} W^o[:6u] + x^{5u} W^o[:4u] + x^{7u} W^o[:2u] + x^{4u} W^o[:7u] \\
& + x^{5u} W^o[:6u] + x^{7u} W^o[:4u] + x^{9u} W^o[:2u] + x^{5u} W^o[:7u] \\
& + x^{6u} W^o[:6u] + x^{8u} W^o[:4u] + x^{10u} W^o[:2u] + x^{7u} W^o[:7u] \\
& + x^{9u} W^o[:5u] + x^{11u} W^o[:3u] + x^{13u} W^o[:u] \\
& + (x^{7u} + x^{8u} + x^{9u} + x^{11u} + x^{12u})R^o.
\end{aligned}$$

Proof. To prove Lemma 2.2 we first construct an automaton for each sequence concerned, and then transform the conditions on infinitely many k 's into finitely many conditions on the states of the automaton. In the following, we will prove that for $T = M_{0,0}^o$, for all integer $k \geq 2$ and $u = 2^{2k}$,

$$\begin{aligned}
T[:14u] &= T[:7u] + x^u T[:6u] + x^{3u} T[:4u] + x^{5u} T[:2u] + x^u T[:7u] + x^{2u} T[:6u] \\
& + x^{4u} T[:4u] + x^{6u} T[:2u] + x^{2u} T[:7u] + x^{3u} T[:6u] + x^{5u} T[:4u] \\
& + x^{7u} T[:2u] + x^{4u} T[:7u] + x^{5u} T[:6u] + x^{7u} T[:4u] + x^{9u} T[:2u] \\
& + x^{5u} T[:7u] + x^{6u} T[:6u] + x^{8u} T[:4u] + x^{10u} T[:2u] + x^{7u} T[:7u] \\
& + x^{9u} T[:5u] + x^{11u} T[:3u] + x^{13u} T[:u].
\end{aligned}$$

The proof of the other 15 cases are similar. We break down the above identity into 7 parts:

$$\begin{aligned}
0 &= x^{7u} T[:u] + x^{5u} T[2u:3u] + x^{3u} T[4u:5u] + x^u T[6u:7u] + T[7u:8u], \\
0 &= x^{8u} T[:u] + x^{7u} T[u:2u] + x^{6u} T[2u:3u] + x^{5u} T[3u:4u]
\end{aligned}$$

$$\begin{aligned}
& + x^{4u}T[4u:5u] + x^{3u}T[5u:6u] + x^{2u}T[6u:7u] + T[8u:9u], \\
0 & = x^{8u}T[u:2u] + x^{6u}T[3u:4u] + x^{4u}T[5u:6u] + T[9u:10u], \\
0 & = x^{10u}T[:u] + x^{8u}T[2u:3u] + x^{6u}T[4u:5u] + x^{4u}T[6u:7u] \\
& \quad + T[10u:11u], \\
0 & = x^{11u}T[:u] + x^{10u}T[u:2u] + x^{9u}T[2u:3u] + x^{8u}T[3u:4u] \\
& \quad + x^{7u}T[4u:5u] + x^{6u}T[5u:6u] + x^{5u}T[6u:7u] + T[11u:12u], \\
0 & = x^{11u}T[u:2u] + x^{9u}T[3u:4u] + x^{7u}T[5u:6u] + T[12u:13u], \\
0 & = x^{13u}T[:u] + x^{11u}T[2u:3u] + x^{9u}T[4u:5u] + x^{7u}T[6u:7u] \\
& \quad + T[13u:14u],
\end{aligned}$$

which can be rewritten as

$$(2.7) \quad 0 = T[[w]_2] + T[[10w]_2] + T[[100w]_2] + T[[110w]_2] + T[[111w]_2],$$

$$0 = T[[w]_2] + T[[1w]_2] + T[[10w]_2] + T[[11w]_2] + T[[100w]_2]$$

$$(2.8) \quad + T[[101w]_2] + T[[110w]_2] + T[[1000w]_2],$$

$$(2.9) \quad 0 = T[[1w]_2] + T[[11w]_2] + T[[101w]_2] + T[[1001w]_2],$$

$$(2.10) \quad 0 = T[[w]_2] + T[[10w]_2] + T[[100w]_2] + T[[110w]_2] + T[[1010w]_2],$$

$$0 = T[[w]_2] + T[[1w]_2] + T[[10w]_2] + T[[11w]_2] + T[[100w]_2]$$

$$(2.11) \quad + T[[101w]_2] + T[[110w]_2] + T[[1011w]_2],$$

$$(2.12) \quad 0 = T[[1w]_2] + T[[11w]_2] + T[[101w]_2] + T[[1100w]_2],$$

$$(2.13) \quad 0 = T[[w]_2] + T[[10w]_2] + T[[100w]_2] + T[[110w]_2] + T[[1101w]_2],$$

for all binary word w of length $2k$ and $w \neq 0^{2k}$, and

$$(2.14) \quad 0 = T[[w]_2] + T[[10w]_2] + T[[100w]_2] + T[[110w]_2] + T[[111w]_2],$$

$$0 = T[[w]_2] + T[[1w]_2] + T[[10w]_2] + T[[11w]_2] + T[[100w]_2]$$

$$(2.15) \quad + T[[101w]_2] + T[[110w]_2] + T[[1000w]_2],$$

$$(2.16) \quad 0 = T[[1w]_2] + T[[11w]_2] + T[[101w]_2] + T[[1001w]_2],$$

$$(2.17) \quad 0 = T[[w]_2] + T[[10w]_2] + T[[100w]_2] + T[[110w]_2] + T[[1010w]_2],$$

$$0 = T[[w]_2] + T[[1w]_2] + T[[10w]_2] + T[[11w]_2] + T[[100w]_2]$$

$$(2.18) \quad + T[[101w]_2] + T[[110w]_2] + T[[1011w]_2],$$

$$(2.19) \quad 0 = T[[1w]_2] + T[[11w]_2] + T[[101w]_2] + T[[1100w]_2],$$

$$(2.20) \quad 0 = T[[w]_2] + T[[10w]_2] + T[[100w]_2] + T[[110w]_2] + T[[1101w]_2],$$

for $w = 0^{2k}$.

First we calculate an 2-automaton that generates T from its minimal polynomial and its first terms. This automaton has 277 states; its transition function and output function can be found in the annex. Let $A(s, w)$ denote the state reached after reading w from right to left starting from the state s , and τ the output function. Define

$$E_{2k} = \{A(i, w) : |w| = 2k, w \neq 0^{2k}\}.$$

Identities (2.7) through (2.20) can be written as

$$0 = \tau(A(s, \epsilon)) + \tau(A(s, 10)) + \tau(A(s, 100)) + \tau(A(s, 110))$$

$$\begin{aligned}
(2.21) \quad & +\tau(A(s, 111)), \\
& 0 = \tau(A(s, \epsilon)) + \tau(A(s, 1)) + \tau(A(s, 10)) + \tau(A(s, 11)) + \tau(A(s, 100)) \\
(2.22) \quad & +\tau(A(s, 101)) + \tau(A(s, 110)) + \tau(A(s, 1000)), \\
(2.23) \quad & 0 = \tau(A(s, 1)) + \tau(A(s, 11)) + \tau(A(s, 101)) + \tau(A(s, 1001)), \\
& 0 = \tau(A(s, \epsilon)) + \tau(A(s, 10)) + \tau(A(s, 100)) + \tau(A(s, 110)) \\
(2.24) \quad & +\tau(A(s, 1010)), \\
& 0 = \tau(A(s, \epsilon)) + \tau(A(s, 1)) + \tau(A(s, 10)) + \tau(A(s, 11)) + \tau(A(s, 100)) \\
(2.25) \quad & +\tau(A(s, 101)) + \tau(A(s, 110)) + \tau(A(s, 1011)), \\
(2.26) \quad & 0 = \tau(A(s, 1)) + \tau(A(s, 11)) + \tau(A(s, 101)) + \tau(A(s, 1100)), \\
& 0 = \tau(A(s, \epsilon)) + \tau(A(s, 10)) + \tau(A(s, 100)) + \tau(A(s, 110)) \\
(2.27) \quad & +\tau(A(s, 1101)),
\end{aligned}$$

for all $s \in E_{2k}$, and

$$\begin{aligned}
(2.28) \quad & 0 = \tau(A(s, \epsilon)) + \tau(A(s, 10)) + \tau(A(s, 100)) + \tau(A(s, 110)) \\
& +\tau(A(s, 111)), \\
& 0 = \tau(A(s, \epsilon)) + \tau(A(s, 1)) + \tau(A(s, 10)) + \tau(A(s, 11)) + \tau(A(s, 100)) \\
(2.29) \quad & +\tau(A(s, 101)) + \tau(A(s, 110)) + \tau(A(s, 1000)), \\
(2.30) \quad & 0 = \tau(A(s, 1)) + \tau(A(s, 11)) + \tau(A(s, 101)) + \tau(A(s, 1001)), \\
& 0 = \tau(A(s, \epsilon)) + \tau(A(s, 10)) + \tau(A(s, 100)) + \tau(A(s, 110)) \\
(2.31) \quad & +\tau(A(s, 1010)), \\
& 0 = \tau(A(s, \epsilon)) + \tau(A(s, 1)) + \tau(A(s, 10)) + \tau(A(s, 11)) + \tau(A(s, 100)) \\
(2.32) \quad & +\tau(A(s, 101)) + \tau(A(s, 110)) + \tau(A(s, 1011)), \\
(2.33) \quad & 0 = \tau(A(s, 1)) + \tau(A(s, 11)) + \tau(A(s, 101)) + \tau(A(s, 1100)), \\
& 0 = \tau(A(s, \epsilon)) + \tau(A(s, 10)) + \tau(A(s, 100)) + \tau(A(s, 110)) \\
(2.34) \quad & +\tau(A(s, 1101)),
\end{aligned}$$

for $s = A(i, 0^{2k})$. We find that $(A(i, 0^{18}), E_{18}) = (A(i, 0^{16}), E_{16})$, so that we only have to verify that identities (2.21) through (2.34) hold for $2 \leq k \leq 9$, which turns out to be true. $\square$

In the following lemma, we express M_{2k} , M_{2k+1} , W_{2k} , and W_{2k+1} in terms of M^e , M^o , W^e , and W^o .

Lemma 2.3. *For all integer $k \geq 2$, and $u = 2^{2k-1}$,*

$$\begin{aligned}
M_{2k} &= M^e[:7u] + x^u M^e[:6u] + x^{3u} M^e[:4u] + x^{5u} M^e[:2u] + x^{7u} R^e, \\
W_{2k} &= W^e[:7u] + x^u W^e[:6u] + x^{3u} W^e[:4u] + x^{5u} W^e[:2u] + x^{7u} R^e.
\end{aligned}$$

For all integer $k \geq 2$, and $u = 2^{2k}$,

$$\begin{aligned}
M_{2k+1} &= M^o[:7u] + x^u M^o[:6u] + x^{3u} M^o[:4u] + x^{5u} M^o[:2u] + x^{7u} R^o, \\
W_{2k+1} &= W^o[:7u] + x^u W^o[:6u] + x^{3u} W^o[:4u] + x^{5u} W^o[:2u] + x^{7u} R^o.
\end{aligned}$$

Proof. We call the four identities in Lemma 2.3 also by the name M_{2k} , W_{2k} , M_{2k+1} , and W_{2k+1} . For $n = 2$, the identities can be verified directly. For $n \geq 2$, we claim

that

$$\begin{aligned} M_{2k} \wedge W_{2k} &\Rightarrow M_{2k+1} \wedge W_{2k+1}, \\ M_{2k+1} \wedge W_{2k+1} &\Rightarrow M_{2k+2} \wedge W_{2k+2}. \end{aligned}$$

We give the proof of

$$M_{2k} \wedge W_{2k} \Rightarrow M_{2k+1},$$

the other ones being similar. Set $u = 2^{2k}$ and $v = 2^{2k-1}$. By definition and induction hypothesis, the left side of identity M_{2k+1} is equal to

$$(2.35) \quad \begin{aligned} W_{2k} M_{2k} &= (W^e[:7v] + x^v W^e[:6v] + x^{3v} W^e[:4v] + x^{5v} W^e[:2v] + x^{7u} R^e) \times (\\ &M^e[:7v] + x^v M^e[:6v] + x^{3v} M^e[:4v] + x^{5v} M^e[:2v] + x^{7u} R^e); \end{aligned}$$

Call this expression *lhs*. Note that both sides of identity M_{2k+1} have the same term of highest degree $x^{14v} R^o$. Therefore we only need to prove that their difference is $O(x^{14v})$. Using Lemma 2.1 it can be seen that the right side of identity M_{2k+1} is congruent, modulo x^{14v} , to

$$\begin{aligned} W^e[:14v] M^e[:14v] + x^{2v} W^e[:12v] M^e[:12v] + x^{6v} W^e[:8v] M^e[:8v] \\ + x^{10v} W^e[:4v] M^e[:4v]. \end{aligned}$$

For all $n \leq 14$, replace the occurrences of $W^e[:n \cdot v]$ and $M^e[:n \cdot v]$ in the above expression by the reduction modulo $x^{n \cdot v}$ of the right side of the corresponding identity in Lemma 2.2 and get a new expression, which we call *rhs*. Define

$$\begin{aligned} X &:= x^v, \\ a_n &:= W^e[n \cdot v : (n+1) \cdot v] / X^n, \\ b_n &:= M^e[n \cdot v : (n+1) \cdot v] / X^n, \\ c &:= R^e. \end{aligned}$$

Using the notation introduced above, we can represent the expressions *lhs* (2.35) and *rhs* as polynomials in $\mathbb{F}_2[a_1, \dots, a_{14}, b_1, \dots, b_{14}, c][X]$. Note that it is not a problem that a_j commutes with b_k while W^e does not commute with M^e , because in the expressions concerned, the products of W^e -terms and M^e -terms are always in the same order. We let the computer do the simplification and check that the difference between these two polynomials is indeed $O(X^{14})$, which completes the proof. $\square$

Proof of Theorem 1.1. We prove the theorem for $\text{CF}(a, b)$; for $\text{CF}(b, a)$, the proof is similar. By Lemma 2.3, we have For all $0 \leq j, k \leq 1$,

$$\begin{aligned} \lim_{n \rightarrow \infty} M_{2n,j,k} &= M_{j,k}^e, \\ \lim_{n \rightarrow \infty} M_{2n+1,j,k} &= M_{j,k}^o, \\ \lim_{n \rightarrow \infty} W_{2n,j,k} &= W_{j,k}^e, \\ \lim_{n \rightarrow \infty} W_{2n+1,j,k} &= W_{j,k}^o. \end{aligned}$$

Let $z = 1/x$. By the convergence theorem and identity (2.2),

$$\text{CF}(a, b) = \frac{M_{0,1}^e(x)}{M_{0,0}^e(x)}.$$

By definition, $\phi(M^e, 0, 1)$ and $\phi(M^e, 0, 0)$ are minimal polynomials of $M_{0,1}^e$ and $M_{0,0}^e$. Therefore

$$P(x, y) = \text{Res}_t (\phi(M^e, 0, 1)(x, t), y^{12}\phi(M^e, 0, 1)(x, t/y))$$

is an annihilating polynomial of $f(x) = M_{0,1}^e/M_{0,0}^e$.

Define

$$Q(x, y) = q_4(x)y^4 + q_3(x)y^3 + q_2(x)y^2 + q_1(x)y + q_0(x),$$

where

$$q_0(x) = x^{20} + x^{19} + x^{16} + x^{11} + x^9 + x^8 + x^7,$$

$$q_1(x) = x^{19} + x^{17} + x^{15} + x^{11} + x^7 + x^3,$$

$$q_2(x) = x^{18} + x^{16} + x^8 + x^6 + 1,$$

$$q_3(x) = x^{19} + x^{17} + x^{15} + x^{11} + x^7 + x^3,$$

$$q_4(x) = x^{19} + x^{18} + x^{16} + x^{12} + x^{11} + x^9 + x^8 + x^7 + x^6.$$

The polynomial $Q(x, y)$ is the candidate for the minimal polynomial of $f(x)$ found by Padé-Hermite approximation. To prove that it is indeed the minimal polynomial of $f(x)$, we only need to prove that it is an irreducible factor of $P(x, y)$ of multiplicity m and $R(x, y) := P(x, y)/Q(x, y)^m$ is not an annihilating polynomial of $f(x)$. We verify the first point directly. For the second point, we find that when we truncate $f(x)$ to order 216, and substitute it for y in $R(z, y)$, we get a series with valuation smaller than 216, which proves that $R(z, y)$ is not an annihilating polynomial of $f(x)$. Finally, $z^{12}Q(1/z, y)$ is the minimal polynomial of $\text{CF}(a, b) = f(1/z)$. $\square$

3. ANNEXE

All 16 polynomials are of the form

$$p_0(x) + p_3(x)y^3 + p_6(x)y^6 + p_9(x)y^9 + p_{12}(x)y^{12}.$$

The coefficients $p_j(x)$ and the 16 initial terms to determine the solutions uniquely are given below.

For $\phi(M^e, 0, 0)$:

$$\begin{aligned} p_0(x) = & x^{114} + x^{112} + x^{110} + x^{104} + x^{102} + x^{98} + x^{92} + x^{88} + x^{86} + x^{80} + x^{78} \\ & + x^{74} + x^{68} + x^{66} + x^{64} + x^{62} + x^{60} + x^{56} + x^{50} + x^{48} + x^{46} + x^{38} \\ & + x^{36}, \end{aligned}$$

$$\begin{aligned} p_3(x) = & x^{106} + x^{104} + x^{100} + x^{98} + x^{96} + x^{94} + x^{92} + x^{88} + x^{86} + x^{76} + x^{72} \\ & + x^{66} + x^{58} + x^{56} + x^{54} + x^{46} + x^{40} + x^{34} + x^{22} + x^{20}, \end{aligned}$$

$$\begin{aligned} p_6(x) = & x^{110} + x^{108} + x^{104} + x^{102} + x^{98} + x^{94} + x^{92} + x^{90} + x^{86} + x^{84} + x^{78} \\ & + x^{68} + x^{64} + x^{62} + x^{60} + x^{58} + x^{54} + x^{50} + x^{48} + x^{46} + x^{44} + x^{40} \\ & + x^{32} + x^{30} + x^{26} + x^{22} + x^{20} + x^{12} + x^6 + 1, \end{aligned}$$

$$\begin{aligned} p_9(x) = & x^{112} + x^{110} + x^{108} + x^{106} + x^{104} + x^{100} + x^{98} + x^{96} + x^{88} + x^{84} \\ & + x^{76} + x^{72} + x^{68} + x^{66} + x^{64} + x^{60} + x^{58} + x^{56} + x^{48} + x^{44} + x^{40} \\ & + x^{38} + x^{34} + x^{24} + x^{22} + x^{14} + x^{12} + x^6, \end{aligned}$$

$$p_{12}(x) = x^{112} + x^{80} + x^{64} + x^{16} + 1,$$

and the initial terms are $[1, 0, 0, 0, 0, 0, 0, 0, 1, 0, 0, 0, 0, 0, 1, 0]$.

For $\phi(M^e, 0, 1)$:

$$\begin{aligned}
p_0(x) &= x^{111} + x^{110} + x^{103} + x^{102} + x^{95} + x^{94} + x^{91} + x^{90} + x^{86} + x^{85} + x^{82} \\
&\quad + x^{81} + x^{79} + x^{78} + x^{63} + x^{62} + x^{61} + x^{60} + x^{59} + x^{58} + x^{55} + x^{53} \\
&\quad + x^{51} + x^{49} + x^{47} + x^{44} + x^{43} + x^{40} + x^{39}, \\
p_3(x) &= x^{109} + x^{108} + x^{106} + x^{105} + x^{103} + x^{102} + x^{101} + x^{97} + x^{96} + x^{94} \\
&\quad + x^{93} + x^{91} + x^{90} + x^{88} + x^{87} + x^{75} + x^{74} + x^{73} + x^{71} + x^{70} + x^{68} \\
&\quad + x^{66} + x^{64} + x^{63} + x^{61} + x^{60} + x^{59} + x^{51} + x^{50} + x^{49} + x^{47} + x^{46} \\
&\quad + x^{44} + x^{42} + x^{40} + x^{39} + x^{37} + x^{36} + x^{35} + x^{29} + x^{28} + x^{27}, \\
p_6(x) &= x^{108} + x^{107} + x^{106} + x^{100} + x^{99} + x^{98} + x^{88} + x^{87} + x^{86} + x^{80} + x^{79} \\
&\quad + x^{78} + x^{76} + x^{75} + x^{74} + x^{72} + x^{71} + x^{70} + x^{68} + x^{67} + x^{66} + x^{56} \\
&\quad + x^{55} + x^{54} + x^{52} + x^{51} + x^{50} + x^{40} + x^{39} + x^{38} + x^{36} + x^{35} + x^{34} \\
&\quad + x^{28} + x^{27} + x^{26} + x^{24} + x^{23} + x^{22} + x^{20} + x^{19} + x^{18}, \\
p_9(x) &= x^{107} + x^{106} + x^{104} + x^{102} + x^{101} + x^{99} + x^{98} + x^{97} + x^{95} + x^{94} + x^{93} \\
&\quad + x^{91} + x^{90} + x^{89} + x^{87} + x^{86} + x^{84} + x^{82} + x^{81} + x^{79} + x^{78} + x^{77} \\
&\quad + x^{73} + x^{72} + x^{70} + x^{68} + x^{67} + x^{63} + x^{62} + x^{60} + x^{58} + x^{57} + x^{49} \\
&\quad + x^{48} + x^{47} + x^{41} + x^{40} + x^{38} + x^{36} + x^{34} + x^{33} + x^{31} + x^{30} + x^{28} \\
&\quad + x^{27} + x^{25} + x^{24} + x^{22} + x^{21} + x^{11} + x^{10} + x^9, \\
p_{12}(x) &= x^{106} + x^{105} + x^{104} + x^{102} + x^{101} + x^{100} + x^{94} + x^{93} + x^{92} + x^{90} \\
&\quad + x^{89} + x^{88} + x^{82} + x^{81} + x^{80} + x^{78} + x^{77} + x^{76} + x^{74} + x^{73} + x^{72} \\
&\quad + x^{66} + x^{65} + x^{64} + x^{62} + x^{61} + x^{60} + x^{58} + x^{57} + x^{56} + x^{50} + x^{49} \\
&\quad + x^{48} + x^{46} + x^{45} + x^{44} + x^{38} + x^{37} + x^{36} + x^{34} + x^{33} + x^{32} + x^{26} \\
&\quad + x^{25} + x^{24} + x^{22} + x^{21} + x^{20} + x^{14} + x^{13} + x^{12} + x^6 + x^5 + x^4 + x^2 \\
&\quad + x + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 1, 1, 0, 1, 1, 0, 1, 0, 1, 0, 1, 0, 0]$.

For $\phi(M^e, 1, 0)$:

$$\begin{aligned}
p_0(x) &= x^{111} + x^{110} + x^{103} + x^{102} + x^{95} + x^{94} + x^{91} + x^{90} + x^{86} + x^{85} + x^{82} \\
&\quad + x^{81} + x^{79} + x^{78} + x^{63} + x^{62} + x^{61} + x^{60} + x^{59} + x^{58} + x^{55} + x^{53} \\
&\quad + x^{51} + x^{49} + x^{47} + x^{44} + x^{43} + x^{40} + x^{39}, \\
p_3(x) &= x^{109} + x^{108} + x^{106} + x^{105} + x^{103} + x^{102} + x^{101} + x^{97} + x^{96} + x^{94} \\
&\quad + x^{93} + x^{91} + x^{90} + x^{88} + x^{87} + x^{75} + x^{74} + x^{73} + x^{71} + x^{70} + x^{68} \\
&\quad + x^{66} + x^{64} + x^{63} + x^{61} + x^{60} + x^{59} + x^{51} + x^{50} + x^{49} + x^{47} + x^{46} \\
&\quad + x^{44} + x^{42} + x^{40} + x^{39} + x^{37} + x^{36} + x^{35} + x^{29} + x^{28} + x^{27}, \\
p_6(x) &= x^{108} + x^{107} + x^{106} + x^{100} + x^{99} + x^{98} + x^{88} + x^{87} + x^{86} + x^{80} + x^{79} \\
&\quad + x^{78} + x^{76} + x^{75} + x^{74} + x^{72} + x^{71} + x^{70} + x^{68} + x^{67} + x^{66} + x^{56}
\end{aligned}$$

$$\begin{aligned}
& + x^{55} + x^{54} + x^{52} + x^{51} + x^{50} + x^{40} + x^{39} + x^{38} + x^{36} + x^{35} + x^{34} \\
& + x^{28} + x^{27} + x^{26} + x^{24} + x^{23} + x^{22} + x^{20} + x^{19} + x^{18}, \\
p_9(x) = & x^{107} + x^{106} + x^{104} + x^{102} + x^{101} + x^{99} + x^{98} + x^{97} + x^{95} + x^{94} + x^{93} \\
& + x^{91} + x^{90} + x^{89} + x^{87} + x^{86} + x^{84} + x^{82} + x^{81} + x^{79} + x^{78} + x^{77} \\
& + x^{73} + x^{72} + x^{70} + x^{68} + x^{67} + x^{63} + x^{62} + x^{60} + x^{58} + x^{57} + x^{49} \\
& + x^{48} + x^{47} + x^{41} + x^{40} + x^{38} + x^{36} + x^{34} + x^{33} + x^{31} + x^{30} + x^{28} \\
& + x^{27} + x^{25} + x^{24} + x^{22} + x^{21} + x^{11} + x^{10} + x^9, \\
p_{12}(x) = & x^{106} + x^{105} + x^{104} + x^{102} + x^{101} + x^{100} + x^{94} + x^{93} + x^{92} + x^{90} \\
& + x^{89} + x^{88} + x^{82} + x^{81} + x^{80} + x^{78} + x^{77} + x^{76} + x^{74} + x^{73} + x^{72} \\
& + x^{66} + x^{65} + x^{64} + x^{62} + x^{61} + x^{60} + x^{58} + x^{57} + x^{56} + x^{50} + x^{49} \\
& + x^{48} + x^{46} + x^{45} + x^{44} + x^{38} + x^{37} + x^{36} + x^{34} + x^{33} + x^{32} + x^{26} \\
& + x^{25} + x^{24} + x^{22} + x^{21} + x^{20} + x^{14} + x^{13} + x^{12} + x^6 + x^5 + x^4 + x^2 \\
& + x + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 1, 1, 0, 1, 1, 0, 1, 0, 1, 0, 1, 0, 0]$.

For $\phi(M^e, 1, 1)$:

$$\begin{aligned}
p_0(x) = & x^{108} + x^{104} + x^{102} + x^{96} + x^{92} + x^{90} + x^{88} + x^{86} + x^{84} + x^{82} + x^{80} \\
& + x^{78} + x^{68} + x^{66} + x^{64} + x^{62} + x^{60} + x^{56} + x^{52} + x^{50} + x^{42}, \\
p_3(x) = & x^{102} + x^{100} + x^{96} + x^{94} + x^{92} + x^{90} + x^{80} + x^{78} + x^{72} + x^{66} + x^{60} \\
& + x^{58} + x^{52} + x^{50} + x^{48} + x^{42} + x^{40} + x^{34} + x^{28} + x^{26} + x^{22} + x^{18}, \\
p_6(x) = & x^{102} + x^{96} + x^{94} + x^{86} + x^{82} + x^{80} + x^{78} + x^{74} + x^{72} + x^{70} + x^{64} \\
& + x^{60} + x^{58} + x^{56} + x^{46} + x^{44} + x^{40} + x^{36} + x^{32} + x^{30} + x^{22} + x^{16} \\
& + x^{14} + x^{12} + x^6 + x^4 + x^2 + 1, \\
p_9(x) = & x^{100} + x^{96} + x^{94} + x^{88} + x^{86} + x^{80} + x^{76} + x^{74} + x^{66} + x^{62} + x^{60} \\
& + x^{58} + x^{54} + x^{42} + x^{40} + x^{38} + x^{36} + x^{32} + x^{28} + x^{26} + x^{22} + x^{20} \\
& + x^{16} + x^{14} + x^{12} + x^{10} + x^8 + x^6, \\
p_{12}(x) = & x^{100} + x^{98} + x^{96} + x^{92} + x^{90} + x^{88} + x^{76} + x^{74} + x^{72} + x^{60} + x^{58} \\
& + x^{56} + x^{44} + x^{42} + x^{40} + x^{36} + x^{34} + x^{32} + x^{20} + x^{18} + x^{16} + x^{12} \\
& + x^{10} + x^8 + x^4 + x^2 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 0, 1, 0, 1, 0, 0, 0, 1, 0, 0, 0]$.

For $\phi(W^e, 0, 0)$:

$$\begin{aligned}
p_0(x) = & x^{108} + x^{100} + x^{98} + x^{94} + x^{90} + x^{88} + x^{82} + x^{72} + x^{64} + x^{62} + x^{54} \\
& + x^{50} + x^{48} + x^{46} + x^{44} + x^{42} + x^{36}, \\
p_3(x) = & x^{102} + x^{100} + x^{98} + x^{92} + x^{88} + x^{84} + x^{78} + x^{76} + x^{74} + x^{72} + x^{70} \\
& + x^{68} + x^{66} + x^{64} + x^{62} + x^{60} + x^{58} + x^{54} + x^{50} + x^{46} + x^{42} + x^{40} \\
& + x^{38} + x^{36} + x^{34} + x^{32} + x^{30} + x^{28} + x^{26} + x^{20},
\end{aligned}$$

$$\begin{aligned}
p_6(x) &= x^{102} + x^{98} + x^{96} + x^{94} + x^{92} + x^{90} + x^{88} + x^{86} + x^{84} + x^{82} + x^{80} \\
&\quad + x^{72} + x^{68} + x^{62} + x^{60} + x^{58} + x^{48} + x^{44} + x^{36} + x^{24} + x^{22} + x^{18} \\
&\quad + x^6 + x^4 + x^2 + 1, \\
p_9(x) &= x^{100} + x^{96} + x^{94} + x^{88} + x^{86} + x^{80} + x^{76} + x^{74} + x^{66} + x^{62} + x^{60} \\
&\quad + x^{58} + x^{54} + x^{42} + x^{40} + x^{38} + x^{36} + x^{32} + x^{28} + x^{26} + x^{22} + x^{20} \\
&\quad + x^{16} + x^{14} + x^{12} + x^{10} + x^8 + x^6, \\
p_{12}(x) &= x^{100} + x^{98} + x^{96} + x^{92} + x^{90} + x^{88} + x^{76} + x^{74} + x^{72} + x^{60} + x^{58} \\
&\quad + x^{56} + x^{44} + x^{42} + x^{40} + x^{36} + x^{34} + x^{32} + x^{20} + x^{18} + x^{16} + x^{12} \\
&\quad + x^{10} + x^8 + x^4 + x^2 + 1,
\end{aligned}$$

and the initial terms are $[1, 0, 0, 0, 0, 0, 1, 0, 1, 0, 0, 0, 0, 0, 0, 0]$.

For $\phi(W^e, 0, 1)$:

$$\begin{aligned}
p_0(x) &= x^{111} + x^{108} + x^{107} + x^{106} + x^{105} + x^{104} + x^{100} + x^{97} + x^{94} + x^{92} \\
&\quad + x^{91} + x^{90} + x^{89} + x^{88} + x^{87} + x^{85} + x^{84} + x^{81} + x^{79} + x^{78} + x^{75} \\
&\quad + x^{74} + x^{72} + x^{71} + x^{70} + x^{69} + x^{68} + x^{64} + x^{63} + x^{60} + x^{55} + x^{47} \\
&\quad + x^{46} + x^{45} + x^{43} + x^{40} + x^{39}, \\
p_3(x) &= x^{109} + x^{108} + x^{106} + x^{105} + x^{103} + x^{102} + x^{101} + x^{97} + x^{96} + x^{94} \\
&\quad + x^{93} + x^{91} + x^{90} + x^{88} + x^{87} + x^{75} + x^{74} + x^{73} + x^{71} + x^{70} + x^{68} \\
&\quad + x^{66} + x^{64} + x^{63} + x^{61} + x^{60} + x^{59} + x^{51} + x^{50} + x^{49} + x^{47} + x^{46} \\
&\quad + x^{44} + x^{42} + x^{40} + x^{39} + x^{37} + x^{36} + x^{35} + x^{29} + x^{28} + x^{27}, \\
p_6(x) &= x^{108} + x^{107} + x^{106} + x^{100} + x^{99} + x^{98} + x^{88} + x^{87} + x^{86} + x^{80} + x^{79} \\
&\quad + x^{78} + x^{76} + x^{75} + x^{74} + x^{72} + x^{71} + x^{70} + x^{68} + x^{67} + x^{66} + x^{56} \\
&\quad + x^{55} + x^{54} + x^{52} + x^{51} + x^{50} + x^{40} + x^{39} + x^{38} + x^{36} + x^{35} + x^{34} \\
&\quad + x^{28} + x^{27} + x^{26} + x^{24} + x^{23} + x^{22} + x^{20} + x^{19} + x^{18}, \\
p_9(x) &= x^{107} + x^{106} + x^{104} + x^{102} + x^{101} + x^{99} + x^{98} + x^{97} + x^{95} + x^{94} + x^{93} \\
&\quad + x^{91} + x^{90} + x^{89} + x^{87} + x^{86} + x^{84} + x^{82} + x^{81} + x^{79} + x^{78} + x^{77} \\
&\quad + x^{73} + x^{72} + x^{70} + x^{68} + x^{67} + x^{63} + x^{62} + x^{60} + x^{58} + x^{57} + x^{49} \\
&\quad + x^{48} + x^{47} + x^{41} + x^{40} + x^{38} + x^{36} + x^{34} + x^{33} + x^{31} + x^{30} + x^{28} \\
&\quad + x^{27} + x^{25} + x^{24} + x^{22} + x^{21} + x^{11} + x^{10} + x^9, \\
p_{12}(x) &= x^{106} + x^{105} + x^{104} + x^{102} + x^{101} + x^{100} + x^{94} + x^{93} + x^{92} + x^{90} \\
&\quad + x^{89} + x^{88} + x^{82} + x^{81} + x^{80} + x^{78} + x^{77} + x^{76} + x^{74} + x^{73} + x^{72} \\
&\quad + x^{66} + x^{65} + x^{64} + x^{62} + x^{61} + x^{60} + x^{58} + x^{57} + x^{56} + x^{50} + x^{49} \\
&\quad + x^{48} + x^{46} + x^{45} + x^{44} + x^{38} + x^{37} + x^{36} + x^{34} + x^{33} + x^{32} + x^{26} \\
&\quad + x^{25} + x^{24} + x^{22} + x^{21} + x^{20} + x^{14} + x^{13} + x^{12} + x^6 + x^5 + x^4 + x^2 \\
&\quad + x + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 1, 0, 1, 0, 0, 0, 0, 1, 0, 0, 0, 0]$.

For $\phi(W^e, 1, 0)$:

$$\begin{aligned}
p_0(x) &= x^{111} + x^{108} + x^{107} + x^{106} + x^{105} + x^{104} + x^{100} + x^{97} + x^{94} + x^{92} \\
&\quad + x^{91} + x^{90} + x^{89} + x^{88} + x^{87} + x^{85} + x^{84} + x^{81} + x^{79} + x^{78} + x^{75} \\
&\quad + x^{74} + x^{72} + x^{71} + x^{70} + x^{69} + x^{68} + x^{64} + x^{63} + x^{60} + x^{55} + x^{47} \\
&\quad + x^{46} + x^{45} + x^{43} + x^{40} + x^{39}, \\
p_3(x) &= x^{109} + x^{108} + x^{106} + x^{105} + x^{103} + x^{102} + x^{101} + x^{97} + x^{96} + x^{94} \\
&\quad + x^{93} + x^{91} + x^{90} + x^{88} + x^{87} + x^{75} + x^{74} + x^{73} + x^{71} + x^{70} + x^{68} \\
&\quad + x^{66} + x^{64} + x^{63} + x^{61} + x^{60} + x^{59} + x^{51} + x^{50} + x^{49} + x^{47} + x^{46} \\
&\quad + x^{44} + x^{42} + x^{40} + x^{39} + x^{37} + x^{36} + x^{35} + x^{29} + x^{28} + x^{27}, \\
p_6(x) &= x^{108} + x^{107} + x^{106} + x^{100} + x^{99} + x^{98} + x^{88} + x^{87} + x^{86} + x^{80} + x^{79} \\
&\quad + x^{78} + x^{76} + x^{75} + x^{74} + x^{72} + x^{71} + x^{70} + x^{68} + x^{67} + x^{66} + x^{56} \\
&\quad + x^{55} + x^{54} + x^{52} + x^{51} + x^{50} + x^{40} + x^{39} + x^{38} + x^{36} + x^{35} + x^{34} \\
&\quad + x^{28} + x^{27} + x^{26} + x^{24} + x^{23} + x^{22} + x^{20} + x^{19} + x^{18}, \\
p_9(x) &= x^{107} + x^{106} + x^{104} + x^{102} + x^{101} + x^{99} + x^{98} + x^{97} + x^{95} + x^{94} + x^{93} \\
&\quad + x^{91} + x^{90} + x^{89} + x^{87} + x^{86} + x^{84} + x^{82} + x^{81} + x^{79} + x^{78} + x^{77} \\
&\quad + x^{73} + x^{72} + x^{70} + x^{68} + x^{67} + x^{63} + x^{62} + x^{60} + x^{58} + x^{57} + x^{49} \\
&\quad + x^{48} + x^{47} + x^{41} + x^{40} + x^{38} + x^{36} + x^{34} + x^{33} + x^{31} + x^{30} + x^{28} \\
&\quad + x^{27} + x^{25} + x^{24} + x^{22} + x^{21} + x^{11} + x^{10} + x^9, \\
p_{12}(x) &= x^{106} + x^{105} + x^{104} + x^{102} + x^{101} + x^{100} + x^{94} + x^{93} + x^{92} + x^{90} \\
&\quad + x^{89} + x^{88} + x^{82} + x^{81} + x^{80} + x^{78} + x^{77} + x^{76} + x^{74} + x^{73} + x^{72} \\
&\quad + x^{66} + x^{65} + x^{64} + x^{62} + x^{61} + x^{60} + x^{58} + x^{57} + x^{56} + x^{50} + x^{49} \\
&\quad + x^{48} + x^{46} + x^{45} + x^{44} + x^{38} + x^{37} + x^{36} + x^{34} + x^{33} + x^{32} + x^{26} \\
&\quad + x^{25} + x^{24} + x^{22} + x^{21} + x^{20} + x^{14} + x^{13} + x^{12} + x^6 + x^5 + x^4 + x^2 \\
&\quad + x + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 1, 0, 1, 0, 0, 0, 0, 1, 0, 0, 0, 0]$.

For $\phi(W^e, 1, 1)$:

$$\begin{aligned}
p_0(x) &= x^{114} + x^{108} + x^{102} + x^{100} + x^{98} + x^{96} + x^{94} + x^{92} + x^{90} + x^{88} + x^{78} \\
&\quad + x^{76} + x^{74} + x^{72} + x^{66} + x^{60} + x^{56} + x^{50} + x^{48} + x^{44} + x^{42}, \\
p_3(x) &= x^{110} + x^{106} + x^{104} + x^{100} + x^{96} + x^{92} + x^{88} + x^{86} + x^{82} + x^{76} + x^{72} \\
&\quad + x^{70} + x^{66} + x^{56} + x^{46} + x^{42} + x^{40} + x^{38} + x^{34} + x^{30} + x^{26} + x^{22} \\
&\quad + x^{20} + x^{18}, \\
p_6(x) &= x^{104} + x^{102} + x^{100} + x^{92} + x^{90} + x^{86} + x^{82} + x^{78} + x^{76} + x^{70} + x^{64} \\
&\quad + x^{62} + x^{50} + x^{48} + x^{46} + x^{42} + x^{40} + x^{38} + x^{32} + x^{22} + x^{20} + x^{18} \\
&\quad + x^6 + 1, \\
p_9(x) &= x^{112} + x^{110} + x^{108} + x^{106} + x^{104} + x^{100} + x^{98} + x^{96} + x^{88} + x^{84}
\end{aligned}$$

$$\begin{aligned}
& + x^{76} + x^{72} + x^{68} + x^{66} + x^{64} + x^{60} + x^{58} + x^{56} + x^{48} + x^{44} + x^{40} \\
& + x^{38} + x^{34} + x^{24} + x^{22} + x^{14} + x^{12} + x^6, \\
p_{12}(x) &= x^{112} + x^{80} + x^{64} + x^{16} + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 0, 0, 0, 1, 0, 0, 0, 1, 0, 1, 0]$.

For $\phi(M^o, 0, 0)$:

$$\begin{aligned}
p_0(x) &= x^{111} + x^{110} + x^{109} + x^{107} + x^{106} + x^{105} + x^{102} + x^{99} + x^{97} + x^{96} \\
& + x^{94} + x^{93} + x^{92} + x^{88} + x^{80} + x^{79} + x^{78} + x^{77} + x^{75} + x^{74} + x^{73} \\
& + x^{71} + x^{69} + x^{68} + x^{66} + x^{65} + x^{62} + x^{61} + x^{58} + x^{52} + x^{51} + x^{49} \\
& + x^{47} + x^{46} + x^{45} + x^{43} + x^{42} + x^{41} + x^{39} + x^{37} + x^{36}, \\
p_3(x) &= x^{110} + x^{107} + x^{106} + x^{103} + x^{102} + x^{100} + x^{99} + x^{96} + x^{91} + x^{90} \\
& + x^{89} + x^{88} + x^{84} + x^{83} + x^{81} + x^{80} + x^{76} + x^{74} + x^{73} + x^{72} + x^{71} \\
& + x^{68} + x^{66} + x^{64} + x^{63} + x^{60} + x^{59} + x^{57} + x^{55} + x^{54} + x^{51} + x^{50} \\
& + x^{48} + x^{46} + x^{43} + x^{42} + x^{41} + x^{40} + x^{38} + x^{30} + x^{28} + x^{27} + x^{25} \\
& + x^{24}, \\
p_6(x) &= x^{110} + x^{107} + x^{106} + x^{105} + x^{104} + x^{102} + x^{101} + x^{98} + x^{95} + x^{94} \\
& + x^{93} + x^{91} + x^{90} + x^{85} + x^{82} + x^{79} + x^{75} + x^{70} + x^{65} + x^{64} + x^{63} \\
& + x^{61} + x^{59} + x^{58} + x^{56} + x^{50} + x^{49} + x^{47} + x^{46} + x^{44} + x^{41} + x^{38} \\
& + x^{36} + x^{34} + x^{33} + x^{28} + x^{27} + x^{22} + x^{21} + x^{20} + x^{18} + x^{16} + x^{15} \\
& + x^{14} + x^6 + x^5 + x^4 + x^2 + x + 1, \\
p_9(x) &= x^{110} + x^{109} + x^{108} + x^{106} + x^{105} + x^{104} + x^{98} + x^{97} + x^{96} + x^{90} \\
& + x^{89} + x^{88} + x^{82} + x^{81} + x^{80} + x^{74} + x^{73} + x^{72} + x^{66} + x^{65} + x^{64} \\
& + x^{58} + x^{57} + x^{56} + x^{50} + x^{49} + x^{48} + x^{46} + x^{45} + x^{44} + x^{30} + x^{29} \\
& + x^{28} + x^{26} + x^{25} + x^{24} + x^{18} + x^{17} + x^{16} + x^{14} + x^{13} + x^{12}, \\
p_{12}(x) &= x^{110} + x^{109} + x^{108} + x^{102} + x^{101} + x^{100} + x^{98} + x^{97} + x^{96} + x^{94} \\
& + x^{93} + x^{92} + x^{78} + x^{77} + x^{76} + x^{62} + x^{61} + x^{60} + x^{38} + x^{37} + x^{36} \\
& + x^{34} + x^{33} + x^{32} + x^{30} + x^{29} + x^{28} + x^{22} + x^{21} + x^{20} + x^{18} + x^{17} \\
& + x^{16} + x^6 + x^5 + x^4 + x^2 + x + 1,
\end{aligned}$$

and the initial terms are $[1, 0, 0, 0, 0, 0, 1, 1, 1, 1, 0, 1, 1, 0, 1, 1]$.

For $\phi(M^o, 0, 1)$:

$$\begin{aligned}
p_0(x) &= x^{108} + x^{106} + x^{105} + x^{104} + x^{103} + x^{101} + x^{99} + x^{97} + x^{94} + x^{92} \\
& + x^{85} + x^{84} + x^{80} + x^{77} + x^{76} + x^{74} + x^{73} + x^{72} + x^{69} + x^{68} + x^{66} \\
& + x^{64} + x^{63} + x^{58} + x^{56} + x^{46} + x^{45} + x^{42} + x^{39}, \\
p_3(x) &= x^{101} + x^{99} + x^{77} + x^{75} + x^{69} + x^{67} + x^{61} + x^{59} + x^{53} + x^{51} + x^{45} \\
& + x^{43} + x^{37} + x^{35} + x^{29} + x^{27}, \\
p_6(x) &= x^{102} + x^{100} + x^{96} + x^{92} + x^{90} + x^{86} + x^{84} + x^{82} + x^{78} + x^{76} + x^{74}
\end{aligned}$$

$$\begin{aligned}
& + x^{66} + x^{64} + x^{62} + x^{58} + x^{56} + x^{52} + x^{48} + x^{44} + x^{42} + x^{38} + x^{36} \\
& + x^{34} + x^{22} + x^{20} + x^{18}, \\
p_9(x) &= x^{103} + x^{101} + x^{91} + x^{89} + x^{87} + x^{85} + x^{75} + x^{73} + x^{71} + x^{69} + x^{59} \\
& + x^{57} + x^{55} + x^{53} + x^{43} + x^{41} + x^{23} + x^{21} + x^{11} + x^9, \\
p_{12}(x) &= x^{104} + x^{102} + x^{98} + x^{94} + x^{90} + x^{86} + x^{84} + x^{80} + x^{78} + x^{74} + x^{70} \\
& + x^{68} + x^{64} + x^{62} + x^{58} + x^{54} + x^{52} + x^{48} + x^{46} + x^{42} + x^{40} + x^{36} \\
& + x^{34} + x^{32} + x^{24} + x^{22} + x^{18} + x^{14} + x^{10} + x^8 + x^4 + x^2 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 1, 1, 0, 1, 1, 0, 0, 0, 0, 0, 0, 0]$.

For $\phi(M^o, 1, 0)$:

$$\begin{aligned}
p_0(x) &= x^{114} + x^{113} + x^{112} + x^{111} + x^{110} + x^{106} + x^{104} + x^{102} + x^{100} + x^{94} \\
& + x^{92} + x^{90} + x^{88} + x^{87} + x^{85} + x^{84} + x^{82} + x^{80} + x^{78} + x^{77} + x^{75} \\
& + x^{71} + x^{69} + x^{68} + x^{67} + x^{64} + x^{63} + x^{61} + x^{60} + x^{58} + x^{57} + x^{56} \\
& + x^{55} + x^{54} + x^{53} + x^{52} + x^{51} + x^{48} + x^{47} + x^{46} + x^{42} + x^{41} + x^{39}, \\
p_3(x) &= x^{113} + x^{109} + x^{107} + x^{105} + x^{103} + x^{99} + x^{89} + x^{85} + x^{83} + x^{79} + x^{77} \\
& + x^{71} + x^{69} + x^{63} + x^{61} + x^{55} + x^{53} + x^{47} + x^{45} + x^{39} + x^{37} + x^{33} \\
& + x^{31} + x^{27}, \\
p_6(x) &= x^{114} + x^{110} + x^{102} + x^{98} + x^{94} + x^{90} + x^{78} + x^{74} + x^{66} + x^{62} + x^{50} \\
& + x^{46} + x^{26} + x^{18}, \\
p_9(x) &= x^{115} + x^{111} + x^{109} + x^{107} + x^{105} + x^{103} + x^{101} + x^{97} + x^{91} + x^{87} \\
& + x^{85} + x^{81} + x^{75} + x^{71} + x^{69} + x^{65} + x^{59} + x^{55} + x^{53} + x^{51} + x^{49} \\
& + x^{47} + x^{45} + x^{41} + x^{35} + x^{31} + x^{29} + x^{27} + x^{25} + x^{23} + x^{21} + x^{19} \\
& + x^{17} + x^{15} + x^{13} + x^9, \\
p_{12}(x) &= x^{116} + x^{112} + x^{100} + x^{92} + x^{84} + x^{80} + x^{76} + x^{68} + x^{64} + x^{60} + x^{44} \\
& + x^{36} + x^{16} + x^{12} + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 1, 0, 1, 0, 0, 0, 0, 0, 0, 0, 0]$.

For $\phi(M^o, 1, 1)$:

$$\begin{aligned}
p_0(x) &= x^{111} + x^{109} + x^{106} + x^{105} + x^{104} + x^{103} + x^{102} + x^{100} + x^{96} + x^{85} \\
& + x^{81} + x^{80} + x^{77} + x^{75} + x^{72} + x^{71} + x^{69} + x^{64} + x^{63} + x^{57} + x^{56} \\
& + x^{55} + x^{54} + x^{52} + x^{51} + x^{50} + x^{49} + x^{48} + x^{45} + x^{43} + x^{42}, \\
p_3(x) &= x^{110} + x^{108} + x^{106} + x^{105} + x^{102} + x^{101} + x^{99} + x^{98} + x^{97} + x^{91} \\
& + x^{90} + x^{89} + x^{88} + x^{80} + x^{79} + x^{77} + x^{76} + x^{75} + x^{73} + x^{69} + x^{68} \\
& + x^{67} + x^{61} + x^{60} + x^{58} + x^{57} + x^{56} + x^{54} + x^{53} + x^{47} + x^{46} + x^{45} \\
& + x^{41} + x^{39} + x^{38} + x^{37} + x^{35} + x^{34} + x^{32} + x^{31} + x^{30} + x^{29} + x^{27} \\
& + x^{26}, \\
p_6(x) &= x^{110} + x^{108} + x^{106} + x^{102} + x^{101} + x^{100} + x^{99} + x^{95} + x^{92} + x^{91}
\end{aligned}$$

$$\begin{aligned}
& + x^{90} + x^{88} + x^{87} + x^{84} + x^{82} + x^{80} + x^{78} + x^{76} + x^{74} + x^{72} + x^{71} \\
& + x^{70} + x^{69} + x^{67} + x^{66} + x^{65} + x^{64} + x^{63} + x^{61} + x^{59} + x^{57} + x^{56} \\
& + x^{55} + x^{54} + x^{52} + x^{51} + x^{49} + x^{47} + x^{46} + x^{44} + x^{42} + x^{39} + x^{35} \\
& + x^{33} + x^{26} + x^{24} + x^{23} + x^{21} + x^{19} + x^{18} + x^{17} + x^{15} + x^{13} + x^{12} \\
& + x^6 + x^5 + x^4 + x^2 + x + 1, \\
p_9(x) &= x^{110} + x^{109} + x^{108} + x^{106} + x^{105} + x^{104} + x^{98} + x^{97} + x^{96} + x^{90} \\
& + x^{89} + x^{88} + x^{82} + x^{81} + x^{80} + x^{74} + x^{73} + x^{72} + x^{66} + x^{65} + x^{64} \\
& + x^{58} + x^{57} + x^{56} + x^{50} + x^{49} + x^{48} + x^{46} + x^{45} + x^{44} + x^{30} + x^{29} \\
& + x^{28} + x^{26} + x^{25} + x^{24} + x^{18} + x^{17} + x^{16} + x^{14} + x^{13} + x^{12}, \\
p_{12}(x) &= x^{110} + x^{109} + x^{108} + x^{102} + x^{101} + x^{100} + x^{98} + x^{97} + x^{96} + x^{94} \\
& + x^{93} + x^{92} + x^{78} + x^{77} + x^{76} + x^{62} + x^{61} + x^{60} + x^{38} + x^{37} + x^{36} \\
& + x^{34} + x^{33} + x^{32} + x^{30} + x^{29} + x^{28} + x^{22} + x^{21} + x^{20} + x^{18} + x^{17} \\
& + x^{16} + x^6 + x^5 + x^4 + x^2 + x + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 0, 0, 1, 1, 1, 0, 1, 1, 0, 0, 1]$.

For $\phi(W^o, 0, 0)$:

$$\begin{aligned}
p_0(x) &= x^{111} + x^{110} + x^{109} + x^{107} + x^{106} + x^{105} + x^{102} + x^{99} + x^{97} + x^{96} \\
& + x^{94} + x^{93} + x^{92} + x^{88} + x^{80} + x^{79} + x^{78} + x^{77} + x^{75} + x^{74} + x^{73} \\
& + x^{71} + x^{69} + x^{68} + x^{66} + x^{65} + x^{62} + x^{61} + x^{58} + x^{52} + x^{51} + x^{49} \\
& + x^{47} + x^{46} + x^{45} + x^{43} + x^{42} + x^{41} + x^{39} + x^{37} + x^{36}, \\
p_3(x) &= x^{110} + x^{107} + x^{106} + x^{103} + x^{102} + x^{100} + x^{99} + x^{96} + x^{91} + x^{90} \\
& + x^{89} + x^{88} + x^{84} + x^{83} + x^{81} + x^{80} + x^{76} + x^{74} + x^{73} + x^{72} + x^{71} \\
& + x^{68} + x^{66} + x^{64} + x^{63} + x^{60} + x^{59} + x^{57} + x^{55} + x^{54} + x^{51} + x^{50} \\
& + x^{48} + x^{46} + x^{43} + x^{42} + x^{41} + x^{40} + x^{38} + x^{30} + x^{28} + x^{27} + x^{25} \\
& + x^{24}, \\
p_6(x) &= x^{110} + x^{107} + x^{106} + x^{105} + x^{104} + x^{102} + x^{101} + x^{98} + x^{95} + x^{94} \\
& + x^{93} + x^{91} + x^{90} + x^{85} + x^{82} + x^{79} + x^{75} + x^{70} + x^{65} + x^{64} + x^{63} \\
& + x^{61} + x^{59} + x^{58} + x^{56} + x^{50} + x^{49} + x^{47} + x^{46} + x^{44} + x^{41} + x^{38} \\
& + x^{36} + x^{34} + x^{33} + x^{28} + x^{27} + x^{22} + x^{21} + x^{20} + x^{18} + x^{16} + x^{15} \\
& + x^{14} + x^6 + x^5 + x^4 + x^2 + x + 1, \\
p_9(x) &= x^{110} + x^{109} + x^{108} + x^{106} + x^{105} + x^{104} + x^{98} + x^{97} + x^{96} + x^{90} \\
& + x^{89} + x^{88} + x^{82} + x^{81} + x^{80} + x^{74} + x^{73} + x^{72} + x^{66} + x^{65} + x^{64} \\
& + x^{58} + x^{57} + x^{56} + x^{50} + x^{49} + x^{48} + x^{46} + x^{45} + x^{44} + x^{30} + x^{29} \\
& + x^{28} + x^{26} + x^{25} + x^{24} + x^{18} + x^{17} + x^{16} + x^{14} + x^{13} + x^{12}, \\
p_{12}(x) &= x^{110} + x^{109} + x^{108} + x^{102} + x^{101} + x^{100} + x^{98} + x^{97} + x^{96} + x^{94} \\
& + x^{93} + x^{92} + x^{78} + x^{77} + x^{76} + x^{62} + x^{61} + x^{60} + x^{38} + x^{37} + x^{36}
\end{aligned}$$

$$\begin{aligned}
& + x^{34} + x^{33} + x^{32} + x^{30} + x^{29} + x^{28} + x^{22} + x^{21} + x^{20} + x^{18} + x^{17} \\
& + x^{16} + x^6 + x^5 + x^4 + x^2 + x + 1,
\end{aligned}$$

and the initial terms are $[1, 0, 0, 0, 0, 0, 1, 1, 1, 1, 0, 1, 1, 0, 1, 1]$.

For $\phi(W^o, 0, 1)$:

$$\begin{aligned}
p_0(x) &= x^{114} + x^{113} + x^{112} + x^{111} + x^{110} + x^{106} + x^{104} + x^{102} + x^{100} + x^{94} \\
&+ x^{92} + x^{90} + x^{88} + x^{87} + x^{85} + x^{84} + x^{82} + x^{80} + x^{78} + x^{77} + x^{75} \\
&+ x^{71} + x^{69} + x^{68} + x^{67} + x^{64} + x^{63} + x^{61} + x^{60} + x^{58} + x^{57} + x^{56} \\
&+ x^{55} + x^{54} + x^{53} + x^{52} + x^{51} + x^{48} + x^{47} + x^{46} + x^{42} + x^{41} + x^{39}, \\
p_3(x) &= x^{113} + x^{109} + x^{107} + x^{105} + x^{103} + x^{99} + x^{89} + x^{85} + x^{83} + x^{79} + x^{77} \\
&+ x^{71} + x^{69} + x^{63} + x^{61} + x^{55} + x^{53} + x^{47} + x^{45} + x^{39} + x^{37} + x^{33} \\
&+ x^{31} + x^{27}, \\
p_6(x) &= x^{114} + x^{110} + x^{102} + x^{98} + x^{94} + x^{90} + x^{78} + x^{74} + x^{66} + x^{62} + x^{50} \\
&+ x^{46} + x^{26} + x^{18}, \\
p_9(x) &= x^{115} + x^{111} + x^{109} + x^{107} + x^{105} + x^{103} + x^{101} + x^{97} + x^{91} + x^{87} \\
&+ x^{85} + x^{81} + x^{75} + x^{71} + x^{69} + x^{65} + x^{59} + x^{55} + x^{53} + x^{51} + x^{49} \\
&+ x^{47} + x^{45} + x^{41} + x^{35} + x^{31} + x^{29} + x^{27} + x^{25} + x^{23} + x^{21} + x^{19} \\
&+ x^{17} + x^{15} + x^{13} + x^9, \\
p_{12}(x) &= x^{116} + x^{112} + x^{100} + x^{92} + x^{84} + x^{80} + x^{76} + x^{68} + x^{64} + x^{60} + x^{44} \\
&+ x^{36} + x^{16} + x^{12} + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 1, 0, 1, 0, 0, 0, 0, 0, 0, 0, 0, 0]$.

For $\phi(W^o, 1, 0)$:

$$\begin{aligned}
p_0(x) &= x^{108} + x^{106} + x^{105} + x^{104} + x^{103} + x^{101} + x^{99} + x^{97} + x^{94} + x^{92} \\
&+ x^{85} + x^{84} + x^{80} + x^{77} + x^{76} + x^{74} + x^{73} + x^{72} + x^{69} + x^{68} + x^{66} \\
&+ x^{64} + x^{63} + x^{58} + x^{56} + x^{46} + x^{45} + x^{42} + x^{39}, \\
p_3(x) &= x^{101} + x^{99} + x^{77} + x^{75} + x^{69} + x^{67} + x^{61} + x^{59} + x^{53} + x^{51} + x^{45} \\
&+ x^{43} + x^{37} + x^{35} + x^{29} + x^{27}, \\
p_6(x) &= x^{102} + x^{100} + x^{96} + x^{92} + x^{90} + x^{86} + x^{84} + x^{82} + x^{78} + x^{76} + x^{74} \\
&+ x^{66} + x^{64} + x^{62} + x^{58} + x^{56} + x^{52} + x^{48} + x^{44} + x^{42} + x^{38} + x^{36} \\
&+ x^{34} + x^{22} + x^{20} + x^{18}, \\
p_9(x) &= x^{103} + x^{101} + x^{91} + x^{89} + x^{87} + x^{85} + x^{75} + x^{73} + x^{71} + x^{69} + x^{59} \\
&+ x^{57} + x^{55} + x^{53} + x^{43} + x^{41} + x^{23} + x^{21} + x^{11} + x^9, \\
p_{12}(x) &= x^{104} + x^{102} + x^{98} + x^{94} + x^{90} + x^{86} + x^{84} + x^{80} + x^{78} + x^{74} + x^{70} \\
&+ x^{68} + x^{64} + x^{62} + x^{58} + x^{54} + x^{52} + x^{48} + x^{46} + x^{42} + x^{40} + x^{36} \\
&+ x^{34} + x^{32} + x^{24} + x^{22} + x^{18} + x^{14} + x^{10} + x^8 + x^4 + x^2 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 1, 1, 0, 1, 1, 0, 0, 0, 0, 0, 0, 0, 0]$.

For $\phi(W^o, 1, 1)$:

$$\begin{aligned}
p_0(x) &= x^{111} + x^{109} + x^{106} + x^{105} + x^{104} + x^{103} + x^{102} + x^{100} + x^{96} + x^{85} \\
&\quad + x^{81} + x^{80} + x^{77} + x^{75} + x^{72} + x^{71} + x^{69} + x^{64} + x^{63} + x^{57} + x^{56} \\
&\quad + x^{55} + x^{54} + x^{52} + x^{51} + x^{50} + x^{49} + x^{48} + x^{45} + x^{43} + x^{42}, \\
p_3(x) &= x^{110} + x^{108} + x^{106} + x^{105} + x^{102} + x^{101} + x^{99} + x^{98} + x^{97} + x^{91} \\
&\quad + x^{90} + x^{89} + x^{88} + x^{80} + x^{79} + x^{77} + x^{76} + x^{75} + x^{73} + x^{69} + x^{68} \\
&\quad + x^{67} + x^{61} + x^{60} + x^{58} + x^{57} + x^{56} + x^{54} + x^{53} + x^{47} + x^{46} + x^{45} \\
&\quad + x^{41} + x^{39} + x^{38} + x^{37} + x^{35} + x^{34} + x^{32} + x^{31} + x^{30} + x^{29} + x^{27} \\
&\quad + x^{26}, \\
p_6(x) &= x^{110} + x^{108} + x^{106} + x^{102} + x^{101} + x^{100} + x^{99} + x^{95} + x^{92} + x^{91} \\
&\quad + x^{90} + x^{88} + x^{87} + x^{84} + x^{82} + x^{80} + x^{78} + x^{76} + x^{74} + x^{72} + x^{71} \\
&\quad + x^{70} + x^{69} + x^{67} + x^{66} + x^{65} + x^{64} + x^{63} + x^{61} + x^{59} + x^{57} + x^{56} \\
&\quad + x^{55} + x^{54} + x^{52} + x^{51} + x^{49} + x^{47} + x^{46} + x^{44} + x^{42} + x^{39} + x^{35} \\
&\quad + x^{33} + x^{26} + x^{24} + x^{23} + x^{21} + x^{19} + x^{18} + x^{17} + x^{15} + x^{13} + x^{12} \\
&\quad + x^6 + x^5 + x^4 + x^2 + x + 1, \\
p_9(x) &= x^{110} + x^{109} + x^{108} + x^{106} + x^{105} + x^{104} + x^{98} + x^{97} + x^{96} + x^{90} \\
&\quad + x^{89} + x^{88} + x^{82} + x^{81} + x^{80} + x^{74} + x^{73} + x^{72} + x^{66} + x^{65} + x^{64} \\
&\quad + x^{58} + x^{57} + x^{56} + x^{50} + x^{49} + x^{48} + x^{46} + x^{45} + x^{44} + x^{30} + x^{29} \\
&\quad + x^{28} + x^{26} + x^{25} + x^{24} + x^{18} + x^{17} + x^{16} + x^{14} + x^{13} + x^{12}, \\
p_{12}(x) &= x^{110} + x^{109} + x^{108} + x^{102} + x^{101} + x^{100} + x^{98} + x^{97} + x^{96} + x^{94} \\
&\quad + x^{93} + x^{92} + x^{78} + x^{77} + x^{76} + x^{62} + x^{61} + x^{60} + x^{38} + x^{37} + x^{36} \\
&\quad + x^{34} + x^{33} + x^{32} + x^{30} + x^{29} + x^{28} + x^{22} + x^{21} + x^{20} + x^{18} + x^{17} \\
&\quad + x^{16} + x^6 + x^5 + x^4 + x^2 + x + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 0, 0, 1, 1, 1, 0, 1, 1, 0, 0, 1]$.

Below is the transition function and output function of an 2-automaton that generates $T = M_{0,0}^o$:

Transition function $(n, j) \mapsto \delta(n, j)$ ($\Lambda(n) := [\delta(n, 0), \delta(n, 1)]$):

n	$\Lambda(n)$	n	$\Lambda(n)$	n	$\Lambda(n)$	n	$\Lambda(n)$	n	$\Lambda(n)$
0	[1, 2]	56	[1, 23]	112	[134, 147]	168	[213, 76]	224	[254, 255]
1	[3, 4]	57	[60, 8]	113	[8, 60]	169	[140, 82]	225	[256, 257]
2	[5, 6]	58	[60, 7]	114	[148, 149]	170	[214, 133]	226	[11, 205]
3	[7, 8]	59	[69, 34]	115	[73, 132]	171	[79, 147]	227	[258, 152]
4	[1, 8]	60	[38, 83]	116	[150, 151]	172	[215, 149]	228	[26, 70]
5	[9, 10]	61	[84, 13]	117	[11, 146]	173	[131, 216]	229	[259, 132]
6	[11, 12]	62	[85, 86]	118	[128, 80]	174	[124, 7]	230	[72, 35]
7	[7, 7]	63	[87, 88]	119	[140, 27]	175	[217, 63]	231	[23, 8]
8	[13, 14]	64	[89, 90]	120	[152, 133]	176	[81, 77]	232	[260, 78]
9	[15, 16]	65	[91, 84]	121	[66, 147]	177	[218, 78]	233	[261, 139]

10	[17, 18]	66	[92, 93]	122	[153, 96]	178	[219, 82]	234	[64, 82]
11	[19, 20]	67	[94, 95]	123	[154, 155]	179	[143, 210]	235	[262, 257]
12	[18, 21]	68	[96, 57]	124	[97, 156]	180	[61, 60]	236	[145, 12]
13	[22, 8]	69	[86, 97]	125	[157, 158]	181	[220, 151]	237	[263, 210]
14	[7, 23]	70	[98, 99]	126	[159, 160]	182	[145, 127]	238	[124, 8]
15	[24, 25]	71	[100, 101]	127	[161, 162]	183	[211, 80]	239	[218, 69]
16	[26, 27]	72	[102, 103]	128	[163, 40]	184	[129, 77]	240	[136, 27]
17	[28, 29]	73	[104, 105]	129	[164, 165]	185	[23, 60]	241	[264, 210]
18	[30, 31]	74	[106, 107]	130	[166, 167]	186	[22, 23]	242	[23, 7]
19	[32, 27]	75	[108, 109]	131	[168, 169]	187	[207, 130]	243	[124, 23]
20	[33, 34]	76	[110, 111]	132	[170, 171]	188	[73, 216]	244	[260, 69]
21	[29, 35]	77	[20, 112]	133	[39, 40]	189	[122, 2]	245	[26, 82]
22	[36, 37]	78	[109, 113]	134	[172, 173]	190	[221, 152]	246	[142, 147]
23	[4, 38]	79	[114, 115]	135	[14, 174]	191	[136, 77]	247	[263, 123]
24	[37, 39]	80	[116, 117]	136	[175, 176]	192	[33, 222]	248	[217, 80]
25	[40, 3]	81	[118, 119]	137	[177, 178]	193	[223, 132]	249	[64, 70]
26	[41, 42]	82	[120, 121]	138	[179, 180]	194	[29, 209]	250	[206, 152]
27	[43, 44]	83	[8, 23]	139	[181, 182]	195	[152, 34]	251	[219, 27]
28	[45, 46]	84	[1, 7]	140	[183, 184]	196	[68, 78]	252	[261, 76]
29	[47, 48]	85	[122, 123]	141	[185, 186]	197	[26, 77]	253	[81, 27]
30	[49, 50]	86	[124, 60]	142	[187, 188]	198	[224, 29]	254	[135, 69]
31	[51, 52]	87	[125, 126]	143	[189, 56]	199	[131, 74]	255	[219, 77]
32	[53, 54]	88	[11, 127]	144	[190, 191]	200	[221, 25]	256	[265, 266]
33	[55, 56]	89	[128, 63]	145	[178, 192]	201	[219, 70]	257	[267, 268]
34	[57, 58]	90	[129, 82]	146	[193, 194]	202	[148, 72]	258	[113, 185]
35	[46, 59]	91	[60, 60]	147	[105, 195]	203	[225, 226]	259	[269, 270]
36	[22, 7]	92	[28, 130]	148	[196, 197]	204	[227, 228]	260	[156, 243]
37	[1, 60]	93	[131, 132]	149	[198, 199]	205	[229, 230]	261	[271, 272]
38	[8, 7]	94	[129, 70]	150	[200, 201]	206	[56, 231]	262	[273, 274]
39	[61, 7]	95	[25, 34]	151	[202, 161]	207	[232, 19]	263	[275, 84]
40	[60, 23]	96	[7, 60]	152	[180, 96]	208	[233, 234]	264	[276, 3]
41	[62, 63]	97	[8, 8]	153	[122, 203]	209	[197, 51]	265	[258, 25]
42	[64, 27]	98	[25, 133]	154	[204, 10]	210	[235, 236]	266	[32, 77]
43	[65, 34]	99	[134, 67]	155	[145, 205]	211	[237, 238]	267	[215, 72]
44	[66, 67]	100	[135, 78]	156	[22, 60]	212	[239, 240]	268	[30, 74]
45	[68, 69]	101	[136, 70]	157	[206, 25]	213	[241, 242]	269	[75, 139]
46	[32, 70]	102	[137, 29]	158	[136, 82]	214	[243, 36]	270	[81, 70]
47	[71, 72]	103	[73, 31]	159	[207, 29]	215	[244, 245]	271	[264, 123]
48	[73, 74]	104	[138, 139]	160	[208, 74]	216	[59, 246]	272	[61, 8]
49	[75, 76]	105	[140, 77]	161	[208, 31]	217	[247, 58]	273	[24, 152]
50	[64, 77]	106	[141, 133]	162	[72, 209]	218	[186, 91]	274	[32, 82]
51	[78, 34]	107	[142, 67]	163	[122, 210]	219	[248, 249]	275	[263, 203]
52	[79, 67]	108	[143, 123]	164	[211, 63]	220	[250, 251]	276	[263, 2]
53	[62, 80]	109	[23, 23]	165	[140, 70]	221	[83, 55]		

54	[81, 82]	110	[144, 126]	166	[212, 72]	222	[174, 242]
55	[61, 23]	111	[145, 146]	167	[131, 31]	223	[252, 253]

Output function $n \mapsto \tau(n)$:

n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$
0	0	40	1	80	1	120	0	160	1	200	1	240	1
1	0	41	0	81	0	121	0	161	1	201	1	241	1
2	0	42	0	82	0	122	0	162	1	202	0	242	0
3	0	43	1	83	1	123	1	163	0	203	1	243	1
4	0	44	0	84	0	124	1	164	1	204	1	244	1
5	0	45	0	85	0	125	0	165	1	205	0	245	0
6	0	46	0	86	1	126	1	166	1	206	0	246	1
7	0	47	0	87	0	127	1	167	1	207	1	247	1
8	1	48	0	88	0	128	0	168	1	208	1	248	1
9	0	49	0	89	0	129	1	169	1	209	0	249	0
10	0	50	0	90	1	130	1	170	1	210	0	250	0
11	0	51	0	91	1	131	1	171	0	211	1	251	1
12	0	52	0	92	0	132	1	172	1	212	1	252	1
13	1	53	0	93	1	133	0	173	1	213	1	253	0
14	0	54	0	94	1	134	1	174	1	214	1	254	0
15	0	55	0	95	1	135	0	175	1	215	1	255	1
16	0	56	0	96	0	136	1	176	0	216	1	256	1
17	0	57	1	97	1	137	1	177	1	217	1	257	1
18	0	58	1	98	1	138	0	178	1	218	1	258	1
19	0	59	1	99	1	139	0	179	0	219	1	259	0
20	0	60	1	100	0	140	1	180	0	220	0	260	1
21	0	61	0	101	1	141	0	181	0	221	1	261	1
22	1	62	0	102	1	142	1	182	1	222	1	262	0
23	0	63	0	103	0	143	0	183	1	223	1	263	1
24	0	64	0	104	0	144	1	184	1	224	0	264	1
25	1	65	1	105	1	145	1	185	0	225	1	265	1
26	0	66	0	106	0	146	1	186	1	226	0	266	0
27	1	67	1	107	1	147	1	187	1	227	1	267	1
28	0	68	0	108	0	148	0	188	0	228	0	268	0
29	0	69	1	109	0	149	0	189	0	229	0	269	0
30	0	70	1	110	1	150	1	190	1	230	1	270	0
31	0	71	0	111	1	151	0	191	1	231	0	271	1
32	0	72	1	112	1	152	0	192	0	232	1	272	0
33	0	73	0	113	1	153	0	193	1	233	1	273	0
34	1	74	0	114	0	154	1	194	0	234	0	274	0
35	0	75	0	115	0	155	1	195	0	235	0	275	1
36	1	76	1	116	1	156	1	196	0	236	1	276	1
37	0	77	0	117	0	157	0	197	0	237	1		
38	1	78	0	118	0	158	1	198	0	238	1		

$$\left| \begin{array}{cc} 39 & 0 \end{array} \right| \left| \begin{array}{cc} 79 & 0 \end{array} \right| \left| \begin{array}{cc} 119 & 1 \end{array} \right| \left| \begin{array}{cc} 159 & 1 \end{array} \right| \left| \begin{array}{cc} 199 & 1 \end{array} \right| \left| \begin{array}{cc} 239 & 1 \end{array} \right| \left| \begin{array}{cc} & \end{array} \right|$$

UNIVERSITÉ DE STRASBOURG, CNRS, IRMA UMR 7501, F-67000 STRASBOURG, FRANCE
Email address: **guoniu.han@unistra.fr**

SCHOOL OF MATHEMATICS AND STATISTICS, HUAZHONG UNIVERSITY OF SCIENCE AND TECHNOLOGY, WUHAN, PR CHINA
Email address: **huyining@hust.edu.cn**