

PROOF OF THE CONTINUED FRACTION CONJECTURE FOR

$$(a, b) = (z^2 + 1, z^5 + z^3 + z^2 + 1)$$

GUO-NIU HAN* AND YINING HU*

ABSTRACT. The continued fraction conjecture claims that the continued fraction defined by the (a, b) -Thue-Morse sequence is algebraic over $\mathbb{F}_2(x)$ for all pairs of distinct elements (a, b) in $\mathbb{F}_2[x] \setminus \mathbb{F}_2$. In this paper we prove the conjecture for $(a, b) = (z^2 + 1, z^5 + z^3 + z^2 + 1)$.

1. INTRODUCTION

Let $\mathbb{F}_2$ be the finite field of cardinality 2. Given a sequence of polynomials $\mathbf{a} = (a_0(z), a_1(z), a_2(z), \dots)$ of elements from $\mathbb{F}_2[z] \setminus \mathbb{F}_2$, we define the infinite continued fraction

$$(1.1) \quad \text{CF}(\mathbf{a}(z)) = \frac{1}{a_0(z) + \frac{1}{a_1(z) + \frac{1}{a_2(z) + \frac{1}{\ddots}}}}$$

as the limit of the finite continued fractions

$$(1.2) \quad \text{CF}_n(\mathbf{a}(z)) = \frac{1}{a_0(z) + \frac{1}{a_1(z) + \frac{1}{\ddots + \frac{1}{a_n(z)}}}} \in \mathbb{F}_2((1/z)).$$

Conversely, each power series in $\mathbb{F}_2[[1/z]]$ without constant term

$$(1.3) \quad c_{-1}z^{-1} + c_{-2}z^{-2} + c_{-3}z^{-3} + \dots$$

admits a continued fraction expansion of form (1.1).

Let a and b be two distinct elements from $\mathbb{F}_2[z] \setminus \mathbb{F}_2$. We consider the continued fraction (1.1) associated with the (a, b) -Thue-Morse sequence

$$\mathbf{t} = (t_0(z), t_1(z), t_2(z), \dots) = (a, b, b, a, b, a, a, b, \dots)$$

that we shall denote

$$(1.4) \quad \text{CF}(a, b) := \text{CF}(\mathbf{t}).$$

Date: runone at : 30/04/2020 03:11:58 .

2010 Mathematics Subject Classification. 11B85, 11J70, 11B50, 11Y65, 05A15.

Key words and phrases. automatic sequence, continued fraction, Thue-Morse sequence.

* This paper was automatically generated from a computer program developed by the authors, with 12.0 minutes CPU time used.

Theorem 1.1. *When $(a, b) = (z^2 + 1, z^5 + z^3 + z^2 + 1)$, the two power series $\text{CF}(a, b)$ and $\text{CF}(b, a)$ are algebraic over $\mathbb{F}_2(z)$, with minimal polynomials of the form*

$$p_4(z)y^4 + p_3(z)y^3 + p_2(z)y^2 + p_1(z)y + p_0(z) = 0$$

where, for $\text{CF}(a, b)$,

$$\begin{aligned} p_0(z) &= z^{13} + z^{12} + z^{10} + z^7 + z^6 + z^4 + z^3 + 1, \\ p_1(z) &= z^{18} + z^{16} + z^{15} + z^{13} + z^{12} + z^{10} + z^9 + z^8 + z^7 + z^6 + z^5 + z^3, \\ p_2(z) &= z^{20} + z^{16} + z^{10} + z^8 + z^6 + 1, \\ p_3(z) &= z^{18} + z^{16} + z^{15} + z^{13} + z^{12} + z^{10} + z^9 + z^8 + z^7 + z^6 + z^5 + z^3, \\ p_4(z) &= z^{16} + z^{13} + z^{12} + z^7 + z^4 + z^3 + 1, \end{aligned}$$

and for $\text{CF}(b, a)$,

$$\begin{aligned} p_0(z) &= z^{13} + z^{10} + z^7 + z^6 + z^4 + z^3 + 1, \\ p_1(z) &= z^{18} + z^{16} + z^{15} + z^{13} + z^{12} + z^{10} + z^9 + z^8 + z^7 + z^6 + z^5 + z^3, \\ p_2(z) &= z^{20} + z^{16} + z^{10} + z^8 + z^6 + 1, \\ p_3(z) &= z^{18} + z^{16} + z^{15} + z^{13} + z^{12} + z^{10} + z^9 + z^8 + z^7 + z^6 + z^5 + z^3, \\ p_4(z) &= z^{16} + z^{13} + z^7 + z^4 + z^3 + 1. \end{aligned}$$

2. PROOF

Define the sequences of polynomials $(P_n(z))_{n \geq 0}$ and $(Q_n(z))_{n \geq 0}$ by

$$\begin{pmatrix} P_n(z) & Q_n(z) \\ P_{n-1}(z) & Q_{n-1}(z) \end{pmatrix} := \begin{pmatrix} t_n(z) & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} t_{n-1}(z) & 1 \\ 1 & 0 \end{pmatrix} \cdots \begin{pmatrix} t_0(z) & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}.$$

By the basic properties of continued fractions, we have

$$(2.1) \quad \text{CF}_n(a, b) = \frac{P_n(z)}{Q_n(z)} \in \mathbb{F}_2((1/z)).$$

As formula (2.1) is not efficient for calculating $\text{CF}_n(a, b)$, a fast method is to be derived and described as follows. Define

$$\begin{aligned} M_n(x) &= x^{\deg(t_{2^n-1})} \begin{pmatrix} t_{2^n-1}(1/x) & 1 \\ 1 & 0 \end{pmatrix} \times x^{\deg(t_{2^n-2})} \begin{pmatrix} t_{2^n-2}(1/x) & 1 \\ 1 & 0 \end{pmatrix} \times \\ &\quad \cdots \times x^{\deg(t_0)} \begin{pmatrix} t_0(1/x) & 1 \\ 1 & 0 \end{pmatrix}, \\ W_n(x) &= x^{\deg(\bar{t}_{2^n-1})} \begin{pmatrix} \bar{t}_{2^n-1}(1/x) & 1 \\ 1 & 0 \end{pmatrix} \times x^{\deg(\bar{t}_{2^n-2})} \begin{pmatrix} \bar{t}_{2^n-2}(1/x) & 1 \\ 1 & 0 \end{pmatrix} \times \\ &\quad \cdots \times x^{\deg(\bar{t}_0)} \begin{pmatrix} \bar{t}_0(1/x) & 1 \\ 1 & 0 \end{pmatrix}, \end{aligned}$$

where $\bar{t}$ is the (b, a) -Thue-Morse sequence. By the properties of the Thue-Morse sequence, we have for $n \geq 0$

$$\begin{aligned} M_{n+1}(x) &= W_n(x) \cdot M_n(x), \\ W_{n+1}(x) &= M_n(x) \cdot W_n(x). \end{aligned}$$

Let $x := 1/z$. For each non-zero polynomial $P(z)$, define $\tilde{P}(x)$ to be $P(1/x)$. Then,

$$\text{CF}_n(a, b) = \frac{P_n(z)}{Q_n(z)} = \frac{\tilde{P}_n(x)}{\tilde{Q}_n(x)} \in \mathbb{F}_2((x)) = \mathbb{F}_2((1/z)).$$

Comparing the definition of $M_n(x)$ with the definition of $P_n(x)$ and $Q_n(x)$, we see that

$$\begin{aligned} M_n(x)_{0,1} &= x^{d_n} \tilde{P}_{2^n-1}(x), \\ M_n(x)_{0,0} &= x^{d_n} \tilde{Q}_{2^n-1}(x), \end{aligned}$$

for some positive integer d_n . Hence,

$$(2.2) \quad \text{CF}_{2^{2^n}-1}(a, b) = \frac{\tilde{P}_{2^{2^n}-1}(x)}{\tilde{Q}_{2^{2^n}-1}(x)} = \frac{M_{2^n}(x)_{0,1}}{M_{2^n}(x)_{0,0}}.$$

Therefore, by the convergence theorem of continued fraction, the algebraicity of $\text{CF}(a, b)$ will be established if it is shown that both $M_{2^n}(x)_{0,1}$ and $M_{2^n}(x)_{0,0}$ converge to algebraic series in $\mathbb{F}_2[[x]]$.

Actually, we will prove that for all $0 \leq i, j \leq 1$ the four sequences $(M_{2^n}(x)_{i,j})_n$, $(M_{2^{n+1}}(x)_{i,j})_n$, $(W_{2^n}(x)_{i,j})_n$, and $(W_{2^{n+1}}(x)_{i,j})_n$ converge to algebraic series in $\mathbb{F}_2[[x]]$. For this purpose, we define four 2×2 matrices M^e, M^o, W^e, W^o as follows: For each $T \in \{M^e, M^o, W^e, W^o\}$ and $0 \leq i, j \leq 1$, $T_{i,j}$ is defined to be the unique solution in $\mathbb{F}_2[[x]]$ of the polynomial $\phi(T, i, j)$ under certain initial conditions; the polynomials $\phi(T, i, j)$ and initial conditions are given in Section 3. We will prove that these four matrices, whose components are algebraic by definition, are the limits of $(M_{2^n}(x))_n$, $(M_{2^{n+1}}(x))_n$, $(W_{2^n}(x))_n$, and $(W_{2^{n+1}}(x))_n$.

Let us explain how the polynomials $\phi(T, i, j)$ and initial conditions are found, and why the solutions exist and are unique. For $0 \leq i, j \leq 1$, the coefficients of the polynomial $\phi(M^e, i, j)$ (resp. $\phi(M^o, i, j)$, $\phi(W^e, i, j)$, and $\phi(W^o, i, j)$) are the Padé-Hermite approximants of type

$$(175, 175, 175, 175, 175)$$

of the vector

$$(1, T^3, T^6, T^9, T^{12}),$$

where $T = M_{12,i,j}$ (resp. $M_{11,i,j}$, $W_{12,i,j}$, and $W_{11,i,j}$), found by the Derksen algorithm. We take the first 8 terms of $M_{12,i,j}$ (resp. $M_{11,i,j}$, $W_{12,i,j}$, and $W_{11,i,j}$) as initial conditions for $\phi(M^e, i, j)$ (resp. $\phi(M^o, i, j)$, $\phi(W^e, i, j)$, and $\phi(W^o, i, j)$). The following fact will be used to ensure that the solution exists and is unique: let $P(x, y) \in \mathbb{F}_2[x, y]$ and for each series $f(x) = \sum_{n=0}^{\infty} a_n x^n \in \mathbb{F}_2[[x]]$ denote the partial sum $\sum_{j=0}^{n-1} a_j x^j$ by $f_n(x)$ for $n \geq 0$. If for some $n \geq 0$ and $a_0, a_1, \dots, a_{n-1} \in \mathbb{F}_2$ $P(x, \sum_{j=0}^{n-1} a_j x^j) = O(x^n)$ and $Q(x, y) := P(x, \sum_{j=0}^{n-1} a_j x^j + x^n y)$ can be written as $x^m \sum_{j=0}^{\infty} q_j(x) y^j$ where $q_j(x)$ are polynomials for $j \geq 0$, $q_1(0) = 1$, and $q_j(0) = 0$ for $j > 1$, then there exists a unique solution $f(x) \in \mathbb{F}_2[[x]]$ of $P(x, f(x)) = 0$ that satisfies the initial condition $f_n(x) = \sum_{j=0}^{n-1} a_j x^j$.

We state two lemmas concerning the four matrices M^e, M^o, W^e, W^o . The first one is about relations between them; the second, about the structure of the each matrix.

Lemma 2.1. *We have*

$$(2.3) \quad M^e = W^o \cdot M^o,$$

$$(2.4) \quad M^o = W^e \cdot M^e,$$

$$(2.5) \quad W^e = M^o \cdot W^o,$$

$$(2.6) \quad W^o = M^e \cdot W^e.$$

Proof. We give the proof of the identity

$$M_{0,0}^e = W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o,$$

the proofs of the others being similar.

First, we compute the minimal polynomials of $W_{0,0}^o M_{0,0}^o$ and $W_{0,1}^o M_{1,0}^o$. We know that

$$P(x, y) = \text{Res}_z (\phi(W^o, 0, 0)(x, z), z^{12} \cdot \phi(M^o, 0, 0)(x, y/z))$$

is an annihilating polynomial of $W_{0,0}^o M_{0,0}^o$. We use Padé-Hermite approximation to find a candidate for the minimal polynomial of $W_{0,0}^o M_{0,0}^o$, that will be called $\phi_0(x, y)$. To prove that $\phi_0(x, y)$ is indeed the minimal polynomial, it suffices to prove that it is an irreducible factor of $P(x, y)$ of multiplicity m and that $Q(x, y) := P(x, y)/\phi_0(x, y)^m$ is not an annihilating polynomial of $W_{0,0}^o M_{0,0}^o$. We verify the first point directly. For the second point, we truncate $W_{0,0}^o M_{0,0}^o$ to order 630 and substitute it for y in $Q(x, y)$. We get a series of valuation less than 630, which proves that $Q(x, y)$ is not an annihilating polynomial of $W_{0,0}^o M_{0,0}^o$. We find the minimal polynomial $\phi_1(x, y)$ of $W_{0,1}^o M_{1,0}^o$ in a similar way.

Now we prove that $\phi(M^e, 0, 0)$ is the minimal polynomial of $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$. We know that

$$S(x, y) = \text{Res}_z (\phi_0(x, z), \phi_1(x, y + z))$$

is an annihilating of $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$. We verify that $\phi(M^e, 0, 0)$ is an irreducible factor of $S(x, y)$ of multiplicity μ , and that $Q(x, y) := S(x, y)/\phi(M^e, 0, 0)^\mu$ is not an annihilating polynomial of $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$. To see the last point, we truncate $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$ to order 770 and substitute it for y in $Q(x, y)$. We get a series of valuation less than 770, and therefore $Q(x, y)$ is not an annihilating polynomial of $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$.

Finally, the first 16 terms of $M_{0,0}^e$ and $W_{0,0}^o M_{0,0}^o + W_{0,1}^o M_{1,0}^o$ coincide. As these first terms determine a unique solution of $\phi(M^e, 0, 0)$, we know that the two series are one and the same. $\square$

Define

$$R^e = \begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix} \quad \text{and} \quad R^o = \begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix}.$$

Lemma 2.2. *For all integers $k \geq 2$ and $u = 2^{2k-1}$, the following identities hold.*

$$\begin{aligned} M^e[14u] &= M^e[7u] + x^{3u} M^e[4u] + x^{3u} M^e[7u] + x^{6u} M^e[4u] + x^{6u} M^e[7u] \\ &\quad + x^{9u} M^e[4u] + x^{8u} M^e[6u] + x^{9u} M^e[5u] + x^{10u} M^e[4u] \\ &\quad + (x^{7u} + x^{10u} + x^{13u}) R^e, \\ W^e[14u] &= W^e[7u] + x^{3u} W^e[4u] + x^{3u} W^e[7u] + x^{6u} W^e[4u] + x^{6u} W^e[7u] \\ &\quad + x^{9u} W^e[4u] + x^{8u} W^e[6u] + x^{9u} W^e[5u] + x^{10u} W^e[4u] \\ &\quad + (x^{7u} + x^{10u} + x^{13u}) R^e. \end{aligned}$$

For all integers $k \geq 2$ and $u = 2^{2k}$, the following identities hold.

$$\begin{aligned}
M^o[:14u] &= M^o[:7u] + x^{3u}M^o[:4u] + x^{3u}M^o[:7u] + x^{6u}M^o[:4u] + x^{6u}M^o[:7u] \\
&\quad + x^{9u}M^o[:4u] + x^{8u}M^o[:6u] + x^{9u}M^o[:5u] + x^{10u}M^o[:4u] \\
&\quad + (x^{7u} + x^{10u} + x^{13u})R^o, \\
W^o[:14u] &= W^o[:7u] + x^{3u}W^o[:4u] + x^{3u}W^o[:7u] + x^{6u}W^o[:4u] + x^{6u}W^o[:7u] \\
&\quad + x^{9u}W^o[:4u] + x^{8u}W^o[:6u] + x^{9u}W^o[:5u] + x^{10u}W^o[:4u] \\
&\quad + (x^{7u} + x^{10u} + x^{13u})R^o.
\end{aligned}$$

Proof. To prove Lemma 2.2 we first construct an automaton for each sequence concerned, and then transform the conditions on infinitely many k 's into finitely many conditions on the states of the automaton. In the following, we will prove that for $T = M_{0,0}^o$, for all integer $k \geq 2$ and $u = 2^{2k}$,

$$\begin{aligned}
T[:14u] &= T[:7u] + x^{3u}T[:4u] + x^{3u}T[:7u] + x^{6u}T[:4u] + x^{6u}T[:7u] + x^{9u}T[:4u] \\
&\quad + x^{8u}T[:6u] + x^{9u}T[:5u] + x^{10u}T[:4u] + (x^{7u} + x^{10u} + x^{13u}).
\end{aligned}$$

The proof of the other 15 cases are similar. We break down the above identity into 7 parts:

$$\begin{aligned}
x^7u &= x^{3u}T[4u:5u] + T[7u:8u], \\
0 &= x^{8u}T[:u] + x^{3u}T[5u:6u] + T[8u:9u], \\
0 &= x^{8u}T[u:2u] + x^{3u}T[6u:7u] + T[9u:10u], \\
x^{10u} &= x^{10u}T[:u] + x^{8u}T[2u:3u] + x^{6u}T[4u:5u] + T[10u:11u], \\
0 &= x^{10u}T[u:2u] + x^{8u}T[3u:4u] + x^{6u}T[5u:6u] + T[11u:12u], \\
0 &= x^{10u}T[2u:3u] + x^{8u}T[4u:5u] + x^{6u}T[6u:7u] + T[12u:13u], \\
x^{13u} &= x^{10u}T[3u:4u] + x^{9u}T[4u:5u] + x^{8u}T[5u:6u] + T[13u:14u],
\end{aligned}$$

which can be rewritten as

$$(2.7) \quad 0 = T[[100w]_2] + T[[111w]_2],$$

$$(2.8) \quad 0 = T[[w]_2] + T[[101w]_2] + T[[1000w]_2],$$

$$(2.9) \quad 0 = T[[1w]_2] + T[[110w]_2] + T[[1001w]_2],$$

$$(2.10) \quad 0 = T[[w]_2] + T[[10w]_2] + T[[100w]_2] + T[[1010w]_2],$$

$$(2.11) \quad 0 = T[[1w]_2] + T[[11w]_2] + T[[101w]_2] + T[[1011w]_2],$$

$$(2.12) \quad 0 = T[[10w]_2] + T[[100w]_2] + T[[110w]_2] + T[[1100w]_2],$$

$$(2.13) \quad 0 = T[[11w]_2] + T[[100w]_2] + T[[101w]_2] + T[[1101w]_2],$$

for all binary word w of length $2k$ and $w \neq 0^{2k}$, and

$$(2.14) \quad 1 = T[[100w]_2] + T[[111w]_2],$$

$$(2.15) \quad 0 = T[[w]_2] + T[[101w]_2] + T[[1000w]_2],$$

$$(2.16) \quad 0 = T[[1w]_2] + T[[110w]_2] + T[[1001w]_2],$$

$$(2.17) \quad 1 = T[[w]_2] + T[[10w]_2] + T[[100w]_2] + T[[1010w]_2],$$

$$(2.18) \quad 0 = T[[1w]_2] + T[[11w]_2] + T[[101w]_2] + T[[1011w]_2],$$

$$(2.19) \quad 0 = T[[10w]_2] + T[[100w]_2] + T[[110w]_2] + T[[1100w]_2],$$

$$(2.20) \quad 1 = T[[11w]_2] + T[[100w]_2] + T[[101w]_2] + T[[1101w]_2],$$

for $w = 0^{2k}$.

First we calculate an 2-automaton that generates T from its minimal polynomial and its first terms. This automaton has 506 states; its transition function and output function can be found in the annex. Let $A(s, w)$ denote the state reached after reading w from right to left starting from the state s , and τ the output function. Define

$$E_{2k} = \{A(i, w) : |w| = 2k, w \neq 0^{2k}\}.$$

Identities (2.7) through (2.20) can be written as

$$(2.21) \quad 0 = \tau(A(s, 100)) + \tau(A(s, 111)),$$

$$(2.22) \quad 0 = \tau(A(s, \epsilon)) + \tau(A(s, 101)) + \tau(A(s, 1000)),$$

$$(2.23) \quad 0 = \tau(A(s, 1)) + \tau(A(s, 110)) + \tau(A(s, 1001)),$$

$$(2.24) \quad 0 = \tau(A(s, \epsilon)) + \tau(A(s, 10)) + \tau(A(s, 100)) + \tau(A(s, 1010)),$$

$$(2.25) \quad 0 = \tau(A(s, 1)) + \tau(A(s, 11)) + \tau(A(s, 101)) + \tau(A(s, 1011)),$$

$$(2.26) \quad 0 = \tau(A(s, 10)) + \tau(A(s, 100)) + \tau(A(s, 110)) + \tau(A(s, 1100)),$$

$$(2.27) \quad 0 = \tau(A(s, 11)) + \tau(A(s, 100)) + \tau(A(s, 101)) + \tau(A(s, 1101)),$$

for all $s \in E_{2k}$, and

$$(2.28) \quad 1 = \tau(A(s, 100)) + \tau(A(s, 111)),$$

$$(2.29) \quad 0 = \tau(A(s, \epsilon)) + \tau(A(s, 101)) + \tau(A(s, 1000)),$$

$$(2.30) \quad 0 = \tau(A(s, 1)) + \tau(A(s, 110)) + \tau(A(s, 1001)),$$

$$(2.31) \quad 1 = \tau(A(s, \epsilon)) + \tau(A(s, 10)) + \tau(A(s, 100)) + \tau(A(s, 1010)),$$

$$(2.32) \quad 0 = \tau(A(s, 1)) + \tau(A(s, 11)) + \tau(A(s, 101)) + \tau(A(s, 1011)),$$

$$(2.33) \quad 0 = \tau(A(s, 10)) + \tau(A(s, 100)) + \tau(A(s, 110)) + \tau(A(s, 1100)),$$

$$(2.34) \quad 1 = \tau(A(s, 11)) + \tau(A(s, 100)) + \tau(A(s, 101)) + \tau(A(s, 1101)),$$

for $s = A(i, 0^{2k})$. We find that $(A(i, 0^{24}), E_{24}) = (A(i, 0^{20}), E_{20})$, so that we only have to verify that identities (2.21) through (2.34) hold for $2 \leq k \leq 12$, which turns out to be true. $\square$

In the following lemma, we express M_{2k} , M_{2k+1} , W_{2k} , and W_{2k+1} in terms of M^e , M^o , W^e , and W^o .

Lemma 2.3. *For all integer $k \geq 2$, and $u = 2^{2k-1}$,*

$$M_{2k} = M^e[:7u] + x^{3u}M^e[:4u] + x^{7u}R^e,$$

$$W_{2k} = W^e[:7u] + x^{3u}W^e[:4u] + x^{7u}R^e.$$

For all integer $k \geq 2$, and $u = 2^{2k}$,

$$M_{2k+1} = M^o[:7u] + x^{3u}M^o[:4u] + x^{7u}R^o,$$

$$W_{2k+1} = W^o[:7u] + x^{3u}W^o[:4u] + x^{7u}R^o.$$

Proof. We call the four identities in Lemma 2.3 also by the name M_{2k} , W_{2k} , M_{2k+1} , and W_{2k+1} . For $n = 2$, the identities can be verified directly. For $n \geq 2$, we claim that

$$M_{2k} \wedge W_{2k} \Rightarrow M_{2k+1} \wedge W_{2k+1},$$

$$M_{2k+1} \wedge W_{2k+1} \Rightarrow M_{2k+2} \wedge W_{2k+2}.$$

We give the proof of

$$M_{2k} \wedge W_{2k} \Rightarrow M_{2k+1},$$

the other ones being similar. Set $u = 2^{2k}$ and $v = 2^{2k-1}$. By definition and induction hypothesis, the left side of identity M_{2k+1} is equal to

$$(2.35) \quad \begin{aligned} W_{2k} M_{2k} &= (W^e[:7v] + x^{3v} W^e[:4v] + x^{7u} R^e) \times (M^e[:7v] + x^{3v} M^e[:4v] \\ &\quad + x^{7u} R^e); \end{aligned}$$

Call this expression *lhs*. Note that both sides of identity M_{2k+1} have the same term of highest degree $x^{14v} R^o$. Therefore we only need to prove that their difference is $O(x^{14v})$. Using Lemma 2.1 it can be seen that the right side of identity M_{2k+1} is congruent, modulo x^{14v} , to

$$W^e[:14v] M^e[:14v] + x^{6v} W^e[:8v] M^e[:8v].$$

For all $n \leq 14$, replace the occurrences of $W^e[:n \cdot v]$ and $M^e[:n \cdot v]$ in the above expression by the reduction modulo $x^{n \cdot v}$ of the right side of the corresponding identity in Lemma 2.2 and get a new expression, which we call *rhs*. Define

$$\begin{aligned} X &:= x^v, \\ a_n &:= W^e[n \cdot v : (n+1) \cdot v] / X^n, \\ b_n &:= M^e[n \cdot v : (n+1) \cdot v] / X^n, \\ c &:= R^e. \end{aligned}$$

Using the notation introduced above, we can represent the expressions *lhs* (2.35) and *rhs* as polynomials in $\mathbb{F}_2[a_1, \dots, a_{14}, b_1, \dots, b_{14}, c][X]$. Note that it is not a problem that a_j commutes with b_k while W^e does not commute with M^e , because in the expressions concerned, the products of W^e -terms and M^e -terms are always in the same order. We let the computer do the simplification and check that the difference between these two polynomials is indeed $O(X^{14})$, which completes the proof. $\square$

Proof of Theorem 1.1. We prove the theorem for $\text{CF}(a, b)$; for $\text{CF}(b, a)$, the proof is similar. By Lemma 2.3, we have For all $0 \leq j, k \leq 1$,

$$\begin{aligned} \lim_{n \rightarrow \infty} M_{2n,j,k} &= M_{j,k}^e, \\ \lim_{n \rightarrow \infty} M_{2n+1,j,k} &= M_{j,k}^o, \\ \lim_{n \rightarrow \infty} W_{2n,j,k} &= W_{j,k}^e, \\ \lim_{n \rightarrow \infty} W_{2n+1,j,k} &= W_{j,k}^o. \end{aligned}$$

Let $z = 1/x$. By the convergence theorem and identity (2.2),

$$\text{CF}(a, b) = \frac{M_{0,1}^e(x)}{M_{0,0}^e(x)}.$$

By definition, $\phi(M^e, 0, 1)$ and $\phi(M^e, 0, 0)$ are minimal polynomials of $M_{0,1}^e$ and $M_{0,0}^e$. Therefore

$$P(x, y) = \text{Res}_t (\phi(M^e, 0, 1)(x, t), y^{12} \phi(M^e, 0, 1)(x, t/y))$$

is an annihilating polynomial of $f(x) = M_{0,1}^e / M_{0,0}^e$.

Define

$$Q(x, y) = q_4(x)y^4 + q_3(x)y^3 + q_2(x)y^2 + q_1(x)y + q_0(x),$$

where

$$\begin{aligned} q_0(x) &= x^{20} + x^{17} + x^{16} + x^{14} + x^{13} + x^{10} + x^8 + x^7, \\ q_1(x) &= x^{17} + x^{15} + x^{14} + x^{13} + x^{12} + x^{11} + x^{10} + x^8 + x^7 + x^5 + x^4 + x^2, \\ q_2(x) &= x^{20} + x^{14} + x^{12} + x^{10} + x^4 + 1, \\ q_3(x) &= x^{17} + x^{15} + x^{14} + x^{13} + x^{12} + x^{11} + x^{10} + x^8 + x^7 + x^5 + x^4 + x^2, \\ q_4(x) &= x^{20} + x^{17} + x^{16} + x^{13} + x^8 + x^7 + x^4. \end{aligned}$$

The polynomial $Q(x, y)$ is the candidate for the minimal polynomial of $f(x)$ found by Padé-Hermite approximation. To prove that it is indeed the minimal polynomial of $f(x)$, we only need to prove that it is an irreducible factor of $P(x, y)$ of multiplicity m and $R(x, y) := P(x, y)/Q(x, y)^m$ is not an annihilating polynomial of $f(x)$. We verify the first point directly. For the second point, we find that when we truncate $f(x)$ to order 216, and substitute it for y in $R(z, y)$, we get a series with valuation smaller than 216, which proves that $R(z, y)$ is not an annihilating polynomial of $f(x)$. Finally, $z^{12}Q(1/z, y)$ is the minimal polynomial of $\text{CF}(a, b) = f(1/z)$. $\square$

3. ANNEXE

All 16 polynomials are of the form

$$p_0(x) + p_3(x)y^3 + p_6(x)y^6 + p_9(x)y^9 + p_{12}(x)y^{12}.$$

The coefficients $p_j(x)$ and the 16 initial terms to determine the solutions uniquely are given below.

For $\phi(M^e, 0, 0)$:

$$\begin{aligned} p_0(x) &= x^{120} + x^{114} + x^{112} + x^{108} + x^{106} + x^{104} + x^{102} + x^{100} + x^{98} + x^{92} \\ &\quad + x^{90} + x^{88} + x^{86} + x^{82} + x^{80} + x^{76} + x^{64} + x^{62} + x^{58} + x^{56} + x^{54} \\ &\quad + x^{50} + x^{46} + x^{44} + x^{40} + x^{36} + x^{32} + x^{30} + x^{24}, \\ p_3(x) &= x^{114} + x^{110} + x^{102} + x^{98} + x^{92} + x^{88} + x^{72} + x^{66} + x^{62} + x^{60} + x^{56} \\ &\quad + x^{54} + x^{52} + x^{50} + x^{48} + x^{42} + x^{40} + x^{38} + x^{36} + x^{34} + x^{26} + x^{22} \\ &\quad + x^{18} + x^{16}, \\ p_6(x) &= x^{114} + x^{112} + x^{108} + x^{102} + x^{100} + x^{90} + x^{86} + x^{78} + x^{74} + x^{70} + x^{68} \\ &\quad + x^{66} + x^{62} + x^{54} + x^{52} + x^{48} + x^{46} + x^{42} + x^{38} + x^{32} + x^{30} + x^{22} \\ &\quad + x^{12} + x^{10} + x^4 + 1, \\ p_9(x) &= x^{114} + x^{110} + x^{106} + x^{104} + x^{102} + x^{96} + x^{94} + x^{86} + x^{84} + x^{80} + x^{70} \\ &\quad + x^{60} + x^{58} + x^{56} + x^{54} + x^{52} + x^{46} + x^{44} + x^{42} + x^{40} + x^{38} + x^{34} \\ &\quad + x^{32} + x^{24} + x^{22} + x^{20} + x^{16} + x^{14} + x^{10} + x^4, \\ p_{12}(x) &= x^{120} + x^{104} + x^{96} + x^{88} + x^{80} + x^{72} + x^{64} + x^{48} + x^{40} + x^{24} + x^{16} + 1 \end{aligned}$$

and the initial terms are $[1, 0, 0, 0, 0, 0, 0, 0, 1, 0, 1, 0, 0, 0, 0, 0]$.

For $\phi(M^e, 0, 1)$:

$$\begin{aligned}
p_0(x) &= x^{111} + x^{108} + x^{103} + x^{102} + x^{100} + x^{96} + x^{94} + x^{91} + x^{90} + x^{86} + x^{85} \\
&\quad + x^{82} + x^{81} + x^{80} + x^{77} + x^{76} + x^{73} + x^{72} + x^{71} + x^{70} + x^{69} + x^{68} \\
&\quad + x^{67} + x^{65} + x^{63} + x^{62} + x^{60} + x^{55} + x^{52} + x^{50} + x^{49} + x^{48} + x^{47} \\
&\quad + x^{45} + x^{44} + x^{42} + x^{41} + x^{40} + x^{39} + x^{37} + x^{35} + x^{34} + x^{33}, \\
p_3(x) &= x^{84} + x^{82} + x^{80} + x^{74} + x^{72} + x^{68} + x^{64} + x^{60} + x^{56} + x^{50} + x^{40} \\
&\quad + x^{34} + x^{32} + x^{26} + x^{20} + x^{18}, \\
p_6(x) &= x^{93} + x^{90} + x^{89} + x^{87} + x^{86} + x^{84} + x^{83} + x^{80} + x^{73} + x^{70} + x^{69} \\
&\quad + x^{67} + x^{66} + x^{65} + x^{64} + x^{63} + x^{62} + x^{60} + x^{59} + x^{56} + x^{45} + x^{42} \\
&\quad + x^{39} + x^{37} + x^{36} + x^{34} + x^{31} + x^{29} + x^{28} + x^{26} + x^{25} + x^{23} + x^{22} \\
&\quad + x^{21} + x^{20} + x^{19} + x^{18} + x^{16} + x^{15} + x^{12}, \\
p_9(x) &= x^{102} + x^{100} + x^{94} + x^{86} + x^{80} + x^{78} + x^{76} + x^{74} + x^{72} + x^{70} + x^{68} \\
&\quad + x^{66} + x^{64} + x^{62} + x^{60} + x^{48} + x^{46} + x^{42} + x^{36} + x^{26} + x^{20} + x^{18} \\
&\quad + x^{16} + x^{10} + x^8 + x^6, \\
p_{12}(x) &= x^{111} + x^{108} + x^{99} + x^{96} + x^{95} + x^{92} + x^{83} + x^{80} + x^{79} + x^{76} + x^{67} \\
&\quad + x^{64} + x^{63} + x^{60} + x^{51} + x^{48} + x^{31} + x^{28} + x^{19} + x^{16} + x^{15} + x^{12} \\
&\quad + x^3 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 1, 0, 1, 0, 1, 0, 1, 1, 0, 1, 0, 0, 0, 1]$.

For $\phi(M^e, 1, 0)$:

$$\begin{aligned}
p_0(x) &= x^{111} + x^{108} + x^{103} + x^{102} + x^{100} + x^{96} + x^{94} + x^{91} + x^{90} + x^{86} + x^{85} \\
&\quad + x^{82} + x^{81} + x^{80} + x^{77} + x^{76} + x^{73} + x^{72} + x^{71} + x^{70} + x^{69} + x^{68} \\
&\quad + x^{67} + x^{65} + x^{63} + x^{62} + x^{60} + x^{55} + x^{52} + x^{50} + x^{49} + x^{48} + x^{47} \\
&\quad + x^{45} + x^{44} + x^{42} + x^{41} + x^{40} + x^{39} + x^{37} + x^{35} + x^{34} + x^{33}, \\
p_3(x) &= x^{84} + x^{82} + x^{80} + x^{74} + x^{72} + x^{68} + x^{64} + x^{60} + x^{56} + x^{50} + x^{40} \\
&\quad + x^{34} + x^{32} + x^{26} + x^{20} + x^{18}, \\
p_6(x) &= x^{93} + x^{90} + x^{89} + x^{87} + x^{86} + x^{84} + x^{83} + x^{80} + x^{73} + x^{70} + x^{69} \\
&\quad + x^{67} + x^{66} + x^{65} + x^{64} + x^{63} + x^{62} + x^{60} + x^{59} + x^{56} + x^{45} + x^{42} \\
&\quad + x^{39} + x^{37} + x^{36} + x^{34} + x^{31} + x^{29} + x^{28} + x^{26} + x^{25} + x^{23} + x^{22} \\
&\quad + x^{21} + x^{20} + x^{19} + x^{18} + x^{16} + x^{15} + x^{12}, \\
p_9(x) &= x^{102} + x^{100} + x^{94} + x^{86} + x^{80} + x^{78} + x^{76} + x^{74} + x^{72} + x^{70} + x^{68} \\
&\quad + x^{66} + x^{64} + x^{62} + x^{60} + x^{48} + x^{46} + x^{42} + x^{36} + x^{26} + x^{20} + x^{18} \\
&\quad + x^{16} + x^{10} + x^8 + x^6, \\
p_{12}(x) &= x^{111} + x^{108} + x^{99} + x^{96} + x^{95} + x^{92} + x^{83} + x^{80} + x^{79} + x^{76} + x^{67} \\
&\quad + x^{64} + x^{63} + x^{60} + x^{51} + x^{48} + x^{31} + x^{28} + x^{19} + x^{16} + x^{15} + x^{12} \\
&\quad + x^3 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 1, 0, 1, 0, 1, 0, 1, 1, 0, 1, 0, 0, 0, 1]$.

For $\phi(M^e, 1, 1)$:

$$\begin{aligned}
p_0(x) &= x^{102} + x^{94} + x^{90} + x^{86} + x^{84} + x^{82} + x^{74} + x^{70} + x^{68} + x^{66} + x^{62} \\
&\quad + x^{60} + x^{50} + x^{48} + x^{46} + x^{44} + x^{42}, \\
p_3(x) &= x^{96} + x^{92} + x^{90} + x^{86} + x^{78} + x^{72} + x^{70} + x^{66} + x^{62} + x^{48} + x^{46} \\
&\quad + x^{44} + x^{42} + x^{40} + x^{38} + x^{34} + x^{32} + x^{26} + x^{24} + x^{20} + x^{18} + x^{12}, \\
p_6(x) &= x^{96} + x^{94} + x^{90} + x^{88} + x^{80} + x^{78} + x^{68} + x^{66} + x^{64} + x^{62} + x^{52} \\
&\quad + x^{48} + x^{44} + x^{42} + x^{40} + x^{38} + x^{34} + x^{30} + x^{26} + x^{22} + x^{18} + x^{14} \\
&\quad + x^8 + x^6 + x^4 + 1, \\
p_9(x) &= x^{96} + x^{92} + x^{90} + x^{88} + x^{84} + x^{82} + x^{80} + x^{76} + x^{70} + x^{64} + x^{58} \\
&\quad + x^{42} + x^{38} + x^{34} + x^{32} + x^{30} + x^{26} + x^{24} + x^{14} + x^4, \\
p_{12}(x) &= x^{102} + x^{96} + x^{86} + x^{80} + x^{70} + x^{64} + x^{54} + x^{48} + x^{22} + x^{16} + x^6 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 1, 0, 0, 0, 1, 0, 0, 0, 0, 0, 1, 0]$.

For $\phi(W^e, 0, 0)$:

$$\begin{aligned}
p_0(x) &= x^{102} + x^{94} + x^{86} + x^{78} + x^{76} + x^{60} + x^{50} + x^{42} + x^{24}, \\
p_3(x) &= x^{96} + x^{92} + x^{90} + x^{86} + x^{84} + x^{80} + x^{78} + x^{70} + x^{68} + x^{66} + x^{62} \\
&\quad + x^{52} + x^{46} + x^{42} + x^{40} + x^{38} + x^{36} + x^{34} + x^{26} + x^{20} + x^{18} + x^{16}, \\
p_6(x) &= x^{96} + x^{94} + x^{88} + x^{82} + x^{76} + x^{70} + x^{66} + x^{62} + x^{58} + x^{52} + x^{46} \\
&\quad + x^{38} + x^{32} + x^{28} + x^{20} + x^{18} + x^{12} + x^6 + x^4 + 1, \\
p_9(x) &= x^{96} + x^{92} + x^{90} + x^{88} + x^{84} + x^{82} + x^{80} + x^{76} + x^{70} + x^{64} + x^{58} \\
&\quad + x^{42} + x^{38} + x^{34} + x^{32} + x^{30} + x^{26} + x^{24} + x^{14} + x^4, \\
p_{12}(x) &= x^{102} + x^{96} + x^{86} + x^{80} + x^{70} + x^{64} + x^{54} + x^{48} + x^{22} + x^{16} + x^6 + 1,
\end{aligned}$$

and the initial terms are $[1, 0, 0, 0, 1, 0, 0, 0, 0, 0, 0, 0, 0, 0, 1, 0]$.

For $\phi(W^e, 0, 1)$:

$$\begin{aligned}
p_0(x) &= x^{111} + x^{108} + x^{103} + x^{102} + x^{100} + x^{96} + x^{94} + x^{93} + x^{91} + x^{89} + x^{87} \\
&\quad + x^{85} + x^{84} + x^{83} + x^{82} + x^{78} + x^{77} + x^{76} + x^{75} + x^{69} + x^{66} + x^{64} \\
&\quad + x^{60} + x^{55} + x^{53} + x^{52} + x^{51} + x^{50} + x^{49} + x^{47} + x^{46} + x^{45} + x^{43} \\
&\quad + x^{42} + x^{40} + x^{36} + x^{33}, \\
p_3(x) &= x^{84} + x^{82} + x^{80} + x^{74} + x^{72} + x^{68} + x^{64} + x^{60} + x^{56} + x^{50} + x^{40} \\
&\quad + x^{34} + x^{32} + x^{26} + x^{20} + x^{18}, \\
p_6(x) &= x^{93} + x^{90} + x^{89} + x^{87} + x^{86} + x^{84} + x^{83} + x^{80} + x^{73} + x^{70} + x^{69} \\
&\quad + x^{67} + x^{66} + x^{65} + x^{64} + x^{63} + x^{62} + x^{60} + x^{59} + x^{56} + x^{45} + x^{42} \\
&\quad + x^{39} + x^{37} + x^{36} + x^{34} + x^{31} + x^{29} + x^{28} + x^{26} + x^{25} + x^{23} + x^{22} \\
&\quad + x^{21} + x^{20} + x^{19} + x^{18} + x^{16} + x^{15} + x^{12}, \\
p_9(x) &= x^{102} + x^{100} + x^{94} + x^{86} + x^{80} + x^{78} + x^{76} + x^{74} + x^{72} + x^{70} + x^{68} \\
&\quad + x^{66} + x^{64} + x^{62} + x^{60} + x^{48} + x^{46} + x^{42} + x^{36} + x^{26} + x^{20} + x^{18}
\end{aligned}$$

$$\begin{aligned}
& + x^{16} + x^{10} + x^8 + x^6, \\
p_{12}(x) = & x^{111} + x^{108} + x^{99} + x^{96} + x^{95} + x^{92} + x^{83} + x^{80} + x^{79} + x^{76} + x^{67} \\
& + x^{64} + x^{63} + x^{60} + x^{51} + x^{48} + x^{31} + x^{28} + x^{19} + x^{16} + x^{15} + x^{12} \\
& + x^3 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 1, 0, 1, 1, 0, 1, 1, 1, 1, 0, 0]$.

For $\phi(W^e, 1, 0)$:

$$\begin{aligned}
p_0(x) = & x^{111} + x^{108} + x^{103} + x^{102} + x^{100} + x^{96} + x^{94} + x^{93} + x^{91} + x^{89} + x^{87} \\
& + x^{85} + x^{84} + x^{83} + x^{82} + x^{78} + x^{77} + x^{76} + x^{75} + x^{69} + x^{66} + x^{64} \\
& + x^{60} + x^{55} + x^{53} + x^{52} + x^{51} + x^{50} + x^{49} + x^{47} + x^{46} + x^{45} + x^{43} \\
& + x^{42} + x^{40} + x^{36} + x^{33}, \\
p_3(x) = & x^{84} + x^{82} + x^{80} + x^{74} + x^{72} + x^{68} + x^{64} + x^{60} + x^{56} + x^{50} + x^{40} \\
& + x^{34} + x^{32} + x^{26} + x^{20} + x^{18}, \\
p_6(x) = & x^{93} + x^{90} + x^{89} + x^{87} + x^{86} + x^{84} + x^{83} + x^{80} + x^{73} + x^{70} + x^{69} \\
& + x^{67} + x^{66} + x^{65} + x^{64} + x^{63} + x^{62} + x^{60} + x^{59} + x^{56} + x^{45} + x^{42} \\
& + x^{39} + x^{37} + x^{36} + x^{34} + x^{31} + x^{29} + x^{28} + x^{26} + x^{25} + x^{23} + x^{22} \\
& + x^{21} + x^{20} + x^{19} + x^{18} + x^{16} + x^{15} + x^{12}, \\
p_9(x) = & x^{102} + x^{100} + x^{94} + x^{86} + x^{80} + x^{78} + x^{76} + x^{74} + x^{72} + x^{70} + x^{68} \\
& + x^{66} + x^{64} + x^{62} + x^{60} + x^{48} + x^{46} + x^{42} + x^{36} + x^{26} + x^{20} + x^{18} \\
& + x^{16} + x^{10} + x^8 + x^6, \\
p_{12}(x) = & x^{111} + x^{108} + x^{99} + x^{96} + x^{95} + x^{92} + x^{83} + x^{80} + x^{79} + x^{76} + x^{67} \\
& + x^{64} + x^{63} + x^{60} + x^{51} + x^{48} + x^{31} + x^{28} + x^{19} + x^{16} + x^{15} + x^{12} \\
& + x^3 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 1, 0, 1, 1, 0, 1, 1, 1, 1, 0, 0]$.

For $\phi(W^e, 1, 1)$:

$$\begin{aligned}
p_0(x) = & x^{120} + x^{114} + x^{112} + x^{106} + x^{104} + x^{102} + x^{98} + x^{96} + x^{86} + x^{84} \\
& + x^{76} + x^{74} + x^{70} + x^{62} + x^{56} + x^{48} + x^{42}, \\
p_3(x) = & x^{114} + x^{110} + x^{96} + x^{88} + x^{78} + x^{74} + x^{70} + x^{68} + x^{64} + x^{58} + x^{54} \\
& + x^{50} + x^{48} + x^{42} + x^{40} + x^{36} + x^{32} + x^{28} + x^{26} + x^{12}, \\
p_6(x) = & x^{114} + x^{112} + x^{98} + x^{92} + x^{90} + x^{88} + x^{86} + x^{84} + x^{64} + x^{58} + x^{56} \\
& + x^{54} + x^{46} + x^{44} + x^{42} + x^{36} + x^{34} + x^{32} + x^{30} + x^{24} + x^{20} + x^{18} \\
& + x^{10} + x^8 + x^4 + 1, \\
p_9(x) = & x^{114} + x^{110} + x^{106} + x^{104} + x^{102} + x^{96} + x^{94} + x^{86} + x^{84} + x^{80} + x^{70} \\
& + x^{60} + x^{58} + x^{56} + x^{54} + x^{52} + x^{46} + x^{44} + x^{42} + x^{40} + x^{38} + x^{34} \\
& + x^{32} + x^{24} + x^{22} + x^{20} + x^{16} + x^{14} + x^{10} + x^4, \\
p_{12}(x) = & x^{120} + x^{104} + x^{96} + x^{88} + x^{80} + x^{72} + x^{64} + x^{48} + x^{40} + x^{24} + x^{16} + 1
\end{aligned}$$

,

and the initial terms are $[0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 1, 0, 0, 0, 0, 0]$.

For $\phi(M^o, 0, 0)$:

$$\begin{aligned}
p_0(x) &= x^{111} + x^{108} + x^{105} + x^{103} + x^{102} + x^{100} + x^{99} + x^{97} + x^{94} + x^{93} \\
&\quad + x^{92} + x^{85} + x^{83} + x^{82} + x^{81} + x^{80} + x^{78} + x^{77} + x^{76} + x^{75} + x^{73} \\
&\quad + x^{70} + x^{69} + x^{68} + x^{67} + x^{66} + x^{64} + x^{62} + x^{60} + x^{59} + x^{56} + x^{54} \\
&\quad + x^{53} + x^{51} + x^{50} + x^{49} + x^{48} + x^{47} + x^{43} + x^{42} + x^{41} + x^{37} + x^{36} \\
&\quad + x^{35} + x^{34} + x^{33} + x^{32} + x^{30} + x^{28} + x^{27} + x^{24}, \\
p_3(x) &= x^{99} + x^{96} + x^{93} + x^{90} + x^{87} + x^{83} + x^{81} + x^{79} + x^{77} + x^{74} + x^{72} \\
&\quad + x^{71} + x^{70} + x^{69} + x^{68} + x^{66} + x^{64} + x^{63} + x^{62} + x^{60} + x^{56} + x^{54} \\
&\quad + x^{52} + x^{49} + x^{47} + x^{46} + x^{44} + x^{43} + x^{41} + x^{39} + x^{32} + x^{30} + x^{29} \\
&\quad + x^{28} + x^{26} + x^{25} + x^{22} + x^{20} + x^{19} + x^{16}, \\
p_6(x) &= x^{105} + x^{103} + x^{102} + x^{100} + x^{97} + x^{96} + x^{94} + x^{93} + x^{92} + x^{91} + x^{89} \\
&\quad + x^{88} + x^{87} + x^{86} + x^{84} + x^{83} + x^{76} + x^{75} + x^{71} + x^{70} + x^{69} + x^{66} \\
&\quad + x^{62} + x^{60} + x^{59} + x^{58} + x^{56} + x^{54} + x^{52} + x^{51} + x^{50} + x^{48} + x^{47} \\
&\quad + x^{46} + x^{45} + x^{43} + x^{42} + x^{40} + x^{39} + x^{37} + x^{35} + x^{31} + x^{30} + x^{29} \\
&\quad + x^{26} + x^{23} + x^{22} + x^{21} + x^{19} + x^{18} + x^{17} + x^{16} + x^{15} + x^{14} + x^{11} \\
&\quad + x^8 + x^3 + 1, \\
p_9(x) &= x^{99} + x^{96} + x^{91} + x^{88} + x^{67} + x^{64} + x^{59} + x^{56} + x^{19} + x^{16} + x^{11} + x^8, \\
p_{12}(x) &= x^{111} + x^{108} + x^{95} + x^{92} + x^{91} + x^{88} + x^{83} + x^{80} + x^{79} + x^{76} + x^{63} \\
&\quad + x^{60} + x^{59} + x^{56} + x^{51} + x^{48} + x^{31} + x^{28} + x^{15} + x^{12} + x^{11} + x^8 \\
&\quad + x^3 + 1,
\end{aligned}$$

and the initial terms are $[1, 0, 0, 0, 0, 1, 0, 0, 1, 1, 0, 0, 0, 1, 1, 0, 1]$.

For $\phi(M^o, 0, 1)$:

$$\begin{aligned}
p_0(x) &= x^{106} + x^{104} + x^{102} + x^{100} + x^{97} + x^{95} + x^{93} + x^{90} + x^{89} + x^{85} + x^{84} \\
&\quad + x^{83} + x^{82} + x^{81} + x^{79} + x^{78} + x^{77} + x^{75} + x^{74} + x^{73} + x^{72} + x^{71} \\
&\quad + x^{70} + x^{66} + x^{63} + x^{61} + x^{58} + x^{57} + x^{56} + x^{50} + x^{48} + x^{44} + x^{41} \\
&\quad + x^{40} + x^{37} + x^{34} + x^{33}, \\
p_3(x) &= x^{79} + x^{76} + x^{75} + x^{73} + x^{72} + x^{71} + x^{70} + x^{69} + x^{68} + x^{67} + x^{66} \\
&\quad + x^{65} + x^{64} + x^{63} + x^{62} + x^{61} + x^{60} + x^{59} + x^{58} + x^{57} + x^{56} + x^{55} \\
&\quad + x^{54} + x^{53} + x^{52} + x^{51} + x^{50} + x^{49} + x^{48} + x^{46} + x^{45} + x^{42} + x^{39} \\
&\quad + x^{36} + x^{35} + x^{33} + x^{32} + x^{31} + x^{30} + x^{29} + x^{28} + x^{27} + x^{26} + x^{25} \\
&\quad + x^{24} + x^{22} + x^{21} + x^{18}, \\
p_6(x) &= x^{88} + x^{86} + x^{82} + x^{80} + x^{68} + x^{66} + x^{62} + x^{58} + x^{56} + x^{40} + x^{38} \\
&\quad + x^{36} + x^{32} + x^{30} + x^{28} + x^{24} + x^{22} + x^{18} + x^{14} + x^{12},
\end{aligned}$$

$$\begin{aligned}
p_9(x) &= x^{97} + x^{94} + x^{89} + x^{86} + x^{65} + x^{62} + x^{57} + x^{54} + x^{17} + x^{14} + x^9 + x^6, \\
p_{12}(x) &= x^{106} + x^{104} + x^{102} + x^{100} + x^{98} + x^{96} + x^{94} + x^{92} + x^{86} + x^{84} + x^{82} \\
&\quad + x^{80} + x^{74} + x^{72} + x^{70} + x^{68} + x^{66} + x^{64} + x^{62} + x^{60} + x^{54} + x^{52} \\
&\quad + x^{50} + x^{48} + x^{26} + x^{24} + x^{22} + x^{20} + x^{18} + x^{16} + x^{14} + x^{12} + x^6 \\
&\quad + x^4 + x^2 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 1, 0, 1, 0, 0, 0, 0, 0, 1, 0, 0, 0, 1, 0]$.

For $\phi(M^o, 1, 0)$:

$$\begin{aligned}
p_0(x) &= x^{120} + x^{112} + x^{111} + x^{108} + x^{105} + x^{104} + x^{103} + x^{101} + x^{100} + x^{94} \\
&\quad + x^{93} + x^{92} + x^{91} + x^{89} + x^{88} + x^{87} + x^{83} + x^{82} + x^{80} + x^{78} + x^{77} \\
&\quad + x^{75} + x^{73} + x^{72} + x^{70} + x^{60} + x^{59} + x^{58} + x^{55} + x^{54} + x^{53} + x^{52} \\
&\quad + x^{47} + x^{44} + x^{43} + x^{42} + x^{41} + x^{37} + x^{33}, \\
p_3(x) &= x^{93} + x^{91} + x^{90} + x^{89} + x^{88} + x^{86} + x^{81} + x^{79} + x^{78} + x^{77} + x^{76} \\
&\quad + x^{74} + x^{61} + x^{59} + x^{58} + x^{57} + x^{56} + x^{54} + x^{53} + x^{51} + x^{50} + x^{48} \\
&\quad + x^{47} + x^{45} + x^{44} + x^{42} + x^{41} + x^{39} + x^{38} + x^{36} + x^{35} + x^{33} + x^{32} \\
&\quad + x^{30} + x^{25} + x^{23} + x^{22} + x^{21} + x^{20} + x^{18}, \\
p_6(x) &= x^{102} + x^{98} + x^{96} + x^{92} + x^{90} + x^{86} + x^{84} + x^{82} + x^{80} + x^{78} + x^{76} \\
&\quad + x^{74} + x^{72} + x^{70} + x^{68} + x^{66} + x^{64} + x^{62} + x^{60} + x^{56} + x^{54} + x^{48} \\
&\quad + x^{46} + x^{42} + x^{40} + x^{38} + x^{36} + x^{32} + x^{30} + x^{26} + x^{24} + x^{22} + x^{20} \\
&\quad + x^{18} + x^{16} + x^{12}, \\
p_9(x) &= x^{111} + x^{109} + x^{108} + x^{106} + x^{103} + x^{101} + x^{100} + x^{99} + x^{98} + x^{97} \\
&\quad + x^{96} + x^{94} + x^{91} + x^{89} + x^{88} + x^{86} + x^{79} + x^{77} + x^{76} + x^{74} + x^{71} \\
&\quad + x^{69} + x^{68} + x^{67} + x^{66} + x^{65} + x^{64} + x^{62} + x^{59} + x^{57} + x^{56} + x^{54} \\
&\quad + x^{31} + x^{29} + x^{28} + x^{26} + x^{23} + x^{21} + x^{20} + x^{19} + x^{18} + x^{17} + x^{16} \\
&\quad + x^{14} + x^{11} + x^9 + x^8 + x^6, \\
p_{12}(x) &= x^{120} + x^{108} + x^{104} + x^{100} + x^{80} + x^{76} + x^{72} + x^{68} + x^{56} + x^{48} + x^{40} \\
&\quad + x^{28} + x^{24} + x^{20} + x^8 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 1, 0, 1, 1, 0, 1, 1, 0, 0, 1, 0]$.

For $\phi(M^o, 1, 1)$:

$$\begin{aligned}
p_0(x) &= x^{111} + x^{108} + x^{105} + x^{103} + x^{102} + x^{100} + x^{97} + x^{96} + x^{94} + x^{92} \\
&\quad + x^{91} + x^{90} + x^{89} + x^{88} + x^{87} + x^{86} + x^{85} + x^{84} + x^{83} + x^{82} + x^{80} \\
&\quad + x^{76} + x^{75} + x^{74} + x^{73} + x^{72} + x^{70} + x^{69} + x^{62} + x^{61} + x^{60} + x^{59} \\
&\quad + x^{58} + x^{56} + x^{55} + x^{54} + x^{52} + x^{50} + x^{48} + x^{47} + x^{44} + x^{43} + x^{42}, \\
p_3(x) &= x^{99} + x^{96} + x^{93} + x^{90} + x^{84} + x^{83} + x^{79} + x^{78} + x^{73} + x^{72} + x^{67} \\
&\quad + x^{65} + x^{61} + x^{59} + x^{58} + x^{57} + x^{53} + x^{52} + x^{51} + x^{50} + x^{49} + x^{48} \\
&\quad + x^{47} + x^{46} + x^{44} + x^{40} + x^{39} + x^{38} + x^{37} + x^{34} + x^{33} + x^{32} + x^{28}
\end{aligned}$$

$$\begin{aligned}
& + x^{27} + x^{25} + x^{24} + x^{23} + x^{22}, \\
p_6(x) &= x^{105} + x^{103} + x^{102} + x^{100} + x^{99} + x^{97} + x^{94} + x^{92} + x^{90} + x^{87} + x^{84} \\
& + x^{80} + x^{76} + x^{75} + x^{73} + x^{71} + x^{67} + x^{65} + x^{64} + x^{63} + x^{58} + x^{55} \\
& + x^{54} + x^{50} + x^{46} + x^{44} + x^{36} + x^{35} + x^{34} + x^{31} + x^{30} + x^{27} + x^{25} \\
& + x^{24} + x^{20} + x^{17} + x^{14} + x^{12} + x^3 + 1, \\
p_9(x) &= x^{99} + x^{96} + x^{91} + x^{88} + x^{67} + x^{64} + x^{59} + x^{56} + x^{19} + x^{16} + x^{11} + x^8, \\
p_{12}(x) &= x^{111} + x^{108} + x^{95} + x^{92} + x^{91} + x^{88} + x^{83} + x^{80} + x^{79} + x^{76} + x^{63} \\
& + x^{60} + x^{59} + x^{56} + x^{51} + x^{48} + x^{31} + x^{28} + x^{15} + x^{12} + x^{11} + x^8 \\
& + x^3 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 0, 0, 1, 0, 0, 1, 0, 0, 1, 1, 1]$.

For $\phi(W^o, 0, 0)$:

$$\begin{aligned}
p_0(x) &= x^{111} + x^{108} + x^{105} + x^{103} + x^{102} + x^{100} + x^{99} + x^{97} + x^{94} + x^{93} \\
& + x^{92} + x^{85} + x^{83} + x^{82} + x^{81} + x^{80} + x^{78} + x^{77} + x^{76} + x^{75} + x^{73} \\
& + x^{70} + x^{69} + x^{68} + x^{67} + x^{66} + x^{64} + x^{62} + x^{60} + x^{59} + x^{56} + x^{54} \\
& + x^{53} + x^{51} + x^{50} + x^{49} + x^{48} + x^{47} + x^{43} + x^{42} + x^{41} + x^{37} + x^{36} \\
& + x^{35} + x^{34} + x^{33} + x^{32} + x^{30} + x^{28} + x^{27} + x^{24}, \\
p_3(x) &= x^{99} + x^{96} + x^{93} + x^{90} + x^{87} + x^{83} + x^{81} + x^{79} + x^{77} + x^{74} + x^{72} \\
& + x^{71} + x^{70} + x^{69} + x^{68} + x^{66} + x^{64} + x^{63} + x^{62} + x^{60} + x^{56} + x^{54} \\
& + x^{52} + x^{49} + x^{47} + x^{46} + x^{44} + x^{43} + x^{41} + x^{39} + x^{32} + x^{30} + x^{29} \\
& + x^{28} + x^{26} + x^{25} + x^{22} + x^{20} + x^{19} + x^{16}, \\
p_6(x) &= x^{105} + x^{103} + x^{102} + x^{100} + x^{97} + x^{96} + x^{94} + x^{93} + x^{92} + x^{91} + x^{89} \\
& + x^{88} + x^{87} + x^{86} + x^{84} + x^{83} + x^{76} + x^{75} + x^{71} + x^{70} + x^{69} + x^{66} \\
& + x^{62} + x^{60} + x^{59} + x^{58} + x^{56} + x^{54} + x^{52} + x^{51} + x^{50} + x^{48} + x^{47} \\
& + x^{46} + x^{45} + x^{43} + x^{42} + x^{40} + x^{39} + x^{37} + x^{35} + x^{31} + x^{30} + x^{29} \\
& + x^{26} + x^{23} + x^{22} + x^{21} + x^{19} + x^{18} + x^{17} + x^{16} + x^{15} + x^{14} + x^{11} \\
& + x^8 + x^3 + 1, \\
p_9(x) &= x^{99} + x^{96} + x^{91} + x^{88} + x^{67} + x^{64} + x^{59} + x^{56} + x^{19} + x^{16} + x^{11} + x^8, \\
p_{12}(x) &= x^{111} + x^{108} + x^{95} + x^{92} + x^{91} + x^{88} + x^{83} + x^{80} + x^{79} + x^{76} + x^{63} \\
& + x^{60} + x^{59} + x^{56} + x^{51} + x^{48} + x^{31} + x^{28} + x^{15} + x^{12} + x^{11} + x^8 \\
& + x^3 + 1,
\end{aligned}$$

and the initial terms are $[1, 0, 0, 0, 1, 0, 0, 1, 1, 0, 0, 0, 1, 1, 0, 1]$.

For $\phi(W^o, 0, 1)$:

$$\begin{aligned}
p_0(x) &= x^{120} + x^{112} + x^{111} + x^{108} + x^{105} + x^{104} + x^{103} + x^{101} + x^{100} + x^{94} \\
& + x^{93} + x^{92} + x^{91} + x^{89} + x^{88} + x^{87} + x^{83} + x^{82} + x^{80} + x^{78} + x^{77} \\
& + x^{75} + x^{73} + x^{72} + x^{70} + x^{60} + x^{59} + x^{58} + x^{55} + x^{54} + x^{53} + x^{52}
\end{aligned}$$

$$\begin{aligned}
& + x^{47} + x^{44} + x^{43} + x^{42} + x^{41} + x^{37} + x^{33}, \\
p_3(x) &= x^{93} + x^{91} + x^{90} + x^{89} + x^{88} + x^{86} + x^{81} + x^{79} + x^{78} + x^{77} + x^{76} \\
& + x^{74} + x^{61} + x^{59} + x^{58} + x^{57} + x^{56} + x^{54} + x^{53} + x^{51} + x^{50} + x^{48} \\
& + x^{47} + x^{45} + x^{44} + x^{42} + x^{41} + x^{39} + x^{38} + x^{36} + x^{35} + x^{33} + x^{32} \\
& + x^{30} + x^{25} + x^{23} + x^{22} + x^{21} + x^{20} + x^{18}, \\
p_6(x) &= x^{102} + x^{98} + x^{96} + x^{92} + x^{90} + x^{86} + x^{84} + x^{82} + x^{80} + x^{78} + x^{76} \\
& + x^{74} + x^{72} + x^{70} + x^{68} + x^{66} + x^{64} + x^{62} + x^{60} + x^{56} + x^{54} + x^{48} \\
& + x^{46} + x^{42} + x^{40} + x^{38} + x^{36} + x^{32} + x^{30} + x^{26} + x^{24} + x^{22} + x^{20} \\
& + x^{18} + x^{16} + x^{12}, \\
p_9(x) &= x^{111} + x^{109} + x^{108} + x^{106} + x^{103} + x^{101} + x^{100} + x^{99} + x^{98} + x^{97} \\
& + x^{96} + x^{94} + x^{91} + x^{89} + x^{88} + x^{86} + x^{79} + x^{77} + x^{76} + x^{74} + x^{71} \\
& + x^{69} + x^{68} + x^{67} + x^{66} + x^{65} + x^{64} + x^{62} + x^{59} + x^{57} + x^{56} + x^{54} \\
& + x^{31} + x^{29} + x^{28} + x^{26} + x^{23} + x^{21} + x^{20} + x^{19} + x^{18} + x^{17} + x^{16} \\
& + x^{14} + x^{11} + x^9 + x^8 + x^6, \\
p_{12}(x) &= x^{120} + x^{108} + x^{104} + x^{100} + x^{80} + x^{76} + x^{72} + x^{68} + x^{56} + x^{48} + x^{40} \\
& + x^{28} + x^{24} + x^{20} + x^8 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 1, 0, 1, 1, 0, 1, 1, 0, 0, 1, 0]$.

For $\phi(W^o, 1, 0)$:

$$\begin{aligned}
p_0(x) &= x^{106} + x^{104} + x^{102} + x^{100} + x^{97} + x^{95} + x^{93} + x^{90} + x^{89} + x^{85} + x^{84} \\
& + x^{83} + x^{82} + x^{81} + x^{79} + x^{78} + x^{77} + x^{75} + x^{74} + x^{73} + x^{72} + x^{71} \\
& + x^{70} + x^{66} + x^{63} + x^{61} + x^{58} + x^{57} + x^{56} + x^{50} + x^{48} + x^{44} + x^{41} \\
& + x^{40} + x^{37} + x^{34} + x^{33}, \\
p_3(x) &= x^{79} + x^{76} + x^{75} + x^{73} + x^{72} + x^{71} + x^{70} + x^{69} + x^{68} + x^{67} + x^{66} \\
& + x^{65} + x^{64} + x^{63} + x^{62} + x^{61} + x^{60} + x^{59} + x^{58} + x^{57} + x^{56} + x^{55} \\
& + x^{54} + x^{53} + x^{52} + x^{51} + x^{50} + x^{49} + x^{48} + x^{46} + x^{45} + x^{42} + x^{39} \\
& + x^{36} + x^{35} + x^{33} + x^{32} + x^{31} + x^{30} + x^{29} + x^{28} + x^{27} + x^{26} + x^{25} \\
& + x^{24} + x^{22} + x^{21} + x^{18}, \\
p_6(x) &= x^{88} + x^{86} + x^{82} + x^{80} + x^{68} + x^{66} + x^{62} + x^{58} + x^{56} + x^{40} + x^{38} \\
& + x^{36} + x^{32} + x^{30} + x^{28} + x^{24} + x^{22} + x^{18} + x^{14} + x^{12}, \\
p_9(x) &= x^{97} + x^{94} + x^{89} + x^{86} + x^{65} + x^{62} + x^{57} + x^{54} + x^{17} + x^{14} + x^9 + x^6, \\
p_{12}(x) &= x^{106} + x^{104} + x^{102} + x^{100} + x^{98} + x^{96} + x^{94} + x^{92} + x^{86} + x^{84} + x^{82} \\
& + x^{80} + x^{74} + x^{72} + x^{70} + x^{68} + x^{66} + x^{64} + x^{62} + x^{60} + x^{54} + x^{52} \\
& + x^{50} + x^{48} + x^{26} + x^{24} + x^{22} + x^{20} + x^{18} + x^{16} + x^{14} + x^{12} + x^6 \\
& + x^4 + x^2 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 1, 0, 1, 0, 0, 0, 0, 0, 1, 0, 0, 0, 1, 0]$.

For $\phi(W^o, 1, 1)$:

$$\begin{aligned}
p_0(x) &= x^{111} + x^{108} + x^{105} + x^{103} + x^{102} + x^{100} + x^{97} + x^{96} + x^{94} + x^{92} \\
&\quad + x^{91} + x^{90} + x^{89} + x^{88} + x^{87} + x^{86} + x^{85} + x^{84} + x^{83} + x^{82} + x^{80} \\
&\quad + x^{76} + x^{75} + x^{74} + x^{73} + x^{72} + x^{70} + x^{69} + x^{62} + x^{61} + x^{60} + x^{59} \\
&\quad + x^{58} + x^{56} + x^{55} + x^{54} + x^{52} + x^{50} + x^{48} + x^{47} + x^{44} + x^{43} + x^{42}, \\
p_3(x) &= x^{99} + x^{96} + x^{93} + x^{90} + x^{84} + x^{83} + x^{79} + x^{78} + x^{73} + x^{72} + x^{67} \\
&\quad + x^{65} + x^{61} + x^{59} + x^{58} + x^{57} + x^{53} + x^{52} + x^{51} + x^{50} + x^{49} + x^{48} \\
&\quad + x^{47} + x^{46} + x^{44} + x^{40} + x^{39} + x^{38} + x^{37} + x^{34} + x^{33} + x^{32} + x^{28} \\
&\quad + x^{27} + x^{25} + x^{24} + x^{23} + x^{22}, \\
p_6(x) &= x^{105} + x^{103} + x^{102} + x^{100} + x^{99} + x^{97} + x^{94} + x^{92} + x^{90} + x^{87} + x^{84} \\
&\quad + x^{80} + x^{76} + x^{75} + x^{73} + x^{71} + x^{67} + x^{65} + x^{64} + x^{63} + x^{58} + x^{55} \\
&\quad + x^{54} + x^{50} + x^{46} + x^{44} + x^{36} + x^{35} + x^{34} + x^{31} + x^{30} + x^{27} + x^{25} \\
&\quad + x^{24} + x^{20} + x^{17} + x^{14} + x^{12} + x^3 + 1, \\
p_9(x) &= x^{99} + x^{96} + x^{91} + x^{88} + x^{67} + x^{64} + x^{59} + x^{56} + x^{19} + x^{16} + x^{11} + x^8, \\
p_{12}(x) &= x^{111} + x^{108} + x^{95} + x^{92} + x^{91} + x^{88} + x^{83} + x^{80} + x^{79} + x^{76} + x^{63} \\
&\quad + x^{60} + x^{59} + x^{56} + x^{51} + x^{48} + x^{31} + x^{28} + x^{15} + x^{12} + x^{11} + x^8 \\
&\quad + x^3 + 1,
\end{aligned}$$

and the initial terms are $[0, 0, 0, 0, 0, 0, 0, 1, 0, 0, 1, 0, 0, 1, 1, 1]$.

Below is the transition function and output function of an 2-automaton that generates $T = M_{0,0}^o$:

Transition function $(n, j) \mapsto \delta(n, j)$ ($\Lambda(n) := [\delta(n, 0), \delta(n, 1)]$):

n	$\Lambda(n)$	n	$\Lambda(n)$	n	$\Lambda(n)$	n	$\Lambda(n)$	n	$\Lambda(n)$
0	[1, 2]	102	[166, 37]	204	[310, 130]	306	[18, 368]	408	[308, 310]
1	[3, 4]	103	[167, 168]	205	[11, 55]	307	[60, 369]	409	[349, 446]
2	[5, 6]	104	[169, 170]	206	[311, 174]	308	[370, 371]	410	[444, 118]
3	[7, 8]	105	[158, 122]	207	[312, 114]	309	[372, 373]	411	[436, 309]
4	[9, 10]	106	[171, 172]	208	[149, 128]	310	[374, 375]	412	[166, 138]
5	[11, 12]	107	[173, 174]	209	[46, 128]	311	[95, 257]	413	[310, 437]
6	[13, 14]	108	[175, 176]	210	[153, 47]	312	[376, 182]	414	[326, 32]
7	[15, 16]	109	[152, 148]	211	[44, 44]	313	[377, 378]	415	[306, 316]
8	[17, 18]	110	[177, 178]	212	[313, 314]	314	[20, 379]	416	[53, 13]
9	[19, 20]	111	[171, 179]	213	[147, 149]	315	[72, 380]	417	[122, 160]
10	[21, 22]	112	[180, 181]	214	[315, 316]	316	[229, 370]	418	[134, 37]
11	[23, 24]	113	[182, 183]	215	[117, 305]	317	[265, 96]	419	[124, 447]
12	[25, 26]	114	[184, 185]	216	[138, 317]	318	[381, 382]	420	[177, 34]
13	[27, 28]	115	[24, 186]	217	[122, 131]	319	[383, 236]	421	[424, 431]
14	[29, 30]	116	[28, 187]	218	[137, 113]	320	[194, 384]	422	[151, 153]
15	[31, 32]	117	[188, 106]	219	[318, 166]	321	[385, 386]	423	[448, 48]
16	[33, 34]	118	[189, 190]	220	[177, 51]	322	[387, 383]	424	[449, 450]

17	[35, 36]	119	[191, 192]	221	[120, 135]	323	[388, 389]	425	[368, 451]
18	[37, 38]	120	[193, 194]	222	[178, 319]	324	[68, 390]	426	[260, 452]
19	[37, 39]	121	[195, 196]	223	[320, 130]	325	[181, 391]	427	[453, 454]
20	[40, 41]	122	[197, 198]	224	[137, 321]	326	[392, 110]	428	[272, 455]
21	[42, 43]	123	[16, 199]	225	[322, 164]	327	[239, 393]	429	[456, 457]
22	[44, 45]	124	[200, 201]	226	[12, 323]	328	[394, 395]	430	[458, 372]
23	[46, 47]	125	[202, 203]	227	[319, 172]	329	[396, 397]	431	[459, 460]
24	[42, 48]	126	[204, 205]	228	[47, 151]	330	[105, 200]	432	[461, 462]
25	[32, 49]	127	[206, 207]	229	[301, 324]	331	[398, 399]	433	[463, 464]
26	[50, 41]	128	[208, 209]	230	[52, 179]	332	[400, 401]	434	[403, 465]
27	[51, 52]	129	[210, 211]	231	[141, 156]	333	[402, 403]	435	[357, 466]
28	[53, 54]	130	[212, 213]	232	[321, 325]	334	[395, 404]	436	[467, 224]
29	[55, 56]	131	[214, 215]	233	[326, 160]	335	[405, 406]	437	[468, 469]
30	[57, 58]	132	[198, 216]	234	[327, 328]	336	[407, 392]	438	[391, 104]
31	[59, 60]	133	[217, 71]	235	[311, 329]	337	[408, 409]	439	[421, 470]
32	[61, 62]	134	[218, 219]	236	[113, 330]	338	[338, 338]	440	[471, 472]
33	[63, 64]	135	[220, 221]	237	[331, 332]	339	[410, 411]	441	[473, 474]
34	[65, 66]	136	[222, 223]	238	[333, 334]	340	[412, 413]	442	[475, 69]
35	[67, 68]	137	[224, 225]	239	[45, 154]	341	[414, 259]	443	[216, 476]
36	[69, 70]	138	[226, 227]	240	[335, 320]	342	[415, 254]	444	[469, 356]
37	[71, 72]	139	[228, 229]	241	[336, 337]	343	[237, 416]	445	[477, 478]
38	[73, 74]	140	[230, 231]	242	[146, 129]	344	[417, 100]	446	[479, 480]
39	[75, 76]	141	[232, 233]	243	[129, 153]	345	[74, 418]	447	[481, 482]
40	[77, 78]	142	[234, 235]	244	[154, 154]	346	[419, 420]	448	[483, 484]
41	[79, 80]	143	[236, 237]	245	[128, 151]	347	[192, 421]	449	[320, 304]
42	[81, 79]	144	[238, 239]	246	[45, 338]	348	[422, 423]	450	[485, 9]
43	[82, 83]	145	[240, 241]	247	[338, 45]	349	[293, 98]	451	[318, 134]
44	[84, 85]	146	[242, 243]	248	[128, 152]	350	[54, 321]	452	[443, 429]
45	[86, 87]	147	[244, 90]	249	[150, 151]	351	[113, 135]	453	[336, 329]
46	[88, 89]	148	[91, 210]	250	[153, 146]	352	[424, 425]	454	[322, 51]
47	[90, 91]	149	[245, 246]	251	[47, 147]	353	[154, 45]	455	[136, 114]
48	[92, 93]	150	[247, 245]	252	[147, 148]	354	[134, 170]	456	[432, 12]
49	[94, 95]	151	[248, 249]	253	[57, 132]	355	[426, 296]	457	[331, 447]
50	[96, 97]	152	[89, 208]	254	[335, 9]	356	[342, 427]	458	[143, 56]
51	[98, 99]	153	[250, 251]	255	[40, 339]	357	[129, 149]	459	[331, 309]
52	[100, 101]	154	[252, 84]	256	[326, 57]	358	[160, 58]	460	[341, 159]
53	[102, 103]	155	[253, 65]	257	[159, 32]	359	[327, 55]	461	[129, 46]
54	[104, 105]	156	[254, 255]	258	[340, 162]	360	[345, 145]	462	[486, 127]
55	[106, 107]	157	[256, 29]	259	[35, 119]	361	[167, 165]	463	[170, 38]
56	[108, 109]	158	[6, 61]	260	[32, 58]	362	[311, 337]	464	[178, 117]
57	[4, 67]	159	[257, 258]	261	[37, 317]	363	[121, 326]	465	[433, 446]
58	[110, 111]	160	[259, 260]	262	[114, 14]	364	[175, 428]	466	[487, 488]
59	[112, 113]	161	[99, 261]	263	[312, 137]	365	[149, 146]	467	[319, 305]
60	[53, 114]	162	[262, 263]	264	[51, 117]	366	[136, 13]	468	[333, 488]

61	[12, 115]	163	[264, 265]	265	[341, 162]	367	[131, 49]	469	[150, 129]
62	[14, 116]	164	[266, 267]	266	[164, 319]	368	[336, 429]	470	[146, 152]
63	[34, 117]	165	[268, 269]	267	[133, 163]	369	[113, 325]	471	[155, 446]
64	[118, 119]	166	[97, 270]	268	[342, 343]	370	[151, 148]	472	[340, 159]
65	[114, 120]	167	[271, 272]	269	[148, 150]	371	[430, 431]	473	[426, 119]
66	[121, 122]	168	[273, 274]	270	[344, 178]	372	[432, 143]	474	[489, 143]
67	[8, 123]	169	[275, 75]	271	[120, 325]	373	[155, 142]	475	[148, 128]
68	[124, 125]	170	[276, 277]	272	[345, 300]	374	[131, 307]	476	[166, 170]
69	[126, 127]	171	[278, 279]	273	[126, 346]	375	[433, 434]	477	[118, 298]
70	[128, 129]	172	[280, 232]	274	[338, 154]	376	[13, 120]	478	[490, 310]
71	[9, 130]	173	[281, 282]	275	[169, 8]	377	[426, 36]	479	[139, 306]
72	[131, 132]	174	[283, 284]	276	[315, 168]	378	[435, 140]	480	[438, 447]
73	[133, 134]	175	[285, 286]	277	[321, 330]	379	[294, 134]	481	[347, 167]
74	[14, 135]	176	[287, 288]	278	[55, 144]	380	[436, 125]	482	[299, 145]
75	[136, 137]	177	[289, 290]	279	[138, 38]	381	[34, 52]	483	[320, 437]
76	[8, 39]	178	[291, 292]	280	[341, 326]	382	[328, 56]	484	[491, 426]
77	[138, 123]	179	[267, 293]	281	[14, 330]	383	[303, 437]	485	[269, 468]
78	[54, 120]	180	[31, 57]	282	[13, 321]	384	[438, 332]	486	[492, 408]
79	[139, 140]	181	[294, 169]	283	[347, 315]	385	[140, 439]	487	[493, 494]
80	[141, 142]	182	[295, 296]	284	[173, 329]	386	[170, 123]	488	[495, 496]
81	[143, 144]	183	[120, 116]	285	[118, 296]	387	[54, 113]	489	[186, 108]
82	[40, 145]	184	[162, 131]	286	[348, 35]	388	[313, 440]	490	[497, 212]
83	[33, 51]	185	[297, 163]	287	[349, 142]	389	[153, 150]	491	[22, 498]
84	[146, 147]	186	[47, 152]	288	[344, 34]	390	[133, 169]	492	[328, 323]
85	[148, 47]	187	[52, 161]	289	[134, 8]	391	[57, 307]	493	[310, 10]
86	[149, 150]	188	[295, 298]	290	[143, 115]	392	[164, 171]	494	[499, 167]
87	[151, 46]	189	[171, 161]	291	[159, 57]	393	[441, 427]	495	[345, 339]
88	[152, 149]	190	[299, 300]	292	[157, 54]	394	[8, 317]	496	[158, 326]
89	[152, 46]	191	[301, 302]	293	[319, 179]	395	[349, 434]	497	[150, 147]
90	[147, 153]	192	[46, 146]	294	[350, 276]	396	[442, 426]	498	[500, 176]
91	[154, 44]	193	[303, 304]	295	[351, 92]	397	[50, 339]	499	[501, 502]
92	[155, 156]	194	[52, 305]	296	[352, 353]	398	[321, 116]	500	[503, 481]
93	[157, 13]	195	[178, 171]	297	[354, 355]	399	[34, 171]	501	[45, 44]
94	[158, 159]	196	[306, 165]	298	[356, 357]	400	[442, 295]	502	[504, 314]
95	[160, 49]	197	[163, 138]	299	[358, 264]	401	[443, 337]	503	[167, 439]
96	[117, 161]	198	[294, 166]	300	[359, 360]	402	[328, 115]	504	[505, 396]
97	[162, 160]	199	[32, 307]	301	[361, 359]	403	[444, 35]	505	[306, 168]
98	[163, 37]	200	[170, 39]	302	[362, 363]	404	[297, 169]		
99	[33, 164]	201	[122, 32]	303	[183, 206]	405	[44, 338]		
100	[140, 165]	202	[308, 303]	304	[364, 365]	406	[445, 324]		
101	[160, 132]	203	[124, 309]	305	[366, 367]	407	[117, 172]		

Output function $n \mapsto \tau(n)$:

n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$
0	0	73	0	146	1	219	1	292	0	365	1	438	0
1	0	74	1	147	0	220	1	293	0	366	1	439	1
2	0	75	1	148	0	221	0	294	0	367	1	440	0
3	0	76	0	149	1	222	1	295	0	368	0	441	1
4	0	77	1	150	0	223	1	296	1	369	0	442	0
5	0	78	0	151	1	224	1	297	1	370	1	443	1
6	0	79	0	152	0	225	0	298	0	371	0	444	0
7	0	80	1	153	1	226	1	299	0	372	1	445	1
8	0	81	0	154	0	227	0	300	1	373	0	446	0
9	0	82	1	155	0	228	0	301	0	374	1	447	0
10	0	83	1	156	1	229	0	302	0	375	1	448	1
11	0	84	1	157	0	230	1	303	0	376	0	449	1
12	1	85	0	158	0	231	1	304	1	377	1	450	0
13	0	86	1	159	1	232	1	305	1	378	1	451	1
14	1	87	1	160	0	233	0	306	0	379	0	452	1
15	0	88	0	161	1	234	1	307	1	380	0	453	0
16	1	89	0	162	1	235	0	308	1	381	1	454	0
17	0	90	0	163	0	236	0	309	1	382	0	455	1
18	0	91	0	164	0	237	1	310	1	383	0	456	1
19	0	92	0	165	0	238	0	311	0	384	0	457	1
20	1	93	0	166	1	239	1	312	0	385	1	458	0
21	0	94	0	167	0	240	1	313	1	386	1	459	1
22	1	95	0	168	1	241	0	314	1	387	0	460	0
23	0	96	0	169	0	242	1	315	1	388	1	461	1
24	0	97	1	170	1	243	1	316	0	389	1	462	0
25	1	98	0	171	1	244	0	317	0	390	0	463	1
26	0	99	1	172	0	245	1	318	1	391	0	464	1
27	0	100	1	173	1	246	1	319	0	392	0	465	1
28	1	101	0	174	0	247	0	320	1	393	1	466	1
29	1	102	1	175	1	248	1	321	1	394	0	467	0
30	0	103	0	176	0	249	0	322	0	395	0	468	0
31	0	104	0	177	1	250	1	323	1	396	0	469	0
32	1	105	0	178	1	251	0	324	1	397	0	470	1
33	1	106	1	179	0	252	0	325	0	398	1	471	0
34	1	107	1	180	0	253	0	326	0	399	1	472	1
35	0	108	1	181	0	254	1	327	1	400	0	473	1
36	1	109	0	182	0	255	1	328	0	401	1	474	0
37	0	110	1	183	0	256	0	329	0	402	0	475	0
38	0	111	1	184	1	257	1	330	0	403	0	476	1
39	1	112	0	185	1	258	1	331	1	404	1	477	1
40	1	113	0	186	0	259	0	332	0	405	1	478	0
41	0	114	1	187	1	260	1	333	0	406	1	479	0
42	0	115	0	188	0	261	0	334	0	407	0	480	0

43	1	116	1	189	1	262	1	335	1	408	1	481	0
44	1	117	0	190	0	263	0	336	0	409	0	482	0
45	1	118	1	191	0	264	0	337	1	410	0	483	1
46	0	119	0	192	0	265	0	338	0	411	0	484	1
47	0	120	0	193	0	266	0	339	0	412	1	485	0
48	0	121	1	194	1	267	0	340	1	413	1	486	0
49	0	122	0	195	1	268	0	341	0	414	0	487	1
50	0	123	1	196	0	269	0	342	0	415	0	488	1
51	0	124	1	197	0	270	0	343	1	416	1	489	0
52	1	125	1	198	0	271	0	344	0	417	0	490	0
53	1	126	1	199	1	272	1	345	1	418	1	491	1
54	0	127	0	200	1	273	1	346	1	419	1	492	0
55	1	128	1	201	0	274	0	347	0	420	1	493	1
56	1	129	1	202	1	275	0	348	1	421	1	494	1
57	0	130	1	203	1	276	1	349	0	422	1	495	1
58	1	131	1	204	1	277	1	350	0	423	1	496	0
59	0	132	0	205	0	278	1	351	0	424	1	497	0
60	1	133	0	206	0	279	1	352	1	425	0	498	0
61	1	134	1	207	0	280	0	353	0	426	1	499	1
62	1	135	1	208	1	281	1	354	1	427	0	500	0
63	1	136	1	209	0	282	0	355	1	428	1	501	1
64	1	137	1	210	1	283	0	356	0	429	1	502	0
65	1	138	1	211	1	284	1	357	1	430	0	503	0
66	1	139	0	212	1	285	1	358	0	431	1	504	0
67	0	140	1	213	0	286	1	359	1	432	1	505	0
68	1	141	1	214	1	287	0	360	1	433	1		
69	1	142	1	215	0	288	0	361	0	434	0		
70	1	143	0	216	1	289	1	362	0	435	1		
71	0	144	0	217	0	290	0	363	1	436	0		
72	1	145	1	218	1	291	1	364	1	437	0		

UNIVERSITÉ DE STRASBOURG, CNRS, IRMA UMR 7501, F-67000 STRASBOURG, FRANCE

Email address: `guoniu.han@unistra.fr`

SCHOOL OF MATHEMATICS AND STATISTICS, HUAZHONG UNIVERSITY OF SCIENCE AND TECHNOLOGY, WUHAN, PR CHINA

Email address: `huyining@hust.edu.cn`