

## TD 7-Immersiones and the geometry of valuations

---

Recall that a morphism of schemes  $f : X \rightarrow Y$  is a **closed immersion** (resp. **open immersion**) if  $|f| : |X| \rightarrow |Y|$  is a homeomorphism onto a closed (resp. open) subset of  $|Y|$  and the map  $O_Y \rightarrow f_*O_X$  is surjective (resp.  $f^{-1}O_Y \rightarrow O_X$  is an isomorphism). If  $Y = \text{Spec}(R)$ , any closed immersion  $f : X \rightarrow Y$  is isomorphic to  $\text{Spec}(R/I) \rightarrow \text{Spec}(R)$  for some ideal  $I$  of  $R$ . An **immersion** is a morphism of schemes  $X \rightarrow Y$  which factors  $X \rightarrow Z \rightarrow Y$ , where  $X \rightarrow Z$  is a closed immersion and  $Z \rightarrow Y$  an open immersion.

A property  $P$  of morphisms of schemes is **stable under base change (resp. composition)** if whenever  $f : X \rightarrow S$  has  $P$  and  $S' \rightarrow S$  is an  $S$ -scheme,  $X \times_S S' \rightarrow S'$  has  $P$  (resp. whenever  $X \rightarrow Y$  and  $Y \rightarrow Z$  have  $P$ , so does  $X \rightarrow Z$ ). Being a closed/open immersion or an immersion is stable under base change and composition. Finally, **iff** is a shortcut for "if and only if"...

### 0.1 Basics on closed immersions

1. Let  $S'$  be an  $S$ -scheme. If  $X \rightarrow S$  is a closed immersion, prove that so is  $X \times_S S' \rightarrow S'$ . If  $f : X \rightarrow Y$  is an  $S$ -morphism and a closed immersion, prove that  $X \times_S S' \rightarrow Y \times_S S'$  is a closed immersion.
2. Let  $\mathcal{I} \subset O_X$  be a quasi-coherent sheaf of ideals<sup>1</sup> and  $V(\mathcal{I}) = \text{Supp}(O_Y/\mathcal{I})$ , a closed subset of  $|Y|$ .
  - a) Letting  $i : V(\mathcal{I}) \rightarrow Y$  the inclusion, prove that there is a unique sheaf of rings  $\mathcal{O}$  on  $V(\mathcal{I})$  for which  $i_*\mathcal{O} = O_Y/\mathcal{I}$ , and that  $V(\mathcal{I}) := (V(\mathcal{I}), \mathcal{O})$  is a scheme with a natural closed immersion  $V(\mathcal{I}) \rightarrow Y$ .
  - b) If  $f : X \rightarrow Y$  is a closed immersion, prove that  $\mathcal{I} = \ker(O_Y \rightarrow f_*O_X)$  is a quasi-coherent sheaf of ideals of  $O_Y$  and there is a unique factorisation  $X \simeq V(\mathcal{I}) \rightarrow Y$  of  $f$ .
3. Let  $Z \rightarrow X$  be a closed immersion, with  $Z$  reduced. Prove that a morphism  $f : Y \rightarrow X$ , with  $Y$  reduced, factors through  $Z \rightarrow X$  iff  $f(Y) \subset \text{Im}(Z \rightarrow X)$  set-theoretically.
4. Let  $T \subset X$  be a closed subset of a scheme  $X$ . Prove that  $\mathcal{I}(T) = \{f \in O_X(U) \mid f(t) = 0 \in k(t), \forall t \in T \cap U\}$  is a quasi-coherent sheaf of ideals of  $O_X$  and that there is a natural bijection  $V(\mathcal{I}) = T$  (this scheme structure on  $T$  is called the **reduced induced scheme structure on  $T$** ).

### 0.2 Diagonal morphism, graphs and equalizers

If  $X$  is an  $S$ -scheme, its **diagonal morphism**  $\Delta_{X/S} : X \rightarrow X \times_S X$  is given on  $T$ -points  $s$  by  $\Delta(s) = (s, s) : X(T) \rightarrow (X \times_S X)(T) = X(T) \times_{S(T)} X(T)$ . If  $f : X \rightarrow Y$  is an  $S$ -morphism of schemes, its **graph** is the morphism  $\Gamma_f : X \rightarrow X \times_S Y$  given on  $T$ -points by  $x \in X(T) \rightarrow (x, f(x)) \in X(T) \times_{S(T)} Y(T) = (X \times_S Y)(T)$ .

1. Prove that  $\Delta_{X/S}$  is a closed immersion if  $X, S$  are affine. Deduce that  $\Delta_{X/S}$  is always an immersion, and it's a closed immersion iff its image is closed. **Warning** :  $\text{Im}(\Delta_{X/S}) \subset \{z \in X \times_S X \mid p(z) = q(z)\}$  ( $p, q$  are the natural projections  $X \times_S X \rightarrow X$ ), but this inclusion may not be an equality.
2. For an  $S$ -morphism  $f : X \rightarrow Y$  prove that  $\Gamma_f$  is the base change of  $\Delta_{Y/S}$  by  $X \times_S Y \rightarrow Y \times_S Y$  (given on  $T$ -points by  $(x, y) \rightarrow (f(x), y)$ ), i.e.<sup>2</sup>  $X \simeq Y \times_{(Y \times_S Y)} (X \times_S Y)$ , in such a way that the natural projections to  $Y$  and  $X \times_S Y$  are  $f$  and  $\Gamma_f$ . Deduce that  $\Gamma_f$  is always an immersion.
3. Let  $f, g : X \rightarrow Y$  be  $S$ -morphisms of schemes. Define  $\text{eq}(f, g) = X \times_{(X \times_S Y)} X$ , where the two maps  $X \rightarrow X \times_S Y$  are  $\Gamma_f$  and  $\Gamma_g$ . Prove that there is a canonical immersion  $\text{eq}(f, g) \rightarrow X$ , inducing a functorial bijection<sup>3</sup>  $\text{eq}(f, g)_S(T) = \{x \in X_S(T) \mid f(x) = g(x)\}$  in the  $S$ -scheme  $T$ . Moreover,  $x \in X$  is in the image of  $\text{eq}(f, g) \rightarrow X$  iff  $f(x) = g(x)$  and the two maps  $k(f(x)) = k(g(x)) \rightarrow k(x)$  coincide.
4. Prove that  $f : X \rightarrow S$  is a monomorphism (i.e.  $X(T) \rightarrow S(T)$  is injective for all  $T$ ) iff  $\Delta_{X/S}$  is an isomorphism, and that  $f$  is universally injective<sup>4</sup> iff  $\Delta_{X/S}$  is surjective.

---

1. i.e. a sheaf of ideals whose restriction to any affine open  $U$  of  $X$  is a quasi-coherent  $O_U$ -module.

2. Seen as  $Y \times_S Y$ -schemes via  $\Delta_{Y/S}$  and via the morphism  $X \times_S Y \rightarrow Y \times_S Y$  above.

3. We also write  $f, g$  for the induced maps  $X_S(T) \rightarrow Y_S(T)$ .

4. i.e.  $X \times_S S' \rightarrow S'$  is injective for all  $S$ -schemes  $S'$ , or equivalently  $X(K) \rightarrow S(K)$  is injective for all fields  $K$ .

### 0.3 The diagonal morphism, separated morphisms

We say that  $f : X \rightarrow S$  is **separated** (resp. **quasi-separated**) if  $\Delta_{X/S}$  is a closed immersion (resp. a **quasi-compact morphism**, i.e. the inverse image of any quasi-compact open subset is quasi-compact).

1. Prove that each of the following statements is equivalent to  $f : X \rightarrow S$  being separated :
  - for any  $S$ -morphisms  $f, g : T \rightarrow X$ ,  $\text{eq}(f, g) \rightarrow T$  is a closed immersion.
  - For any  $S$ -morphism  $f : T \rightarrow X$  the graph  $\Gamma_f$  is a closed immersion.
2. a) Prove that a morphism of affine schemes is separated. Also, any immersion is separated.  
 b) Prove that being separated (resp. quasi-separated) is stable under base change and composition. Moreover, an  $S$ -morphism of schemes  $f : X \rightarrow Y$  is separated (resp. quasi-separated) if  $X \rightarrow S$  is so.
3. a) Let  $f, g : X \rightarrow Y$  be  $S$ -morphisms of schemes such that  $f|_U = g|_U$  for an open dense subscheme  $U$  of  $X$ . If  $Y \rightarrow S$  is separated and  $X$  is reduced, prove that  $f = g$ .  
 b) Let  $f, g : Z := \text{Spec}(k[X, Y]/(XY, Y^2)) \rightarrow \text{Spec}(k[T]/T^2)$  be induced by the maps on rings sending  $T$  to 0 (resp.  $Y$ ). Prove that there is an open dense subset  $U \subset Z$  such that  $f|_U = g|_U$ , yet  $f \neq g$ .
4. a) Prove that  $X \rightarrow \text{Spec}(R)$  is separated if and only if  $U \cap V$  is affine and  $O_X(U) \otimes_R O_X(V) \rightarrow O_X(U \cap V)$  is surjective for any affine opens  $U, V$  of  $X$ . Deduce that  $\mathbf{P}^n \rightarrow \text{Spec}(\mathbf{Z})$  is separated.  
 b) If  $X \rightarrow S$  and  $S \rightarrow \text{Spec}(\mathbf{Z})$  are separated, then  $U \cap V$  is affine and a closed subscheme of  $U \times V$  for any affine opens  $U, V$  of  $X$ . If  $S \rightarrow \text{Spec}(\mathbf{Z})$  is separated and  $f : X \rightarrow S$  is any morphism, then  $U \cap f^{-1}(V)$  is affine for any open affine subschemes  $V \subset S$  and  $U \subset X$ .

### 0.4 Geometry of valuation rings

A **valuation ring** is an integral domain  $V$  such that for all nonzero  $x \in \text{Frac}(V)$  we have  $x \in V$  or  $1/x \in V$ .

1. Prove that any valuation ring is normal (i.e. integrally closed in its field of fractions) and local.
2. (hard) Let  $K$  be a field and let  $A \subset K$  be a local subring such that  $\text{Frac}(A) = K$ .
  - a) Prove that  $A$  is a valuation ring if and only if for any local ring  $B$  with  $A \subset B \subset K$  and such that  $A \rightarrow B$  is a local map we have  $A = B$ . **Hint** : use your favorite commutative algebra book!
  - b) Deduce that there is a valuation ring  $V$  such that  $A \subset V \subset K$  and such that  $A \rightarrow V$  is a local map. If  $X$  is a scheme and  $x, x' \in X$  we say that  $x$  is a **specialization of  $x'$**  and write  $x' \rightsquigarrow x$  if  $x \in \overline{\{x'\}}$ . A morphism  $f : X \rightarrow S$  is **specializing** if any specialization of  $f(x')$  (with  $x' \in X$ ) is of the form  $f(x)$  for a specialization  $x$  of  $x'$ .
3. a) If  $X$  is a scheme and  $x' \rightsquigarrow x$  in  $X$ , prove that there is a valuation ring  $A$  with  $\text{Frac}(A) = k(x')$  and a morphism  $\text{Spec}(A) \rightarrow X$  sending the closed point to  $x$  and the generic point to  $x'$ .  
 b) If  $f : X \rightarrow S$  is a quasi-compact morphism of schemes<sup>5</sup>, then  $f(X)$  is closed in  $S$  iff  $f(X)$  is stable under specialization (i.e. if  $s' \rightsquigarrow s$  and  $s' \in f(X)$  then  $s \in f(X)$ ), and  $f$  is closed iff  $f$  is specializing.
4. (**valuative criteria**) Let  $f : X \rightarrow S$  be a morphism of schemes.
  - a) Prove that  $X \times_S S' \rightarrow S'$  is specializing for any  $S$ -scheme  $S'$  iff  $X_S(V) \rightarrow X_S(\text{Frac}(V))$  is surjective for any valuation ring  $V$  with a map  $\text{Spec}(V) \rightarrow S$ . If  $f$  is quasi-compact, this is also equivalent to  $f$  being **universally closed** (i.e.  $X \times_S S' \rightarrow S'$  is closed for all  $S$ -schemes  $S'$ ).
  - b) Prove that  $f : X \rightarrow S$  is separated iff  $f$  is quasi-separated and  $X_S(V) \rightarrow X_S(\text{Frac}(V))$  is injective for any valuation ring  $V$  with a map  $\text{Spec}(V) \rightarrow S$ .

### 0.5 Properness

A morphism of schemes  $f : X \rightarrow S$  is called **proper** if  $f$  is of finite type<sup>6</sup>, separated and universally closed.

1. a) Prove that being proper is stable under base change and composition. Moreover,  $f : X \rightarrow S$  is proper if and only if  $f^{-1}(U_i) \rightarrow U_i$  are proper, for an open covering  $S = \cup_i U_i$ .  
 b) Let  $f : X \rightarrow Y$  be an  $S$ -morphism, with  $Y \rightarrow S$  separated. If  $X \rightarrow S$  is proper, then  $f$  is proper.  
 c) "The image of a proper scheme is proper" : if  $f : X \rightarrow Y$  is a surjective  $S$ -morphism, with  $X \rightarrow S$  proper and  $Y \rightarrow S$  separated and of finite type, then  $Y \rightarrow S$  is proper.

5. i.e. the inverse image of any quasi-compact open is quasi-compact.

6. i.e.  $f$  is quasi-compact and  $O_S(V) \rightarrow O_X(U)$  is of finite type whenever  $U, V$  are affine opens of  $X, S$  such that  $f(U) \subset V$ .

2. a) (**valuative criterion**) Prove that  $f$  is proper iff  $f$  is quasi-separated, of finite type and  $X_S(V) \rightarrow X_S(\text{Frac}(V))$  is bijective for any valuation ring  $V$  with a map  $\text{Spec}(V) \rightarrow S$ .  
 b) If  $R$  is a Dedekind domain and  $X \rightarrow \text{Spec}(R)$  is proper, prove that  $X(R) \rightarrow X(\text{Frac}(R))$  is bijective.
3. a) Prove that any closed immersion, as well as  $\mathbf{P}^n \rightarrow \text{Spec}(\mathbf{Z})$  is proper.  
 b) Prove that  $\mathbf{A}^1 = \text{Spec}(\mathbf{Z}[T]) \rightarrow \text{Spec}(\mathbf{Z})$  is not proper.  
 c) (hard) If  $f : \text{Spec}(A) \rightarrow \text{Spec}(B)$  is universally closed (e.g. proper), then  ${}^7 B \rightarrow A$  is integral.
4. (hard) Let  $k$  be an algebraically closed field and  $X \rightarrow \text{Spec}(k)$  a proper morphism, with  $X$  reduced and connected. Prove that  $O_X(X) = k$ . **Hint** : see  $f$  as a morphism  $X \rightarrow \mathbf{A}_k^1 \subset \mathbf{P}_k^1$ .

## 0.6 Proper normal curves over a field

This exercise is hard and fully uses the results in exercises 4,5. Fix a field  $k$ . A **curve** over  $k$  is a reduced, irreducible scheme  $C$  of dimension 1, with a morphism of finite type  $C \rightarrow \text{Spec}(k)$ . Its function field  $K(C) = O_{C,\eta}$  ( $\eta \in C$  being the generic point of  $C$ ) has transcendence degree 1 over  $k$ . Call  $C$  **normal** if  $O_{C,x}$  is a discrete valuation ring (i.e. a noetherian valuation ring, equivalently<sup>8</sup> normal) for all  $x \in C \setminus \{\eta\}$ .  **$C$  will always denote a normal curve over  $k$  and  $K$  will always denote an extension of  $k$  of transcendence degree 1.** For such  $K/k$ , its **Riemann-Zariski space**  $RZ(K/k)$  is the set of valuation rings  $V$  such that  $k \subset V \subset K$  and  $\text{Frac}(V) = K$ , with the topology for which a nonempty  $U \subset RZ(K/k)$  is open if  $K \in U$  (note that  $K \in RZ(K/k)$ ) and  $RZ(K/k) \setminus U$  is finite.

1. a) Prove that  $\{K\}$  is dense in  $RZ(K/k)$ , and any other point of  $RZ(K/k)$  is closed. Moreover, for any  $f \in K$  the set  $\{V \in RZ(K/k) | f \in V\}$  is open. Deduce that a nonempty  $U \subset RZ(K/k)$  is open if and only if  $U$  is a union of sets of the form  $\{V \in RZ(K/k) | f_1, \dots, f_n \in V\}$ , with  $f_1, \dots, f_n \in K$ . **Hint** : if  $V \in RZ(K/k)$  and  $f \notin V$ , study the integral closure of  $k[1/f]$  in  $K$ .  
 b) Prove that if  $V \in RZ(K/k)$ , then either  $V = K$  or  $V$  is a discrete valuation ring.
2. Let  $C$  be a normal curve over  $k$  and let  $K = K(C)$  be its function field.  
 a) Prove that  $x \rightarrow O_{C,x}$  induces a continuous open map  $\iota : |C| \rightarrow RZ(K/k)$ .  
 b) Prove that  $C \rightarrow \text{Spec}(k)$  is separated (resp. proper) if and only if  $\iota$  is an open embedding (resp. a homeomorphism). **Hint** : use several times the valuative criteria.
3. Fix  $K/k$  of transcendence degree 1 and let  $X = RZ(K/k)$ , a topological space. Define a pre-sheaf of rings  $O_X$  on  $X$  by setting  $O_X(U) = \bigcap_{V \in U} V \subset K$  for  $U \subset X$  nonempty.  
 a) Prove that  $O_X$  is a sheaf of  $k$ -algebras on  $X$ , and that  $O_{X,V} \simeq V$  for all  $V \in X$ , in particular  $X := (X, O_X)$  is a locally ringed space.  
 b) Let  $C$  be a normal curve over  $k$ , such that  $K(C) = K$ . Prove that there is a unique morphism of locally ringed spaces  $f : C \rightarrow X$  compatible with the natural maps to  $\text{Spec}(k)$ . Moreover, this map is an open immersion if  $C \rightarrow \text{Spec}(k)$  is separated.  
 c) Prove that  $X$  is a normal curve over  $k$ ,  $X \rightarrow \text{Spec}(k)$  is proper and  $K(X) \simeq K$ . Conversely, if  $C$  is a normal curve over  $k$ , there is a natural isomorphism of  $k$ -schemes  $C \simeq RZ(K(C))$ .

---

7. It is actually enough to assume that  $\text{Spec}(A[T]) \rightarrow \text{Spec}(B[T])$  is closed.

8. Though this is not really trivial...